

Plant Pest Risk Assessment for Event H7-1 Sugar Beet

Monsanto Company (St. Louis, MO) and KWS SAAT AG (Einbeck, Germany), petitioned APHIS (APHIS number 03-323-01p) for a determination that genetically engineered (GE) sugar beet (*Beta vulgaris* L. ssp. *vulgaris*) event H7-1 is unlikely to pose a plant pest risk (Monsanto and KWS SAAT AG 2004) and, therefore, should no longer be a regulated article under APHIS' 7 Code of Federal Regulations (CFR) part 340. APHIS administers 7 CFR part 340 under the authority of the plant pest provisions of the Plant Protection Act of 2000¹. This plant pest risk assessment was conducted to determine if H7-1 is unlikely to pose a plant pest risk.

The initial request was received from Monsanto and KWS SAAT AG in 2003. Upon completing a Plant Pest Risk Assessment, an Environmental Assessment (EA) and issuing a Finding of No Significant Impact (FONSI) (70 FR 13007-13008, Docket No. 04-075-2), APHIS advised the public of its determination, effective March 4, 2005, that the Monsanto and KWS SAAT AG (2004) sugar beet event H7-1 did not pose a plant pest risk and therefore was no longer considered a regulated article under APHIS regulations in 7 CFR part 340. A complaint was filed in January, 2008, challenging APHIS' decision to grant nonregulated status to event H7-1. On September 21, 2009, the US District Court for the Northern District of California ruled that APHIS should have prepared an Environmental Impact Statement before making a decision to grant nonregulated status to event H7-1 (*Center for Food Safety et al. vs. Thomas Vilsack et al.*). On August 13, 2010, the Court vacated APHIS's decision to grant nonregulated status to the event H7-1 sugar beet varieties making them subject to the Plant Protection Act of 2000 (PPA) and 7 CFR part 340 once again, and remanded the matter back to the agency to determine regulatory actions, if any, that should be imposed upon event H7-1 until the completion of the EIS and a new decision could be made by APHIS as to whether it would be appropriate to grant full nonregulated status to event H7-1.

APHIS received a supplemental request from Monsanto and KWS SAAT AG (2010) to amend the petition for non-regulated status submitted in 2003 (Petition 03-323-01p) pursuant to the regulatory scheme of 7 CFR part 340 (Reding 2010). These petitioners requested that APHIS grant "partial deregulation" or similar administrative action to authorize the continued cultivation of event H7-1 subject to the interim measures proposed by APHIS in the lawsuit challenging its 2005 determination of non-regulated status of event H7-1. On February 8, 2011, APHIS advised the public of its decision to "partially deregulate" event H7-1 sugar beets as an interim action until the court-ordered EIS was completed. That "partial deregulation" described a system in which root crop production of event H7-1 sugar beets could occur under a Compliance Agreement with USDA/APHIS and growing of event H7-1 for seed production could occur under APHIS' permitting system. The conditions under which commercial H7-1 root production is carried out are enforceable under the mandatory compliance agreements. A Plant Pest Risk

¹ Section 403 (14) of the Plant Protection Act (7USC Sec 7702(14) defines plant pest as: "Plant Pest - The term "plant pest" means any living stage of any of the following that can directly or indirectly injure, cause damage to, or cause disease in any plant or plant product: (A) A protozoan. (B) A nonhuman animal. (C) A parasitic plant. (D) A bacterium. (E) A fungus. (F) A virus or viroid. (G) An infectious agent or other pathogen. (H) Any article similar to or allied with any of the articles specified in the preceding subparagraphs."

Assessment for root production was prepared to consider the plant pest risk potential of H7-1 sugar beet root crop under the conditions of the interim action.

H7-1 was grown in the U.S. under APHIS permits or notifications from 1998 through 2005. From 2005 to 2010, H7-1 was grown widely without APHIS regulatory restrictions. Since February 2011, H7-1 sugar beets have been grown commercially for root production under APHIS-mandated compliance agreements and for seed production under APHIS issued permits. This Plant Pest Risk Assessment was conducted to determine whether sugar beet event H7-1 is unlikely to pose a plant pest risk under full deregulation.

Sugar beet line H7-1 was produced by transformation of a sugar beet line (3S0057) with *Agrobacterium tumefaciens*. Because *A. tumefaciens* is a plant pest and some of the regulatory sequences (figwort mosaic virus promoter) used to facilitate expression of the gene and the sequences that code for the protein (*cp4epsps* gene from *A. tumefaciens*) in sugar beet were derived from plant pests, H7-1 has been considered a regulated article under APHIS regulations at 7 CFR part 340. Potential impacts to be addressed in this risk assessment are those that pertain to the use of H7-1 and its progeny in the absence of confinement. APHIS utilizes data and information from current scientific literature, in addition to information submitted by the applicant, to determine if H7-1 is unlikely to pose a plant pest risk. If APHIS determines that a GE organism is not a plant pest, then APHIS has no regulatory authority over that organism. Thus, if APHIS determines that H7-1 sugar beet does not pose a plant pest risk, then H7-1 sugar beet would no longer be subject to the plant pest provisions of the Plant Protection Act or to the regulatory requirements of 7 CFR part 340, and therefore, APHIS must grant it non-regulated status.

Potential impacts analyzed in this Plant Pest Risk Assessment are those that pertain to plant pest risk characteristics. APHIS regulation 7 CFR 340.6(c) specifies the information a petitioner must include in a petition for nonregulated status. APHIS will evaluate information submitted by the applicant related to plant pest risk characteristics, disease and pest susceptibilities, expression of the gene product, new enzymes, or changes to plant metabolism, weediness of the regulated article, any impacts on the weediness of any other plant with which it can interbreed, and transfer of genetic information to organisms with which it cannot interbreed.

Pursuant to NEPA, the EIS for this petition for full deregulation considers whether agricultural or cultivation practices associated with H7-1 may result in impacts on the environment. A thorough assessment of H7-1's effects on the environment including on non-target and beneficial organisms, and threatened and endangered species is included in the EIS.

Development of H7-1 Herbicide Tolerant² Sugar beet

² The applicant has described H7-1 sugar beet as “herbicide tolerant” and historically APHIS has also referred to GE plants with diminished herbicide sensitivity as “herbicide tolerant”. However, the phenotype would fall under the Weed Science Society of America’s (WSSA) definition of “herbicide resistance” since H7-1 has an inherited ability to survive and reproduce following exposure to a dose of herbicide normally lethal to the wild type variety (WSSA 1998). By the WSSA definition, “resistance [to an herbicide] may be naturally occurring or induced by such techniques as genetic engineering or selection of variants produced by tissue culture or mutagenesis.” Herbicide

Origin of Cultivated Beets

Sugar beets (*Beta vulgaris* (L.) ssp. *vulgaris*), a member of the genus *Beta* (Chenopodiaceae), are cultivated worldwide. The species *B. vulgaris* includes many different crop varieties including sugar beets, fodder beets, Swiss chard, and table beets. *Beta vulgaris* L. ssp. *maritima* (L.) Arcang. wild sea beet, is regarded as the progenitor of the *Beta* crop species (fodder beet, sugar beet, Swiss chard and table or red beet) and found widely distributed over the Mediterranean basin and the Near East (OECD 2001).

Plant Description and Flowering Characteristics

Sugar beet is a biennial species (typically will complete its lifecycle in 2 years) that is grown as an annual for sugar production. Since the root crop is usually harvested during the first year while still in the vegetative phase, flowers rarely develop. The plant grows as a rosette in its first year and can be quite variable in size (from 1-6 ft. tall). If allowed to overwinter in a moderate climate, it will “bolt” the second year and produce a flower stalk and seeds (OECD 2001). Being a biennial, the plant typically flowers during the late spring to summer of the 2nd year of growth, usually sending up a single unbranched flower stalk. Flowers begin to release pollen about 4-6 weeks after the flower stalk emerges and continue to release pollen for another 2-4 weeks. This biennial plant life cycle of reduced flowering in the first year has been selected for in sugar beets and vegetable beet varieties. Typically the level of bolting in cultivated sugar beet is reduced to 0.01 – 1 percent (Ingram 2000; Darmency 2009; OECD 2001). However, the presence of the bolting genes and environmental triggers such as day-length, temperature or the amount of cold during the winter may induce flowering during the first year (Boudry 2002; Büttner 2010; Van Dijk 1997).

Sugar beet pollen is mostly carried by wind with insects playing a lesser role (OECD 2001). Exact percentages of outcrossing vary considerably at different locations based on numerous environmental conditions. Wind, however, is considered to be the primary vector for pollination.

H7-1 is a genetically engineered sugar beet line that was developed to increase tolerance to the herbicide glyphosate. Glyphosate was first introduced as an herbicide under the trade name of Roundup[®] by Monsanto in 1975. Glyphosate is a systemic, post-emergence herbicide widely used on both agricultural commodities (food uses) and non-agriculture sites (Cerdeira 2006).

The management of weeds in conventional sugar beet fields can be an expensive, labor intensive, and sometimes complicated operation. Often farmers use pre-emergent herbicides that will stop weed seeds from germinating. However, this requires application of the pre-emergent herbicides on the entire field because farmers cannot predict where, on their fields, weeds will emerge. With H7-1, growers have the option of applying herbicide after weeds have germinated and only in the areas of the field where there are weeds.

tolerance, by the WSSA definition, only applies to plant species with an “inherent ability” to survive and reproduce after herbicide treatment.

These sugar beets were genetically engineered to be glyphosate tolerant by inserting a gene (from *Agrobacterium* sp. strain CP4) that codes for the enzyme 5-enolpyruvylshikimate-3-phosphate synthase (EPSPS) into the sugar beet genome. This gene, along with its regulatory sequences, was introduced into these sugar beets via an *Agrobacterium*-mediated transformation protocol. This is a well-characterized procedure that has been widely used for over two decades for introducing various genes directly into plant genomes.

APHIS authorized the first field testing of these sugar beets from 1998 through 2002. Event H7-1 sugar beets have been evaluated extensively to confirm that they exhibit the desired agronomic characteristics and do not present a plant pest risk. The field tests were conducted in agricultural settings under physical and reproductive confinement conditions.

Description of the Modification

Sugar beet H7-1 was produced by transformation using disarmed *Agrobacterium tumefaciens*. Sugar beet line 3S0057 (KWS SAAT AG proprietary line) cotyledons were infected with *A. tumefaciens* strain CP4 containing plasmid PV-BVGT08 (Monsanto and KWS SAAT AG 2004, pp. 20-21). This technique using disarmed *Agrobacterium* followed by selection (Howard 1990) has a history of use and has been used for transformation of a variety of plant tissues for over 20 years. Plants containing the introduced DNA were selected based on growth in the presence of glyphosate.

The plasmid PV-BVGT08 contains a single expression cassette flanked by the right and left border sequences from the *Agrobacterium* Ti plasmid (Monsanto and KWS SAAT AG 2004, p. 23). Data supplied in the petition (Monsanto and KWS SAAT AG 2004, pp. 29-47) support the conclusion that event H7-1 contains the following sequences:

5-enolpyruvylshikimate-3-phosphate synthase (EPSPS)

- The 35S gene promoter from a figwort mosaic virus (Gowda 1989; Sanger 1990) is constitutively active in plants (Sheperd 1987; Richins 1987; Gowda 1989; Sanger 1990).
- The *ctp2* N-terminal chloroplast transit peptide (CTP) sequence from the *Arabidopsis thaliana cp4 epsps* coding region (Timko 1988) that targets the protein to the chloroplast.
- The *cp4 epsps* gene from *Agrobacterium* sp. strain CP4 (Padgett 1995) has been sequenced and encodes a 47.6 kDa protein consisting of a single polypeptide of 455 amino acids (Padgett 1996).
- 3' untranslated region (transcriptional terminator and polyadenylation site) of the *rbcS E9* gene from *Pisum sativum* (Coruzzi 1984; Morelli 1985).

Data was provided and reviewed by APHIS that demonstrates stable integration and inheritance of the *cp4 epsps* gene and its associated regulatory sequences over several breeding generations (Monsanto and KWS SAAT AG 2004). Statistical analyses show that glyphosate tolerance is inherited as a dominant trait in a typical Mendelian manner.

Potential for H7-1 to Become Invasive and/or a Weed

In the U.S., *Beta vulgaris* L. spp. *vulgaris*, is not listed as a weed by the Weed Science Society of America (2010) nor is it listed as a noxious weed species by the U.S. Federal government (7 CFR part 360;

http://www.aphis.usda.gov/plant_health/plant_pest_info/weeds/downloads/weedlist-2010doc.pdf). In order to assess the potential for H7-1 to become invasive and/or a weed, APHIS looked to whether H7-1's reproductive characteristics and survival biology were any different from the characteristics of conventional sugar beet varieties. One measure of a plant's potential to be invasive is the potential for the plant to produce "volunteers." Sugar beets possess few of the characteristics of plants that are notable of successful weeds (Baker 1965; Keeler 1989). As part of a bilateral agreement between the United States and Canada, USDA/APHIS and the Canadian Food Inspection Agency (CFIA) have generated documents that outline basic data requirements for developers of genetically engineered plants. One of these documents, Appendix II, outlines the environmental characterization data requirements for unconfined releases. These data and the data required in a petition for non-regulated status are designed to inform APHIS of any differences between the characteristics, including growth and seed characteristics that influence reproductive and survival biology of the transgenic plant compared to its non-transgenic counterpart.

Volunteer plants rarely occur in fields planted with a sugar beet root crop. This is because bolters in root crop fields are rare and generally do not produce seed and unharvested roots are incapable of surviving the winters in the Northern sugar beet producing regions and the summers in the Imperial Valley (USDA APHIS, 2011). In fields used to produce sugar beet seed, volunteer plants can occur in subsequent years. These plants can be controlled by mechanical means or the use of several other registered herbicides besides glyphosate (Crop Protection Chemical Reference 1996).

APHIS assessed whether H7-1 is any more likely to become a weed than the isogenic non-transgenic sugar beet line. The assessment encompasses a thorough consideration of the basic biology of sugar beet and an evaluation of the unique characteristics of H7-1 under field conditions. Monsanto conducted field trials of H7-1 between 1998 and 2002 at 98 field trial sites in the U.S. (Monsanto and KWS SAAT AG 2004, p. 62) and Europe (Monsanto and KWS SAAT AG 2004, p. 70). APHIS considered data relating to plant vigor, bolting, seedling emergence, seed germination, seed dormancy and other characteristics that might relate to increased weediness (Monsanto and KWS SAAT AG 2004, pp. 70-77). Field trial data indicated that H7-1 does not exhibit characteristics that would cause it to be weedier than the parental sugar beet line. Additionally, no characteristics relating to disease or insect resistance that might affect weediness were noted that were consistent over all trial locations. H7-1 sugar beet is still susceptible to the typical insect and disease pests of sugar beet.

These results on growth characteristics, seed production and germination indicate that H7-1 is not different from its comparators in these respects. There is no indication that H7-1 possesses a selective advantage that would result in increased weediness or invasiveness. H7-1 lacks the ability to persist as a troublesome weed, and there would be no significant impact on current

weed management practices for sugar beet cultivation, because alternative methods exist that can be used to kill weeds in the absence of using glyphosate (OECD 2001).

Potential for Gene Flow and Gene Introgression from H7-1 into Sexually Compatible Relatives

APHIS evaluated the potential for gene introgression to occur from H7-1 to sexually compatible wild relatives and considered whether such introgression would result in weediness in those resulting hybrids. The centers of origin for *Beta vulgaris* is generally believed to be in the Mediterranean or Near East region and no *Beta* species are known to be native to the U.S. (OECD 2001; Panella 2007).

Sugar beet is strongly self-incompatible (particularly tetraploid plants) although self-fertilizing plants exist in nearly every population (OECD 2001). Outcrossing is very important in developing hybrid seed every year for commercial production. The use of cytoplasmic male sterile lines is used extensively for this work.

Distribution of Beta species in the United States

Beet species that have been introduced into the United States include *B. procumbens* Chr. Sm., *B. vulgaris* ssp. *vulgaris*, *B. vulgaris* ssp. *maritima* and *B. macrocarpa* Guss. (USDA ARS 2011). The distribution of *B. procumbens* is limited to the state of Pennsylvania (USDA ARS 2011).

The only location where feral beets are recorded as having a genuine presence in the environment is in California. The California wild beets belong to two different taxa, *B. vulgaris* and *B. macrocarpa*, and have at least three different origins (Bartsch 2002). They evolved from escaped cultivated Swiss chard or red beet, from *B. macrocarpa*, or from hybridization of *B. vulgaris* with introduced *B. macrocarpa*. Research by Bartsch and Ellstrand (1999) identified that many populations of wild beets in California are actually feral varieties of Swiss chard and table beets (Bartsch 2003). Wild beets are found from the San Francisco Bay area to the Mexican border (Bartsch 1999) (Figures 1 and 2).




Figure 1. Distribution of *B. vulgaris* (Calflora 2011)




Figure 2. Distribution of *B. macrocarpa* (Calflora 2011)

Potential for H7-1 to hybridize with Beta species that occur in the U.S.

Hybrids between sugar beet and *B. procumbens* normally die at the seedling stage (Savitsky 1975; OECD 2001). Therefore, hybridization of H7-1 with the wild *B. procumbens* in Pennsylvania is unlikely to occur due to species incompatibility.

Another *Beta* species, *Beta macrocarpa*, grows in the Imperial Valley of California (Figure 2) where sugar beet root production occurs. Hybrids between sugar beet and *B. macrocarpa* are also not likely to occur. *B. macrocarpa* show a high degree of self-fertilization and are known to occupy the same or overlapping geographic areas without interbreeding (Lange 1999). *B. macrocarpa* usually flowers earlier than sugar beet. Bartsch speculated that *B. macrocarpa* can

cross with sugar beet bolters when flowering times overlap (Bartsch 2002) based on evidence of introgression from isozyme analysis. Lewellen, however, found that *B. macrocarpa* x *B. vulgaris* crosses in greenhouses under ideal conditions were usually unsuccessful and the progeny were genetically unstable, further supporting the conclusion that hybrids between sugar beet and *B. macrocarpa* are unlikely (Lewellen 2003).

Where cultivated sugar beets are grown in close proximity to wild *B. vulgaris*, they can freely hybridize when flowering times overlap (Boudry 1993). In California, feral populations of *Beta vulgaris*, thought to be escaped table beet or Swiss chard, have established along the coast and pose a minor weed problem (Bartsch 1999; Johnson 1958; McFarlane 1975). These plants are not serious weed problems except when they are present in table beet, Swiss chard, and sugar beet fields (Johnson 1958; Panella 2003). Confirmed populations of *Beta vulgaris* are found in the San Francisco Bay area and along the coast where sugar beets are no longer cultivated. Wild *B. vulgaris* would be capable of crossing with flowering sugar beet plants, however, wild populations have not been confirmed in the Imperial Valley (Calflora, 2011, Kelly Richardson, personal communication). These beets are easily controlled with herbicides or mechanical cultivation in non-*Beta* crops. Control of weed beet in *Beta* crop fields is more difficult because young plants resemble *Beta* crops making it difficult to remove them by mechanical means and they are tolerant of the herbicides used for weed control in beet crops.

Escape of the engineered trait into weed beet populations is possible if grown in close proximity to weedy or feral beets. APHIS concludes that the potential of the glyphosate tolerance trait moving from H7-1 to other sexually compatible *Beta* species in the United States is low, because of lack of ability to hybridize or proximity to sexually compatible plants. If the trait did move into wild species, these could not be controlled with glyphosate in H7-1 sugar beets. As glyphosate is not used for post-emergent control of wild beets in other *Beta* crops and other herbicides are effective to control wild beets in non-*Beta* crops, the potential for glyphosate tolerant weed beets to cause problems in H7-1 sugar beet fields or other crops that are resistant to glyphosate would be limited. Even if these plants become tolerant to glyphosate, there are other registered herbicides that can be used to kill them and other methods of control can still be used (OECD 2001).

Potential for Transfer of Genetic Information to Organisms with which H7-1 Cannot Interbreed

Horizontal gene transfer and expression of DNA from a plant species to bacteria is unlikely to occur (Keese 2008). First, many bacteria (or parts thereof) that are closely associated with plants have been sequenced, including *Agrobacterium* and *Rhizobium* (Kaneko 2000; Kaneko 2002; Wood 2001). There is no evidence that these organisms contain genes derived from plants. Second, FDA has evaluated horizontal gene transfer from the use of antibiotic resistance marker genes, and concluded that the likelihood of transfer of antibiotic resistance genes from plant genomes to microorganisms in the gastrointestinal tract of humans or animals, or in the environment, is remote (FDA 1998). Therefore, APHIS concludes that horizontal gene transfer is unlikely to occur and thus poses no plant pest risk.

Potential for H7-1 to have Altered Disease and Pest Susceptibilities

APHIS assessed whether H7-1 is likely to have significantly altered disease and pest susceptibility. This assessment encompassed a thorough consideration of the introduced trait and disease and pest susceptibility data from H7-1 field trials.

The *Agrobacterium* transformed plants used in the generation of H7-1 were treated with an antibiotic to kill the *Agrobacterium* cells (Monsanto and KWS SAAT AG 2004, p. 19). Furthermore, DNA sequences derived from plant pests that were incorporated into H7-1 do not result in the production of infectious agents or disease symptoms in plants, and so it is unlikely that H7-1 could pose a plant pest risk. The description of the genetic modifications, including genetic elements, expression of the gene product and their functions for H7-1 has been summarized above.

Sugar beets are susceptible to a wide variety of insect pests and plant diseases (Cattanach 1991). Both qualitative and quantitative data addressing disease susceptibility, insect damage and overall agronomic performance were collected in order to assess possible effects from introduction of the *cp4 epsps* gene and its associated regulatory sequences. H7-1 was field tested in the U.S. over five years (1998-2002) at ninety-eight field trial sites representing a wide range of environmental conditions (Monsanto and KWS SAAT AG 2004, p. 62). Plant disease was noted in thirty-six of the ninety-eight trial sites (Monsanto and KWS SAAT AG 2004, p. 63-65). Susceptibility to powdery mildew (*Erysiphe betae*), *Cercospora* leaf spot (*Cercospora beticola*), *Rhizoctonia* root rot (*Rhizoctonia solani*), fungal seedling disease (including *Pythium*, *Rhizoctonia*, and *Aphanomyces*, curly top virus and *Rhizomania* (Beet Necrotic Yellow Vein Virus – BNYVV) was similar between H7-1 and conventional sugar beet varieties in all but six field trial sites. Observations found that H7-1 exhibited reduced susceptibility to powdery mildews at three trial sites, increased susceptibility at one site, and no difference at nine sites where the disease was present. A slight increased susceptibility (10%) to *Cercospora* leaf spot was noted in two sites, while no difference was identified at eleven sites. These differing levels of resistance and susceptibility observed between H7-1 and comparator sugar beet plants is likely due to the non-GE part of the genetic background in the H7-1 plants, because there were no trends in disease susceptibility with event H7-1 when differences were observed. European trials conducted in Germany and France over 2 years, using regionally adapted conventional sugar beet lines and genetically similar lines to H7-1, noted no differences in susceptibility to 10 different sugar beet pests (Monsanto and KWS SAAT AG 2004, pp. 69-70).

Nursery trials tested the performance of different sugar beet plant varieties including H7-1 when challenged (artificial or natural infection) with plant pathogens (Monsanto and KWS SAAT AG 2004). During the 2000 and 2001 growing seasons, sugar beet nursery trials were conducted with H7-1 and conventional sugar beet varieties to assess disease resistance (to *Cercospora* leaf spot, *Aphanomyces* root rot, and curly top and *Rhizoctonia* root). H7-1 infection levels were found to be within the range of ratings observed for the conventional registered varieties. Greenhouse trials using *Fusarium* and *Rhizoctonia* isolates have found that some of the isolates produced greater disease severity on H7-1 treated with glyphosate (Hanson 2003; Larson 2006). Other researchers have suggested that it may be difficult to predict field results from greenhouse/laboratory experiments (Estok 1989; Wan 1998). Subsequent field studies did not show increased incidence to these root diseases (Khan 2010).

The major insect and nematode pests in the U.S. (sugar beet root aphid (*Pemphigus populivenerae*), sugar beet root maggot (*Tetanops myopaeformis*), sugar beet cyst nematode (*Heterodera schachtii*) and root knot nematode (*Meloidogyne arenaria*, *M. incognita*, *M. javanica* and *M. hapla*) were monitored during the U.S. field trials (Dewar 2006, Monsanto and KWS SAAT AG 2004, pp. 66-68). No significant differences between H7-1 and conventional sugar beets were observed in any of the 98 field trials.

Al-Kaff (1998) noted gene silencing effects when transgenic plants have been infected by a virus with DNA sequence homology to a portion of the introduced genes. None of the viral diseases of beet is related to figwort mosaic virus (<http://image.fs.uidaho.edu/vide/refs.htm> and Whitney 1986) (a caulimovirus and from which the promoter for the *cp4 epsps* gene originates) so silencing of the *cp4 epsps* gene should not occur.

In addition to field studies on agronomic parameters, Monsanto and KWS SAAT AG (2004, pp. 77-84) analyzed sugar beets for compositional changes as part of their submission to FDA in the consultation process. While FDA uses these data as indicators of possible nutritional changes, APHIS views them as a general indicator of possible unintended changes. Compositional analyses evaluating carbohydrates, proteins, fiber, fat, sugars and eighteen amino acids (a total of 55 statistical comparisons) identified seven statistically different values compared with the near isogenic control line. None of the values for seed composition characteristics were outside the range of natural variability of conventional sugar beet. Therefore, the composition of H7-1 is not biologically different than conventional sugar beets. Based on the known functions and mechanisms of actions of this protein (summarized in the petition), this protein is not expected to directly alter susceptibility to plant pathogens. Thus H7-1 is expected to be susceptible to the same pests as conventional sugar beet.

Conclusion

APHIS has prepared this plant pest risk assessment in order to determine if event H7-1 sugar beet is unlikely to pose a plant pest risk under full deregulation. Based on the information provided by the applicant, consideration of current scientific literature and the lack of plant pest risk from the inserted genetic material, weedy characteristics, atypical responses to disease or plant pests in the field, effects on non-targets or beneficial organisms in the agro-ecosystem, and horizontal gene transfer, APHIS has concluded that event H7-1 sugar beet is unlikely to pose a plant pest risk.

References

- Al-Kaff**, N.S., Covey, S.N., Kreike, M.M., Page, A.M., Pinder, R., Dale, P.J. (1998) Transcriptional and Posttranscriptional Plant Gene Silencing in Response to a Pathogen. *Science* 279: 2113-2115
- Baker**, H.G. (1965) Characteristics and Modes of Origin of Weeds. *In*: The Genetics of Colonizing Species. pp. 147-172. Baker, H. G., Stebbins, G. L. (eds.). Academic Press, New York and London

- Bartsch, D., Ellstrand, N.C.** (1999) Genetic evidence for the origin of Californian wild beets (genus *Beta*). *Theoretical and Applied Genetics* 99:1120-1130
- Bartsch, D., Driessen, S., Gathmann, A., Hoffmann, A., Lehnen, M., Muecher, T., Saeglitz, C., Wehres, U., Schuphan, I.** (2002) Monitoring the environmental consequences of gene flow from transgenic sugar beet. *In Gene Flow Workshop, The Ohio State University*, pp. 78-93
- Bartsch, D., Cuguen, J., Biancardi, E., Sweet, J.** (2003) Environmental implications of gene flow from sugar beet to wild beet -- current status and future research needs. *Environmental-Biosafety-Research* 2:105-115
- Boudry, P., Moerchen, M., Saumitou-Laprade, P., Vernet, P., van Dijk, H.** (1993) The origin and evolution of weed beets: Consequence of the breeding and release of herbicide-resistant transgenic sugar beets. *Theoretical and Applied Genetics* 87:471-478
- Boudry, P., McCombie, H., Van Dijk, H.** (2002) Vernalization requirement of wild beet *Beta vulgaris* spp. *maritima*: among population variation and its adaptive significance. *Journal of Ecology* 90:693-703.
- Buttner, B., Abou-Elwafa, S., Zhang, W., Jung, C., Müller** (2010) A survey of EMS-induced biennial *Beta vulgaris* mutants reveals a novel bolting locus which is unlinked to the bolting gene *B*. *Theor. App. Genet.* 121:1117-1131.
- Cattanach, A.W., Dexter, A.G., Oplinger, E.S.** (1991) Sugarbeets. *Alternative Field Crops Manual*. University of Wisconsin-Extension. *Available at:* <http://www.hort.purdue.edu/newcrop/afcm/sugarbeet.html>. (Accessed: September 1, 2011).
- Calflora** (2011) Search for Plants. *Available at:* < <http://www.calflora.org>>. (Accessed: June 27, 2011).
- Cerdeira, A.L., Duke, S.O.** (2006) The current status and environmental impacts of glyphosate-resistant crops: A review. *J. Environmental Quality* 35:1633-1658
- Coruzzi, G., Broglie, R., Edwards, C., Chua, N.** (1984) Tissue-specific and light-regulated expression of a pea nuclear gene encoding the small subunit of ribulose-1,5-bisphosphate carboxylase. *EMBO J.* 3:1671-1679
- Crop Protection Chemical Reference** (1996) Chemical and Pharmaceutical Press, New York.
- Darmency, H., Klein, E.K., De Garanbé, T.G., Gouyon, P.H., Richard-Molard, M., Muchembled, C.** (2009) Pollen Dispersal in Sugar Beet Production Fields. *Theoretical and Applied Genetics* 118:1083–1092.
- Dewar, A.M., Cooke, D.A.** (2006) Chapter 13: Pests. Pgs. 316–358. *In: Sugar Beet*. Draycott, A. P. (ed.). Oxford, United Kingdom: Blackwell Publishing Ltd. World Agricultural Series: 474 pgs.

- Estok, D., B. Freedman, Boyle, D. (1989)** Effects of the herbicides 2,4-D, glyphosate, Hexazinone, and Triclopyr on the growth of three species of ectomycorrhizal fungi. *Bull. Environ. Contam. Toxicol.* 42:835-839.
- FDA (1998)** Guidance for Industry: Use of Antibiotic Resistance Marker Genes in Transgenic Plants. pp. 28
- Gowda, S., Wu, F.C., Sheperd, R.J. (1989)** Identification of promoter sequences for the major RNA transcripts of figwort mosaic and peanut chlorotic streak viruses (caulimovirus group). *J. Cell. Biochem* 13D (supplement):301
- Hanson, L. (2003)** Effect of treating sugar beet with the herbicide glyphosate on *Fusarium* yellows. Proceeding of the American Society of Agronomy, Crop Science Society of America, and Soil Science Society of America. Published Sept. 2, 2003
- Howard, E., Citovsky, V., Zambryski, P. (1990)** The T-complex of *Agrobacterium tumefaciens*. *In: Plant Gene Transfer*. Beachy, R.N., Lamb, C.J. (eds.), Alan R. Liss, Inc., New York, New York. pp. 1-12.
- Ingram, J. (2000)** Separation Distances Required to Ensure Cross-pollination is Below Specified Limits in Non-seed Crops of Sugar Beet, Maize and Oilseed Rape. *Plant Varieties and Seeds* 13:181
- Johnson, R. T., Burtch, L.M. (1958)** The problem of wild annual sugar beets in California. *Jour. Am. Soc. of Sugar Beet Tech.* 10: 311-317
- Kaneko, T. et al. (2000)** Complete genome structure of the nitrogen-fixing symbiotic bacterium *Mesorhizobium loti*. *DNA Research* 7:331-338
- Kaneko, T. et al. (2002)** Complete genomic sequence of nitrogen-fixing symbiotic bacterium *Bradyrhizobium japonicum* USDA110. *DNA Research* 9:189-197
- Keeler, K. (1989)** Can genetically engineered crops become weeds? *Bio/Technology* 7:1134-1139
- Keese, P. (2008)** Risks from GMOs due to horizontal gene transfer. *Environ. Biosafety Res.* 7:123-149
- Khan, M.F. (2010)** Introduction of glyphosate-tolerant sugar beet in the United State. *Outlooks on Pest Management* 21:38-41
- Lange, W., Brandenburg, W.A., De Bock, T.S.M. (1999)** Taxonomy and cultonomy of beet (*Beta vulgaris* L.). *Botanical Journal of the Linnean Society* 130:81-96

- Larson, R.L., Hill, A.L., Fenwick, A., Kniss, A.R., Hanson, L.E., Miller, S.D.** (2006) Influence of glyphosate on *Rhizoctonia* and *Fusarium* root rot in sugar beet. *Pest Management Science* 62:1182-1192
- Lewellen, R.T., Liu, H.-Y., Wintermantel, W.M., Sears, J.L.** (2003) Inheritance of beet necrotic yellow vein virus (BNYVV) systemic infection in crosses between sugarbeet and *Beta macrocarpa*. 1st Joint IIRB-ASSBT Congress 149-160. Feb. 27-Mar. 1, 2003. San Antonio, TX.
- McFarlane, F.S.** (1975) Naturally occurring hybrids between sugar beet and *Beta macrocarpa* in the Imperial Valley of California. *Journal of American Society of Sugar Beet Technologists* 18:245-251
- Monsanto and KWS SAAT AG** (2004) Petition for Determination of Nonregulated Status for Roundup Ready® Sugar Beet Event H7-1. Submitted by R. Schneider, Regulatory Affairs Manager. Monsanto Company, St. Louis, MO and G. Strittmatter, Head of Research and Breeding. KWS SAAT AG, Einbeck, Germany. *Available at:* <http://www.aphis.usda.gov/biotechnology/not_reg.html>. (Accessed: June 27, 2011).
- Morelli, G., Nagy, F., Fraley, R.T., Rogers, S.G., Chua, N.H.** (1985) A short conserved sequence is involved in the light-inducibility of a gene encoding ribulose 1,5-bisphosphate carboxylase small subunit of pea. *Nature* 315:200-204
- OECD** (2001) Consensus Document on the biology of *Beta vulgaris* L. (Sugar Beet). In Series on Harmonization of Regulatory Oversight in Biotechnology, p. 40
- Padgett, S.R., Kolacz, K.H., Delannay, X., Re, D.B., Lavalley, B.J., Tinius, C.N., Rhodes, Y.I., Otero, W.K., Barry, G.F., Eichholtz, D.A., Peschke, V.M., Nida, D.L., Taylor, N.B., Kishore, G.M.** (1995) Development, identification, and characterization of a glyphosate-tolerant soybean line. *Crop Science* 35:1451-1461
- Padgett, S.R., Re, D.B., Barry, G.F., Eichholtz, D.E., Delannay, X., Fuchs, R.L., Kishore, G.M., Fraley, R.T.** (1996) New weed control opportunities: development of soybeans with a Roundup ready™ gene. pp. 54-84. *In* Agricultural, Environmental, Economic, Regulatory, and Technical Aspects. Duke, S.O. (ed). CRC Press Inc. 4:54-84
- Panella, L.** (2003) USDA/ARS Research Geneticist and Sugarbeet Germplasm Committee Chair, Personal communication to J.R. Stander, Betaseed, Inc. (Appendix 2 of the Petition).
- Panella, L., Lewellen, R.T.** (2007) Broadening the genetic base of sugar beet: introgression from wild relatives. *Euphytica* 154:383-400
- Parker, I.M., Bartsch, D.** (1996) Recent advances in ecological biosafety research on the risks of transgenic plants: a transcontinental perspective. In T Tomiuk, K Wöhrmann, A Sentker, eds., *Transgenic organisms: Biological and Social Implications*. Birkhause Verlag, Basel, pp. 147-161

- Reding, H.K.,** von dem Bussche, P., Hofmann, P. (2010) Letter to Michael C. Gregoire, USDA APHIS, July 29, 2010. Attached to Environmental Report submitted by Monsanto in support of petition 03-323-01p.
- Richins, R.D.,** Scholthof, H.B., Sheperd, R.J. (1987) Sequence of figwort mosaic virus DNA (Caulimovirus Group). *Nucl. Acids Res.* 15:8451-8466
- Sanger, M.,** Daubert, S., Goodman, R.M. (1990) Characteristics of a strong promoter from figwort mosaic virus: comparison with the analogous 35S promoter from cauliflower mosaic virus and the regulated mannopine synthase promoter. *Plant Molecular Biology* 14:433-443
- Sheperd, R.J.,** Richins, J.F., Duffus, J.F., Handley, M.K. (1987) Figwort mosaic virus: properties of the virus and its adaptation to a new host. *Phytopathology* 77:1668-1673
- Savitsky, H.** (1975) Hybridization between *Beta vulgaris* and *B. procumbens* and transmission of nematode (*Heterodera schachtii*) resistance to sugarbeet. *Can. J. Genet. Cytol.* 17:197-209.
- Timko, M.P.,** Herdies, L., de Alameida, E., Cashmore, A.R., Leemans, J., Krebbers, E. (1988) Genetic engineering of nuclear-encoded components of the photosynthetic apparatus of *Arabidopsis*. pp. 279-295. *In: The Impact of Chemistry on Biotechnology – A Multidisciplinary Discussion.* M. Phillips, S.P. Shoemaker, R.D. Middlekauff, R.M. Ottenbrite, editors. ACS Books, Washington, D.C.
- USDA APHIS, 2011.** Glyphosate-tolerant H7-1 Sugar Beets: Request for Nonregulated Status. Draft Environmental Impact Statement.
- USDA ARS,** United States Department of Agriculture Agricultural Research Service. 2010. National Genetic Resources Program Germplasm Resources Information Network - (GRIN) [Online Database]. National Germplasm Resources Laboratory, Beltsville, Maryland. Available at: <<http://www.ars-grin.gov/cgi-bin/npgs/acc/display.pl?1441384>>. (Accessed: June 27, 2011).
- Van Dijk, H.,** Boudry, P., McCombie, H., Vernet, P. (1997) Flowering Time in Wild beet (*Beta vulgaris* ssp. *maritima*) along a Latitudinal Cline. *Acta Oecology* 18: 47–60.
- Wan, M.T.,** Rahe, J.E., Watts, R.G. (1998) A new technique for determining the sublethal toxicity of pesticides to the vesicular-arbuscular mycorrhizal fungus *Glomus intraradices*. *Environ. Toxicol. Chem.* 17(7):14-21
- Whitney, E.D.,** Duffus, J.E. (1986) Compendium of Beet Diseases and Insects. pp. 26-32
- Wood, D.W., et al.** (2001) The genome of the natural genetic engineer *Agrobacterium tumefaciens* C58. *Science.* 294:2317-2323

WSSA (Weed Science Society of America) (1998) “Herbicide Resistance” and “Herbicide Tolerance” defined. Weed Technology. 12: 789