



Pale Cyst Nematode (*Globodera pallida*) Eradication Program- Idaho Falls, Idaho

July 2010 Report

Background

Pale cyst nematodes (PCN), *Globodera pallida*, are soil-borne organisms that do not infest potato tubers. The pests infest feeder roots, where the females attach, feed, and become sedentary. Nematodes reproduce sexually. Females form cysts containing 200 to 600 eggs, which can stay dormant for up to 30 years while the eggs inside remain viable. On host plants, large numbers of PCN can cause wilting, stunted growth, poor root development, and early plant death. If left uncontrolled, PCN can reduce yields up to 80 percent in potato fields. Even with only minor symptoms showing on the foliage, PCN can significantly reduce tuber size. PCN spread primarily by the transport of cysts in soil. This may occur with the movement of soil on farming, construction, and other equipment; infested soil adhering to seed potatoes and other regulated crops; and any other items or means of transport such as water.

On April 19, 2006, officials of USDA's Animal and Plant Health Inspection Service (APHIS) and the Idaho State Department of Agriculture (ISDA) announced the detection of PCN, a major pest of potato crops. This was the first detection of the pest in the United States. The nematode cysts were detected during a routine survey of tare soil at an ISDA grading facility in eastern Idaho. Subsequent 2006 surveying to determine the possible origin and distribution of the pest in Idaho confirmed seven PCN-infested fields totaling 911 acres, all within a one mile radius in Bingham and Bonneville Counties, Idaho. The PCN-infested fields and an area surrounding the fields were placed under a Federal Domestic Quarantine Order and parallel State Rule in August 2006, establishing restrictions on movement of certain regulated articles from Idaho in order to prevent the spread of PCN.

As a result of continued intensive soil sampling in 2007 and 2008, an additional two PCN-infested fields were found in Bingham County, Idaho. The nine PCN-infested fields all continue to be within a one mile radius and the fields associated with them through shared tenancy, farming practices, equipment, and/or shared borders have been extensively surveyed and regulated. Since program inception, a total of 30,753 acres have been regulated due to their infestation or association with an infested field. Non-infested, associated fields have been eligible for federal deregulation following a sequence of soil surveys with no PCN detections. To date, 29,327 acres have been released from federal regulation. Currently, 1,426 acres remain regulated, of which 1,100 are PCN-infested.

Eradication treatments of PCN-infested fields have been ongoing since the spring of 2007. Eradication treatments have included methyl bromide fumigation, Telone II fumigation, and biofumigant plantings. Testing of the soil in infested fields as per the Guidelines indicate the average viability of eggs within the PCN cysts have declined by more than 95% since eradication treatments began.

A description of the current PCN regulated area can be found at:

http://www.aphis.usda.gov/plant_health/plant_pest_info/potato/pcn-maps.shtml

The current Federal PCN rule revised as of January 1, 2010:

http://www.aphis.usda.gov/plant_health/plant_pest_info/potato/downloads/pcndocs/7cfr-10.txt

Survey Information

Type of survey	Idaho soil samples collected	
	July 2010	Since program inception
Detection	0	96,804
Delimiting	0	128,058
Eradication	236	52,586
Total	236	277,448

Identification and Diagnostics

Type of survey	Samples processed by the Idaho PCN Laboratory			Results		
	July 2010	2010 Year to date	Since program inception	July 2010 results	2010 Year to date	Results since program inception
Detection	0	39,104	69,934	Negative	Negative	Negative
Delimiting	0	2,511	117,613	Negative	Negative	Negative ¹
Eradication	711	2,831	49,145	N/A	N/A	N/A
Total	711	44,446	236,692			

¹Except for samples confirmed for the nine infested fields

Type of survey	Samples processed by the Idaho Food Quality Assurance Laboratory	
	Since program inception	Results
Detection	35,524	Negative
Delimiting	31,097	Negative ¹
Total	66,621	

¹Except for samples confirmed for the nine infested fields

Program Research

No report for July

Eradication Activities

There were no eradication treatments in the infested fields in July.

Telone II application to the PCN-infested fields is scheduled for mid-August, 2010. The amended solicitation for bids will close 8/5. Telone II was applied in the late summer of 2007 and 2008. There was no Telone II application in 2009 due to a world-wide shortage of this chemical.

Biofumigants with nematicidal activity were planted in the infested fields in the summers of 2007 (oil radish) and 2009 (arugula).

Regulatory Actions

The PCN-regulated area was not amended in July.

Regulatory Treatments

Treatment type	Regulatory Treatments (# of pieces of equipment)		
	July 2010	2010 Year to date	Since program inception ¹
Pressure Washed	41	232	>6,500
Steam Sanitized	1	84	>800
Total	42	316	>7,300

¹A review of regulatory data is underway while new data collection and management tools are developed.

Regulatory Documentation

Documentation type	Regulatory Documentation		
	July 2010	2010 Year to date	Since program inception ¹
Certificate (PPQ 540)	9	143	>5,300
Limited Permit (PPQ 530)	29	170	>1,100
Active compliance agreements	0	6	136

¹A review of regulatory data is underway while new data collection and management tools are developed.

Impacts on Commerce

In response to the initial PCN detection in 2006, Canada, Mexico and Korea shut off importation of potatoes from Idaho, while Japan cut off importation of potatoes from the entire U.S. The Mexican and Canadian export markets have both been re-opened with the exception of potatoes from PCN-regulated areas. Both require PCN soil surveys from origin fields. The Korean market was reopened in June, 2010 with the exception of potatoes originating from Bingham and Bonneville Counties, ID. The Japanese market remains closed to Idaho potatoes but negotiations are actively underway to regain market access. Because of extensive field surveys conducted throughout production areas in Idaho, all of which have been negative beyond the nine infested fields, the general opinion by our trading partners is that potatoes produced outside regulated areas do not pose the biological risk for introduction of PCN.

Communication and Outreach

There is a regional sewer project being built in Shelley which will encompass many surrounding communities. The project plans to lay a main collection pipe along Country Club Road, which has

infested fields on each side of the road. Regulatory and sanitation issues have been brought to the attention of project principals. Work is scheduled to begin near the infested fields in early August.

The next stakeholder update is due out in September, 2010. Stakeholder updates are available at:
http://www.aphis.usda.gov/plant_health/plant_pest_info/potato/pcn_stakeholder.shtml