



United States  
Department of  
Agriculture

Animal and Plant  
Health Inspection  
Service

September 21, 2012

Version 3



## Weed Risk Assessment for *Oplismenus hirtellus* (L.) P. Beauv. subsp. *undulatifolius* (Ard.) U. Scholz (Poaceae) – Wavyleaf basketgrass



*Oplismenus hirtellus* subsp. *undulatifolius* infestation (photo by Kerrie L. Kyde). Insets from left to right show foliage (photo by K. L. Kyde), flower (photo by K. L. Kyde), and seeds (photo by Garrett Waugaman). Source for images : [www://invasives.org](http://www://invasives.org).

### Agency Contact:

Plant Epidemiology and Risk Analysis Laboratory  
Center for Plant Health Science and Technology

Plant Protection and Quarantine  
Animal and Plant Health Inspection Service  
United States Department of Agriculture  
1730 Varsity Drive, Suite 300  
Raleigh, NC 27606

**Introduction** Plant Protection and Quarantine (PPQ) regulates noxious weeds under the authority of the Plant Protection Act (7 U.S.C. § 7701-7786, 2000) and the Federal Seed Act (7 U.S.C. § 1581-1610, 1939). A noxious weed is “any plant or plant product that can directly or indirectly injure or cause damage to crops (including nursery stock or plant products), livestock, poultry, or other interests of agriculture, irrigation, navigation, the natural resources of the United States, the public health, or the environment” (7 U.S.C. § 7701-7786, 2000). We use weed risk assessment (WRA)—specifically, the PPQ WRA model<sup>1</sup>—to evaluate the risk potential of plants, including those newly detected in the United States, those proposed for import, and those emerging as weeds elsewhere in the world.

Because our WRA model is geographically and climatically neutral, it can be used to evaluate the baseline invasive/weed potential of any plant species for the entire United States or any area within it. We use a climate matching tool in our WRAs to evaluate those areas of the United States that are suitable for the establishment of the plant. We also use a Monte Carlo simulation to evaluate the consequences of uncertainty on the outcome of the risk assessment. For more information on the PPQ WRA process, please refer to the document, *Introduction to the PPQ Weed Risk Assessment Process*, which is available upon request.

---

***Oplismenus hirtellus* (L.) P. Beauv. subsp. *undulatifolius* (Ard.) U. Scholz –  
Wavyleaf basketgrass**

---

**Species** Family: Poaceae

**Information** Initiation: Al Tasker, APHIS PPQ National Noxious Weed Program Manager, requested that the Plant Epidemiology and Risk Analysis Laboratory (PERAL) Weed Team conduct a weed risk assessment for *Oplismenus hirtellus* subsp. *undulatifolius*, wavyleaf basketgrass. This request was initiated after he was informed of an infestation in Maryland. Although *O. hirtellus* is sold by the nursery industry, molecular work conducted by Sharon Talley, PPQ Center for Plant Health and Science Technology (CPHST), confirmed that this was not the same taxon being sold by the nursery industry (Tasker, 2010).

Foreign distribution: *Oplismenus hirtellus* subsp. *undulatifolius* occurs in temperate, subtropical, and tropical areas of the world (Scholz, 1981). It occurs in many African countries (including Madagascar), temperate and tropical Asia, Australia, and parts of Europe (Scholz, 1981; USDA ARS NGRP, 2009).

U.S. distribution and status: First reported in the United States in 1996 in a state park in Maryland, *O. hirtellus* subsp. *undulatifolius* has since been found in at least seven locations in Maryland (Westbrooks and Imlay, 2009) and four in Virginia (Akerson, 2010). It is reported as having been present in Hawaii for over 150 years (Snow and Lau, 2010), but there is some debate as to the identity of the subspecies in Hawaii (Talley, 2012). Maryland and Virginia have not taken regulatory action against this taxon, although considerable effort has been made at state and local levels to combat it. Eradication and control efforts include Early Detection Rapid Response (EDRR) strategies (Kyde and Marose, 2008), and ongoing removal by county and volunteer weed crews (MDDNR, n.d.). In pure

---

<sup>1</sup> Koop, A., L. Fowler, L. Newton, and B. Caton. 2012. Development and validation of a weed screening tool for the United States. *Biological Invasions* 14(2):273-294. DOI:10.1007/s10530-011-0061-4

stands, the herbicide glyphosate is used to kill the plants, while hand-removal is used when plants are growing among native species.

WRA area: The United States, including its territories and possessions.

---

1. *Oplismenus hirtellus* subsp. *undulatifolius* analysis

**Establishment/Spread Potential** In Maryland, *O. hirtellus* subsp. *undulatifolius* has demonstrated a high capacity to establish and spread; since its initial detection in 1996, it has spread to at least seven locations (Westbrooks and Imlay 2009). The most notable characteristics contributing to its high establishment/spread risk score include the following: self reproduction (Scholz, 1981); a short generation time; and sticky propagules that promote dispersal (Kyde, 2010; Scholz, 1981; Talley, 2010). This taxon's rapid spread in Maryland is likely due to unintentional dispersal by bikers, dog walkers, and hikers who use infested areas. For example, a recent article in the Baltimore Sun reports "After a recent hike through Patapsco Valley State Park, Baltimore teacher Greg Schnitzlein's jaw dropped as he watched his two dogs emerge from the woods looking, as he says, like Chia Pets, every inch of their fur slathered in sproutlike seeds" (Rosen, 2011). Although we found little information about *O. hirtellus* subsp. *undulatifolius*, beyond what is known in the United States, this risk element had a relatively low level of uncertainty due to contributions from an expert in PPQ (Talley, 2010).

Risk score = 15                      Uncertainty index = 0.11


**Impact Potential** Once *O. hirtellus* subsp. *undulatifolius* becomes established, it spreads rapidly through wooded natural areas (Westbrooks and Imlay, 2009). It forms dense carpets in the forest understory (Beauchamp, 2012; Kyde, 2010), at times creating a new layer of plant vegetation in habitats (Kyde, 2010). Understory species richness and *O. hirtellus* subsp. *undulatifolius* cover are strongly and negatively correlated (Beauchamp, 2012). Research is still ongoing, however, to determine whether it invades areas that already have low volumes of understory vegetation, or whether it outcompetes existing native vegetation (Beauchamp, 2012). This taxon affects recreational use of infested areas because people do not want themselves or their pets to get covered in seeds (Kyde, 2010; Rosen, 2011). *Oplismenus hirtellus* subsp. *undulatifolius* has been observed in lawns in Virginia (Hughes, 2010). The U.S. Fish and Wildlife Service helps private landowners control this taxon in suburban areas to maintain suitable wildlife on their property (Kyde, 2010). This risk element had a high level of uncertainty due to a lack of knowledge on what impacts this taxon may have in anthropogenic and production systems.

Risk score = 3.3                      Uncertainty index = 0.27

**Geographic Potential** We estimate that about 30 percent of the United States is suitable for the establishment of *O. hirtellus* subsp. *undulatifolius* (Fig. 1). That predicted distribution is based on the taxon's known distribution elsewhere in the world and includes point-referenced localities and areas of occurrence. The map for *O. hirtellus* subsp. *undulatifolius* represents the joint distribution of USDA Plant Hardiness Zones 6-13, areas with 30-100+ inches of annual precipitation, and the following Köppen-Geiger climate classes: tropical rainforest, tropical savannah, mediterranean, humid subtropical, marine west coast, humid continental warm summers, and humid continental cool summers.

**Entry Potential** Because *Oplismenus hirtellus* subsp. *undulatifolius* is already established in the United States (Westbrooks and Imlay 2009), assessment of this risk element was not necessary.

**Figure 1.** Predicted distribution of *Oplismenus hirtellus* subsp. *undulatifolius* in the United States. Map insets for Alaska, Hawaii, and Puerto Rico are not to scale.




## 2. Results and Conclusion

Model Probabilities: P(Major Invader) = 80.0%  
P(Minor Invader) = 19.3%  
P(Non-Invader) = 8.0%


Risk Result = High Risk

Secondary Screening = Not Applicable

**Figure 2.** *Oplismenus hirtellus* subsp. *undulatifolius* risk score (black box) relative to the risk scores of species used to develop and validate the WRA model (other symbols). See Appendix A for the complete assessment.



**Figure 3.** Monte Carlo simulation results (N=5000) for uncertainty around the risk scores for *Oplismenus hirtellus* subsp. *undulatifolius*<sup>a</sup>.



<sup>a</sup>The blue “+” symbol represents the medians of the simulated outcomes. The smallest box contains 50 percent of the outcomes, the second 95 percent, and the largest 99 percent.

### 3. Discussion

The result of the weed risk assessment for *O. hirtellus* subsp. *undulatifolius* is High Risk with a relatively low level of uncertainty (Figs. 2 and 3). This taxon forms dense mats, spreads rapidly through wooded areas (Westbrooks and Imlay, 2009), and renders areas undesirable for recreational use (Kyde, 2010). A strong negative correlation exists between understory species richness and *O. hirtellus* subsp. *undulatifolius* cover (Beauchamp, 2012). A taskforce was established in 2008 to focus on mapping and eradication of known infestations (Westbrooks and Imlay, 2009). The taskforce has developed a public awareness campaign to help detect and report infestations before they become widespread. Land managers believe that, if left unchecked, *O. hirtellus* subsp. *undulatifolius* will have a devastating impact on the deciduous forests of eastern North America for many decades (Westbrooks and Imlay, 2009). Plant managers, state and local native plant societies, Native American tribes, and others wrote an open letter to Congress requesting funding to combat *O. hirtellus* subsp. *undulatifolius* in Maryland and Virginia and prevent its spread into other states (Ford and Imlay, 2012). This taxon may be elevated to the species rank in the future (Thompson, 2012).

### 4. Literature Cited

- 7 U.S.C. § 1581-1610. 1939. The Federal Seed Act, Title 7 United States Code § 1581-1610.
- 7 U.S.C. § 7701-7786. 2000. The Federal Seed Act, Title 7 United States Code § 7701-7786.
- Akerson, J. 2010. Wavyleaf basketgrass in Ovaka Virginia.
- Basnet, D. B. 2005. Some common weed flora in forest plantation of Darjeeling Hill, West Bengal, India. in A. Kuman, (ed.). Biodiversity & Conservation. A.P.H. Publishing Company, New Delhi.
- Beauchamp, V. B. 2012. *Oplismenus* WRA. Email from Vanessa Beauchamp, Towson University, to Leslie Newton, PERAL, on July 20, 2012. (Archived at PERAL).
- Burrows, G. E., and R. J. Tylr. 2001. Toxic Plants of North America (First). Iowa State University Press, Ames.
- Culley, T. M., S. G. Weller, and A. K. Sakai. 2002. The evolution of wind pollination in angiosperms. *Trends in Ecology & Evolution* 17(8):361-369.
- Dave's Garden. 2008. Dave's Garden. <http://davesgarden.com>. (Archived at PERAL).
- Enomoto, T. 2003. Weeds of Japan. Okayama University, The Research Institute for Bioresources, Laboratory of Wild Plant Science. Last accessed August 28, 2008, [http://www.rib.okayama-u.ac.jp/wild/zassou\\_table.htm](http://www.rib.okayama-u.ac.jp/wild/zassou_table.htm). (Archived at PERAL).
- Ford, A., and M. Imlay. 2012. Open letter to Congress: Control of Asiatic wavyleaf basketgrass, *Oplismenus hirtellus undulatifolius*. Letter written (February 3, 2012) to members of Congress on behalf of the Virginia Native Plant Society, the Maryland Native Plant Society, and 24 other organizations requesting funding to contain *O. hirtellus* subsp. *undulatifolius* in Maryland and Virginia. Available at <http://ipetrus.blogspot.com/2012/02/open-letter-to-congress-control-of.html>.
- Holm, L. G., J. V. Pancho, J. P. Herberger, and D. L. Plucknett. 1979. A Geographical Atlas of World Weeds. Krieger Publishing Company, Malabar, Florida, U.S.A. 391 pp.

- Hughes, J. 2010. Wavyleaf Basketgrass Task Force. Personal communication to S. Talley on June 30, 2010, from Jake Hughes, Shenandoah National Park, regarding the threat of wavyleaf basketgrass upon T & E species.
- Kyde, K. L. 2010. Field investigation and greenhouse experimentation of *Oplismenus hirtellus* spp. *undulatifolius*. Personal communication to S. Talley on March 4, 2010, from K. Kyde, Maryland Department of Natural Resources, regarding the report by Sharon Talley, PPQ CPHST, entitled Field investigation and greenhouse experimentation of *Oplismenus hirtellus* ssp. *undulatifolius*. (Archived at PERAL).
- Kyde, K. L. 2012. *Oplismenus* WRA. Email from Kerrie Kyde, Maryland Department of Natural Resources, to Tony Koop, PERAL, on 6/26/2012. (Archived at PERAL).
- Kyde, K. L., and B. H. Marose. 2008. Wavyleaf basketgrass in Maryland: an early detection rapid response program in progress. Poster presented at the 2008 Northeastern Weed Science Society meeting.
- Martin, P. G., and J. M. Dowd. 1990. A protein sequence study of the dicotyledons and its relevance to the evolution of the legumes and nitrogen fixation. *Australian Systematic Botany* 3:91-100.
- McCullum, S. 2012. *Oplismenus hirtellus* subsp. *undulatifolius* WRA. Email from Shiloh McCullum, University of Alabama - Huntsville, to Leslie Newton, PERAL, on 7/6/2012. (Archived at PERAL).
- MDDNR. n.d. Wavyleaf basketgrass: management of wavyleaf basketgrass in Maryland. Maryland Department of Natural Resources (MDDNR), Annapolis, MD. Last accessed April 17, 2012, [http://www.dnr.state.md.us/wildlife/Plants\\_Wildlife/WLBG/wl\\_basketgrass\\_management.asp](http://www.dnr.state.md.us/wildlife/Plants_Wildlife/WLBG/wl_basketgrass_management.asp).
- Peterson, P. M., E. E. Terrell, E. C. Uebel, C. A. Davis, H. Scholz, and R. J. Soreng. 1999. *Oplismenus hirtellus* subspecies *undulatifolius*, a new record for North America. *Castanea* 64(1999):201-202.
- Randall, J. M. 2007. The introduced flora of Australia and its weed status. CRC for Australian Weed Management, Department of Agriculture and Food, Western Australia, Australia. 528 pp.
- Ricketts, T. H., E. Dinerstein, D. M. Olson, C. J. Loucks, W. Elchbaum, D. DellaSala, K. Kavanagh, P. Hedao, P. T. Hurley, K. M. Carney, R. Abell, and S. Walters. 1999. *Terrestrial Ecoregions of North America: A Conservation Assessment*. Island Press, Washington, D.C. 485 pp.
- Rosen, J. 2011. Maryland has become ground zero for a new invasive threat, wavyleaf basketgrass. *The Baltimore Sun*, October 22, 2011.
- Scholz, U. 1981. Monographia der Gattung *Oplismenus* (Gramineae). *Phanerogamarum monographiae* 13(1981):1-213.
- Snow, N., and A. Lau. 2010. Notes on grasses (Poaceae) in Hawaii. *Bishop Museum Occasional Papers* 107:46-60.
- Talley, S. 2010. Field observation, export review, and greenhouse experimentation of *Oplismenus hirtellus* subsp. *undulatifolius*. USDA APHIS PPQ CPHST, Ft. Collins.
- Talley, S. 2012. *Oplismenus hirtellus*. Email from Sharon Talley, APHIS BRS, to Tony Koop, PERAL, on 8/7/2012. (Archived at PERAL).
- Tasker, A. 2010. WLBG (wavyleaf basketgrass). Personal communication to Larry Fowler, PPQ CPHST, on the initiation of a weed risk assessment for wavyleaf basketgrass from AI Tasker, APHIS PPQ. (Archived at PERAL).
- Thompson, J. P. 2012. *Oplismenus hirtellus* ssp. *undulatifolius* (Ard.) U. Scholz.

- Datasheet for CABI Invasive Species Compendium.  
University of Minnesota. 2008. Plant Information Online. University of Minnesota.  
<https://plantinfo.umn.edu/arboretum/default.asp>. (Archived at PERAL).
- USDA APHIS PPQ CPHST. 2007. NPAG Report: *Oplismenus hirtellus* spp.  
*undulatifolius* (Ard.) U. Scholtz, Wavyleaf basketgrass. USDA APHIS  
PPQ CPHST, Raleigh. 3 pp.
- USDA ARS NGRP. 2009. World Economic Plants in GRIN (Germplasm  
Resources Information Network). United States Department of Agriculture,  
Agricultural Research Service, National Germplasm Resources Laboratory  
(NGRP). Beltsville. Last accessed 09/03/2009. <http://www.ars-grin.gov/cgi-bin/npgs/html/taxecon.pl?language=en>. (Archived at PERAL).
- USFWS. 2010. Threatened and endangered species system (TESS).  
[http://ecos.fws.gov/tess\\_public/pub/SpeciesReport.do?dsource=plants](http://ecos.fws.gov/tess_public/pub/SpeciesReport.do?dsource=plants).  
(Archived at PERAL).
- Walker, R. 2003. Parasitic Plants Database.  
[http://www.omnisterra.com/bot/pp\\_home.cgi](http://www.omnisterra.com/bot/pp_home.cgi). (Archived at PERAL).
- Westbrooks, R., and M. Imlay. 2009. Wavyleaf basketgrass - a new invader of  
deciduous forests in Maryland and Virginia. <http://www.se-eppc.org/southcarolina/WLBG.pdf>. (Archived at PERAL).

**Appendix A.** Weed risk assessment for *Oplismenus hirtellus* subsp. *undulatifolius* (Poaceae). The following information was obtained from the species' risk assessment, which was conducted using Microsoft Excel. The information shown below was modified to fit on the page. The original Excel file, the full questions, and the guidance to answer the questions are available upon request.

Question ID	Answer - Uncertainty	Score	Notes (and references)
<b>Establishment/Spread Potential</b>			
ES-1 (Invasiveness elsewhere)	f - negl	5	<i>Oplismenus hirtellus</i> subsp. <i>undulatifolius</i> is reported as having naturalized outside its native range in Australia (Randall, 2007). This rapidly spreading understory grass is invading the states of Maryland and Virginia (Kyde, 2010; Westbrook and Imlay, 2009). Once established, it spreads rapidly through wooded natural areas... (Westbrook and Imlay, 2009). The sticky awns allow the grass seeds to easily spread by adhering to passing animals, people, and vehicles (Westbrook and Imlay, 2009). This subspecies is reported as having been present in Hawaii for over 150 years (Snow and Lau, 2010), although there is some debate as to whether the subspecies in Hawaii is really subsp. <i>undulatifolius</i> (Talley, 2012). Alternate answers for the Monte Carlo simulation are both "e".
ES-2 (Domesticated to reduce weed potential)	n - low	0	<i>Oplismenus hirtellus</i> subsp. <i>undulatifolius</i> is not known to be cultivated. A closely related taxon, <i>O. hirtellus</i> subsp. <i>varigatus</i> , is sold as an ornamental grass by the horticulture industry (Dave's Garden, 2008; University of Minnesota, 2008). It is not clear if this cultivar has been bred for reduced weediness potential. From DNA sequence comparisons of variable regions, the horticultural variety appears to be a hybrid and is believed to be sterile (Talley, 2010).
ES-3 (Weedy congeners)	y - mod	1	<i>Oplismenus compositus</i> is recorded as a common weed of agriculture in Borneo and Malaysia (Holm et al., 1979) and is listed as a major weed in forest plantations in West Bengal, India (Basnet, 2005).
ES-4 (Shade Tolerance)	y - negl	1	<i>Oplismenus</i> species are exclusively grasses of forest understory and are found in areas of moderate to dense shade (Scholz, 1981). <i>Oplismenus hirtellus</i> subsp. <i>undulatifolius</i> is found in shady forests (Scholz, 1981). Generally <i>O. hirtellus</i> subsp. <i>undulatifolius</i> does not grow beyond the treeline (Kyde, 2010). <i>O. hirtellus</i> subsp. <i>undulatifolius</i> grows well under 70% shade in the greenhouse (Talley, 2010).
ES-5 (Climbing or smothering growth form)	n - negl	0	<i>Oplismenus hirtellus</i> subsp. <i>undulatifolius</i> is a stoloniferous grass (Peterson et al., 1999; Scholz, 1981) and has not been observed to have a climbing or smothering habit (Kyde, 2010; Talley, 2010).
ES-6 (Dense Thickets)	y - negl	2	<i>Oplismenus hirtellus</i> subsp. <i>undulatifolius</i> forms dense carpets in the forest understory (Beauchamp, 2012; Kyde, 2010). It is further described as "covering all our woodland habitats like astroturf, killing all the other herbaceous plants" (Ford and Imlay, 2012).
ES-7 (Aquatic)	n - negl	0	<i>Oplismenus hirtellus</i> subsp. <i>undulatifolius</i> is a terrestrial plant (Scholz, 1981) and is not an obligate aquatic plant. <i>O. hirtellus</i> subsp. <i>undulatifolius</i> has been observed growing up to the edge of streams but not in the stream (Beauchamp, 2012; Kyde, 2010).

ES-8 (Grass)	y - negl	1	<i>Oplismenus hirtellus</i> subsp. <i>undulatifolius</i> is a grass and belongs to the Poaceae (Peterson et al., 1999; Scholz, 1981).
ES-9 (N2-fixer)	n - negl	0	There is no evidence that any members of the family Poaceae have nitrogen-fixing capabilities (Martin and Dowd, 1990).
ES-10 (Viable seeds)	y - negl	1	Most species of the genus <i>Oplismenus</i> produce fertile caryopses (Scholz, 1981). <i>O. hirtellus</i> subsp. <i>undulatifolius</i> produces abundant seed; germination rates of <i>O. hirtellus</i> subsp. <i>undulatifolius</i> seeds from their naturalized range in the U.S. can be as high as 97%, when the caryopsis coat is removed to prevent dormancy (Talley, 2010). Recruitment from seedlings is evident from seedlings appearing in late spring and early summer (Kyde, 2010). Soil collected in the fall produced seedlings the following spring (Beauchamp, 2012).
ES-11 (Self-compatible)	y - mod	1	Reproduction in most <i>Oplismenus</i> species is likely apomictic (Scholz, 1981). Florets have both male and female parts (Scholz, 1981). Most grasses are wind or self pollinated. Abundant fertile seeds of <i>O. hirtellus</i> subsp. <i>undulatifolius</i> are produced from greenhouse grown plants that have little to no exposure to pollinators or wind (Talley, 2010).
ES-12 (Special Pollinators)	n - negl	0	No pollinators for <i>Oplismenus hirtellus</i> subsp. <i>undulatifolius</i> have been observed in the field (Kyde, 2010). Because abundant fertile seeds were produced in the greenhouse from plants with little to no exposure to pollinators, pollinators do not appear necessary for effective reproduction (Talley, 2010). Reproduction in most <i>Oplismenus</i> species is likely apomictic (Scholz, 1981). Most grasses (Poaceae) and sedges (Cyperaceae) are wind pollinated (Culley et al., 2002).
ES-13 (Min generation time)	b - low	1	<i>Oplismenus hirtellus</i> subsp. <i>undulatifolius</i> is capable of regenerating from seeds and vegetatively through stolons (Kyde, 2010; Scholz, 1981; Talley, 2010). Seeds can germinate in a few days and it takes about 4 months for seedling to mature enough to produce viable seeds (Talley, 2010). Mature plants that are vegetatively propagated can produce seeds within one month or less (Talley, 2010). Under greenhouse conditions healthy mature plants can produce seeds all year (Talley, 2010). In early spring in MD, <i>O. hirtellus</i> subsp. <i>undulatifolius</i> emerges from overwintering roots. Seed production peaks in August and seedlings emerge the following late spring (Kyde, 2010). Alternate answers for the Monte Carlo simulation are "a" and "c".
ES-14 (Prolific reproduction)	n - low	-1	Rough calculations for plants in the field have been estimated at 1250 seeds/m <sup>2</sup> /yr (Kyde, 2012) and in greenhouse trials at 3,155 seeds/m <sup>2</sup> /yr (McCollum, 2012). Greenhouse grown plants have not been estimated to produce more than 5000 seeds/ m <sup>2</sup> /yr (Talley, 2010).
ES-15 (Unintentional dispersal)	y - negl	1	Seeds of <i>O. hirtellus</i> subsp. <i>undulatifolius</i> are wrapped in a visid awn that aids in its epizoochoric dispersal; seeds readily adheres to human hair, skin, and wool (Scholz, 1981). Seeds with awns also stick to clothing, leather, and rubber (Kyde, 2010; Talley, 2010). Mountain bikers have claimed that seeds can spread via mountain biking activities (Kyde, 2010). Seeds readily stick to the fur of dogs (Rosen, 2011).

Weed Risk Assessment for *Oplismenus hirtellus* subsp. *undulatifolius*

ES-16 (Trade contaminant)	y - low	2	<i>Oplismenus hirtellus</i> subsp. <i>undulatifolius</i> is found in forested areas and can stick to timber and be transported during months when fruit is ripe (Kyde, 2010).
ES-17 (#Natural dispersal vectors)	1	-2	0
ES-17a (Wind dispersal)	n - negl		<i>Oplismenus hirtellus</i> subsp. <i>undulatifolius</i> caryopses do not have wind-specialized awns or other structures that would facilitate wind dispersal; <i>O. hirtellus</i> subsp. <i>undulatifolius</i> caryopses are an oval shape that is not conducive to wind dispersal (Talley, 2010).
ES-17b (Water dispersal)	n - negl		Viable <i>O. hirtellus</i> subsp. <i>undulatifolius</i> seeds are not buoyant and therefore are not adapted to dispersal by water (Talley, 2010).
ES-17c (Bird dispersal)	? - max		Unknown. <i>O. hirtellus</i> subsp. <i>undulatifolius</i> seeds stick to almost anything and can potentially stick to the feet and feathers of ground foraging birds (Beauchamp, 2012; Talley, 2010); however we have no data to support this.
ES-17d (Animal external dispersal)	y - negl		<i>O. hirtellus</i> subsp. <i>undulatifolius</i> caryopses are spread by animals externally via a sticky awn (Scholz, 1981).
ES-17e (Animal internal dispersal)	n - high		There is no evidence that <i>O. hirtellus</i> subsp. <i>undulatifolius</i> is eaten by animals such as deer and birds (Beauchamp, 2012; Kyde, 2010). However, no formal studies have evaluated whether deer use the plant as forage and the caryopsis coat could allow some seeds to pass undigested (Talley, 2010).
ES-18 (Seed bank)	? - max	0	Unknown.
ES-19 (Tolerance to loss of biomass)	n - mod	-1	There is no evidence that <i>O. hirtellus</i> subsp. <i>undulatifolius</i> can withstand mutilation or fire. Hand pulling appears to be an effective form of control (Kyde, 2010).
ES-20 (Herbicide resistance)	n - low	0	<i>Oplismenus hirtellus</i> subsp. <i>undulatifolius</i> can be killed though herbicide treatments with a 1-2% solution of glyphosate (Westbrooks and Imlay, 2009). Herbicide trials with eight different herbicides demonstrated that each herbicide has some effect on <i>O. hirtellus</i> subsp. <i>undulatifolius</i> survival (Kyde, 2010).
ES-21 (# Cold hardiness zones)	8	0	
ES-22 (# Climate types)	7	2	
ES-23 (# Precipitation bands)	8	1	
<b>Impact Potential</b>			
<b>General Impacts</b>			
Imp-G1 (Allelopathic)	n - low	0	An extensive literature review did not reveal any evidence that any member of the <i>Oplismenus</i> genus is allelopathic. Other plants appear to grow well with <i>O. hirtellus</i> subsp. <i>undulatifolius</i> in soil and in hydroponic solutions (Talley, 2010). Additionally, no evidence has been revealed through field observations (Kyde, 2010).
Imp-G2 (Parasitic)	n - negl	0	An extensive literature review did not reveal any evidence that any member of the <i>Oplismenus</i> genus is parasitic. <i>Oplismenus hirtellus</i> subsp. <i>undulatifolius</i> belongs to the Poaceae, and there are no known grasses that are parasitic. <i>O. hirtellus</i> subsp. <i>undulatifolius</i> produces chlorophyll and the plant does not require nutrients from other plants (Talley, 2010). No evidence through field observations (Kyde, 2010). No evidence after searching the online parasitic plant database (Walker, 2003).

<b>Impacts to Natural Systems</b>			
Imp-N1 (Ecosystem processes)	n - mod	0	No evidence that <i>O. hirtellus</i> subsp. <i>undulatifolius</i> changes the physical properties of the ecosystem.
Imp-N2 (Community structure)	y - low	0.2	<i>O. hirtellus</i> subsp. <i>undulatifolius</i> forms a dense carpet layer in the forest understory and has been found to grow underneath Japanese stilt grass (Beauchamp, 2012; Kyde, 2010), thus creating either a new canopy layer where stiltgrass does not occur or an additional layer where stiltgrass does occur. Once it becomes established, <i>O. hirtellus</i> subsp. <i>undulatifolius</i> spreads rapidly through wooded natural areas; it is reported to prevent regeneration of native hardwood tree species (Westbrooks and Imlay, 2009), however, this has not been fully documented (Beauchamp, 2012).
Imp-N3 (Community composition)	y - high	0.2	Once it becomes established, this taxon spreads rapidly through wooded natural areas; it is reported to crowd out and prevent regeneration of native species (Westbrooks and Imlay, 2009). There is a strong negative correlation between understory species richness and wavyleaf basketgrass cover (Beauchamp, 2012). There is still some debate as to whether it is invading areas that already have low volumes of understory vegetation or whether it is outcompeting existing native vegetation; research is underway to determine the true competitive nature of this species (Beauchamp, 2012). The large image on the cover page for this assessment is a photograph of a forest floor covered with a blanket of wavyleaf basketgrass (source: Kerry Kyde, <a href="http://www.invasives.org">www.invasives.org</a> ).
Imp-N4 (T&E species)	y - low	0.1	There are 22 federally threatened or endangered plants in Maryland and Virginia (USFWS, 2010). Jake Hughes, National Park Service, reported to Sharon Talley, PPQ CPHST, that <i>Isotria medeoloides</i> (small whorled pogonia) is a federally threatened species occurring in Virginia that could be negatively impacted by <i>Oplismenus hirtellus</i> subsp. <i>undulatifolius</i> in their state—both occur in the same habitats (Hughes, 2010).
Imp-N5 (Globally outstanding ecoregions)	y - mod	0.1	There are globally-outstanding ecoregions on the east coast of the United States where wavyleaf basketgrass is growing and could spread (Ricketts et al., 1999). Given the impacts described above, it is likely to impact these ecoregions.
Imp-N6 (Natural systems weed)	c - negl	0.6	There are active control programs to manage <i>O. hirtellus</i> subsp. <i>undulatifolius</i> in Patapsco Valley State Park, Shenandoah National park, Rosaryville State Park, Little Paint Branch Park, and Hernwood Sanitary Landfill (Kyde, 2010). Alternate answers for the uncertainty simulation are both "b".
<b>Impact to Anthropogenic areas (cities, suburbs, roadways)</b>			
Imp-A1 (Affects property, civilization, ...)	n - mod	0	No evidence.
Imp-A2 (Recreational use)	y - low	0.1	Hunters do not want to bow hunt in infested areas because <i>O. hirtellus</i> subsp. <i>undulatifolius</i> seeds sticks to clothing. Some people do not want to walk dogs in infested areas because <i>O. hirtellus</i> subsp. <i>undulatifolius</i> seeds stick to dog fur (Kyde, 2010; Rosen, 2011).
Imp-A3 (Affects ornamental plants)	? - max		<i>Oplismenus hirtellus</i> subsp. <i>undulatifolius</i> has been observed in lawns in Virginia (Hughes, 2010) and is controlled there (Kyde, 2010).

Imp-A4 (Anthropogenic weed)	c - low	0.4	MassWildlife Landowner Incentive Program (LIP) funded through the U. S. Fish and Wildlife Service helps private landowners control <i>O. hirtellus</i> subsp. <i>undulatifolius</i> in suburban areas to maintain suitable wildlife habitat on their property (Kyde, 2010). Alternate answers for the Monte Carlo simulation are both "b".
<b>Impact to Production systems (agriculture, nurseries, forest plantations, orchards, etc.)</b>			
Imp-P1 (Crop yield)	? - max		<i>Oplismenus undulatifolius</i> (synonym) is a weed of rice in Japan (Enomoto, 2003) and listed as an agricultural weed (Randall, 2007). Though it occurs in agricultural cropping systems, it is not listed as a Principal or Serious weed by any country; that is, it has not reduced crop yield significantly (Holm et al., 1979).
Imp-P2 (Commodity Value)	? - max		No evidence for a poorly studied taxon. <i>O. undulatifolius</i> (synonym) is a weed of rice in Japan (Enomoto, 2003) and listed as an agricultural weed (Randall, 2007).
Imp-P3 (Affects trade)	n - mod	0	There is no evidence that <i>O. hirtellus</i> subsp. <i>Undulatifolius</i> or any of its conspecifics or congeners impacts trade due to regulatory restrictions (USDA APHIS PPQ CPHST, 2007).
Imp-P4 (Irrigation)	? - max		<i>O. hirtellus</i> subsp. <i>undulatifolius</i> is not well studied and no studies have been found that reports its ability to significantly compete for water.
Imp-P5 (Animal toxicity)	n - mod	0	There is no evidence that <i>O. hirtellus</i> subsp. <i>undulatifolius</i> or any of its congeners is toxic to animals (Burrows and Tyrll, 2001).
Imp-P6 (Production system weed)	c - low	0.6	<i>O. hirtellus</i> subsp. <i>undulatifolius</i> impacts forests in the United States and is actively being controlled (Kyde, 2010). <i>O. hirtellus</i> subsp. <i>undulatifolius</i> may prevent the establishment of tree seedlings (Kyde, 2010). <i>O. undulatifolius</i> is a weed of rice in Japan (Enomoto, 2003) and listed as an agricultural weed (Randall, 2007). Because forests are considered production systems, answering "c". Alternate answers for the Monte Carlo simulation are both "b".
<b>Geographic Potential</b>			
<b>Plant cold hardiness zones</b>			
Geo-Z1 (Zone 1)	n - negl	N/A	No evidence.
Geo-Z2 (Zone 2)	n - negl	N/A	No evidence.
Geo-Z3 (Zone 3)	n - negl	N/A	No evidence.
Geo-Z4 (Zone 4)	n - negl	N/A	No evidence.
Geo-Z5 (Zone 5)	n - high	N/A	No evidence, though some point source data came close to occurring in this range.
Geo-Z6 (Zone 6)	y - low	N/A	Italy, Japan.
Geo-Z7 (Zone 7)	y - negl	N/A	US-Maryland, Japan-Tohoku.
Geo-Z8 (Zone 8)	y - negl	N/A	Japan-Kinki.
Geo-Z9 (Zone 9)	y - negl	N/A	China-Yunnan, Taiwan.
Geo-Z10 (Zone 10)	y - negl	N/A	China-Taiwan.
Geo-Z11 (Zone 11)	y - negl	N/A	Australia-Queensland, Papua New Guinea.
Geo-Z12 (Zone 12)	y - negl	N/A	Papua New Guinea, Sri Lanka.
Geo-Z13 (Zone 13)	y - negl	N/A	Papua New Guinea, Sri Lanka.
<b>Koppen-Geiger climate classes</b>			
Geo-C1 (Tropical rainforest)	y - negl	N/A	Australia-Queensland, Papua New Guinea.
Geo-C2 (Tropical savanna)	y - low	N/A	Africa-Uganda, Australia-Queensland.
Geo-C3 (Steppe)	n - low	N/A	No evidence.
Geo-C4 (Desert)	n - negl	N/A	No evidence.
Geo-C5 (Mediterranean)	y - high	N/A	Spain.

Geo-C6 (Humid subtropical)	y - negl	N/A	US-Maryland.
Geo-C7 (Marine west coast)	y - negl	N/A	New Zealand, Australia-New South Wales.
Geo-C8 (Humid cont. warm sum.)	y - low	N/A	Japan, South Korea.
Geo-C9 (Humid cont. cool sum.)	y - low	N/A	Italy, Japan-Shikoku.
Geo-C10 (Subarctic)	n - negl	N/A	No evidence.
Geo-C11 (Tundra)	n - negl	N/A	No evidence.
Geo-C12 (Icecap)	n - negl	N/A	No evidence.
<b>10-inch precipitation bands</b>			
Geo-R1 (0-10")	n - negl	N/A	No evidence.
Geo-R2 (10-20")	n - low	N/A	No evidence.
Geo-R3 (20-30")	n - high	N/A	No evidence.
Geo-R4 (30-40")	y - low	N/A	Spain-Cataluna; <i>Oplismenus hirtellus</i> subsp. <i>undulatifolius</i> is native to moist parts of Europe in Northern Italy and Southern Switzerland (and southeast Asia) that receive between 950 mm (37 in) and 2000 mm (79 in) of rain (Scholz, 1981).
Geo-R5 (40-50")	y - negl	N/A	US-Maryland; <i>Oplismenus hirtellus</i> subsp. <i>undulatifolius</i> is native to moist parts of Europe in Northern Italy and Southern Switzerland (and southeast Asia) that receive between 950 mm (37 in) and 2000 mm (79 in) of rain (Scholz, 1981).
Geo-R6 (50-60")	y - negl	N/A	Japan-Kanto; <i>Oplismenus hirtellus</i> subsp. <i>undulatifolius</i> is native to moist parts of Europe in Northern Italy and Southern Switzerland (and southeast Asia) that receive between 950 mm (37 in) and 2000 mm (79 in) of rain (Scholz, 1981).
Geo-R7 (60-70")	y - negl	N/A	Japan-Kanto, Italy-Lombardi; <i>Oplismenus hirtellus</i> subsp. <i>undulatifolius</i> is native to moist parts of Europe in Northern Italy and Southern Switzerland (and southeast Asia) that receive between 950 mm (37 in) and 2000 mm (79 in) of rain (Scholz, 1981).
Geo-R8 (70-80")	y - negl	N/A	Japan-Chubu; <i>Oplismenus hirtellus</i> subsp. <i>undulatifolius</i> is native to moist parts of Europe in Northern Italy and Southern Switzerland (and southeast Asia) that receive between 950 mm (37 in) and 2000 mm (79 in) of rain (Scholz, 1981).
Geo-R9 (80-90")	y - negl	N/A	Papua New Guinea.
Geo-R10 (90-100")	y - negl	N/A	Papua New Guinea.
Geo-R11 (100"+)	y - negl	N/A	Papua New Guinea.
<b>Entry Potential</b>			
Ent-1 (Already here)	y - negl	1	<i>O. hirtellus</i> subsp. <i>undulatifolius</i> is invading the states of Maryland and Virginia (Akerson, 2010; Kyde, 2010; Westbrook and Imlay, 2009). <i>O. hirtellus</i> subsp. <i>undulatifolius</i> was first reported in the U.S.A. in 1996 (Peterson et al., 1999). This subspecies is reported as having been in Hawaii for over 150 years (Snow and Lau, 2010), but there is some debate as to the identity of the subspecies in Hawaii (Talley, 2012).
Ent-2 (Proposed for entry)	-	N/A	
Ent-3 (Human value & cultivation/trade status)	-	N/A	
<b>Ent-4 (Entry as a Contaminant)</b>			
Ent-4a (In MX, CA, Central Amer., Carib., or China)	-	N/A	
Ent-4b (Propagative material)	-	N/A	
Ent-4c (Seeds)	-	N/A	
Ent-4d (Ballast water)	-	N/A	
Ent-4e (Aquaria)	-	N/A	
Ent-4f (Landscape products)	-	N/A	

Weed Risk Assessment for *Oplismenus hirtellus* subsp. *undulatifolius*

Ent-4g (Container, packing, trade goods)	-	N/A
Ent-4h (Commodities for consumption)	-	N/A
Ent-4i (Other pathway)	-	N/A
Ent-5 (Natural dispersal)	-	N/A