



U.S. Department of Agriculture
Animal and Plant Health Inspection Service
Wildlife Services

U.S. Government Publication

Winter and Summer Home Ranges of American White Pelicans (*Pelecanus erythrorhynchos*) Captured at Loafing Sites in the Southeastern United States

Author(s): D. Tommy King, Justin Fischer, Bronson Strickland, W. David Walter, Fred L. Cunningham and Guiming Wang

Source: Waterbirds, 39(3):287-294.

Published By: The Waterbird Society

DOI: <http://dx.doi.org/10.1675/063.039.0308>

URL: <http://www.bioone.org/doi/full/10.1675/063.039.0308>

BioOne (www.bioone.org) is a nonprofit, online aggregation of core research in the biological, ecological, and environmental sciences. BioOne provides a sustainable online platform for over 170 journals and books published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Web site, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/page/terms_of_use.

Usage of BioOne content is strictly limited to personal, educational, and non-commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

Winter and Summer Home Ranges of American White Pelicans (*Pelecanus erythrorhynchos*) Captured at Loafing Sites in the Southeastern United States

D. TOMMY KING^{1,*}, JUSTIN FISCHER², BRONSON STRICKLAND³, W. DAVID WALTER⁴,
FRED L. CUNNINGHAM¹ AND GUIMING WANG³

¹U.S. Department of Agriculture, Wildlife Services, National Wildlife Research Center, P.O. Box 6099, Mississippi State, Mississippi, 39762, USA

²U.S. Department of Agriculture, Wildlife Services, National Wildlife Research Center, Ft. Collins, Colorado, 80521, USA

³Department of Wildlife, Fisheries, and Aquaculture, Mississippi State University, Mississippi State, Mississippi, 39762, USA

⁴U.S. Geological Survey, Pennsylvania Cooperative Fish and Wildlife Research Unit, 403 Forest Resources Building, The Pennsylvania State University, University Park, Pennsylvania, 16802, USA

*Corresponding author; E-mail: Tommy.King@aphis.usda.gov

Abstract.—Satellite telemetry was used to investigate summer and winter home ranges for resident and migrant American White Pelicans (*Pelecanus erythrorhynchos*) captured in the southeastern United States between 2002 and 2007. Home range utilization distributions were calculated using 50% and 95% kernel density estimators with the plug-in bandwidth selector. Mean summer home ranges (95%) varied from 177 to 4,710 km² and mean winter home ranges (95%) ranged from 185 to 916 km². Mean 50% and 95% home ranges of adult American White Pelicans during summer tended to be larger than those during winter, whereas mean 50% and 95% home ranges of immature pelicans during summer tended to be smaller than those during winter. Home ranges for all American White Pelicans encompassed the latitude range of 24°–55° N, including wintering, stop over, and nesting habitat. These data provide baseline movement and home range data for future studies of American White Pelican ecology. Received 21 June 2014, accepted 8 August 2014.

Key words.—American White Pelican, aquaculture, home range, kernel density estimation, movements, *Pelecanus erythrorhynchos*, plug-in bandwidth, satellite telemetry.

Waterbirds 39(3): 287–294, 2016

The American White Pelican (*Pelecanus erythrorhynchos*; hereafter pelican) is a charismatic and well known species in North America (Anderson and King 2005; Keith 2005), but little is actually known about its general movements, home ranges, and core use areas (Knopf and Evans 2004). Using VHF (Very High Frequency) telemetry, King and Werner (2001) documented daily activity budgets of pelicans captured near aquaculture facilities in the southeastern United States. Logistical constraints involved with the utilization of VHF telemetry, however, precluded its use to effectively address many local, regional, and continental questions concerning pelican ranges and movements (King and Werner 2001). Breeding bird surveys and band recoveries of pelicans have partially identified dispersal patterns (Houston 1972; Strait and Sloan 1975; King and Grewe 2001; Anderson and Anderson

2005) and the location of North American colonies (King and Anderson 2005), but seasonal and regional movements of these wide-ranging birds are of considerable added interest. Recent advances in satellite telemetry have allowed monitoring of animal movements over expansive areas (Lindberg and Walker 2007; Robinson *et al.* 2009; Hebblewhite and Haydon 2010). Shannon *et al.* (2002) used satellite telemetry to document American White Pelican soaring flight times, and Izhaki *et al.* (2002) used satellite telemetry to describe the migratory and ranging behavior of immature Great White Pelicans (*P. onocrotalus*). Further advances have led to the incorporation of a Global Positioning System (GPS) into satellite transmitters, allowing for very precise location data and intensive monitoring of animal movements. For example, Avery *et al.* (2011) used GPS-technology to document home ranges of

Black Vultures (*Coragyps atratus*) and Turkey Vultures (*Cathartes aura*) in South Carolina, USA, and their migration to central Florida, USA.

To our knowledge, the home range of American White Pelicans has not been previously estimated. The objectives of this study were to: 1) estimate seasonal home ranges and core use areas for pelicans captured in the southeastern United States; and 2) determine effects of season, age, and sex on home range sizes of pelicans. Data from this study provide novel information on local, regional, and continental movements of American White Pelicans and can also be interpreted on a behavioral basis.

METHODS

Study Area

We captured pelicans at loafing sites near aquaculture-intensive areas in Alabama, Arkansas, Louisiana, and

Mississippi, USA (Fig. 1) using rocket nets and modified foot-hold traps (King *et al.* 1998). We fitted pelicans with backpack 70-g solar-powered GPS satellite transmitters (PTT-100, Microwave Telemetry; Dunstan 1972); transmitters were programmed to record one location per hour for the duration of the study and were < 3% of the individual's body weight. We estimated age (≥ 3 years old = adult; < 3 years old = immature) using plumage and eye and skin color characteristics (D. T. King, unpubl. data) and determined sex by using culmen length (Dorr *et al.* 2005) for each captured pelican. We released all captured birds at the trap site.

The summer season included locations from 1 June-31 August, and the winter season included locations from 1 November-15 April. A pelican's home range for a given season/year was computed only if there were > 50 locations and if those locations were logged during > 50% of the time period. We identified 1 June the year following capture as the date when a bird tagged as a 2-year-old immature would be considered an adult. We defined migration as seasonal movements from wintering areas in the southern United States to breeding and summering areas in the north central United States and southern Canada. Because a few individuals exhibited nomadic behavior when flying north in the spring, but never reached breeding colonies, we identified 36°




Figure 1. American White Pelican capture locations in Alabama, Arkansas, Louisiana, and Mississippi, 19 April 2002-4 April 2007.

30' N latitude as the dividing line and 15 June as the cutoff date for individuals to either be classified as residents or migrants.

Statistical Analysis

We used kernel density estimation (KDE) to estimate summer and winter 50% (i.e., core area) and 95% (i.e., overall use area) utilization distributions for pelicans. We used the second generation plug-in bandwidth selection method because our GPS data set had numerous locations (i.e., 50-24,000 per season) that often caused first generation methods to fail (Gitzen *et al.* 2006; Walter *et al.* 2011). First generation methods such as least squares cross-validation failed to converge with a large number of identical or very tight clustered points (Hemson *et al.* 2005; Gitzen *et al.* 2006; Pellerin *et al.* 2008) or had a tendency to grossly over-smooth the data (default or reference bandwidth; Worton 1995; Seaman and Powell 1996; Hemson *et al.* 2005). The plug-in bandwidth selection method is data-based, uses an equation to directly estimate the 'ideal' bandwidth, and estimates pilot bandwidths to tune the performance of the covariance estimator (Jones *et al.* 1996; Duong and Hazelton 2003; Duong 2007). We calculated KDE plug-in home ranges in the R language for statistical computing (R Development Core Team 2012) using the ks package (Duong 2007).

We used mixed-effect models to detect effects of season, age, and sex on home range sizes of pelicans with animal ID as a random effect variable to account for temporal autocorrelation because of repeated observations of the same individual (R Development Core Team 2012). If the estimate of variance for animal ID was 0.00, we used analysis of variance to compare home range sizes by age, sex, season, and their interactions at the significance level of 0.05. In preliminary analyses, effects of year were not significant ($P > 0.10$). Therefore, we pooled our data over years in subsequent analyses.

We tested for effects of fixed-effect variables (age, sex, and season) and their interactions with: 1) all possible combinations of two-factor interactions; and 2) three-factor interaction. We conducted backward model selection at the significance level of 0.05. We conducted multiple comparisons of least square means (lsmeans)

or sliced lsmeans of home ranges with the Tukey adjustment using the SAS procedure MIXED with the slice option (Littell *et al.* 2006). We conducted linear model or mixed-model analysis for 50% and 95% KDE home range sizes, with home range sizes log transformed for the normality assumption.

RESULTS

From 19 April 2002-4 April 2007, we captured 39 pelicans (3-17 per year) and fitted them with transmitters. We captured 34 males (5 adults, 29 immatures) and 5 immature females. These birds were tracked from 19 April 2002-29 October 2007. Transmitters ($n = 39$) operated an average of 17.3 months (± 1.5 SE; Range = 1-52), excluding birds that were known to be killed (shot under depredation permits at aquaculture facilities) within 3 weeks of release. We obtained an average of 2,299 (± 271.8 SE) locations per bird, some of which were represented by captured immatures reaching adulthood (male: $n = 17$; female: $n = 4$).

Mean 50% and 95% home ranges of adult pelicans during summer tended to be larger than those during winter, whereas mean 50% and 95% home ranges of immature pelicans during summer tended to be smaller than those during winter (Table 1). Estimated mean 95% home ranges for all pelicans covered the latitude range of 24°-55° N, including wintering, stop over, and nesting habitat. Mean 50% summer home ranges for migrating pelicans were located in the northern part of the species' range (i.e., summer nesting habitat; Fig. 2). The

Table 1. Mean summer and winter home ranges of American White Pelicans using kernel density estimation and the plug-in bandwidth selector for 50% and 95% utilization distributions (km² (\pm SE)). Pelicans were captured in the southeastern United States from April 2002-October 2007. Each season-year was considered a separate range ($n = 178$).

Sex	Age	Season	50%	95%
Male	Adult	Summer	2,818 \pm 1,418	28,487 \pm 10,147
		Winter	916 \pm 228	12,632 \pm 3,140
	Immature	Summer	326 \pm 124	5,275 \pm 2,175
		Winter	681 \pm 162	11,410 \pm 2,994
Female	Adult	Summer	4,711 \pm 2,794	65,880 \pm 28,866
		Winter	186 \pm 29	1,329 \pm 248
	Immature	Summer	177 \pm 66	2,089 \pm 557
		Winter	342 \pm 79	4,960 \pm 814



Figure 2. Extent of 50% and 95% fixed-kernel density estimates (KDE) with the plug-in bandwidth selector of summer home ranges for migrating American White Pelicans captured in the southeastern United States during 2002-2007.

extents of mean 50% and 95% home ranges of summer residents (i.e., non-migrants) were similar to those of winter home ranges (Figs. 3 and 4).

50% Kernel Home Range

The estimate of variance for animal ID was 0.00 when including animal ID as a random effect. Therefore, we compared 50% kernel home ranges using analysis of variance. No comparisons of 50% kernel home range sizes by age, sex, and season were significant ($P > 0.10$).

95% Kernel Home Range

In the mixed model, the age-sex-season interaction was significant ($F_{4, 75} = 4.82$, $P = 0.002$). Female adults had larger home ranges in summer than in winter ($t_{75} = 3.28$, Tukey

adjusted $P = 0.03$), with the least square mean of log transformed 95% kernel home range sizes being 10.4 and 7.06, respectively. Moreover, female adults had larger home range sizes than did male immatures during summer ($t_{75} = 3.6$, adjusted $P = 0.01$), with the least square mean of log transformed 95% kernel home range sizes being 10.4 and 7.21, respectively. Female home range sizes did not differ from those of adult males.

DISCUSSION

Migratory birds make seasonal movements between wintering and summer nesting habitats to take advantage of available resources in different locations to meet habitat and nutritional requirements (Pulido 2007; Ramenofsky and Wingfield 2007). Mean 50% kernel home ranges (i.e., core home




Figure 3. Extent of 50% and 95% fixed-kernel density estimates (KDE) with the plug-in bandwidth selector of summer home ranges for resident American White Pelicans captured in the southeastern United States during 2002-2007.

ranges) of pelicans were spatially unconnected (Figs. 2 and 4). The northern boundary of winter 95% kernel home ranges probably represents the northern boundary of pelican wintering habitat (Fig. 4). For migrating pelicans, the lower latitude limit of the range of summer core or 50% kernel home ranges was approximately the same as that of known breeding colonies (Fig. 2; King and Anderson 2005). Therefore, our estimated summer and winter core home ranges probably delineate the lower latitude limit of nesting habitat and upper latitude limit of wintering habitat respectively, providing insight for future studies of habitat selection and requirements by nesting and wintering American White Pelicans.

Habitat, food abundance, and bird reproductive condition may affect bird movement distance and home range sizes (McNab 1963; Weimerskirch *et al.* 1994). More abun-

dant food resources may result in smaller home ranges of animals (McNab 1963). Home ranges of adult pelicans during winter tended to be smaller than those during summer (Table 1). Additionally, home range size of female adults in summer was greater than that in winter. Wintering pelicans feed on fish at commercial aquaculture facilities, such as catfish (*Ictalurus punctatus*) ponds, in the southeastern United States (King *et al.* 2010). Catfish aquaculture in the southeastern United States was estimated to produce a total of 1,626,401,709 kg of fish during our study period (National Agricultural Statistics Service 2013). Abundant fish at commercial aquaculture facilities improves the body condition of pelicans (King *et al.* 2010) and may reduce winter home range sizes of pelicans. During the breeding season, egg incubation and provision feeding by adults are energetically demanding with nesting birds often




Figure 4. Extent of 50% and 95% fixed-kernel density estimates (KDE) with the plug-in bandwidth selector of winter home ranges for American White Pelicans captured in the southeastern United States during 2002-2007.

making long foraging trips to meet these energetic demands (Behle 1958; Johnson and Sloan 1978; Weimerskirch *et al.* 1994; Findholt and Anderson 1995).

When compared to peak years, the aquaculture industry in the USA has undergone a 54.6% reduction in total catfish water surface acres and a 55.6% reduction in the number of commercial catfish farms due to foreign imports and rising feed costs (Hanson and Sites 2014). It has been speculated that the remaining commercial catfish farms will suffer increased predation losses and disease risk as pelicans concentrate their foraging efforts on fewer facilities, which could also restrict pelican winter home ranges to smaller areas. This information will be important to catfish farmers to help them better understand the threat that changes in American White Pelican distribution represents, allowing farmers to concentrate control and harassment efforts during the ap-

propriate months to increase efficiency and reduce predation losses. These data can be used as baseline movement and home range data for future studies of American White Pelican ecology.

ACKNOWLEDGMENTS

We thank S. Baruch-Mordo for assistance with the R software plug-in programming. We thank the staff of the USDA/WS/NWRC Mississippi Field Station for assisting in pelican capture and transmitter attachment. We also thank M. Tobin and two anonymous reviewers for providing helpful comments on this manuscript. This research was conducted under U.S. Department of Agriculture, National Wildlife Research Center, Animal Care and Use Committee study protocol QA-1018; USGS, BRD Federal Bird Marking and Salvage Permit No. 20873; Mississippi Department of Wildlife, Fisheries, and Parks Administrative Scientific Collecting Permit; Alabama Department of Conservation and Natural Resources Scientific Collecting Permit No. 2031; Arkansas Game and Fish Commission Scientific Collection Permit No. 3010; and Louisiana Department of Wildlife and Fisheries Scientific Collecting Permit Number LNHP-01-029.

LITERATURE CITED

- Anderson, D. W. and D. T. King. 2005. Introduction: biology and conservation of the American White Pelican. *Waterbirds* 28 (Special Publication 1): 1-8.
- Anderson, J. G. T. and K. B. Anderson. 2005. An analysis of band returns of the American White Pelican, 1922-1981. *Waterbirds* 28 (Special Publication 1): 55-60.
- Avery, M. L., J. S. Humphrey, T. S. Daughtery, J. W. Fischer, M. P. Miles, E. A. Tillman, W. E. Bruce and W. D. Walter. 2011. Vulture flight behavior and implications for aircraft safety. *Journal of Wildlife Management* 75: 1581-1587.
- Behle, W. H. 1958. *The bird life of Great Salt Lake*. University of Utah Press, Provo, Utah.
- Dorr, B. S., D. T. King, J. B. Harrel, P. Gerard and M. G. Spalding. 2005. The use of culmen length to determine sex of American White Pelicans (*Pelecanus erythrorhynchos*). *Waterbirds* 28 (Special Publication 1): 102-106.
- Dunstan, T. C. 1972. A harness for radio-tagging raptorial birds. *Inland Bird Banding News* 44: 4-8.
- Duong, T. 2007. ks: kernel density estimation and kernel discriminant analysis for multivariate data in R. *Journal of Statistical Software* 21: 1-16.
- Duong, T. and M. L. Hazelton. 2003. Plug-in bandwidth matrices for bivariate kernel density estimation. *Nonparametric Statistics* 15: 17-30.
- Findholt, S. L. and S. H. Anderson. 1995. Foraging areas and feeding habitat selection of American White Pelicans (*Pelecanus erythrorhynchos*) nesting at Pathfinder Reservoir, Wyoming. *Colonial Waterbirds* 18: 47-57.
- Gitzen, R. A., J. J. Millsbaugh and B. J. Kernohan. 2006. Bandwidth selection for fixed-kernel analysis of animal utilization distributions. *Journal of Wildlife Management* 70: 1334-1344.
- Hanson, T. and D. Sites. 2014. 2013 U.S. catfish database. Fisheries and Allied Aquaculture Department Series No. 1, Alabama Agriculture Experiment Station, Auburn, Alabama.
- Hebblewhite, M. and D. T. Haydon. 2010. Distinguishing technology from biology: a critical review of the use of GPS telemetry data in ecology. *Philosophical Transactions of the Royal Society B* 365: 2303-2312.
- Hemson, G., P. Johnson, A. South, R. Kenward, R. Ripley and D. MacDonald. 2005. Are kernels the mustard? Data from global positioning system (GPS) collars suggests problems for kernel home-range analyses with least-squares cross-validation. *Journal of Animal Ecology* 74: 455-463.
- Houston, C. S. 1972. Recent Saskatchewan banding of the white pelican. *Blue Jay* 30: 24-27.
- Izhaki, I., M. Shmueli, Z. Arad, Y. Steinberg and A. Crivelli. 2002. Satellite tracking of migratory and ranging behavior of immature Great White Pelicans. *Waterbirds* 25: 295-304.
- Johnson, R. F. and N. F. Sloan. 1978. White pelican production and survival of young at Chase Lake National Wildlife Refuge, North Dakota. *Wilson Bulletin* 90: 346-352.
- Jones, M. C., J. S. Marron and S. J. Sheather. 1996. A brief survey of bandwidth selection for density estimation. *Journal of the American Statistical Association* 91: 401-407.
- Keith, J. O. 2005. An overview of the American White Pelican. *Waterbirds* 28 (Special Publication 1): 9-17.
- King, D. T. and A. H. Grewe, Jr. 2001. Movements and mortality of American White Pelicans banded at Marsh Lake, Minnesota. *North American Bird Bander* 26: 57-60.
- King, D. T. and S. J. Werner. 2001. Daily activity budgets and population size of American White Pelicans wintering in south Louisiana and the delta region of Mississippi. *Waterbirds* 24: 250-254.
- King, D. T. and D. W. Anderson. 2005. Recent population status of the American White Pelican: a continental perspective. *Waterbirds* 28 (Special Publication 1): 48-54.
- King, D. T., J. L. Belant, J. B. Harrel and J. F. Glahn. 2010. Superabundant food at catfish aquaculture facilities improves body condition in American White Pelicans. *Waterbirds* 33: 221-227.
- King, D. T., J. D. Paulson, D. J. LeBlanc and K. Bruce. 1998. Two capture techniques for American White Pelicans and Great Blue Herons. *Colonial Waterbirds* 21: 258-260.
- Knopf, F. L. and R. M. Evans. 2004. American White Pelican (*Pelecanus erythrorhynchos*). No. 57 in *The Birds of North America Online* (A. Poole, Ed.). Cornell Lab of Ornithology, Ithaca, New York. <http://bna.birds.cornell.edu/bna/species/057>, accessed 11 September 2012.
- Lindberg, M. S. and J. Walker. 2007. Satellite telemetry in avian research and management: sample size considerations. *Journal of Wildlife Management* 71: 1002-1009.
- Littell, R. C., G. A. Milliken, W. W. Stroup and R. D. Wolfinger. 2006. *SAS for mixed models*. SAS Press, Cary, North Carolina.
- McNab, B. K. 1963. Bioenergetics and determination of home range size. *American Naturalist* 97: 133-140.
- National Agricultural Statistics Service. 2013. Catfish processing. U.S. Department of Agriculture, Agricultural Statistics Board, National Agricultural Statistics Service, Washington, D.C. <http://usda.mannlib.cornell.edu/usda/current/CatfProc/CatfProc-03-20-2013.pdf>, accessed 14 June 2013.
- Pellerin, M., S. Said and J. M. Gaillard. 2008. Roe deer *Capreolus capreolus* home-range sizes estimated from VHF and GPS data. *Wildlife Biology* 14: 101-110.
- Pulido, F. 2007. Phenotypic changes in spring arrival: evolution, phenotypic plasticity, effects of weather and condition. *Climate Research* 35: 5-23.
- R Development Core Team. 2012. R: a language and environment for statistical computing v. 2.1.4.3. R Foundation for Statistical Computing, Vienna, Austria. <http://www.R-project.org/>, accessed 5 January 2013.
- Ramenofsky, M. and J. C. Wingfield. 2007. Regulation of migration. *Bioscience* 57: 135-143.
- Robinson, W. D., M. S. Bowlin, I. Bisson, J. Shamoun-Baranes, K. Thorup, R. H. Diehl, T. H. Kunz, S. Mabey

- and D. W. Winkler. 2009. Integrating concepts and technologies to advance the study of bird migration. *Frontiers in Ecology and the Environment* 8: 354-361.
- Seaman, D. E. and R. A. Powell. 1996. An evaluation of the accuracy of kernel density estimators for home range analysis. *Ecology* 77: 2075-2085.
- Shannon, H. D., G. S. Young, M. A. Yates, M. R. Fuller and W. S. Seegar. 2002. American White Pelican soaring flight times and altitudes relative to changes in thermal depth and intensity. *Condor* 104: 679-683.
- Strait, L. E. and N. F. Sloan. 1975. Movements and mortality of juvenile white pelicans from North Dakota. *Wilson Bulletin* 87: 54-59.
- Walter, W. D., J. W. Fischer, S. Baruch-Mordo and K. C. VerCauteren. 2011. What is the proper method to delineate home range of an animal using today's advanced GPS telemetry systems: the initial step. Pages 249-268 in *Modern Telemetry* (O. Krejcar, Ed.). InTech, DOI: 10.5772/24660. <http://www.intechopen.com/books/modern-telemetry/what-is-the-proper-method-to-delineate-home-range-of-an-animal-using-today-s-advanced-gps-telemetry->, accessed 20 August 2012.
- Weimerskirch, H., O. Chastel, L. Ackermann and T. Chaurand. 1994. Alternate long and short foraging trips in pelagic seabird parents. *Animal Behaviour* 47: 472-476.
- Worton, B. J. 1995. Using Monte Carlo simulation to evaluate kernel-based home range estimators. *Journal of Wildlife Management* 59: 794-800.