



United States Department of Agriculture

Animal and
Plant Health
Inspection
Service

Veterinary
Services

February 2020

Influenza A Virus in Swine Surveillance

Fiscal Year 2020 Quarterly Report

Surveillance Summary for First Quarter Fiscal Year 2020: October 1 to December 31, 2019

Report Summary¹

- This report covers the first quarter (Q1) of fiscal year 2020 from October 1 to December 31, 2019.
- Where relevant, the report also includes previous years' data for historical perspective.
- The report provides data from both national and regional levels.
- In fiscal year 2020 Q1, there were 896 samples submitted for influenza A virus (IAV) surveillance in swine from 942 accessions.
- H1N2 was the predominant subtype reported in USDA data in Q1 FY2020.
- Over the past 7 quarters, H1N2 was the main subtype in Regions 1, 3 and 5 while Region 2 and Region 4 most frequently isolated H1N1 and H3N2, respectively. For regions recorded as "unknown," H1N2 and H3N2 were the most frequent subtypes.
- Limited accessions from a region can skew data and lead to misinterpretation. Therefore, less inference can be applied to results from Regions 3, 4, and 5.
- All IAV-S submissions are voluntary and based on clinical case submissions to veterinary diagnostic labs. These data are not a statistically representative sampling of the U.S. swine population.
- Due to the voluntary nature of this surveillance, the information in this report cannot be used to determine regional and/or national incidence, prevalence, or other epidemiological measures, but it may help identify IAV-S trends.

Introduction

This report, based on data received as of July 15, 2020, provides a brief update on the status of national surveillance for IAV in swine for producers, swine practitioners, diagnosticians and the public. Summaries in this report may differ from those provided in past reports due to the regular addition of data from participating laboratories. Reporting months are based on the month the sample was collected. The IAV-S surveillance program is voluntary and, as a result, the accessions and samples

¹ In November 2016, VS modernized the process that prepares and stages laboratory results data for reporting. Consequently, VS recognizes there is a small difference in previously reported summary numbers for IAV-S surveillance. The results in this report reflect updated and corrected numbers achieved with the modernized data process.

submitted represent a subset of the swine population. Submitted samples should only be collected from animals displaying influenza-like illness. Due to its voluntary nature, this surveillance system does not entirely represent the total U.S. domestic swine population. Therefore, the data cannot be used to determine IAV-S prevalence or other epidemiologic measures in the swine population. However, the data may help identify trends of influenza in swine.

When the submitter does not report relevant information, data are recorded as “unknown.” Summaries in this report may differ from those provided in past reports due to the ongoing addition of data from participating labs. Reporting months are based on the month when the sample was collected.

A laboratory accession is generally a set of samples collected at a single premises on a single day and received at the laboratory. A maximum of 10 samples of any kind is allowed per accession for reimbursement under the USDA IAV-S system. However, no more than five of the 10 samples may be oral fluid for any given accession. This does not prevent additional samples from being tested at the owner’s expense. While a nasal swab or lung tissue sample represents a single animal within the herd, a single oral fluid sample may represent one to two pens of animals in a herd. A positive sample status is based on the screening real-time reverse transcriptase polymerase chain reaction (rRT-PCR). The subtype result is based on the rRT-PCR-based subtyping assays. Virus isolation (VI) and sequencing in the NAHLN labs are only attempted on rRT-PCR positives meeting criteria with sequences deposited into GenBank, the public sequence database. On a monthly basis, USDA also performs whole genomic sequencing (WGS) on a selected subset of virus isolates received through surveillance and deposits those sequences into Genbank. On a quarterly basis, a phylogenetic analysis is performed; phylogenetic analyses are based on all successful USDA surveillance sequencing results, with sequences deposited into GenBank®, the public sequence database.

Program Updates

There have been recent changes to the USDA’s webpages that provide information on IAV-S and IAV-S surveillance. Please visit the new webpage and navigate around to find the information you may be seeking as well as the IAV-S quarterly reports. The new web address is:

<https://www.aphis.usda.gov/aphis/ourfocus/animalhealth/animal-disease-information/swine-disease-information/influenza-a-virus>

The focus of IAV-S surveillance remains on acquiring viruses. The National Animal Health Laboratory Network (NAHLN) has several submission options to ensure unusual viruses identified by methods other than those approved for NAHLN testing can be submitted into the program. An updated version of the IAV-S NAHLN testing guidelines and instruction sheet can be found at:

https://www.aphis.usda.gov/animal_health/animal_dis_spec/swine/downloads/appendix_c_testing_guidelines.pdf

and

https://www-author.aphis.usda.gov/animal_health/animal_dis_spec/swine/downloads/iav-s-algorithm-instructions.pdf

IAV-S Surveillance Objectives

USDA’s National Surveillance Plan for Swine Influenza Virus in Pigs (July 2010) describes the current surveillance system for IAV in swine in detail. The surveillance objectives are to:

1. Monitor genetic evolution of endemic IAV in swine to better understand endemic and emerging influenza virus ecology;
2. Make influenza isolates from swine available for research and establish a data management system to facilitate genetic analysis of these isolates and related information; and
3. Select proper isolates for the development of relevant diagnostic reagents, updated diagnostic assays, and vaccine seed stock products.

Objective 1. Monitoring Genetic Evolution of Endemic IAV in Swine to Better Understand Endemic and Emerging Influenza Virus Ecology

Objective 1 is met through the submission of diagnostic laboratory samples to the surveillance system, collection of the viruses that are isolated from the samples, and analysis of the hemagglutinin (HA) and neuraminidase (NA) sequences that are generated at the NAHLN laboratories. Each month, selected viruses undergo whole genomic sequencing by the National Veterinary Services Laboratories (NVSL). Phylogenetic analysis of the genetic sequences submitted through the surveillance program is provided through an interagency agreement with the USDA’s Agricultural Research Service (ARS) National Animal Disease Center (NADC).

National Surveillance Data Summary

From fiscal year 2010 through fiscal year 2016, the total number of accessions and samples submitted increased. Changes initiated in fiscal year 2016 resulted in decreased laboratory accessions and samples, but a higher percentage of accessions resulting in a virus isolate that can be sequenced and analyzed. In fiscal year 2020, 942 samples have been tested from 896 accessions (Figure 1). Figure 2 shows the overall increasing trends in total accessions, rRT-PCR and VI positive accessions, and subtyped accessions.




Figure 1. Number of IAV-S laboratory accessions and samples tested in swine, fiscal year 2012 through fiscal year 2020 Q1.




Figure 2. Accessions submitted, subtyped accessions, rRT-PCR positive accessions, and virus isolation positive accessions over time with trend lines for influenza A virus in swine, fiscal year 2011 to fiscal year 2020 Q1.

Figure 3 shows the number of subtype detections in fiscal year 2020 Q1. The total number of samples subtyped was 380, including H1N1 (n=130), H1N2 (n=135), H3N2 (N=106) and mixed (N=17). H3N1 was not isolated.




Figure 3. Number of influenza A subtype detections in swine fiscal year 2020 Q1.

Figure 4 breaks down accessions by rRT-PCR subtype from fiscal year 2011 Q1 to fiscal year 2020 Q1. H1N1 was the predominant subtype detected in 2012, 2013, 2014 and 2018. H1N2 was detected most often in 2011, 2015, 2016, 2017, 2019 and 2020.

Influenza A Virus in Swine Surveillance Quarterly Report for fiscal year 2020, Quarter 1




Figure 4. Number of subtypes, fiscal year 2012 through fiscal year 2020 Q1.

Figure 5 displays the number of times VI was conducted in blue, the number of times VI was conducted and was positive in purple, and the number of viral isolates submitted to GenBank in green. Since the implementation of the June 2016 modifications to the program, almost all VIs attempted now yield a virus and the sequences are submitted to Genbank for analysis. Due to an unresolvable data processing coding error, samples sequenced and sent to Genbank during FY2019 Q1 appear greatly reduced in Figure 5. As with other fiscal years, the samples sent during that timeframe closely follow the number of attempted virus isolations as well as the number of positive virus isolation samples.




Figure 5. Number of influenza A virus isolations attempted, number of virus isolations that were positive, and the number of viruses submitted to GenBank from fiscal year 2011 to fiscal year 2020 Q1.

Laboratory accessions were evaluated by age-class for the first quarter. The most common subtype isolated among grow/finish age class was H1N1. Among nursery and suckling age classes, H1N2 was the predominant isolated subtype. The sow/boar class had limited testing, with four isolates of H1N2 and

two isolates of H3N2. Among isolates for which the age class was unknown or not recorded, both H1N1 and H1N2 were the predominant subtype with 15 isolates each (Table 1). Excluding specimen types with less than 10 percent of total submissions, samples collected from lung and nasal specimens (swabs or wipes) were the most successful at providing a virus isolate which enables sequence submission to GenBank (Table 2).

Table 1. Number of positive accessions tested for influenza A virus in swine by age class and subtype fiscal year 2020, Q1.

Age class	Number of accessions with subtype reported	Number of H1N1	Number of H1N2	Number of H3N1	Number of H3N2	Number of Mixed
Suckling	62	13	27	0	19	3
Nursery	160	54	63	0	37	7
Sow/Boar	6	0	4	0	2	0
Grow/finish	113	51	25	0	34	4
Unknown	42	15	15	0	13	3

Table 2. Number of positive accessions* tested for influenza A virus in swine by specimen type and subtype fiscal year 2020 Q1.**

Specimen Type	Number of accessions with subtype reported	Percent of subtyped accessions with positive virus isolation	Number of H1N1	Number of H1N2	Number of H3N1	Number of H3N2	Number of Mixed	Number of samples sequenced and sent to GenBank
Lung	327	88%	122	110	0	88	8	287
Nasal	33	88%	4	16	0	8	5	26
Oral Fluid	15	80%	1	5	0	5	4	9
Other	2	100%	0	1	0	1	0	2

*Accessions may include samples with multiple specimen types. In these cases, individual accessions are counted in more than one specimen type category.

** Other includes specimen types recorded as swab, mixed tissue, or unknown.

Regional surveillance data

In this section, we present data in five different regions (Figure 6). These regions are based on former USDA administrative districts only and do not represent specific industry distribution. Submissions are voluntary, as is any identifying information accompanying the submission (except the State of animal origin), and therefore no sampling strategies can be applied to the regions.



Figure 6. A map of the regions for national IAV-S surveillance.

Summary of Regional Data from ARS

Table 3. Summary of predominant subtypes in each region in a 2 year window (fiscal year 2018 Q1 through fiscal year 2020 Q1).

Region	Total number	HA/NA subtype
1	264	Gamma H1/Classical N1 Delta2 H1/1998-N2 2010.1 H1/2002-N2
2	1,275	Gamma H1/Classical N1 2010.1 H3/2002-N2 Delta2 H1/1998-N2
3	159	Alpha H1/2002-N2 Gamma H1/Classical N1 2010.1 H3/2002-N2
4	172	2010.1 H3/2002-N2 IV-A H3/2002-N2 Gamma H1/Classical N1
5	38	Low participation 2010.1 H3/2002 N2 Pandemic H1/Pandemic N1 Alpha H1/2002-N2

Most Predominant HA/NA phylo-types overall: H1N1 (Gamma H1/Classical N1), H3N2 (2010.1 H3/2002-N2) and H1N2 (Delta2 H1/1998-N2)

Figure 7 shows the distribution of rRT-PCR subtyped accessions among the five regions for Q3 fiscal year 2018 through Q1 fiscal year 2020. Regions 1, 3 and 5 demonstrate H1N2 as the predominant subtype while Region 2 and Region 4 isolated H1N1 and H3N2 most frequently, respectively. For regions recorded as “unknown,” H1N2 and H3N2 were the most frequent subtypes.



Figure 7. Percentage of influenza A subtyped accessions for swine by region for fiscal year 2018 Q3 to fiscal year 2020 Q1.

Regional phylogenetic analysis

Phylogenetic analysis of sequences from the IAV-S surveillance system

Phylogenetic analysis of gene sequences of IAV in swine is conducted to further examine the genetic changes that occur in HA, NA, and Matrix (M) genes of this rapidly changing virus. Through collaboration with ARS, a dataset^{2,3} of 376 isolates with published sequences in GenBank was characterized by phylogenetic analysis for the Q1 fiscal year 2020 report. This analysis provides information on the genetic diversity and evolution patterns of influenza in swine and allows for inferences about population and/or vaccine immunity.

The following series of bar charts parse the data into an approximately 2-year window by quarters and subtypes for each region, followed by charts further describing the phylogenetic clades of H1 and H3 subtypes. Regional charts depicting the various combinations of HA and NA are in Appendix 1.

² Participating NAHLN labs included M gene sequencing in their testing until July 2016 because the 2009 H1N1 M gene was the predominant circulating gene.

³ The ARS dataset is comprised of IAV-S surveillance isolate sequences that were posted in Genbank. This represents only a subset of the complete IAV-S surveillance dataset that includes PCR diagnostic test-based results as well as sequencing results. Therefore, ARS dataset results, such as subtype percentages, differ from the complete IAV-S dataset results provided in other sections of this report.




Figure 8. Temporal distribution of Influenza A virus subtype by region for Q2 fiscal year 2018 to Q1 fiscal year 2020

Figure 8 demonstrates the four subtypes H1N1, H1N2, H3N2 and mixed subtypes across the five regions. Regions 1 and 2 reported the most submissions, with a mixture of mostly H1N1, H1N2, and H3N2. Limited accessions from a region can skew data and lead to misinterpretation and therefore, less inference can be applied to results from Regions 3, 4, and 5.

National phylogenetic HA gene information

HA genes from H1 subtype viruses are classified as alpha, beta, gamma, delta-1, delta-2, or pandemic H1N1 2009 (H1N1pdm09) phylogenetic clades based on a previously published nomenclature system. Similarly, H3 subtype viruses are classified as Cluster IV, Cluster IV-A, Cluster IV-B, Cluster IV-C, Cluster IV-D, Cluster IV-E, Cluster IV-F, or human-like. In the H1 subtypes (Figure 9), gamma viruses are the predominant H1 with delta2 and alpha frequently detected.

H1 phylogenetic-clades by Region




Figure 9. Temporal distribution of H1 phylogenetic clades by region for fiscal year 2018 Q2 to fiscal year 2020 Q1.

H3 phylogenetic-clades by Region




Figure 10. Temporal distribution of H3 phylogenetic clades by region for fiscal year 2018 Q2 to fiscal year 2020 Q1.

In Q1 fiscal year 2020, there were 45 detections of H3s (Figure 10). 2010.2 H3 was detected in Arkansas, Iowa, Illinois, Indiana and Oklahoma paired with human N2 with on Iowa paired with 2002 N2. Cluster I

H3 was detected in Alabama, Iowa, Illinois, Indiana, Kansas, Minnesota, Missouri, Nebraska, South Dakota and Tennessee. Human-like_2010.1 H3 was the predominant H3.

National phylogenetic NA gene information

Whole genome patterns with HA/NA pairs were reported and the dominant patterns are in descending order below with most frequent pattern by HA/NA pair:

- TTTTPT: H3.2010.1/N2-02, H1.gamma/cN1, H1.delta1a/N2-02,
- TTTTPT: H1.delta2/N2-98, H1.gamma/cN1, H1.delta1b/N2-02,
- TTPPPT: H1.gamma/cN1, H1.alpha/N2-02, H1.delta1a/N2-02.

Leading vaccine constellation is TVVTPT, with other patterns detected. Both the N1 and N2 subtypes are found in circulating swine viruses. Classical N1 continued to be the dominant cluster at 82.5 percent; and the 2002-lineage N2 represents 65 percent of N2 collections.

Objective 2. Make Influenza Isolates from Swine Available for Research and Establish a Data Management System to Facilitate Genetic Analysis of these Isolates and Related Information

A primary goal of IAV swine surveillance is to share selected virus isolates obtained through the surveillance system with public health, animal health, and academic researchers to facilitate genetic analysis and research on viruses of interest. The NVSL Diagnostic Virology Laboratory maintains a repository of the viruses submitted into the surveillance system and provides these viruses upon request.

In the first quarter of fiscal year 2020, the NVSL Diagnostic Virology Laboratory provided 33 isolates to two government, two academic, two pharmaceutical, and two international institutions. NVSL received 328 isolates into the repository (Table 4). Table 5 reports the total number of isolates available in the repository by subtype for sharing.

Table 4. Virus isolates received in repository.

Virus isolates in the repository	
2020	277
2019	1,055
2018	994
2017	844
2016	1,046
2015	883
2014	765
TOTAL TO DATE	5,864

Objective 3. Select Proper Isolates for Development of Relevant Diagnostic Reagents, Updating Diagnostic Assays, and Vaccine Seed Stock Products

USDA makes IAV-S isolates available in the public domain for further research. ARS-NADC conducts research on isolates obtained from the repository and sequences generated from the surveillance system. Genetic sequencing from the USDA program that is reported to GenBank is available for private corporations, government entities, academia, and other scientific community partners for research and vaccine strain selection and efficacy testing. NVSL and ARS staff are consulted as subject matter experts when necessary.

Table 5. Total number of subtyped isolates available through repository.

Subtyped isolates available through repository	
H3N2	1,932
H3N1	16
H1N1	2,563
H1N2	2,396
Mixed	302
TOTAL	7,209

Conclusion

The IAV voluntary surveillance system in swine continues to provide insight into the genetic makeup of circulating influenza A virus in limited populations of commercial pigs. Genetic information and virus isolates are made publicly available for further research and possible vaccine strain selection and efficacy testing. Influenza A virus in swine remains a dynamic virus with high levels of genetic variability in the hemagglutinin and neuraminidase genes.

Appendix 1. Regional Charts of HA and NA Combinations by Percentage

The following charts present the percentages of combinations of HA and NA on the national and regional scales based on ARS-NADC phylogenetic analyses. The results are reported from January 2018 to December 2019. These “heat maps” represent the percentage of combinations by using a color gradient where a deeper gradient color represents a greater percentage occurrence for a particular HA-NA combination. HA clusters are listed on the left vertical axis of the chart and NA clusters are listed on the bottom horizontal axis. Line up the HA cluster with the corresponding NA cluster to determine the occurrence of that particular combination.



Region 1. Total HA & NA combinations –264

Region 1

Percentage of HA and NA combinations – Jan 2018 to Dec 2019




Total HA & NA combinations – 264



Region 2. Total HA & NA combinations –1,278

Region 2

Percentage of HA and NA combinations – Jan 2018 to Dec 2019




Total HA & NA combinations – 1278



Region 3. Total HA & NA combinations - 159

Region 3




Total HA & NA combinations – 159



Region 4. Total HA & NA combinations – 172

Region 4

Percentage of HA and NA combinations – Jan 2018 to Dec 2019




Total HA & NA combinations – 172



Region 5. Total HA & NA combinations – 38

Region 5

Percentage of HA and NA combinations – Jan 2018 to Dec 2019



Total HA & NA combinations – 38