



United States Department of Agriculture

---

Animal and  
Plant Health  
Inspection  
Service

Veterinary  
Services

January 2018

# Influenza A Virus in Swine Surveillance

## Fiscal Year 2017 Quarterly Report

Surveillance Summary for Fourth Quarter FY 2017:

July 1 – September 30, 2017

---

### Report Summary<sup>1</sup>

- This report covers the fourth quarter (Q4) of fiscal year (FY) 2017, from July 1, 2017– September 30, 2017.
- Where relevant, the report also includes previous years' data for historical perspective.
- The report provides data from both national and regional levels.
- In FY 2017 Q4, 6,508 samples were submitted for IAV-S surveillance from 1,980 accessions.
- H1N2 was the predominant subtype.
- Over the past 8 quarters, H1N2 predominated in Regions 1, 2, 3, 4 and 5. When Regions are recorded as "unknown," H1N2 predominates as well (see map in regional section).
- Limited accessions from a region can skew data and lead to misinterpretation. Therefore, less inference can be applied to results from Regions 3, 4, and 5.
- All IAV-S submissions are voluntary and based on clinical case submissions to veterinary diagnostics labs. These data are not a statistically representative sampling of the U.S. swine population.
- Due to the voluntary nature of this surveillance, the information in this report cannot be used to determine regional and/or national incidence, prevalence, or other epidemiological measures, but it may help identify IAV-S trends.

### Introduction

This report, based on data received as of December 28, 2017, provides a brief update on the status of national surveillance for IAV in swine for producers, swine practitioners, diagnosticians, and the public. Summaries in this report may differ from those provided in past reports due to the regular addition of data from participating laboratories. Reporting months are based on the month the sample was collected. The USDA-APHIS web site provides general information about the IAV-S surveillance program at [https://www.aphis.usda.gov/aphis/ourfocus/animalhealth/animal-disease-information/swine-disease-information/ct\\_siv\\_surveillance](https://www.aphis.usda.gov/aphis/ourfocus/animalhealth/animal-disease-information/swine-disease-information/ct_siv_surveillance)

---

<sup>1</sup> In November 2016, VS modernized the process that prepares and stages laboratory results data for reporting. Consequently, VS recognizes there is a small difference in previously reported summary numbers for IAV-S surveillance. The results in this report reflect updated and corrected numbers achieved with the modernized data process.

The IAV-S surveillance program is voluntary and, as a result, the accessions and samples submitted represent a subset of the swine population. Submitted samples should only be collected from animals displaying influenza-like illness. Due to its voluntary nature, this surveillance system does not entirely represent the total U.S. domestic swine population. Therefore, the data cannot be used to determine IAV-S prevalence or other epidemiologic measures in the swine population. However, the data may help identify trends in influenza in swine.

When the submitter does not report relevant information, data are recorded as “unknown.” Summaries in this report may differ from those provided in past reports due to the ongoing addition of data from participating labs. Reporting months are based on the month when the sample was collected.

A laboratory accession is generally a set of samples collected at a single premises on a single day and received at the laboratory. While a nasal swab or lung tissue sample represents a single animal within the herd, a single oral fluid sample may represent one to two pens of animals in a herd. A positive sample status is based on the screening matrix real-time reverse transcriptase polymerase chain reaction (rRT-PCR). The subtype is based on the rRT-PCR- subtyping assays and sequencing if the sample is sequenced. Virus isolation (VI) and sequencing are only attempted on rRT-PCR positives meeting criteria listed in the testing algorithm posted on the web ([https://www.aphis.usda.gov/animal\\_health/animal\\_dis\\_spec/swine/downloads/appendix\\_c\\_testing\\_guidelines.pdf](https://www.aphis.usda.gov/animal_health/animal_dis_spec/swine/downloads/appendix_c_testing_guidelines.pdf)). Phylogenetic analyses are based on successful sequencing results, with sequences deposited into GenBank, the public sequence database.

## Program Updates

Virus isolates acquired through USDA’s IAV-S surveillance are sent to the National Veterinary Services Laboratories (NVSL) in Ames, Iowa, and deposited into the IAV-S virus repository. Testing labs post the genomic sequences from the isolates into GenBank®, a publically accessible data base maintained by the National Institutes for Health.

USDA swine surveillance influenza A virus isolates can be identified in GenBank® by their eight-digit barcode designations in the virus name, e.g. A/swine/NY/A01104005/2011(H3N2). The information in GenBank® will usually list “USDA Swine Surveillance” or similar in the *Consortium* field. The majority of available isolates will have H, M, and N sequenced, but some will also have full genome sequencing completed. The repository contains over 3,000 viruses. These viruses are made available to the public upon request.

NVSL recently posted instructions for repository IAV-S isolate requests. In order to obtain isolates from the repository:

- Contact the repository microbiologist by email at [nvsl.dvl.bpa@aphis.usda.gov](mailto:nvsl.dvl.bpa@aphis.usda.gov) or telephone at (515) 337-7551.
- Provide a list of barcode IDs (e.g. A01104005) for the sought-after viruses to assess the availability of each isolate. Many isolates are ready to ship, but some will need to be propagated before shipping.
- Once the request list has been finalized, visit the following website for information on permit requirements and placing your order:  
[https://www.aphis.usda.gov/aphis/ourfocus/animalhealth/lab-info-services/sa\\_reagents/ct\\_reagent\\_order](https://www.aphis.usda.gov/aphis/ourfocus/animalhealth/lab-info-services/sa_reagents/ct_reagent_order)
  - The reagent code number for repository isolates is SIV-REP.
  - The reagent name is the barcode ID.
  - A user fee is charged per vial of virus ordered.

These instructions are posted on the USDA website here:

[https://www.aphis.usda.gov/aphis/ourfocus/animalhealth/lab-info-services/sa\\_reagents/ct\\_reagents](https://www.aphis.usda.gov/aphis/ourfocus/animalhealth/lab-info-services/sa_reagents/ct_reagents)

The link to the PDF of the instructions can be downloaded from here:

[https://www.aphis.usda.gov/animal\\_health/lab\\_info\\_services/downloads/OrderingIAV-SRepositoryIsolates.pdf](https://www.aphis.usda.gov/animal_health/lab_info_services/downloads/OrderingIAV-SRepositoryIsolates.pdf)

## **IAV-S Surveillance Objectives**

USDA's National Surveillance Plan for Swine Influenza Virus in Pigs (July 2010) describes the current surveillance system for IAV-S in detail. The surveillance objectives are to:

1. Monitor genetic evolution of endemic IAV in swine to better understand endemic and emerging influenza virus ecology;
2. Make influenza isolates from swine available for research and establish a data management system to facilitate genetic analysis of these isolates and related information; and
3. Select proper isolates for the development of relevant diagnostic reagents, updated diagnostic assays, and vaccine seed stock products.

### **Objective 1. Monitoring genetic evolution of endemic IAV in swine to better understand endemic and emerging influenza virus ecology**

Objective 1 is met through the submission of diagnostic laboratory samples to the surveillance system, collection of the viruses that are isolated from the samples, and analysis of the HA and NA sequences that are generated at the NAHLN laboratories and posted into GenBank®. Each month selected viruses undergo whole genomic sequencing by NVSL. Phylogenetic analysis of the genetic sequences is provided through an interagency agreement with the USDA's Agricultural Research Service (ARS) National Animal Disease Center (NADC).

### **National Surveillance Data Summary**

From FY 2011 through FY 2016, the total number of accessions and samples submitted rose over time. Changes initiated in FY 2016 have resulted in decreased laboratory accessions and samples, but have resulted in a higher percentage of accessions that yield a virus that can be sequenced and analyzed. For FY 2017's fourth quarter, 6,508 samples were tested from 1,980 accessions (Figure 2) for a fiscal year-to-date total of 20,246 samples and 6,657 accessions. Figure 3 shows the overall trends in total accessions, PCR-positive accessions, subtyped accessions, and VI positive accessions.

Influenza A Virus in Swine Surveillance Quarterly Report for FY 2017, Quarter 4




Figure 2. Number of IAV-S laboratory accessions and samples tested in swine, FY 2011 through FY 2017 Q4




Figure 3. Accessions submitted, subtyped accessions, positive accessions, and VI positive accessions over time with trend lines, FY 2011 through FY 2017 Q4

Figure 4 shows the number of subtype detections in FY 2017 Q4. The total number of samples subtyped was 177, including 50 H1N1, 76 H1N2, 46 H3N2, 0 H3N1, and 5 mixed.




Figure 4. Number of subtype detections in FY 2017 Q4

Figure 5 breaks down accessions by rRT-PCR subtype from FY 2011 to FY 2017 Q4. H1N1 remains the major subtype over the course of the surveillance; however, H1N2 and H3N2 detections have increased substantially since 2012 and H1N2 jumped to the most common subtype through the first, second, third, and fourth quarters of FY 2017.





Figure 5. Number of subtypes, FY 2011 through FY 2017 Q4

Figure 6 displays the number of VIs attempted, the number of those attempts that were positive, and the number of positive VIs that are submitted to GenBank.



**Figure 6. Number of virus isolations attempted, positive virus isolations, and GenBank submissions from FY 2013 through FY 2017 Q4**

When accessions were evaluated by age-class for the third quarter, the following observations were noted. H1N2 was the most common subtype among suckling and nursery pigs. H1N1 was the most common subtype among grower/finishers. Sow/boar had limited testing, with only two samples, both of which were H1N2. Among accessions for which the age class was unknown or not recorded, H1N1 was the predominant subtype (Table 1). Samples collected from nasal/ nasal swabs are the most successful at providing positive virus isolation and submission to GenBank (Table 2).

**Table 1.** Number of positive accessions tested for IAV-S by age class and by viral subtype, Q4 FY 2017.

Age Class (group)	Number of accessions with subtype reported	Number of H1N1	Number of H1N2	Number of H3N1	Number of H3N2	Number of Mixed
Suckling	29	0	20	0	8	1
Nursery	62	14	25	0	20	3
Grower/Finisher	47	20	16	0	10	1
Sow/Boar	2	0	2	0	0	0
Not Recorded/Unknown	37	16	13	0	8	0

**Table 2.** Number of positive accessions tested for IAV-S by specimen type and by viral subtype, Q4 FY 2017.

Specimen Type (group)	Number of accessions with subtype reported	Percent of subtyped accessions with positive virus isolation	Number of H1N1	Number of H1N2	Number of H3N1	Number of H3N2	Number of Mixed	Number of samples sequenced and sent to GenBank
Lung	144	91%	42	61	0	36	5	129
Nasal or Nasal Swab	23	100%	5	9	0	9	0	23
Oral Fluids	10	60%	3	6	0	1	0	7
Other Specimens	0		0	0	0	0	0	0


\*Accessions may include samples with multiple specimen types. In these cases, individual accessions are counted in more than one specimen type category.

\*\*Other includes specimen types recorded as swab, mixed tissue, or unknown.

## Regional surveillance data

In this section, we present data in five different regions (Figure 6) to parse the analysis across regions. These regions are based on current USDA administrative districts for simplicity; these divisions do not represent specific industry distribution. Submissions are voluntary, as is any identifying information accompanying the submission (except the State of animal origin), and therefore no sampling strategies can be applied to the regions.

**Figure 6. A map of the regions for national IAV-S surveillance**



### Summary of Regional Data from ARS

**Table 3. Summary of predominant subtypes in each region for FY 2015 Q4 through FY 2017 Q4**

**Most Predominant HA/NA phylo-types overall:**

H1N1 (Gamma H1/Classical N1)    H1N2 (Delta1a H1/2002-N2)  
 H1N2 (Delta2 H1/1998-N2)    H3N2 (IV-A H3/2002-N2)  
 H3N2 (hu-like\_2010 H3/2002-N2)    H1N2 (Delta1b H1/2002-N2)

**Region 1 (Total HA/NA: 334)**

Gamma H1/Classical N1  
 IV-A H3/2002-N2  
 Delta2 H1/1998-N2  
 Alpha H1/2002-N2  
 Low frequency but consistent detections of IV-B H3/2002-N2, pdm H1/pdm N1, delta2 H1/2002-N2, delta1a H1/2002-N2, human-like\_2010 H3/2002-N2, IV-B H3/classical N1

**Region 4 (Total HA/NA: 129)**

Delta1b H1/2002-N2  
 IV-A H3/2002-N2  
 Delta1a H1/2002-N2  
 Gamma H1/Classical N1  
 Pdm H1/Pdm N1  
 Beta H1/2002-N2  
 Low frequency but consistent detections of Alpha H1/2002-N2, Hu-like\_2010 H3/2002-N2, gamma H1/2002-N2

**Region 2 (Total HA/NA: 1,060)**

Most diversity of all regions  
 Gamma H1/Classical N1  
 Delta1 H1a/2002-N2  
 Hu-like\_2010 H3/2002-N2  
 Delta2 H1/1998-N2  
 IV-A H3/2002-N2  
 Low frequency but consistent detections of IV-E H3/2002-N2, IV-B H3/2002-N2, pdm H1/pdm N1, alpha H1/2002-N2, delta1b H1/2002-N2

**Region 5 (Total HA/NA: 7)**

Low participation  
 Three Delta 1b H1/2002-N2  
 Two Alpha H1/2002-N2  
 One Hu-like\_2010 H3/2002 N1  
 One IV-A H3/2002-N2

**Region 3 (Total HA/NA: 126)**

Delta1a H1/2002-N2  
 Gamma H1/Classical N1  
 IV-A H3/2002-N2  
 Hu-like\_2010 H3/2002-N2  
 Low frequency detections of Beta H1/Classical N1, Pdm H1/Pdm N1, Delta1b H1/2002-N2



Figure 8 shows the distribution of rRT-PCR subtyped accessions among the five regions for Q4 FY 2015 through Q4 FY 2017. Regions 1, 2, 3, 4, and 5 demonstrate H1N2 as the predominant subtype. For regions recorded as “unknown,” H1N2 was also the predominant subtype.




Figure 8. Percentage of subtyped accessions by region for FY 2015 Q4 through FY2017 Q4


## Regional phylogenetic analysis

### Phylogenetic analysis of sequences from the IAV-S surveillance system

Phylogenetic analysis of gene sequences of the influenza A virus in swine is conducted to further examine the genetic changes that occur in HA, NA, and M genes of this rapidly changing virus. Through collaboration with ARS, a dataset<sup>2</sup> of 151 isolates with published sequences in GenBank<sup>®</sup> was characterized by phylogenetic analysis in Q4 FY 2017. This analysis provides information on the genetic diversity and evolution patterns of influenza in swine and allows for inferences about population and/or vaccine immunity.

The following series of bar charts parse the data into an approximate 2-year window by quarters and subtypes for each region, followed by charts further parsing the H1 and H3 subtypes into phylogenetic clades. Regional charts depicting the various combinations of HA and NA are available in Appendix 1.

<sup>2</sup> The ARS dataset is comprised of IAV-S surveillance isolate sequences that were posted in Genbank. This represents only a subset of the complete IAV-S surveillance dataset that includes PCR diagnostic test-based results as well as sequencing results. Therefore, ARS dataset results, such as subtype percentages, differ from the complete IAV-S dataset results provided in other sections of this report.



**Figure 9. Virus type by region 2-year summary Q1 FY 2016 to Q4 FY 2017**

Figure 9 demonstrates the four subtypes H1N1, H1N2, H3N1, H3N2, and mixed subtypes across the five regions. There was also an H4N6 reported in FY 2016 Q1. Regions 1 and 2 reported the most submissions, with a mixture of mostly H1N1, H1N2, and H3N2. Limited accessions from a region can skew data and lead to misinterpretation and therefore, less inference can be applied to results from Regions 3, 4, and 5.

**National phylogenetic HA gene information**

HA genes from H1 subtype viruses are classified as alpha, beta, gamma, delta-1, delta-2, or pandemic H1N1 2009 (H1N1pdm09) phylogenetic clades based on a previously published nomenclature system. Similarly, H3 subtype viruses are classified as Cluster IV, Cluster IV-A, Cluster IV-B, Cluster IV-C, Cluster IV-D, Cluster IV-E, Cluster IV-F, or human-like. In the H1 subtypes (see Figure 10), there continued to be detections of alpha with two aa deletions (n=5). Delta 1a, delta 2, and gamma viruses are the predominant H1.





Figure 10. H1 phylo-cluster by region – 2-year summary Q1 FY 2016 to Q4 FY 2017



**Figure 11. H3 phylo-cluster by region, 2-year summary Q1 FY 2016 to Q4 FY 2017**

In Q4 FY 2017, there were detections of human-like H3 in Iowa, Illinois, Ohio, and Wisconsin. Out of 23 H3s in Q4 FY 2017, 18 are human-like H3s. Human-like H3 is the predominant H3. A new human-like H3 spillover associated with the H3 circulating in humans in 2016 appeared in IAV-S surveillance data in 2017.

**National phylogenetic NA gene information**

NA gene information remained the same in Q4 FY 2017. Both the N1 and N2 subtypes are found in circulating swine viruses. Classical N1 continued to be the dominant cluster. The 2002-lineage N2 represents 82 percent of N2 collections.

## Objective 2. Make influenza isolates from swine available for research and establish a data management system to facilitate genetic analysis of these isolates and related information

A primary goal of IAV-S surveillance is to share selected virus isolates obtained through the surveillance system with public health, animal health, and academic researchers to facilitate genetic analysis and research on viruses of interest. The NVSL Diagnostic Virology Laboratory maintains a repository of the viruses submitted into the surveillance system and provides these viruses upon request.

In the fourth quarter of FY 2017, the NVSL Diagnostic Virology Laboratory provided 71 isolates to four institutions: one governmental, two pharmaceutical, and one international (government). NVSL received 144 isolates into the repository in Q4 for a total of 844 in FY 2017 (Table 4). Table 5 reports the total number of isolates available in the repository by subtype for sharing.

**Table 4. Virus isolates received in repository**

Virus isolates in the repository	
2017 YTD	844
2016	1,046
2015	883
2014	765
2013	820
<b>TOTAL TO DATE</b>	<b>4,358</b>

## Objective 3. Select proper isolates for development of relevant diagnostic reagents, updating diagnostic assays, and vaccine seed stock products

USDA makes IAV-S isolates available in the public domain for further research. ARS-NADC conducts research on isolates obtained from the repository and sequences generated from the surveillance system. Genetic sequencing reported to GenBank is available for private corporations, government entities, academia, and other scientific community partners for research and vaccine strain selection and efficacy testing. NVSL and ARS staff are consulted as subject matter experts when necessary.

**Table 5. Total number of subtyped isolates available through repository**

Subtyped isolates available through repository	
H3N2	1,627
H3N1	15
H1N1	2,084
H1N2	1,969
Mixed	316
<b>TOTAL</b>	<b>6,011</b>

## **Conclusion**

The IAV-S voluntary surveillance system continues to provide insight into the genetic makeup of circulating influenza virus in limited populations of commercial pigs. Genetic information and virus isolates are made publicly available for further research and possible vaccine strain selection and efficacy testing. Influenza A virus in swine remains a dynamic virus with high levels of genetic variability in the hemagglutinin and neuraminidase genes.


## **Appendix 1. Regional Charts of HA and NA Combinations by Percentage**

The following charts present the percentages of combinations of HA and NA on the national and regional scales based on ARS-NADC phylogenetic analyses. The results are reported from October 2015 to September 2017. These “heat maps” represent the percentage of combinations by using a color gradient where a deeper gradient represents a greater percentage occurrence for a particular HA-NA combination. HA clusters are listed on the left vertical axis of the chart and NA clusters are listed on the bottom horizontal axis. Line up the HA cluster with the corresponding NA cluster to determine the percentage of occurrence of that particular combination.



Total HA & NA combinations – 1,658

Percentage of HA and NA combinations – Oct 2015 to Sept 2017








**Region 1. Total HA & NA combinations – 334**


Percentage of HA and NA combinations – October 2015 to September 2017





**Region 2. Total HA & NA combinations – 1,060**


Percentage of HA and NA combinations – October 2015 to September 2017





**Region 3. Total HA & NA combinations – 126**


Percentage of HA and NA combinations – October 2015 to September 2017





**Region 4.** Total HA & NA combinations – 129

Percentage of HA and NA combinations – October 2015 to September 2017





**Region 5. Total HA & NA combinations –7**

Percentage of HA and NA combinations – October 2015 to September 2017

