

**FOOT-AND-MOUTH DISEASE
STANDARD OPERATING PROCEDURES:
1. OVERVIEW OF ETIOLOGY AND ECOLOGY**

FAD PReP

**Foreign Animal Disease
Preparedness & Response Plan**



**United States
Department of
Agriculture**

United States Department of Agriculture • Animal and Plant Health Inspection Service • Veterinary Services

File name:	FADPREP_FMD_E&E_SOP	
Lead section:	Preparedness and Incident Coordination	Version number: 2.0
Effective date:	November 2016	Review date: November 2018

The Foreign Animal Disease Preparedness and Response Plan (FAD PReP) Standard Operating Procedures (SOPs) provide operational guidance for responding to an animal health emergency in the United States.

These draft SOPs are under ongoing review. This document was last updated in **November 2016**. Please send questions or comments to:

National Preparedness and Incident Coordination Center
 Veterinary Services
 Animal and Plant Health Inspection Service
 U.S. Department of Agriculture
 4700 River Road, Unit 41
 Riverdale, Maryland 20737
 Fax: (301) 734-7817
 E-mail: FAD.PReP.Comments@aphis.usda.gov

While best efforts have been used in developing and preparing the FAD PReP SOPs, the U.S. Government, U.S. Department of Agriculture (USDA), Animal and Plant Health Inspection Service, and other parties, such as employees and contractors contributing to this document, neither warrant nor assume any legal liability or responsibility for the accuracy, completeness, or usefulness of any information or procedure disclosed. The primary purpose of these FAD PReP SOPs is to provide operational guidance to those government officials responding to a foreign animal disease outbreak. It is only posted for public access as a reference.

The FAD PReP SOPs may refer to links to various other Federal and State agencies and private organizations. These links are maintained solely for the user's information and convenience. If you link to such site, please be aware that you are then subject to the policies of that site. In addition, please note that USDA does not control and cannot guarantee the relevance, timeliness, or accuracy of these outside materials. Further, the inclusion of links or pointers to particular items in hypertext is not intended to reflect their importance, nor is it intended to constitute approval or endorsement of any views expressed, or products or services offered, on these outside websites, or the organizations sponsoring the websites.

Trade names are used solely for the purpose of providing specific information. Mention of a trade name does not constitute a guarantee or warranty of the product by USDA or an endorsement over other products not mentioned.

USDA prohibits discrimination in all its programs and activities on the basis of race, color, national origin, sex, religion, age, disability, political beliefs, sexual orientation, or marital or family status. (Not all prohibited bases apply to all programs.) Persons with disabilities who require alternative means for communication of program information (Braille, large print, audiotape, etc.) should contact USDA's TARGET Center at (202) 720-2600 (voice and telecommunications device for the deaf [TDD]).

To file a complaint of discrimination, write USDA, Director, Office of Civil Rights, Room 326-W, Whitten Building, 1400 Independence Avenue SW, Washington, DC 20250-9410 or call (202) 720-5964 (voice and TDD). USDA is an equal opportunity provider and employer.

Contents

1.1	Introduction.....	1-2
	1.1.1 Goals	1-2
	1.1.2 Further Information.....	1-2
1.2	Purpose.....	1-3
1.3	Etiology.....	1-3
	1.3.1 Name	1-3
	1.3.2 Virus Characteristics	1-3
	1.3.3 Morphology.....	1-4
	1.3.4 FMD Serotypes and Strains	1-4
1.4	Ecology	1-5
	1.4.1 General Overview	1-5
	1.4.2 Susceptible Species	1-5
	1.4.3 Carrier State	1-6
	1.4.4 Introduction and Transmission of FMDV	1-8
	1.4.5 Incubation Period	1-11
	1.4.6 Morbidity and Mortality	1-12
1.5	Environmental Persistence of FMDV	1-13
	1.5.1 Environmental Persistence of FMDV in Excretions and Surroundings	1-14
	1.5.2 Environmental Persistence of FMDV in Milk and Dairy Products	1-14
	1.5.3 Environmental Persistence of FMDV in Meat and Meat Products.....	1-15
	1.5.4 Environmental Persistence of FMDV in Wool and Hides.....	1-15
1.6	OIE Procedures for the Inactivation of FMDV.....	1-15
	1.6.1 Meat	1-16
	1.6.2 Wool and Hair.....	1-16
	1.6.3 Milk and Cream for Human Consumption	1-17
	1.6.4 Milk for Animal Consumption.....	1-17
	Attachment 1.A References and Selected Resources	1-18
	Attachment 1.B Abbreviations.....	1-23

Foot-and-Mouth Disease

Etiology & Ecology Quick Summary

Disease

Foot-and-mouth disease (FMD), fiebre aftosa, fièvre aphteuse, Maulund-Klauenseuche.

Mortality & Morbidity

Very high morbidity; high mortality in young animals.

Susceptible Species

Domestic cloven hoofed animals (cattle, swine, sheep, goats) and wildlife (deer, bison, pronghorn antelope, feral swine).

Zoonotic Potential?

Not a threat to public health.

Carriers

Ruminants may become carriers of FMD; an estimated 50 percent of cattle may become carriers.

Transmission

Pigs excrete large amounts of virus through respiration. Virus spreads readily on contaminated equipment and other fomites. Windborne transmission can occur.

Persistence in the Environment

Survives longer at cold temperatures. Susceptible to acid and alkaline pH. Can remain infective for long periods of time where organic matter is present.

Animal Products and By-Products

FMD virus can survive for months in chilled/frozen bone marrow, lymph nodes, and can survive in sausages and other cured meats. Typical pasteurization may not completely inactivate FMD from milk.

1.1 Introduction

Foot-and-mouth disease (FMD) is a highly contagious viral disease that affects domestic cloven-hoofed animals (cattle, swine, sheep, and goats) and more than 70 wildlife species (deer, bison, pronghorn antelope, and feral swine). The disease is characterized by fever, vesicular (blister-like) lesions, and subsequent erosions on the surfaces of the mouth, tongue, nostrils, muzzle, feet, and teats.

FMD is one of the most contagious livestock diseases and endemic in many areas of the world; introduction of FMD into the United States is a serious concern. All secretions, excretions, and tissues are contagious, and the virus may be present in respirations, milk, and semen. The virus enters a new susceptible animal either orally (especially swine) or via the respiratory tract (especially cattle). Aerosol transmission is the major means of animal-to-animal spread within premises. Though FMD virus (FMDV) is not typically considered a zoonotic disease, and is not a threat to public health, there is evidence demonstrating humans can carry the virus mechanically in their nasal passages.^{1,2,3}

FMD is not normally characterized by high death rates in adult animals, but it can produce morbidity rates of almost 100 percent, resulting in severe losses to productivity. It is probable that all animals in a susceptible population will be affected. Although it does not tend to kill adults, high mortality rates may be observed in young animals.^{4,5,6}

1.1.1 Goals

As a preparedness goal, the Animal and Plant Health Inspection Service (APHIS) will provide etiology and ecology summaries for FMD, and update these summaries at regular intervals.

As a response goal, the Unified Command and stakeholders will have a common set of etiology and ecology definitions and descriptions, to ensure proper understanding of FMD when establishing or revising goals, objectives, strategies, and procedures.

1.1.2 Further Information

This document is intended to be an overview, focusing on FMD in domestic livestock. Additional resources on FMD are listed in Attachment 1.A. In addition to this *FMD Standard Operating Procedure (SOP): Overview of Etiology and Ecology*, information can also be found

¹ Sellers, R.F., et al. (1970). Inhalation, persistence and dispersal of foot-and-mouth disease virus by man. *Journal of Hygiene* (London), 68(4), 565–573. DOI: 10.1017/s0022172400042492.

² Amass, S., et al. (2003). Procedures for preventing the transmission of foot-and-mouth disease virus to pigs and sheep by personnel in contact with infected pigs. *Veterinary Record*, 153(5), 137–140. DOI: 10.1136/vr.153.5.137.

³ Amass, S., Pacheco, J., Mason, P., Pacheco, J., Miller, C., Ramirez, A., Clark, L., Ragland, D., Schneider, J., & Kenyon, S. (2004). Procedures for preventing transmission of foot-and-mouth disease virus (O/TAW/97) by people. *Veterinary Microbiology*, 103(3–4), 143–149. DOI: 10.1016/j.vetmic.2004.07.020.

⁴ Musser, J.M. (2004). A practitioner's primer on foot-and-mouth disease. *Journal of the American Veterinary Medical Association*, 224(8), 1261–1268. DOI: 10.2460/javma.2004.224.1261.

⁵ Kitching, R.P. & Alexandersen, S. (2002). Clinical variation in foot and mouth disease: pigs. *Scientific and Technical Review of the Office International des Epizooties*, 21(3), 513–518. DOI: 10.20506/rst.21.3.1367.

⁶ Kitching, R.P. & Hughes, G.J. (2002). Clinical variation in foot and mouth disease: sheep and goats. *Scientific and Technical Review of the Office International des Epizooties*, 21(3), 505–512. DOI: 10.20506/rst.21.3.1342.

in the *FMD Response Plan: The Red Book*. Case definitions and laboratory criteria are provided from Surveillance, Design, and Analysis in the APHIS Center for Epidemiology and Animal Health.

This document does not comprehensively discuss vaccination, or its effects on immunity. For more specific technical information related to vaccination, as well as a detailed review of scientific literature relating to FMD, and please see the *National Animal Health Emergency Management System (NAHEMS) Guidelines: Vaccination for Contagious Diseases, Appendix A: FMD*.

These documents are available on the APHIS Foreign Animal Disease Preparedness and Response (FAD PReP) website: www.aphis.usda.gov/fadprep.

1.2 Purpose

This document provides responders and stakeholders with a common understanding of the disease agent.

1.3 Etiology

1.3.1 Name

This disease is called foot-and-mouth disease, fiebre aftosa, fievre aphteuse, and maulund-klauenseuche.

1.3.2 Virus Characteristics

According to the International Committee on Taxonomy of Viruses, this disease has the following characteristics:⁷

- Family: Picornaviridae
 - Genus: *Aphthovirus*, containing four species:
 - Foot-and-mouth disease virus
 - Bovine rhinitis A virus
 - Bovine rhinitis B virus
 - Equine rhinitis A virus
- Baltimore Classification: Group IV (+) ssRNA.^{8,9}

⁷ International Committee on Taxonomy of Viruses. (2015). Foot-and-Mouth Disease. Retrieved from <http://ictvonline.org>.

⁸ Adams, R.L.P., et al. (1992). The structure of viruses. *The biochemistry of the nucleic acids* (p. 75–76). United Kingdom: Chapman & Hall. DOI: 10.1007/978-94-011-2290-0.

⁹ Shors, T. (2013). Baltimore classification. In *Understanding viruses* (56). Burlington, MA: Jones & Bartlett Learning.


1.3.3 Morphology

FMD is a single-stranded ribonucleic acid (RNA) virus, approximately 30 nanometers in diameter, with icosahedral symmetry. There are 60 copies of four structural proteins, VP1, VP2, VP3, and VP4. These proteins encapsidate a single strand of positive-sense RNA^{10,11,12}

1.3.4 FMD Serotypes and Strains

There are seven serotypes of FMD: O, A, C, Asia-1, and South African Territories (SAT) 1, SAT 2, and SAT 3. Type O is the most prevalent serotype worldwide, followed by type A.¹³ Within these serotypes, more than 60 strains have been identified.¹⁴ Figure 1-1 illustrates the typical distribution of serotypes in the world.^{15,16}

Figure 1-1. Distribution of FMD Serotypes



¹⁰ Belsham, G. & Bøtner, A. (2015). Use of recombinant capsid proteins in the development of a vaccine against foot-and-mouth disease virus (FMDV). *Virus Adaptation and Treatment*, 7, 11–23. DOI: 10.2147/VAAT.S55351.

¹¹ Gay, C.G., et al. (2010). Foot-and-Mouth Disease (FMD): Gap Analysis Workshop Report. In *Agricultural Research Service*. Retrieved from <http://go.usa.gov/kCqF>.

¹² Goodwin, S., et al. (2009). Foot-and-mouth disease virus assembly: processing of recombinant capsid precursor by exogenous protease induces self-assembly of pentamers in vitro in a myristoylation-dependent manner. *Journal of Virology*, 83(21), 11275–11282. DOI: 10.1128/JVI.01263-09.

¹³ Gay, C.G., et al. (2010). Foot-and-Mouth Disease (FMD): Gap Analysis Workshop Report. In *Agricultural Research Service*. Retrieved from <http://go.usa.gov/kCqF>.

¹⁴ Center for Food Security and Public Health (CFSPH). (2014). Foot-and-Mouth Disease. In *Technical Factsheet*. Retrieved from www.cfsph.iastate.edu.

¹⁵ World Organization for Animal Health (OIE). (2016). Foot-and-Mouth Disease Distribution Maps, 2013–2016. In *World Animal Health Information Database (WAHID) World Animal Health Information System (WAHIS) Interface*. Retrieved from http://www.oie.int/wahis_2/public/wahid.php/Diseaseinformation/Diseasedistributionmap.

¹⁶ Knowles, N.J., et al. (2005). Pandemic strain of foot-and-mouth disease virus serotype O. *Emerging Infectious Diseases*, 11(12), 1887–1893. DOI: 10.32001/eid1112.050908.

FMDV is highly susceptible to change, resulting from errors in RNA replication, recombination, and host selection.¹⁷ There is no cross-protection between the distinct serotypes.¹⁸ Protection within serotypes varies based on the antigenic similarity of the strains. Subsequently, any vaccine must be carefully matched with the field strain to be effective. For more information on FMD vaccination matching, please see the *FAD PReP/NAHEMS Guidelines: Vaccination for Contagious Diseases, Appendix A: FMD* found at www.aphis.usda.gov/fadprep.

1.4 Ecology

1.4.1 General Overview

FMD is currently endemic in areas of Africa, Asia, Eastern Europe, the Middle East, and South America.¹⁹ North America (the United States, Canada, and Mexico), Central America, Western Europe, Australia, and New Zealand are all free of FMD.²⁰ From 2013 to 2016, FMD outbreaks have occurred in countries including (but not limited to) Egypt, China, Vietnam, North and South Korea, Russia, South Africa, Iraq, Nepal, Cambodia, Venezuela, and Botswana.²¹ The last FMD outbreak in the United States was in 1929.²²

1.4.2 Susceptible Species

FMDV infects cloven-hooved mammals (order *Artiodactyla*), as well as a few species in other orders. Livestock susceptible to FMD, common in the United States, include:

- cattle,
- pigs,
- sheep, and
- goats.

In addition, deer, bison, and elk are also susceptible to the virus. Wild pigs, antelope, African buffalo, Bactrian camels, and giraffe are also all susceptible species. Llamas and alpacas have been infected experimentally, but do not seem to be highly susceptible to natural infection. Other animals like rats, mice, guinea pigs, and armadillos have all been experimentally infected.^{23,24,25}

¹⁷ OIE. (2013). Foot and Mouth Disease. In *Technical Disease Card*. Retrieved from <http://www.oie.int>.

¹⁸ OIE. (2013). Foot and Mouth Disease. In *Technical Disease Card*. Retrieved from <http://www.oie.int>.

¹⁹ OIE. (2013). Foot and Mouth Disease. In *Technical Disease Card*. Retrieved from <http://www.oie.int>.

²⁰ For a list of the regions considered to be free of Foot-and-Mouth Disease by U.S. Department of Agriculture (USDA) APHIS, please see <https://www.aphis.usda.gov/aphis/ourfocus/animalhealth/animal-and-animal-product-import-information/import-live-animals/ct-foot-and-mouth-and-rinderpest>.

²¹ OIE. (2016). Foot-and-Mouth Disease Distribution Maps, 2013–2016. In *World Animal Health Information Database (WAHID) World Animal Health Information System (WAHIS) Interface*. Retrieved from http://www.oie.int/wahis_2/public/wahid.php/Diseaseinformation/Diseasedistributionmap.

²² CFSPH. (2014). Foot-and-Mouth Disease. In *Technical Factsheet*. Retrieved from www.cfsph.iastate.edu.


²³ Musser, J.M. (2004). A practitioner's primer on foot-and-mouth disease. *Journal of the American Veterinary Medical Association*, 224(8), 1261–1268. DOI: 10.2460/javma.2004.224.1261.

²⁴ OIE. (2013). Foot and Mouth Disease. In *Technical Disease Card*. Retrieved from <http://www.oie.int>.

²⁵ Lubroth, J., et al. (1990). Foot-and-mouth disease virus in the llama (*Lama glama*): diagnosis, transmission, and susceptibility. *Journal of Veterinary Diagnostic Investigation*, 2(3), 197–203. DOI: 10.1177/104063879000200308.

Strains may have a predilection for one animal species over another.²⁶ Figure 1-2 presents the density of FMD susceptible livestock species in the world.

Figure 1-2. Density of FMD Susceptible Animals Livestock Species, i.e., Combined Cattle, Pigs, Sheep, and Goats



Source: Knight-Jones, T. and Rushton, J. (n.d.).

1.4.3 Carrier State

A carrier is defined as an animal in which there is persistence of FMDV or the viral genome in the pharyngeal region for at least 28 days post-infection.^{27,28} Generally speaking, a “carrier” is defined in epidemiological terms as an animal that is infected and can disseminate the infection in the absence of symptoms. However, with FMD, carrier animals may or may not be able to transmit infection.²⁹ African buffalo are the only carrier animals that have been demonstrated to

²⁶ CFSPH. (2014). Foot-and-Mouth Disease. In *Technical Factsheet*. Retrieved from www.cfsph.iastate.edu.

²⁷ Suttmoller, P., et al. (1968). The epizootiological importance of foot-and-mouth disease carriers. I: Experimentally produced foot-and-mouth disease carriers in susceptible and immune carriers. *Archives of Virology*, 23(3), 227–235. DOI: 10.1007/BF01241895.

²⁸ Suttmoller, P. & Casas-Olascoaga, R. (2002). Unapparent foot and mouth disease infection (sub-clinical infections and carriers): implications for control. *Scientific and Technical Review of the Office International des Epizooties*, 21(3), 519–529. DOI: 10.20506/rst.21.3.1366.

²⁹ Suttmoller, P. & Casas-Olascoaga, R. (2002). Unapparent foot and mouth disease infection (sub-clinical infections and carriers): implications for control. *Scientific and Technical Review of the Office International des Epizooties*, 21(3), 519–529. DOI: 10.20506/rst.21.3.1366.

be able to definitively transmit SAT FMDV to other buffalo and potentially cattle.^{30,31,32,33} Serotypes A, O, Asia-1, and C do not seem to be transmitted from carriers to naïve animals. For more information on carrier transmission, please see the *FAD PReP/NAHEMS Guidelines: Vaccination for Contagious Diseases, Appendix A: FMD* found at www.aphis.usda.gov/fadprep.

In addition to African buffalo, cattle, sheep, goats, and water buffalo can also become carriers; however, there is not sufficient evidence to suggest a carrier state exists in pigs.³⁴ In comparison to cattle and buffalo, sheep and goats are carriers for a shorter period of time—typically between 1–5 months.³⁵ On the other hand, while most cattle carry FMDV for 6 months or less, there is evidence of cattle carrying FMDV for more than 3 years.³⁶ Persistent infections have also been identified in some experimentally infected wildlife species, including white-tailed deer and kudu.³⁷

The epidemiological significance of carrier animals in FMD outbreaks of cattle is uncertain. There is no evidence that sheep or goats can transmit FMDV as carriers to naïve animals. In addition, the prevalence of carrier animals in a herd and factors influencing this prevalence—regardless of whether vaccination is employed—requires more research.^{38,39} The *NAHEMS Guidelines: Vaccination for Contagious Diseases, Appendix A: FMD* contains more information on the known epidemiological role and prevalence of carrier animals, impact of vaccination on carrier status, and detection of carriers via diagnostic testing.

³⁰ Alexandersen, S., et al. (2002). Aspects of the persistence of foot-and-mouth disease virus in animals—the carrier problem. *Microbes and Infection*, 4(10), 1099–1110. DOI: 10.1016/S1286-4579(02)01634-9.

³¹ Suttmoller, P. & Casas-Olascoaga, R. (2002). Unapparent foot and mouth disease infection (sub-clinical infections and carriers): implications for control. *Scientific and Technical Review of the Office International des Epizooties*, 21(3), 519–529. DOI: 10.20506/rst.21.3.1366.

³² Condy, J.B., et al. (1985). The duration of the foot-and-mouth disease virus carrier state in African buffalo (i) in the individual animal and (ii) in a free-living herd. *Comparative Immunology, Microbiology & Infectious Diseases*, 8(3–4), 259–265. DOI: 10.1016/0147-9571(85)90004-9.

³³ Anderson, E.C., et al. (1979). The pathogenesis of foot-and-mouth disease in the African buffalo (*Syncerus caffer*) and the role of this species in the epidemiology of the disease in Kenya. *Journal of Comparative Pathology*, 89(4), 541–549. DOI: 10.1016/0021-9975(79)90045-8.

³⁴ Kitching, R.P. (2002). Identification of foot and mouth disease virus carrier and subclinically infected animals and differentiation from vaccinated animals. *Scientific and Technical Review of the Office International des Epizooties*, 21(3), 531–538. DOI: 10.20506/rst.21.3.1365.

³⁵ Suttmoller, P. & Casas-Olascoaga, R. (2002). Unapparent foot and mouth disease infection (sub-clinical infections and carriers): implications for control. *Scientific and Technical Review of the Office International des Epizooties*, 21(3), 519–529. DOI: 10.20506/rst.21.3.1366.

³⁶ Mushayabasa, S. & Tapedzesa, G. (2015). Modeling the effects of multiple intervention strategies on controlling foot-and-mouth disease. *BioMed Research International*, 2015, 584234.

³⁷ Suttmoller, P. & Casas-Olascoaga, R. (2002). Unapparent foot and mouth disease infection (sub-clinical infections and carriers): implications for control. *Scientific and Technical Review of the Office International des Epizooties*, 21(3), 519–529. DOI: 10.20506/rst.21.3.1366.

³⁸ Tenzin, et al. (2008). Rate of foot-and-mouth disease virus transmission by carriers quantified from experimental data. *Risk Analysis*, 28(2), 303–309. DOI: 10.1111/j.1539-6924.2008.01020.x.

³⁹ Donaldson A.I. & Kitching, R.P. (1989). Transmission of foot-and-mouth disease by vaccinated cattle following natural challenge. *Research in Veterinary Science*, 46(1), 9–14.

1.4.4 Introduction and Transmission of FMDV

FMD is highly contagious. FMDV is typically introduced via contact with infected animals, their secretions, excretions, or fomites, or products contaminated with FMDV. FMD can also be introduced into a naïve animal population by feeding contaminated meat, milk, or garbage.^{40,41} Conveyances may be responsible for transmitting the disease between an infected and an uninfected premises.⁴² In past FMD outbreaks in both the United Kingdom (1967–68) and Denmark (1982), FMD spread was also epidemiologically linked to the routes of bulk milk tankers, though the precise transmission mechanism was not identified.⁴³ Insects and birds may also be mechanical vectors; no biological vector of the virus has been identified.⁴⁴

Cattle typically become infected through aerosolized virus.⁴⁵ Pigs usually become infected by eating virus-contaminated food, or through direct contact with the vesicular lesions of other animals.^{46,47} Pigs also can excrete large quantities of the virus through respiration, infecting susceptible animals.⁴⁸ As such, pigs are considered key amplifiers of the virus.⁴⁹

1.4.4.1 Live Animals and Virus Shedding

Animal to animal contact is a common mode of transmission. FMDV is shed in all secretions and excretions, including saliva, milk, semen, feces, urine, and ruptured vesicular fluid.⁵⁰ Pigs produce nearly 3,000-fold more of FMDV in respiratory secretions than either cattle or sheep.^{51,52} Table 1-1 provides the approximate amount of virus excretion for cattle and pigs to

⁴⁰ Musser, J.M. (2004). A practitioner's primer on foot-and-mouth disease. *Journal of the American Veterinary Medical Association*, 224(8), 1261–1268. DOI: 10.2460/javma.2004.224.1261.

⁴¹ Donaldson, A.I. & Alexandersen, S. (2002). Predicting the spread of foot and mouth disease by airborne virus. *Scientific and Technical Review of the Office International des Epizooties (Paris)*, 21(31), 569–575. DOI: 10.20506/rst.21.3.1362.

⁴² Alexandersen, S., et al. (2003). The pathogenesis and diagnosis of foot-and-mouth disease. *Journal of Comparative Pathology*, 129(1), 1–36. DOI: 10.1016/S0021-9975(03)00041-0.

⁴³ Donaldson, A.I. (1997). Risks of spreading foot and mouth disease through milk and dairy products. *Scientific and Technical Review of the Office International des Epizooties*, 16(1), 117–124. DOI: 10.20506/rst.16.1.1013.

⁴⁴ Hugh-Jones, M.E. & Wright, P.B. (1970). Studies on the 1967–8 foot-and-mouth disease epidemic: The relation of weather to the spread of disease. *Journal of Hygiene*, 68(2), 253–271. DOI: 10.1017/s0022172400028722.

⁴⁵ Kitching, R.P. & Hughes, G.J. (2002). Clinical variation in foot-and-mouth disease: cattle. *Scientific and Technical Review of the Office International des Epizooties*, 21(3), 499–504. DOI: 10.20506/rst.21.3.1343.

⁴⁶ Kitching, R.P. & Alexandersen, S. (2002). Clinical variation in foot and mouth disease: pigs. *Scientific and Technical Review of the Office International des Epizooties*, 21(3), 513–518. DOI: 10.20506/rst.21.3.1367.

⁴⁷ Alexandersen, S., et al. (2003). The pathogenesis and diagnosis of foot-and-mouth disease. *Journal of Comparative Pathology*, 129(1), 1–36. DOI: 10.1016/S0021-9975(03)00041-0.

⁴⁸ Alexandersen, S., et al. (2003). The pathogenesis and diagnosis of foot-and-mouth disease. *Journal of Comparative Pathology*, 129(1), 1–36. DOI: 10.1016/S0021-9975(03)00041-0.

⁴⁹ Amass, S., et al. (2003). Procedures for preventing the transmission of foot-and-mouth disease virus to pigs and sheep by personnel in contact with infected pigs. *Veterinary Record*, 153(5), 137–140. DOI: 10.1136/vr.153.5.137.

⁵⁰ Alexandersen, S., et al. (2003). The pathogenesis and diagnosis of foot-and-mouth disease. *Journal of Comparative Pathology*, 129(1), 1–36. DOI: 10.1016/S0021-9975(03)00041-0.

⁵¹ Thomson, G. R. (1996). The role of carrier animals in the transmission of foot and mouth disease. In *OIE comprehensive reports on technical items presented to the International Committee or to Regional Commissions* (p. 87–103). Retrieved from <http://www.oie.int/doc/ged/D3014.PDF>.

⁵² Donaldson, A.I. (1983). Quantitative data on airborne foot-and-mouth disease virus; its production, carriage, and deposition. In *Philosophical Transactions of the Royal Society of London, Series B*, 302(1111), 529–534. DOI: 10.1098/rstb.1983.0072.

indicate the significant difference in their levels of virus shedding. The amount of virus excreted by various species will also vary based on the serotype and strain of the FMDV.⁵³

Table 1-1. Approximate Virus Shedding

Species	Approximate Amount (per 24 hours)	Type of Excretion
Cattle ⁵⁴	Log ₁₀ 6.7 TCID ₅₀	Milk
Cattle ⁵⁵	Log ₁₀ 5.1 TCID ₅₀	Respiratory
Pigs ⁵⁶	From Log ₁₀ 5.8 to 7.6 TCID ₅₀	Respiratory
Sheep ⁵⁷	Log ₁₀ 4.3 TCID ₅₀	Respiratory

Note: TCID₅₀ = 50 percent tissue culture infectious dose.

Because cattle are more likely to be infected through inhalation, and pigs shed a significant amount of FMDV via respiration, highly concentrated herds of infected pigs in close proximity to naïve cattle herds poses a significant risk of transmission from pigs to cattle.⁵⁸

1.4.4.2 Air/Windborne Transmission

Airborne FMDV can result from a large number of infected pigs, resulting in plumes of aerosolized virus in the atmosphere.⁵⁹ Cattle, because they inhale more air and are easily infected through respiration, are the species frequently infected when FMDV is airborne.⁶⁰ Under specific climate conditions (particularly downwind), aerosolized FMDV produced by infected pigs can travel a significant distance infecting cattle from 20 kilometers ([km] up to 300 km away and infecting sheep from 10–100 km away), as was predicted with simulation models in the United Kingdom.^{61,62}

⁵³ Donaldson, A.I., et al. (1970). Further investigations on the airborne excretion of foot-and-mouth disease virus. *Journal of Hygiene*, 68(4), 557–564. DOI: 10.1017/s0022172400042480.

⁵⁴ Kitching, R.P. & Hughes, G.J. (2002). Clinical variation in foot-and-mouth disease: cattle. *Scientific and Technical Review of the Office International des Epizooties*, 21(3), 499–504. DOI: 10.20506/rst.21.3.1343.

⁵⁵ Kitching, R.P. & Hughes, G.J. (2002). Clinical variation in foot-and-mouth disease: cattle. *Scientific and Technical Review of the Office International des Epizooties*, 21(3), 499–504. DOI: 10.20506/rst.21.3.1343.

⁵⁶ Kitching, R.P. & Alexandersen, S. (2002). Clinical variation in foot and mouth disease: pigs. *Scientific and Technical Review of the Office International des Epizooties*, 21(3), 513–518. DOI: 10.20506/rst.21.3.1367.

⁵⁷ Alexandersen, S., et al. (2003). The pathogenesis and diagnosis of foot-and-mouth disease. *Journal of Comparative Pathology*, 129(1), 1–36. DOI: 10.1016/S0021-9975(03)00041-0.

⁵⁸ Donaldson, A.I. & Alexandersen, S. (2002). Predicting the spread of foot and mouth disease by airborne virus. *Scientific and Technical Review of the Office International des Epizooties (Paris)*, 21(31), 569–575. DOI: 10.20506/rst.21.3.1362.

⁵⁹ Morris, R.S., et al. (2002). Decision-support tools for foot and mouth disease control. *Scientific and Technical Review of the Office International des Epizooties*, 21(3), 557–567.

⁶⁰ Alexandersen, S., et al. (2003). The pathogenesis and diagnosis of foot-and-mouth disease. *Journal of Comparative Pathology*, 129(1), 1–36. DOI: 10.1016/S0021-9975(03)00041-0.

⁶¹ Alexandersen, S., et al. (2003). The pathogenesis and diagnosis of foot-and-mouth disease. *Journal of Comparative Pathology*, 129(1), 1–36. DOI: 10.1016/S0021-9975(03)00041-0.

⁶² Donaldson, A.I. & Alexandersen, S. (2002). Predicting the spread of foot and mouth disease by airborne virus. *Scientific and Technical Review of the Office International des Epizooties (Paris)*, 21(31), 569–575. DOI: 10.20506/rst.21.3.1362.

Many different factors influence how well FMD aerosolizes, and how far aerosolized virus may spread. As already mentioned, the species of the infected animal is significant, as pigs excrete more virus through respiration than cattle or sheep. In addition, the amount of virus emitted into the air will be impacted by the stage of the disease in the infected animal, as well as the number and concentration of infected animals, and the strain of the virus.⁶³ In terms of climatic conditions, a relative humidity of 55 percent or more, with low and steady winds, is favorable for FMDV spread via aerosol.^{64,65,66} The virus also seems to spread significantly better over water than over land.⁶⁷

1.4.4.3 Fomite Transmission

FMDV is readily transmitted through vehicles, equipment, boots, clothing, and other fomites. In an FMD outbreak, the movement of fomites is a critical transmission pathway which must be addressed, particularly because FMDV can persist on fomites for an extended period of time with persistence based on many factors, including decreased temperatures.⁶⁸

1.4.4.4 Personnel

It has been demonstrated that humans can harbor FMDV in their nasal passages for up to approximately 28 hours and throats for short periods of time as well.⁶⁹ Under experimental conditions, personnel did transmit infection to naïve animals.⁷⁰ As such, during FMD outbreaks, it has been directed that responders have a “down period” or waiting time in moving between premises to avoid the possibility of transmitting FMDV to susceptible, uninfected animals.^{71,72} This time period is determined in an incident by relevant regulatory officials. Evidence suggests

⁶³ Donaldson, A.I. & Alexandersen, S. (2002). Predicting the spread of foot and mouth disease by airborne virus. *Scientific and Technical Review of the Office International des Epizooties (Paris)*, 21(31), 569–575. DOI: 10.20506/rst.21.3.1362.

⁶⁴ Alexandersen, S., et al. (2003). The pathogenesis and diagnosis of foot-and-mouth disease. *Journal of Comparative Pathology*, 129(1), 1–36. DOI: 10.1016/S0021-9975(03)00041-0.

⁶⁵ Donaldson, A.I. & Alexandersen, S. (2002). Predicting the spread of foot and mouth disease by airborne virus. *Scientific and Technical Review of the Office International des Epizooties (Paris)*, 21(31), 569–575. DOI: 10.20506/rst.21.3.1362.

⁶⁶ Donaldson, A.I. (1972). The influence of relative humidity on the aerosol stability of different strains of foot and mouth disease virus suspended in saliva. *Journal of General Virology*, 15(1), 25–33. DOI: 10.1099/0022-1317-15-1-25.

⁶⁷ Morris, R.S., et al. (2002). Decision-support tools for foot and mouth disease control. *Scientific and Technical Review of the Office International des Epizooties*, 21(3), 557–567.

⁶⁸ Cottral, G.E. (1969). Persistence of foot-and-mouth disease virus in animals, their products and the environment. *Scientific and Technical Review of the Office International des Epizooties*, 71(3–4), 549–568.

⁶⁹ Sellers, R.F., et al. (1970). Inhalation, persistence and dispersal of foot-and-mouth disease virus by man. *Journal of Hygiene (London)*, 68(4), 565–573. DOI: 10.1017/s0022172400042492.

⁷⁰ Sellers, R.F., et al. (1970). (1971). Transfer of foot-and-mouth disease virus in the nose of man from infected to non-infected animals. *Veterinary Record*, 89(16), 447–449. DOI: 10.1136/vr.89.16.447-a.

⁷¹ Amass, S., et al. (2003). Procedures for preventing the transmission of foot-and-mouth disease virus to pigs and sheep by personnel in contact with infected pigs. *Veterinary Record*, 153(5), 137–140. DOI: 10.1136/vr.153.5.137.

⁷² Amass, S., et al. (2004). Procedures for preventing transmission of foot-and-mouth disease virus (O/TAW/97) by people. *Veterinary Microbiology*, 103(3–4), 143–149. DOI: 10.1016/j.vetmic.2004.07.020.

that with appropriate biosecurity and cleaning and disinfection measures, the necessity for extended down periods is decreased.^{73,74,75}

1.4.4.5 Wildlife

It is unclear what role wildlife would play in disease transmission if there was an FMD outbreak in the United States, Red, fallow, and roe deer were all susceptible to FMDV when exposed in the laboratory, though severity of clinical signs varied.⁷⁶ However, in the 2001 epidemic in the United Kingdom, evidence suggests deer were not epidemiologically important in the spread of FMD.^{77,78} Transmission of FMDV from feral swine may also be a possibility in a U.S. FMD outbreak.⁷⁹

1.4.5 Incubation Period

The incubation period for FMD is typically 2–14 days, and is defined by the World Organization for Animal Health (OIE) as 14 days for the purpose of the *OIE Terrestrial Animal Health Code* (2016).^{80,81} During the beginning phases in the prevalence of FMDV, the incubation period may be as short as 24 hours.⁸² How fast clinical signs appear depends on the dose of the virus, species of the animal, as well as the route of infection.^{83,84} Animals may shed FMDV before the appearance of clinical signs.^{85,86}

⁷³ Amass, S., et al. (2003). Procedures for preventing the transmission of foot-and-mouth disease virus to pigs and sheep by personnel in contact with infected pigs. *Veterinary Record*, 153(5), 137–140. DOI: 10.1136/vr.153.5.137.

⁷⁴ Donaldson, A.I. (2008). Foot-and-mouth disease and the use of face masks. *Veterinary Record*, 162(5), 163. DOI: 10.1136/vr.162.5.163.

⁷⁵ Amass, S., et al. (2003). Procedures for preventing the transmission of foot-and-mouth disease virus to pigs and sheep by personnel in contact with infected pigs. *Veterinary Record*, 153(5), 137–140. DOI: 10.1136/vr.153.5.137.

⁷⁶ Forman, A.J. & Gibbs, E.P.J. (1974). Studies with foot-and-mouth disease virus in British deer (red, fallow, and roe): 1. Clinical Disease. *Journal of Comparative Pathology*, 84(2), 215–220. DOI: 10.1016/0021-9975(74)90062-0.

⁷⁷ McVicar, J.W., et al. (1974). Foot-and-mouth disease in white tailed deer: clinical signs and transmission in the laboratory. In *Proceedings of the 78th annual meeting of the United States Animal Health Association* (p. 169–180).

⁷⁸ Davies, G. (2002). The foot and mouth disease epidemic in the United Kingdom, 2001. *Comparative Immunology, Microbiology & Infectious Diseases*, 25(5–6), 331–343. DOI: 10.1016/S0147-9571(02)00030-9.

⁷⁹ Mohamed, F., et al. (2011). Foot-and-mouth disease in feral swine: Susceptibility and transmission. *Transboundary Emerging Diseases*, 58(4), 358–371. DOI: 10.1111/j.1865-1682.2011.01213.x.

⁸⁰ OIE. (2013). Foot and Mouth Disease. In *Technical Disease Card*. Retrieved from <http://www.oie.int>.

⁸¹ OIE. (2016). Foot-and-Mouth Disease. In *Terrestrial Animal Health Code* (8.8.). Retrieved from <http://www.oie.int>.

⁸² Yang, P.C., Chu, R.M., Chung, W.B., & Sung, H.T. (1999). Epidemiological characteristics and financial costs of the 1997 foot-and-mouth disease epidemic in Taiwan. *Veterinary Record*, 145(25), 731–734. DOI: 10.1136/vr.145.25.731.

⁸³ Kitching, R.P. & Hughes, G.J. (2002). Clinical variation in foot and mouth disease: sheep and goats. *Scientific and Technical Review of the Office International des Epizooties*, 21(3), 505–512. DOI: 10.20506/rst.21.3.1342.

⁸⁴ Kitching, R.P. & Alexandersen, S. (2002). Clinical variation in foot and mouth disease: pigs. *Scientific and Technical Review of the Office International des Epizooties*, 21(3), 513–518. DOI: 10.20506/rst.21.3.1367.

⁸⁵ Gay, C.G., et al. (2010). Foot-and-Mouth Disease (FMD): Gap Analysis Workshop Report. In *Agricultural Research Service*. Retrieved from <http://go.usa.gov/kCqF>.

⁸⁶ Orsel, K., et al. (2009) Foot and mouth disease virus transmission during the incubation period of the disease in piglets, lambs, calves, and dairy cows. *Preventive Veterinary Medicine*, 88(2), 158–163. DOI: 10.1016/j.prevetmed.2008.09.001.

1.4.5.1 Infectious Dose

Different animal species vary in their susceptibility to FMDV, which is also influenced by the route of infection. Table 1-2 lists the infectious dose and mode of infection for key animals, given the primary mode of transmission for that species.

Table 1-2. Infectious Dose and Mode of Transmission

Species	Infectious Dose	Common Mode
Cattle ⁸⁷	May be as low as 20 TCID ₅₀	Inhalation
Sheep and goats ⁸⁸	May be as low as 20 TCID ₅₀	Direct contact
Pigs ⁸⁹	Approximately log10 ⁵ TCID ₅₀	Ingestion

Note: TCID₅₀ = 50 percent tissue culture infectious dose.

1.4.6 Morbidity and Mortality

The morbidity and mortality of FMD varies depending on the species affected, as well as the serotype and strain of the virus. Morbidity is significant, and can approach 100 percent.⁹⁰ Mortality is typically low in adult animals (1 to 5 percent), though higher mortality rates are typically observed in very young animals, usually from acute myocarditis.⁹¹

1.4.6.1 Clinical Signs

FMDV is typically characterized by high morbidity, evidenced by characteristic vesicles on the oral and nasal mucosa, teats, coronary bands, and interdigital spaces. However, before vesicles appear and a decreased appetite progresses fever develops.⁹² The clinical signs can vary based on the serotype and strain of the FMDV. Generally speaking, sheep and goats typically have milder clinical signs than cattle.⁹³ The following sections provide more detail on the clinical signs in cattle, pigs, and sheep and goats.

1.4.6.1.1 Cattle

Cattle usually present with fever, anorexia, shivering, and reduction in milk production for approximately 2–3 days, before vesicular lesions are observable on mucous membranes, interdigital spaces, and on the coronary band. The vesicles will rupture in approximately 1 day,

⁸⁷ Kitching, R.P. & Hughes, G.J. (2002). Clinical variation in foot-and-mouth disease: cattle. *Scientific and Technical Review of the Office International des Epizooties*, 21(3), 499–504. DOI: 10.20506/rst.21.3.1343.

⁸⁸ Kitching, R.P. & Hughes, G.J. (2002). Clinical variation in foot and mouth disease: sheep and goats. *Scientific and Technical Review of the Office International des Epizooties*, 21(3), 505–512. DOI: 10.20506/rst.21.3.1342.

⁸⁹ Kitching, R.P. & Alexandersen, S. (2002). Clinical variation in foot and mouth disease: pigs. *Scientific and Technical Review of the Office International des Epizooties*, 21(3), 513–518. DOI: 10.20506/rst.21.3.1367.

⁹⁰ OIE. (2013). Foot and Mouth Disease. In *Technical Disease Card*. Retrieved from <http://www.oie.int>.

⁹¹ Musser, J.M. (2004). A practitioner's primer on foot-and-mouth disease. *Journal of the American Veterinary Medical Association*, 224(8), 1261–1268. DOI: 10.2460/javma.2004.224.1261.

⁹² Musser, J.M. (2004). A practitioner's primer on foot-and-mouth disease. *Journal of the American Veterinary Medical Association*, 224(8), 1261–1268. DOI: 10.2460/javma.2004.224.1261.

⁹³ Musser, J.M. (2004). A practitioner's primer on foot-and-mouth disease. *Journal of the American Veterinary Medical Association*, 224(8), 1261–1268. DOI: 10.2460/javma.2004.224.1261.

and recovery occurs in 8–15 days. Excessive salivation is often observed in cattle and milk yields are reduced by 80 percent.^{94,95,96}

1.4.6.1.2 Pigs

In pigs, severe lesions typically are observed on the feet, as well as on the snout, udder, as well as hock and elbow. Excessive salivation is less likely in pigs than in cattle, and lesions in the mouth are milder than those observed elsewhere.^{97,98}

1.4.6.1.3 Sheep and Goats

Fewer clinical signs are seen in sheep and goats. Mouth lesions are often not obvious, though lesions can develop on heel bulbs and on the coronary band.⁹⁹ Sheep and goats may be important in transmission, as infection presents with mild clinical signs and may not be as immediately recognized.¹⁰⁰

1.5 Environmental Persistence of FMDV

FMDV is moderately stable in the environment.¹⁰¹ The FMD virus is susceptible to both acid and alkaline pH. However, under certain conditions, it can maintain infectivity in the environment for extended periods of time. Presence of organic matter increases its persistence.

Table 1-3. Resistance of FMDV to Physical and Chemical Action

Action	Resistance
Temperature	Preserved by refrigeration and freezing. Progressively inactivated by temperatures above 50 °C. Heating meat to a minimum core temperature of 70 °C for at least 30 minutes inactivates the virus.
pH	Quickly inactivated by pH <6.0 or >9.0.
Disinfectants	Inactivated by sodium hydroxide (2%), sodium carbonate (4%), citric acid (0.2%), acetic acid (2%), sodium hypochlorite (3%), potassium peroxymonosulfate/sodium chloride (1%), and chlorine dioxide. Resistant to iodophores, quaternary ammonium compounds, and phenol, especially in the presence of organic matter.

⁹⁴ OIE. (2013). Foot and Mouth Disease. In *Technical Disease Card*. Retrieved from <http://www.oie.int>.

⁹⁵ Musser, J.M. (2004). A practitioner's primer on foot-and-mouth disease. *Journal of the American Veterinary Medical Association*, 224(8), 1261–1268. DOI: 10.2460/javma.2004.224.1261.

⁹⁶ Knight-Jones, T. & Rushton, J. (2013). The economic impacts of foot and mouth disease—What are they, how big are they and where do they occur? *Preventive Veterinary Medicine*, 112 (3–4), 161–173. DOI: 10.1016/j.prevetmed.2013.07.013.

⁹⁷ Musser, J.M. (2004). A practitioner's primer on foot-and-mouth disease. *Journal of the American Veterinary Medical Association*, 224(8), 1261–1268. DOI: 10.2460/javma.2004.224.1261.

⁹⁸ Kitching, R.P. & Alexandersen, S. (2002). Clinical variation in foot and mouth disease: pigs. *Scientific and Technical Review of the Office International des Epizooties*, 21(3), 513–518. DOI: 10.20506/rst.21.3.1367.

⁹⁹ Kitching, R.P. & Hughes, G.J. (2002). Clinical variation in foot and mouth disease: sheep and goats. *Scientific and Technical Review of the Office International des Epizooties*, 21(3), 505–512. DOI: 10.20506/rst.21.3.1342.

¹⁰⁰ Alexandersen, S., et al. (2003). The pathogenesis and diagnosis of foot-and-mouth disease. *Journal of Comparative Pathology*, 129(1), 1–36. DOI: 10.1016/S0021-9975(03)00041-0.

¹⁰¹ Alexandersen, S., et al. (2003). The pathogenesis and diagnosis of foot-and-mouth disease. *Journal of Comparative Pathology*, 129(1), 1–36. DOI: 10.1016/S0021-9975(03)00041-0.

Action	Resistance
Survival	Survives in lymph nodes and bone marrow at neutral pH, but destroyed in muscle at pH <6.0, i.e., after rigor mortis. Survives in frozen bone marrow or lymph nodes. Residual virus survives in milk and milk products during regular pasteurization, but is inactivated by ultra-high temperature pasteurization. Survives drying but may persist for days to weeks in organic matter under moist and cool temperatures. Can persist in contaminated fodder and the environment for up to 1 month, depending on the temperature and pH conditions.

Source: OIE Technical Disease Card for FMD, 2013.

Sunlight does not have a significant effect on FMDV infectivity.¹⁰² FMDV can be found in the bone marrow, lymph nodes, and certain organs of deceased animals for extended periods of time because the pH does not decline sufficiently after death.¹⁰³

1.5.1 Environmental Persistence of FMDV in Excretions and Surroundings

Under certain environmental conditions, FMDV can survive for significant periods on organic material. Reported survival times of FMDV include up to 20 weeks on hay, 4 weeks in cow's hair in temperate temperatures, 14 days in dry feces, 39 days in urine, 6 months in slurry (winter), and 28 days in soil (fall).¹⁰⁴ FMDV has also been recovered from bovine semen, and can maintain virus titers if frozen at -50 °C for up to 320 days.¹⁰⁵

1.5.2 Environmental Persistence of FMDV in Milk and Dairy Products

Raw milk and milk products have the potential to cause infection in animals if not properly treated. Even with a typical pasteurization process, milk and milk products can still be infective to naïve animals.¹⁰⁶ Typical pasteurization processes do not activate FMDV.

In milk, experiments have demonstrated that FMDV¹⁰⁷

- survived in skim milk after heating to 72 °C for 2 minutes,
- survived in whole milk after heating to 72 °C for 5 minutes, and
- survived in cream after heating to 93 °C for 15 seconds.

¹⁰² Alexandersen, S., et al. (2003). The pathogenesis and diagnosis of foot-and-mouth disease. *Journal of Comparative Pathology*, 129(1), 1–36. DOI: 10.1016/S0021-9975(03)00041-0.

¹⁰³ Alexandersen, S., et al. (2003). The pathogenesis and diagnosis of foot-and-mouth disease. *Journal of Comparative Pathology*, 129(1), 1–36. DOI: 10.1016/S0021-9975(03)00041-0.

¹⁰⁴ Alexandersen, S., et al. (2003). The pathogenesis and diagnosis of foot-and-mouth disease. *Journal of Comparative Pathology*, 129(1), 1–36. DOI: 10.1016/S0021-9975(03)00041-0.

¹⁰⁵ Cottral, G.E. (1969). Persistence of foot-and-mouth disease virus in animals, their products and the environment. *Scientific and Technical Review of the Office International des Epizooties*, 71(3–4), 549–568.

¹⁰⁶ Tomasula, P.M. & Konstance, R.P. (2004). The survival of foot-and-mouth disease virus in raw and pasteurized milk and milk products. *Journal of Dairy Science*, 87(4), 1115–1121. DOI: 10.3168/jds.S0022-0302(04)73258-0.

¹⁰⁷ Blackwell, J.H. & Hyde, J.L. (1976). Effect of heat on foot-and-mouth disease virus (FMDV) in the components of milk from FMDV-infected cows. *Journal of Hygiene*, 77(1), 77–83. DOI: 10.1017/s0022172400055534.

Additionally, here are some examples of the experimental resistance of FMDV in cheese:¹⁰⁸

- Survived in Camembert cheese for 21 days at 2 °C.
- Survived processing for Cheddar cheese prepared from heated milk (but 30 days of curing inactivated the virus).
- Survived processing and curing for Cheddar cheese prepared from unheated milk for 60 days.

FMDV can also survive in other products, such as butter and butter oil, for extended periods (at least 45 days with storage at 4 °C).¹⁰⁹ However, FMDV has not been detected in whey constituents after processing.¹¹⁰

1.5.3 Environmental Persistence of FMDV in Meat and Meat Products

FMDV can maintain infectivity for months in chilled or frozen lymph nodes (beef), liver, and blood. FMDV was detected at 11 days in beef frozen after slaughter and at 4 months in frozen liver.¹¹¹ In salted bacon, FMDV was detected at 190 days and in pork sausages at 56 days.¹¹²

1.5.4 Environmental Persistence of FMDV in Wool and Hides

FMDV can survive on many materials. FMDV has been identified in wool 5-days post-exposure, but was not recovered at measureable amounts from wool stored at 37 °C after 40–96 hours, or wool stored at 18 °C after 10–14 days, based on the strain.¹¹³ In hides, FMDV has been detected for 32 days or longer, depending on the humidity and temperature of drying and storage. For example, hides salted and stored at 4 °C still had detectable FMDV at 352 days; hides dried at 20 °C, 40 percent relative humidity had detectable FMDV at 42 days.¹¹⁴

1.6 OIE Procedures for the Inactivation of FMDV

The following sections detail OIE recommended inactivation procedures for FMDV in various animal products. For details on products listed here, please see the *OIE Terrestrial Animal Health Code* (2016).¹¹⁵

¹⁰⁸ Blackwell, J.H. (1976). Survival of Foot-and-Mouth Disease Virus in Cheese. *Journal of Dairy Science*, 59(9), 1574–1579. DOI: 10.3168/jds.S0022-0302(76)84407-4.

¹⁰⁹ Blackwell, J.H. (1978). Persistence of foot-and-mouth disease virus in butter and butter oil. *Journal of Dairy Science*, 45(2), 283–285. DOI: 10.1017/S0022029900016460.

¹¹⁰ Blackwell, J.H. (1978). Potential transmission of foot-and-mouth disease in whey constituents. *Journal of Food Protection*, 41(8), 631–633. Retrieved from <http://www.ingentaconnect.com/content/iafp/jfp/1978/00000041/00000008/art00006>.

¹¹¹ Ryan, E., et al. (2008). Foot-and-mouth disease virus concentrations in products of animal origin. *Transboundary Emerging Diseases*, 55(2), 89–98. DOI: 10.1111/j.1865-1682.2007.01004.x.

¹¹² Ryan, E., et al. (2008). Foot-and-mouth disease virus concentrations in products of animal origin. *Transboundary Emerging Diseases*, 55(2), 89–98. DOI: 10.1111/j.1865-1682.2007.01004.x.

¹¹³ McColl, K.A., et al. (1995). The persistence of foot-and-mouth disease virus on wool. *The Australian Veterinary Journal*, 72(8), 286–292. DOI: 10.1111/j.1751-0813.1995.tb03556.x.

¹¹⁴ Gailiunas, P. & Cottral, G.E. (1967). Survival of foot and mouth disease virus in bovine hides. *American Journal of Veterinary Research*, 28(125), 1047–1053.

¹¹⁵ OIE. (2016). Foot-and-Mouth Disease. In *Terrestrial Animal Health Code* (8.8.). Retrieved from <http://www.oie.int>.

1.6.1 Meat

Article 8.8.31 in the 2016 *OIE Terrestrial Animal Health Code* provides the following procedures for the inactivation of FMDV in meat and meat products:

1. Canning

Meat and meat products are subjected to heat treatment in a hermetically sealed container to reach an internal core temperature of at least 70 °C for a minimum of 30 minutes or to any equivalent treatment which has been demonstrated to inactivate FMDV.

2. Thorough cooking

Meat, previously deboned and defatted, and meat products are subjected to heat treatment so that an internal temperature of 70 °C for a minimum of 30 minutes.

After cooking, they should be packed and handled in such a way that it cannot be exposed to a source of FMDV.

3. Drying after salting

When *rigor mortis* is complete, the meat must be deboned, salted with cooking salt (NaCl) and 'completely dried'. It must not deteriorate at ambient temperature.

'Completely dried' is defined in terms of the ration between water and protein which must not be greater than 2.25:1 or a water activity (A_w) that is not greater than 0.85.

1.6.2 Wool and Hair

Article 8.8.32 in the 2016 *OIE Terrestrial Animal Health Code* provides the following procedures for the inactivation of FMDV in wool and hair:

For the inactivation of viruses present in wool and hair for industrial use, one of the following procedures should be used:

1. industrial washing, which consists of the immersion of the wool in a series of baths of water, soap, and sodium hydroxide (soda) or potassium hydroxide (potash);
2. chemical depilation by means of slaked lime or sodium sulphide;
3. fumigation in formaldehyde in a hermetically sealed chamber for at least 24 hours;
4. industrial scouring which consists of the immersion of wool in a water-soluble detergent held at 60–70 °C;
5. storage of wool at 18 °C for four weeks, or 4 °C for four months or 37 °C for eight days.

1.6.3 Milk and Cream for Human Consumption

Article 8.8.35 in the 2016 *OIE Terrestrial Animal Health Code* provides the following procedures for the inactivation of FMDV in milk and cream for human consumption:

For the inactivation of FMDV present in milk and cream for human consumption, one of the following procedures should be used:

1. a process applying a minimum temperature of 132 °C for at least one second (ultra-high temperature [UHT]), or
2. if the milk has a pH less than 7.0, a process applying a minimum temperature of 72 °C for at least 15 seconds (high temperature-short time pasteurization [HTST]), or
3. if the milk has a pH of 7.0 or over, the HTST process applied twice.

1.6.4 Milk for Animal Consumption

Article 8.8.36 in the 2016 *OIE Terrestrial Animal Health Code* provides the following procedures for the inactivation of FMDV in milk for animal consumption:

For the inactivation of FMDV present in milk for animal consumption, one of the following procedures should be used:

1. the HTST process applied twice; or
2. HTST combined with another physical treatment, e.g. maintaining a pH 6 for at least one hour or additional heating to at least 72 °C combined with desiccation; or
3. UHT combined with another physical treatment referred to in point 2 above.

Attachment 1.A References and Selected Resources

- Adams, R.L.P., Knowler, J., & Leader, D. (1992). The structure of viruses. *The biochemistry of the nucleic acids* (p.75–76). United Kingdom: Chapman & Hall. DOI: 10.1007/978-94-011-2290-0.
- Alexandersen, S., Zhang, Z., & Donaldson, A.I. (2002). Aspects of the persistence of foot-and-mouth disease virus in animals—the carrier problem. *Microbes and Infection*, 4(10), 1099–1110. DOI: 10.1016/S1286-4579(02)01634-9.
- Alexandersen, S., Zhang, Z., Donaldson, A.I., & Garland, A.J. (2003). The pathogenesis and diagnosis of foot-and-mouth disease. *Journal of Comparative Pathology*, 129(1), 1–36. DOI: 10.1016/S0021-9975(03)00041-0.
- Amass, S., Pacheco, J., Mason, P., Pacheco, J., Miller, C., Ramirez, A., Clark, L., Ragland, D., Schneider, J., & Kenyon, S. (2004). Procedures for preventing transmission of foot-and-mouth disease virus (O/TAW/97) by people. *Veterinary Microbiology*, 103(3–4), 143–149. DOI: 10.1016/j.vetmic.2004.07.020.
- Amass, S., Pacheco, J., Mason, P., Schneider, J., Alvarez, R., Clark, L., & Ragland, D. (2003). Procedures for preventing the transmission of foot-and-mouth disease virus to pigs and sheep by personnel in contact with infected pigs. *Veterinary Record*, 153(5), 137–140. DOI: 10.1136/vr.153.5.137.
- Anderson, E.C., Doughty, W.J., Anderson, J., & Paling, R. (1979). The pathogenesis of foot-and-mouth disease in the African buffalo (*Syncerus caffer*) and the role of this species in the epidemiology of the disease in Kenya. *Journal of Comparative Pathology*, 89(4), 541–549. DOI: 10.1016/0021-9975(79)90045-8.
- Belsham, G. & Bøtner, A. (2015). Use of recombinant capsid proteins in the development of a vaccine against foot-and-mouth disease virus (FMDV). *Virus Adaptation and Treatment*, 7, 11–23. DOI: 10.2147/VAAT.S55351.
- Blackwell, J.H. (1976). Survival of Foot-and-Mouth Disease Virus in Cheese. *Journal of Dairy Science*, 59(9), 1574–1579. DOI: 10.3168/jds.S0022-0302(76)84407-4.
- Blackwell, J.H. (1978). Persistence of foot-and-mouth disease virus in butter and butter oil. *Journal of Dairy Science*, 45(2), 283–285. DOI: 10.1017/S0022029900016460.
- Blackwell, J.H. (1978). Potential transmission of foot-and-mouth disease in whey constituents. *Journal of Food Protection*, 41(8), 631–633. Retrieved from <http://www.ingentaconnect.com/content/iafp/jfp/1978/00000041/00000008/art00006>.
- Blackwell, J.H. & Hyde, J.L. (1976). Effect of heat on foot-and-mouth disease virus (FMDV) in the components of milk from FMDV-infected cows. *Journal of Hygiene*, 77(1), 77–83. DOI: 10.1017/s0022172400055534.

-
- Center for Food Security and Public Health (CFSPH). (2014). Foot-and-Mouth Disease. In *Technical Factsheet*. Retrieved from www.cfsph.iastate.edu.
- Condy, J.B., Hedger, R.S., Hamblin, C., & Barnett, I.T. (1985). The duration of the foot-and-mouth disease virus carrier state in African buffalo (i) in the individual animal and (ii) in a free-living herd. *Comparative Immunology, Microbiology & Infectious Diseases*, 8(3–4), 259–265. DOI: 10.1016/0147-9571(85)90004-9.
- Cottral, G.E. (1969). Persistence of foot-and-mouth disease virus in animals, their products and the environment. *Scientific and Technical Review of the Office International des Epizooties*, 71(3–4), 549–568.
- Davies, G. (2002). The foot and mouth disease epidemic in the United Kingdom, 2001. *Comparative Immunology, Microbiology & Infectious Diseases*, 25(5–6), 331–343. DOI: 10.1016/S0147-9571(02)00030-9.
- Donaldson, A.I. (1972). The influence of relative humidity on the aerosol stability of different strains of foot and mouth disease virus suspended in saliva. *Journal of General Virology*, 15(1), 25–33. DOI: 10.1099/0022-1317-15-1-25.
- Donaldson, A.I. (1983). Quantitative data on airborne foot-and-mouth disease virus; its production, carriage, and deposition. In *Philosophical Transactions of the Royal Society of London, Series B*, 302(1111), 529–534. DOI: 10.1098/rstb.1983.0072.
- Donaldson, A.I. (1997). Risks of spreading foot and mouth disease through milk and dairy products. *Scientific and Technical Review of the Office International des Epizooties*, 16(1), 117–124. DOI: 10.20506/rst.16.1.1013.
- Donaldson, A.I. (2008). Foot-and-mouth disease and the use of face masks. *Veterinary Record*, 162(5), 163. DOI: 10.1136/vr.162.5.163.
- Donaldson, A.I. & Alexandersen, S. (2002). Predicting the spread of foot and mouth disease by airborne virus. *Scientific and Technical Review of the Office International des Epizooties (Paris)*, 21(31), 569–575. DOI: 10.20506/rst.21.3.1362.
- Donaldson, A.I., Herniman, K.A., Parker, J., & Sellers, R.F. (1970). Further investigations on the airborne excretion of foot-and-mouth disease virus. *Journal of Hygiene*, 68(4), 557–564. DOI: 10.1017/s0022172400042480.
- Donaldson A.I. & Kitching, R.P. (1989). Transmission of foot-and-mouth disease by vaccinated cattle following natural challenge. *Research in Veterinary Science*, 46(1), 9–14.
- Forman, A.J. & Gibbs, E.P.J. (1974). Studies with foot-and-mouth disease virus in British deer (red, fallow, and roe): 1. Clinical Disease. *Journal of Comparative Pathology*, 84(2), 215–220. DOI: 10.1016/0021-9975(74)90062-0.
- Gailiunas, P. & Cottral, G.E. (1967). Survival of foot and mouth disease virus in bovine hides. *American Journal of Veterinary Research*, 28(125), 1047–1053.

-
- Gay, C.G., Borca, M., Capozzo, A., De Clercq, K., Elsken, L., Espinoza, A., Hammond, H., Konig, G., Leon, E., Mattion, N., Metwally, S., Perez-Filgueira, M., Perez, A., Rieder, E., Smitsaart, E., Vosloo, W., Art, J., Brake, D., Duque, H., Golde, W., Grubman, M., Krug, P., Lombard, M., McIntosh, M., & Rodriguez, L. (2010). Foot-and-Mouth Disease (FMD): Gap Analysis Workshop Report. In *Agricultural Research Service*. Retrieved from <http://go.usa.gov/kCqF>.
- Goodwin, S., Tuthill, T.J., Arias, A., Killington, R.A., & Rowlands, D.J. (2009). Foot-and-mouth disease virus assembly: processing of recombinant capsid precursor by exogenous protease induces self-assembly of pentamers in vitro in a myristoylation-dependent manner. *Journal of Virology*, 83(21), 11275–11282. DOI: 10.1128/JVI.01263-09.
- Hugh-Jones, M.E. & Wright, P.B. (1970). Studies on the 1967–8 foot-and-mouth disease epidemic: The relation of weather to the spread of disease. *Journal of Hygiene*, 68(2), 253–271. DOI: 10.1017/s0022172400028722.
- International Committee on Taxonomy of Viruses. (2015). Foot-and-Mouth Disease. Retrieved from <http://ictvonline.org>.
- Kitching, R.P. (2002). Identification of foot and mouth disease virus carrier and subclinically infected animals and differentiation from vaccinated animals. *Scientific and Technical Review of the Office International des Epizooties*, 21(3), 531–538. DOI: 10.20506/rst.21.3.1365.
- Kitching, R.P. & Alexandersen, S. (2002). Clinical variation in foot and mouth disease: pigs. *Scientific and Technical Review of the Office International des Epizooties*, 21(3), 513–518. DOI: 10.20506/rst.21.3.1367.
- Kitching, R.P. & Hughes, G.J. (2002). Clinical variation in foot-and-mouth disease: cattle. *Scientific and Technical Review of the Office International des Epizooties*, 21(3), 499–504. DOI: 10.20506/rst.21.3.1343.
- Kitching, R.P. & Hughes, G.J. (2002). Clinical variation in foot and mouth disease: sheep and goats. *Scientific and Technical Review of the Office International des Epizooties*, 21(3), 505–512. DOI: 10.20506/rst.21.3.1342.
- Knight-Jones, T. & Rushton, J. (2013). The economic impacts of foot and mouth disease—What are they, how big are they and where do they occur? *Preventive Veterinary Medicine*, 112(3–4), 161–173. DOI: 10.1016/j.prevetmed.2013.07.013.
- Knowles, N.J., Samuel, A.R., Davies, P.R., Midgley, R.J., & Valarcher, J.F. (2005). Pandemic strain of foot-and-mouth disease virus serotype O. *Emerging Infectious Diseases*, 11(12), 1887–1893. DOI: 10.32001/eid1112.050908.
- Lubroth, J., Yedloutschnig, R.J., Culhane, V.K., & Mikiciuk, P.E. (1990). Foot-and-mouth disease virus in the llama (*Lama glama*): diagnosis, transmission, and susceptibility. *Journal of Veterinary Diagnostic Investigation*, 2(3), 197–203. DOI: 10.1177/104063879000200308.

-
- McColl, K.A., Westbury, H.A., Kitching, R.P., & Lewis, V.M. (1995). The persistence of foot-and-mouth disease virus on wool. *The Australian Veterinary Journal*, 72(8), 286–292. DOI: 10.1111/j.1751-0813.1995.tb03556.x.
- McVicar, J.W., Suttmoller, P., Ferris, D.H., & Campbell, C.H. (1974). Foot-and-mouth disease in white tailed deer: clinical signs and transmission in the laboratory. In *Proceedings of the 78th annual meeting of the United States Animal Health Association* (p. 169–180).
- Mohamed, F., Swafford, S., Petrowski, H., Bracht, A., Schmit, B., Fabian, A., Pacheco, J.M., Hartwig, E., Berninger, M., Carrillo, C., Mayr, G., Moran, K., Kavanaugh, D., Leibrecht, H., White, W., Metwally, S. (2011). Foot-and-mouth disease in feral swine: Susceptibility and transmission. *Transboundary Emerging Diseases*, 58(4), 358–371. DOI: 10.1111/j.1865-1682.2011.01213.x.
- Morris, R.S., Sanson, R.L., Stern, M.W., Stevenson, M., Wilesmith, J.W. (2002). Decision-support tools for foot and mouth disease control. *Scientific and Technical Review of the Office International des Epizooties*, 21(3), 557–567.
- Mushayabasa, S. & Tapedzesa, G. (2015). Modeling the effects of multiple intervention strategies on controlling foot-and-mouth disease. *BioMed Research International*, 2015, 584234.
- Musser, J.M. (2004). A practitioner’s primer on foot-and-mouth disease. *Journal of the American Veterinary Medical Association*, 224(8), 1261–1268. DOI: 10.2460/javma.2004.224.1261.
- Orsel, K., Bouma, A., Dekker, A., Stegeman, J.A., & de Jong, M.C. (2009) Foot and mouth disease virus transmission during the incubation period of the disease in piglets, lambs, calves, and dairy cows. *Preventive Veterinary Medicine*, 88(2), 158–163. DOI: 10.1016/j.prevetmed.2008.09.001.
- Ryan, E., Mackay, D., & Donaldson, A. (2008). Foot-and-mouth disease virus concentrations in products of animal origin. *Transboundary Emerging Diseases*, 55(2), 89–98. DOI: 10.1111/j.1865-1682.2007.01004.x.
- Sellers, R.F., Donaldson, A.I., & Herniman, K.A. (1970). Inhalation, persistence and dispersal of foot-and-mouth disease virus by man. *Journal of Hygiene (London)*, 68(4), 565–573. DOI: 10.1017/s0022172400042492.
- Sellers, R.F., Herniman, K.A., & Mann, J.A. (1971). Transfer of foot-and-mouth disease virus in the nose of man from infected to non-infected animals. *Veterinary Record*, 89(16), 447–449. DOI: 10.1136/vr.89.16.447-a.
- Shors, T. (2013). Baltimore classification. In *Understanding viruses* (56). Burlington, MA: Jones & Bartlett Learning.

-
- Sutmoller, P. & Casas-Olascoga, R. (2002). Unapparent foot and mouth disease infection (sub-clinical infections and carriers): implications for control. *Scientific and Technical Review of the Office International des Epizooties*, 21(3), 519–529. DOI: 10.20506/rst.21.3.1366.
- Sutmoller, P., McVicar, J.W., & Cottral, G.E. (1968). The epizootiological importance of foot-and-mouth disease carriers. I: Experimentally produced foot-and-mouth disease carriers in susceptible and immune carriers. *Archives of Virology*, 23(3), 227–235. DOI: 10.1007/BF01241895.
- Tenzin, Dekker, A., Vernooij, H., Bouma, A., & Stegeman, A. (2008). Rate of foot-and-mouth disease virus transmission by carriers quantified from experimental data. *Risk Analysis*, 28(2), 303–309. DOI: 10.1111/j.1539-6924.2008.01020.x.
- Thomson, G. R. (1996). The role of carrier animals in the transmission of foot and mouth disease. In *OIE comprehensive reports on technical items presented to the International Committee or to Regional Commissions* (p. 87–103). Retrieved from <http://www.oie.int/doc/ged/D3014.PDF>.
- Tomasula, P.M. & Konstance, R.P. (2004). The survival of foot-and-mouth disease virus in raw and pasteurized milk and milk products. *Journal of Dairy Science*, 87(4), 1115–1121. DOI: 10.3168/jds.S0022-0302(04)73258-0.
- United States Department of Agriculture (USDA), Animal and Plant Health Inspection Service (APHIS), National Center for Import and Export. (2011). Countries/Regions Considered Free of Foot-and-Mouth Disease. Retrieved from [https://www.aphis.usda.gov/wps/portal/aphis/ourfocus/animalhealth/sa_import_into_us/sa_entry_requirements/ct_foot_and_mouth_and_rinderpest!/ut/p/a0/04_Sj9CPykssy0xPLMnMz0vMAfGjzOK9_D2MDJ0MjDzdgy1dDTz9wtx8LXzMjf09TPQLsh0VAZdihIg!/.](https://www.aphis.usda.gov/wps/portal/aphis/ourfocus/animalhealth/sa_import_into_us/sa_entry_requirements/ct_foot_and_mouth_and_rinderpest!/ut/p/a0/04_Sj9CPykssy0xPLMnMz0vMAfGjzOK9_D2MDJ0MjDzdgy1dDTz9wtx8LXzMjf09TPQLsh0VAZdihIg!/)
- World Organization for Animal Health (OIE). (2013). Foot and Mouth Disease. In *Technical Disease Card*. Retrieved from <http://www.oie.int>.
- World Organization for Animal Health (OIE). (2016). Foot-and-Mouth Disease. In *Terrestrial Animal Health Code* (8.8.). Retrieved from <http://www.oie.int>.
- World Organization for Animal Health (OIE). (2016). Foot-and-Mouth Disease Distribution Maps, 2013–2016. In *World Animal Health Information Database (WAHID) World Animal Health Information System (WAHIS) Interface*. Retrieved from http://www.oie.int/wahis_2/public/wahid.php/Diseaseinformation/Diseasedistributionmap.
- Yang, P.C., Chu, R.M., Chung, W.B., & Sung, H.T. (1999). Epidemiological characteristics and financial costs of the 1997 foot-and-mouth disease epidemic in Taiwan. *Veterinary Record*, 145(25), 731–734. DOI: 10.1136/vr.145.25.731.

Attachment 1.B Abbreviations

APHIS	Animal and Plant Health Inspection Service
CFSPH	Center for Food Security and Public Health
FAD PReP	Foreign Animal Disease Preparedness and Response Plan
FMD	foot-and-mouth disease
FMDV	foot-and-mouth disease virus
HTST	high temperature-short time (pasteurization)
NAHEMS	National Animal Health Emergency Management System
OIE	World Organization for Animal Health
RNA	ribonucleic acid
SAT	South African Territories (serotypes of FMD)
SOP	standard operating procedure
TCID	tissue culture infectious dose
TDD	telecommunications device for the deaf
UHT	ultra-high temperature (pasteurization)
USDA	United States Department of Agriculture
WAHID	World Animal Health Information Database
WAHIS	World Animal Health Information System