



United States
Department of
Agriculture

Animal and
Plant Health
Inspection
Service


Veterinary
Services

National
Animal Health
Monitoring
System

May 2007



Cattle and Calves Nonpredator Death Loss in the United States, 2005



The U.S. Department of Agriculture (USDA) prohibits discrimination in all its programs and activities on the basis of race, color, national origin, age, disability, and where applicable, sex, marital status, familial status, parental status, religion, sexual orientation, genetic information, political beliefs, reprisal, or because all or part of an individual's income is derived from any public assistance program. (Not all prohibited bases apply to all programs.) Persons with disabilities who require alternative means for communication of program information (Braille, large print, audiotape, etc.) should contact USDA's TARGET Center at (202) 720-2600 (voice and TDD). To file a complaint of discrimination, write to USDA, Director, Office of Civil Rights, 1400 Independence Avenue, S.W., Washington, D.C. 20250-9410, or call (800) 795-3272 (voice) or (202) 720-6382 (TDD). USDA is an equal opportunity provider and employer.

Mention of companies or commercial products does not imply recommendation or endorsement by the USDA over others not mentioned. USDA neither guarantees nor warrants the standard of any product mentioned. Product names are mentioned solely to report factually on available data and to provide specific information.

USDA:APHIS:VS:CEAH
NRRC Building B, M.S. 2E7
2150 Centre Avenue
Fort Collins, CO 80526-8117
970.494.7000
E-mail: NAHMS@aphis.usda.gov
<http://www.aphis.usda.gov/nahms>
#N462.0507

Acknowledgments

This report was a cooperative effort between two U.S. Department of Agriculture (USDA) Agencies: the National Agricultural Statistics Service (NASS) and the Animal and Plant Health Inspection Service (APHIS).

Thanks to the NASS enumerators who contacted cattle producers and collected the data. Their hard work and dedication were invaluable. Thanks also to the personnel at the USDA–APHIS–Veterinary Services' Centers for Epidemiology and Animal Health for their efforts in generating and distributing this report.

All participants are to be commended, particularly the producers whose voluntary efforts made this report possible.



Larry Granger
Director
Centers for Epidemiology and Animal Health

Suggested bibliographic citation for this report:

USDA. 2007. Cattle and Calves Nonpredator Death Loss in the United States, 2005

USDA-APHIS-VS, CEAH, National Animal Health Monitoring System, Fort Collins, CO #N462.0507

Contacts for further information:

Questions or comments on the Cattle and Calves Death Loss, 2005 study methods or requests for additional data analysis: Dr. David Dargatz 970.494.7000
Information on reprints or other NAHMS reports: Mr. Brandon O'Neal 970.494.7000

E-mail: NAHMS@aphis.usda.gov

Table of Contents

Introduction 1

Terms Used In This Report 2

Section I: Population Estimates—Number of Head 3

A. U.S. Demographics 3

1. Inventory—January 1, 1996-2006 3
2. Operations—1995-2005 3
3. Calf crop—1996–2006 6

B. Regional Demographics 6

1. Cattle and calves inventory—January 1, 1992, 1996, 2001, and 2006 6
2. Beef and dairy cow inventory—January 1, 1992, 1996, 2001, and 2006 7
3. Calf crop—1991, 1995, 2000, and 2005 8

C. U.S. Annual Death Loss 8

1. Loss from all causes—1991-2005 8
2. U.S. nonpredator and predator death loss (excluding Alaska)—1991, 1995, 2000, and 2005 10
3. Regional predator and nonpredator death loss—1991, 1995, 2000, and 2005 12
4. Regional predator and nonpredator death loss—1995, 2000, and 2005 14

Section II: Population Estimates—Nonpredator Losses 18

A. Operation Classification 18

1. Operation classification process 18
2. Operation characteristics 18

B. Nonpredator Cattle Death Loss 21

1. Losses as a percentage of inventory—2005 21
2. Cause of loss—1991, 1995, 2000, and 2005 26
3. Cause of loss by size of operation—2005 32
4. Cause of loss by region—2005 36

C. Nonpredator Calf Death Loss 40

1. Losses as a percentage of calf crop—2005 40
2. Cause of loss—1991, 1995, 2000, and 2005 46
3. Cause of loss by size of operation—2005 53
4. Cause of loss by region—2005 57

Section III: Methodology 61

A. Reliability of Cattle and Calf Death Loss Estimates 61

1. Survey procedures 61
2. Estimating procedures 61
3. Reliability 61

Appendix: U.S. Populations and Operations 62

Introduction

Each January, the U.S. Department of Agriculture's (USDA) National Agricultural Statistics Service (NASS) collects cattle data on inventory, calf crop, and total death losses for State and national estimating programs. Inventory and calf crop estimates (number of head) are published in January, while cattle and calves death losses (number of head) are published annually in April via the Meat Animals: Production, Disposition and Income report. For the January 1992, 1996, 2001, and 2006 surveys, USDA's Animal and Plant Health Inspection Service (APHIS) provided funding for a detailed breakdown of total losses for producer-attributed cause of loss occurring the previous year. Death losses (number of head) by cause for predator and nonpredator losses were estimated and subsequently published by NASS (Cattle Death Loss, May 5, 2006).

This report provides additional analysis of cattle and calves losses in 2005 from all causes. Death losses by size group and region are also provided, with special emphasis on nonpredator causes of loss. When possible, the 1991, 1995, and 2000 death losses are provided for comparison.

Terms Used In This Report

Beef operation: If an operation had more beef cows than milk cows and milk cows were fewer than five head, the operation was classified as a beef operation.

Cattle: All cows, bulls, steers, and heifers 500 lb or more.

Calves: All bulls, steers, and heifers less than 500 lb.

Dairy operation: If an operation had more milk cows than beef cows and beef cows were fewer than five head, the operation was classified as a dairy operation.

Mixed operation: If an operation had more than five beef cows and more than five milk cows, or if the number of beef cows equaled the number of milk cows, the operation was classified as a mixed operation.

Other operation: Operation that did not meet the criteria to be classified as beef, dairy, or mixed operation.

N/A: Not available

Regions: The United States was divided into six regions for the purposes of providing data on smaller geographic areas. Alaska is not included because cause-of-loss estimates are not generated for the State.

North Central: Iowa, Minnesota, Nebraska, North Dakota, South Dakota

Northeast: Connecticut, Delaware, Illinois, Indiana, Kentucky, Maine, Maryland, Massachusetts, Michigan, New Hampshire, New Jersey, New York, Ohio, Pennsylvania, Rhode Island, Vermont, Virginia, West Virginia, Wisconsin

Northwest: Idaho, Montana, Oregon, Washington, Wyoming

South Central: Arkansas, Kansas, Missouri, Oklahoma, Texas

Southeast: Alabama, Florida, Georgia, Louisiana, Mississippi, North Carolina, South Carolina, Tennessee

Southwest: Arizona, California, Colorado, Hawaii, Nevada, New Mexico, Utah

Operation size: Total number of cows for beef, dairy, and mixed operations.

For beef operations size groups are: 1-49, 50-99, 100-499, and 500 or more head.

For dairy operations size groups are: 1-29, 30-49, 50-99, 100-199, 200-499, and 500 or more head.

For mixed operations size groups are: 1-99, 100-499, and 500 or more head.

For other operations (total number of cattle and calves) size groups are: 1-99 head, 100-499, 500-999, and 1,000 or more head.

Section I: Population Estimates—Number of Head

A. U.S. Demographics

1. Inventory—January 1, 1996-2006¹

Number of cattle and calves, by year:

January 1 Number (x 1,000)*											
Year											
Class	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006
All cattle and calves	103,548	101,656	99,744	99,115	98,199	97,298	96,723	96,100	94,888	95,438	96,702
Cattle 500 lb or more	85,164	83,830	82,343	81,825	81,383	81,082	80,971	80,555	79,679	80,188	81,237
Beef cows	35,319	34,458	33,885	33,750	33,575	33,398	33,134	32,983	32,861	32,915	32,994
Dairy cows	9,420	9,318	9,199	9,128	9,183	9,172	9,106	9,142	8,990	9,005	9,063
Heifers, steers, and bulls	40,426	40,054	39,259	38,946	38,625	38,511	38,731	38,430	37,828	38,268	39,180
Calves	18,384	17,826	17,401	17,290	16,816	16,216	15,753	15,545	15,210	15,250	15,465

*Numbers may not add up due to rounding

2. Operations—1995-2005

a. Number of operations with cattle and calves, beef cows, and milk cows, by year:

Number Operations											
Year											
	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005
Cattle and calves	1,190,630	1,176,700	1,148,050	1,114,650	1,095,560	1,076,370	1,049,170	1,036,430	1,013,570	989,460	982,510
Beef cows	897,660	885,980	872,840	855,460	844,170	831,270	814,520	808,110	792,050	774,930	770,170
Milk cows	139,670	130,980	123,700	117,145	110,855	105,055	97,460	91,240	86,360	81,520	78,300

¹NASS multiple reports

b. Percentage of operations with beef cows, by size of operation and by year:

Percent Operations											
Year											
Size of Operation (Number of Beef Cows)	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005
1 to 49	79.8	79.4	79.3	79.3	79.0	78.8	78.5	78.4	78.3	77.6	77.5
50 to 99	11.8	12.0	12.0	11.9	12.0	12.0	12.1	12.2	12.1	12.4	12.3
100 to 499	7.8	8.0	8.1	8.2	8.4	8.5	8.7	8.7	8.9	9.3	9.5
500 or more	0.6	0.6	0.6	0.6	0.6	0.7	0.7	0.7	0.7	0.7	0.7
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0

c. Percentage of operations with dairy cows, by size of operation and by year:

Percent Operations											
Year											
Size of Operation (Number of Dairy Cows)	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005
1 to 29	34.5	32.9	31.6	30.8	29.7	29.3	29.0	28.9	29.0	29.2	28.7
30 to 49	22.2	22.3	22.1	21.8	21.7	21.1	20.4	19.8	19.5	19.0	19.0
50 to 99	28.1	28.7	29.0	29.1	29.6	29.7	29.8	30.0	29.9	29.5	29.6
100 to 199	10.2	10.7	11.3	11.8	11.9	12.2	12.6	12.7	12.7	12.8	12.8
200 or more	5.0	5.4	6.0	6.5	7.1	7.7	8.2	8.6	8.9	9.5	9.9
200 to 499	N/A	N/A	4.1	4.4	4.8	5.1	5.3	5.4	5.5	5.8	6.0
500 or more	N/A	N/A	1.9	2.1	2.3	2.6	2.9	3.2	3.4	3.7	3.9
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0

d. Percentage of beef cow inventory, by size of operation and by year:

Percent Inventory											
Year											
Size of Operation (Number of Beef Cows)	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005
1 to 49	31.2	30.8	30.4	30.5	29.9	29.5	29.0	29.0	29.1	28.1	27.9
50 to 99	19.2	19.6	19.4	18.8	19.1	19.1	19.1	19.2	19.0	19.1	19.0
100 to 499	35.3	35.4	35.9	36.1	36.6	36.7	37.0	37.3	37.5	38.3	38.5
500 or more	14.3	14.2	14.3	14.6	14.4	14.7	14.9	14.5	14.4	14.5	14.6
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0

e. Percentage of dairy cow inventory, by size of operation and by year:

Percent Inventory											
Year											
Size of Operation (Number of Dairy Cows)	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005
1 to 29	4.0	4.0	3.5	3.6	3.2	2.9	2.7	2.4	2.3	2.1	2.0
30 to 49	13.0	12.0	11.5	10.5	10.2	9.1	8.0	7.4	6.9	6.6	6.4
50 to 99	28.0	27.0	26.0	24.3	23.3	22.0	20.8	19.6	18.8	17.8	17.1
100 to 199	20.0	20.0	20.0	19.3	18.4	18.1	17.2	16.4	15.7	15.1	14.6
200 or more	35.0	37.0	39.0	42.3	44.9	47.9	51.3	54.2	55.3	58.4	59.9
200 to 499	N/A	N/A	14.6	15.5	16.3	16.6	16.3	15.9	15.4	15.5	15.4
500 or more	N/A	N/A	24.4	26.8	28.6	31.3	35.0	38.3	40.9	42.9	44.5
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0

3. Calf crop—1995–2005

Calf crop and calf crop per 100 cows on hand, by year:

	Year										
	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005
Calf crop (1,000 Head)	40,264	39,823	38,961	38,812	38,796	38,631	38,300	38,224	37,903	37,505	37,575
Calf crop per 100 cows on hand (January 1)	90.1	89.0	89.0	90.1	90.5	90.3	90.0	90.5	90.0	89.6	89.6

B. Regional Demographics

1. Cattle and calves inventory—January 1, 1992, 1996, 2001, and 2006

NASS does not conduct a cause-of-loss survey in Alaska; therefore Alaska is excluded from regional demographics tables and nonpredator loss estimates. Overall, the national inventory of cattle and calves remained relatively stable from 1992 to 2006, with 96.7 million head present on January 1, 2006.

Number of cattle and calves and number of cattle over 500 lb, by region and by year:

Region	January 1—Number (x 1,000)							
	Cattle and Calves				Cattle Over 500 lb			
	1992	1996	2001	2006	1992	1996	2001	2006
North Central	18,000	19,110	18,830	18,120	15,248	16,555	16,884	16,390
Northeast	18,638	18,184	16,111	16,185	14,598	14,259	12,903	12,852
Northwest	8,300	8,920	8,600	8,460	7,320	8,018	7,796	7,734
South Central	30,480	33,610	31,500	32,410	24,565	27,390	25,650	26,920
Southeast	10,760	11,710	9,925	9,460	8,152	8,845	7,506	7,194
Southwest	11,270	12,004	12,320	12,051	9,331	10,089	10,334	10,134
Total*	97,548	103,538	97,286	96,686	79,214	85,156	81,073	81,224

*Excludes Alaska, so total of regions does not match published United States total.

2. Beef and dairy cow inventory—January 1, 1992, 1996, 2001, and 2006

From 1992 to 2006, U.S. inventory of beef cows fluctuated between 33 million and 35 million head, while dairy cow inventory decreased from 9.7 million to 9.0 million head. In 2006, the South Central region had more beef cows than any other region (12.2 million head), while the Northeast region had more dairy cows than any other region (3.8 million).

Number of beef and dairy cows that had calved, by region and by year:

January 1—Number (x 1,000)								
Region	Beef Cows				Dairy Cows			
	1992	1996	2001	2006	1992	1996	2001	2006
North Central	5,618.0	6,072.0	6,151.0	6,029.0	12,320.0	1,078.0	949.0	806.0
Northeast	3,832.3	3,926.6	3,510.5	3,695.0	4,787.4	4,417.1	4,017.6	3,831.5
Northwest	3,641.0	3,874.0	3,721.0	3,543.0	549.0	626.0	714.0	857.0
South Central	11,484.0	12,520.0	11,895.0	12,185.0	856.0	820.0	715.0	645.0
Southeast	5,262.0	5,760.0	5,077.0	4,885.0	763.0	640.0	543.0	421.0
Southwest	3,167.0	3,153.0	3,040.0	2,650.4	1,540.0	1,838.0	2,178.0	2,501.6
Total*	33,004.3	35,315.6	33,394.5	32,987.4	9,727.4	9,419.1	9,170.6	9,062.1

*Excludes Alaska, so total of regions does not match published United States total.

3. Calf crop—1991, 1995, 2000, and 2005

About 30 percent of the 37.6 million calves born in 2005 were born in the South Central region.

Calendar year calf crop, by region and by year:

Calf Crop (x 1,000)				
Year				
Region	1991	1995	2000	2005
North Central	6,570	6,770	6,730	6,420
Northeast	7,823	7,576	6,813	6,569
Northwest	3,945	4,280	4,410	4,410
South Central	11,070	11,910	11,330	11,450
Southeast	5,205	5,475	4,880	4,560
Southwest	3,968	4,249	4,464	4,361
Total*	38,581	40,261	38,627	37,570

*Excludes Alaska, so total of regions does not match published United States total.

C. U.S. Annual Death Loss**1. Loss from all causes—1991-2005**

A total of 1.7 million cattle and 2.3 million calves were lost to all causes during 2005. These totals represent 2.1 and 6.2 percent of cattle inventory and calf crop, respectively. While the percentage of cattle inventory lost remained stable, the percentage of calf crop lost declined from 7.0 percent in 1991 to 6.2 percent in 2005.


a. Number of cattle and calf death losses due to all causes, by year:

Number (x 1,000)															
Year															
	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005
Cattle	1,540	1,677	1,702	1,589	1,645	1,761	1,847	1,668	1,658	1,711	1,722	1,710	1,710	1,711	1,718
Calves	2,707	2,689	2,921	2,681	2,739	2,811	2,829	2,541	2,455	2,387	2,487	2,366	2,320	2,292	2,335
Total	4,247	4,366	4,623	4,270	4,384	4,572	4,676	4,209	4,113	4,098	4,209	4,076	4,030	4,003	4,053

b. Cattle death losses as a percentage of January 1 inventory of cattle over 500 lb on January 1 of the following year, by year:

Percent Inventory														
Year														
1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005
1.9	2.1	2.0	1.9	1.9	2.1	2.2	2.0	2.0	2.1	2.1	2.1	2.1	2.1	2.1


Cattle Death Losses as a Percentage of January 1 Inventory of Cattle Over 500 lb on January 1 of the Following Year, by Year



c. Calf death losses as a percentage of calf crop, by year:

Percent Calf Crop														
Year														
1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005
7.0	6.9	7.4	6.7	6.8	7.1	7.3	6.5	6.3	6.2	6.5	6.2	6.1	6.1	6.2

Calf Death Losses as a Percentage of Calf Crop, by Year



2. U.S. nonpredator and predator death loss (excluding Alaska)—1991², 1995², 2000², and 2005

Since 1991, nonpredator causes accounted for the majority of cattle and calf losses; 95.3 percent of losses in 2005 were attributed to nonpredator causes. As expected, predators accounted for a larger percentage of calf losses than cattle losses (6.7 and 2.0 percent, respectively).

a. Number and percentage of cattle and calf death losses, by cause and by year:

Cause	Year							
	1991		1995		2000		2005	
	Number (x 1,000)	Pct.	Number (x 1,000)	Pct.	Number (x 1,000)	Pct.	Number (x 1,000)	Pct.
Nonpredator	4,143.4	97.6	4,263.6	97.3	3,952.0	96.4	3,861.0	95.3
Predator	102.0	2.4	118.8	2.7	144.6	3.6	190.0	4.7
Total	4,245.4	100.0	4,382.4	100.0	4,096.6	100.0	4,051.0	100.0

b. Number and percentage of cattle death losses, by cause and by year:

Cause	Year							
	1991		1995		2000		2005	
	Number (x 1,000)	Pct.	Number (x 1,000)	Pct.	Number (x 1,000)	Pct.	Number (x 1,000)	Pct.
Nonpredator	1523.7	99.0	1,622.7	98.7	1,689.6	98.8	1,683.0	98.0
Predator	15.4	1.0	21.4	1.3	20.5	1.2	34.0	2.0
Total	1,539.1	100.0	1,644.1	100.0	1,710.1	100.0	1,717.0	100.0

c. Number and percentage of calf death losses, by cause and by year:

Cause	Year							
	1991		1995		2000		2005	
	Number (x 1,000)	Pct.	Number (x 1,000)	Pct.	Number (x 1,000)	Pct.	Number (x 1,000)	Pct.
Nonpredator	2,619.7	96.8	2,640.9	96.5	2,262.4	94.8	2,178.0	93.3
Predator	86.6	3.2	97.4	3.5	124.1	5.2	156.0	6.7
Total	2,706.3	100.0	2,738.3	100.0	2,386.5	100.0	2,334.0	100.0

²NASS estimates of total death loss were revised in subsequent publications. However, estimates of predator and nonpredator death loss were not revised. The ratio of predator to nonpredator losses in the initial publication was maintained and applied to the revised total loss estimate to generate the losses due to predators and nonpredators in tables a, b, and c. For the 1991 estimates, cattle losses were revised from 1,583,800 to 1,539,100, and calf losses were revised from 2,786,300 to 2,706,300. For the 1995 estimates, cattle losses were revised from 1,635,100 to 1,644,100, and calf losses were revised from 2,745,300 to 2,738,300. For the 2000 estimates, cattle losses were revised from 1,721,100 to 1,710,100, and calf losses were revised from 2,409,500 to 2,386,500.

3. Regional predator and nonpredator death loss—1991, 1995, 2000, and 2005

a. Number of cattle death losses from all causes, by region and by year:

Number Losses (x 1,000)				
Year				
Region	1991	1995	2000	2005
North Central	264.0	277.0	281.0	267.0
Northeast	312.1	311.1	291.1	300.0
Northwest	103.0	115.0	124.0	117.0
South Central	498.0	522.0	625.0	635.0
Southeast	183.0	201.0	174.0	174.0
Southwest	179.0	218.0	215.0	224.0
Total*	1,539.1	1,644.1	1,710.1	1,717.0

*Excludes Alaska, so total of regions does not match published United States total.

In 2005, the Northwest and North Central regions lost a smaller percentage of cattle (1.5 and 1.6 percent, respectively) than the other regions.

b. Cattle death losses as a percentage of January 1 inventory of cattle over 500 lb on January 1 of the following year, by year:

Percent Inventory				
Year				
Region	1991	1995	2000	2005
North Central	1.7	1.7	1.7	1.6
Northeast	2.1	2.2	2.3	2.3
Northwest	1.4	1.4	1.6	1.5
South Central	2.0	1.9	2.4	2.4
Southeast	2.2	2.3	2.3	2.4
Southwest	1.9	2.2	2.1	2.4
Total	1.9	1.9	2.1	2.1

c. Number of calf death losses from all causes, by region and by year:

Number Losses (x 1,000)				
Year				
Region	1991	1995	2000	2005
North Central	493.0	553.0	425.0	392.0
Northeast	717.3	673.3	538.5	543.0
Northwest	204.0	216.0	225.0	205.0
South Central	689.0	675.0	620.0	638.0
Southeast	297.0	313.0	266.0	238.0
Southwest	306.0	308.0	312.0	318.0
Total*	2,706.3	2,738.3	2,386.5	2,334.0

*Excludes Alaska, so total of regions does not match published United States total.

The Northeast region lost the largest percentage of calf crop (8.2 percent) in 2005, compared all other regions.

d. Calf death losses as a percentage of calf crop, by region and by year:

Percent Calf Crop				
Year				
Region	1991	1995	2000	2005
North Central	7.5	8.2	6.3	6.1
Northeast	9.2	8.9	7.9	8.2
Northwest	5.2	5.0	5.1	4.8
South Central	6.2	5.7	5.5	5.5
Southeast	5.7	5.7	5.5	5.2
Southwest	7.7	7.2	7.0	7.2
Total	7.0	6.8	6.2	6.2


4. Regional predator and nonpredator death loss—1995, 2000, and 2005

In all regions, nonpredator causes accounted for the majority of cattle losses. With the exception of the North Central and South Central regions, the percentages of losses due to nonpredator causes decreased in all regions from 1995 to 2005. In the Southeast region, the percentage of losses due to nonpredator causes decreased from 98.5 percent in 1995 to 94.9 percent in 2005.

a. Number and percentage of cattle death losses by cause—by region and by year:

Region	Cause	Year					
		1995		2000		2005	
		Number (x1,000)	Percent	Number (x1,000)	Percent	Number (x1,000)	Percent
North Central	Nonpredator	274.8	99.2	280.2	99.7	265.6	99.5
	Predator	2.2	0.8	0.8	0.3	1.4	0.5
	Total	277.0	100.0	281.0	100.0	267.0	100.0
Northeast	Nonpredator	308.3	99.1	288.2	99.0	294.7	98.2
	Predator	2.8	0.9	2.9	1.0	5.3	1.8
	Total	311.1	100.0	291.1	100.0	300.0	100.0
Northwest	Nonpredator	113.5	98.7	121.9	98.3	114.1	97.5
	Predator	1.5	1.3	2.1	1.7	2.9	2.5
	Total	115.0	100.0	124.0	100.0	117.0	100.0
South Central	Nonpredator	514.2	98.5	619.7	99.2	625.0	98.4
	Predator	7.8	1.5	5.3	0.8	10.0	1.6
	Total	522.0	100.0	625.0	100.0	635.0	100.0
Southeast	Nonpredator	198.0	98.5	170.5	98.0	165.1	94.9
	Predator	3.0	1.5	3.5	2.0	8.9	5.1
	Total	201.0	100.0	174.0	100.0	174.0	100.0
Southwest	Nonpredator	214.1	98.2	208.8	97.1	218.5	97.5
	Predator	3.9	1.8	6.2	2.9	5.5	2.5
	Total	218.0	100.0	215.0	100.0	224.0	100.0

Percentage of 2005 Cattle Death Losses, by Cause and by Region




In all regions in 2005, nonpredator causes accounted for the majority of calf losses. However, the percentage of total losses due to nonpredator causes ranged from 86.4 percent in the Southeast region to 97.5 percent in the North Central region.

b. Number and percentage of calf death losses by cause—by region and by year:

Region	Cause	Year					
		1995		2000		2005	
		Number (x1,000)	Percent	Number (x1,000)	Percent	Number (x1,000)	Percent
North Central	Nonpredator	545.3	98.6	416.9	98.1	382.1	97.5
	Predator	7.7	1.4	8.1	1.9	9.9	2.5
	Total	553.0	100.0	425.0	100.0	392.0	100.0
Northeast	Nonpredator	662.5	98.4	526.1	97.7	520.3	95.8
	Predator	10.8	1.6	12.4	2.3	22.7	4.2
	Total	673.3	100.0	538.5	100.0	543.0	100.0
Northwest	Nonpredator	206.1	95.4	210.1	93.3	191.4	93.4
	Predator	9.9	4.6	14.9	6.7	13.6	6.6
	Total	216.0	100.0	225.0	100.0	205.0	100.0
South Central	Nonpredator	636.5	94.3	581.6	93.8	581.0	91.1
	Predator	38.5	5.7	38.4	6.2	57.0	8.9
	Total	675.0	100.0	620.0	100.0	638.0	100.0
Southeast	Nonpredator	299.5	95.7	242.4	91.1	205.6	86.4
	Predator	13.5	4.3	23.6	8.9	32.4	13.6
	Total	313.0	100.0	266.0	100.0	238.0	100.0
Southwest	Nonpredator	292.3	94.9	284.5	91.2	297.6	93.6
	Predator	15.7	5.1	27.5	8.8	20.4	6.4
	Total	308.0	100.0	312.0	100.0	317.0	100.0

Percentage of 2005 Calf Death Losses, by Cause and by Region



Section II: Population Estimates—Nonpredator Losses

A. Operation Classification

1. Operation classification process

This section reports death loss and cause of loss on three operation types: beef, dairy, and mixed operations (see p. 2 for definition of operation types). NASS collects inventory numbers of beef cows, milk cows, and heifers on each operation. However, animal breed is not specified in death-loss numbers. To estimate death loss, operations were classified based on inventory numbers of beef and dairy cows.

2. Operation characteristics

Nearly two-thirds (65.6 percent) of the January 1, 2006, inventory of cattle and calves was on beef operations. Over 98 percent of beef cows were on beef operations, and 1.4 percent were on mixed operations. Ninety percent of the dairy cow inventory was on dairy operations, and 9.9 percent was on mixed operations.

a. Percentage of January 1, 2006, inventory for all cattle and calves, beef cows, and dairy cows, by operation type:

Operation Type	Percent Inventory					
	All Cattle and Calves		Beef Cows		Dairy Cows	
	Percent	Std. Error	Percent	Std. Error	Percent	Std. Error
Beef	65.6	(0.8)	98.6	(0.1)	0.1	(0.0)
Dairy	15.6	(0.4)	0.0	(0.0)	90.0	(1.7)
Mixed	2.6	(0.4)	1.4	(0.1)	9.9	(1.7)
Other	16.2	(0.8)	0.0	(--)	0.0	(--)
Total	100.0		100.0		100.0	

Across all operation types, nonpredator causes accounted for the majority of cattle losses in 2005.

b. Percentage of 2005 cattle death losses, by cause and by operation type:


Percent Losses								
Operation Type								
	Beef		Dairy		Mixed		Other	
Cause	Pct.	Std. Error	Pct.	Std. Error	Pct.	Std. Error	Pct.	Std. Error
Nonpredator	96.6	(0.3)	99.7	(0.1)	99.6	(0.1)	99.3	(0.2)
Predator	3.4	(0.3)	0.3	(0.1)	0.4	(0.1)	0.7	(0.2)
Total	100.0		100.0		100.0		100.0	

Just over 90 percent of calf losses on beef operations were due to nonpredator causes, compared to at least 97 percent of losses on dairy, mixed, and other operations (99.0, 97.0, and 98.9 percent of losses, respectively).

c. Percentage of 2005 calf death losses, by cause and by operation type:

Percent Losses								
Operation Type								
	Beef		Dairy		Mixed		Other	
Cause	Pct.	Std. Error	Pct.	Std. Error	Pct.	Std. Error	Pct.	Std. Error
Nonpredator	90.1	(0.4)	99.0	(0.1)	97.0	(0.9)	98.9	(0.3)
Predator	9.9	(0.4)	1.0	(0.1)	3.0	(0.9)	1.1	(0.3)
Total	100.0		100.0		100.0		100.0	

Percentage of 2005 Calf Losses, by Cause and by Operation Type



Half of cattle losses (51.4 percent) and 61.0 percent of calf losses in the United States occurred on beef operations, which is most likely a reflection of the fact that the majority of U.S. cattle and calves are on beef operations.

d. Percentage of nonpredator cattle and calf death losses, by operation type:

Operation Type	Percent Nonpredator Loss			
	Cattle		Calf	
	Percent	Std. Error	Percent	Std. Error
Beef	51.4	(1.4)	61.0	(1.5)
Dairy	24.9	(0.8)	21.0	(0.7)
Mixed	3.2	(0.6)	3.6	(0.4)
Other	20.5	(1.7)	14.4	(1.9)
Total	100.0		100.0	

B. Nonpredator Cattle Death Loss

1. Losses as a percentage of inventory—2005

The percentage of cattle inventory lost to nonpredator causes ranged from 1.5 percent in the Northwest region to 2.6 percent in the South Central region.

a. Percentage of cattle inventory lost,* by region:

Region	Percent Inventory	
	Percent	Standard Error
North Central	1.8	(0.1)
Northeast	2.3	(0.1)
Northwest	1.5	(0.1)
South Central	2.6	(0.1)
Southeast	2.2	(0.1)
Southwest	2.3	(0.1)

*2005 death loss divided by end-of-year (January 1, 2006) inventory of cattle 500 lb or greater X 100.

Overall, 1.7 percent of cattle inventory on beef operations was lost to nonpredator causes. Across all regions, operations with 1 to 49 operations lost a higher percentage of inventory to nonpredator causes than operations with 500 or more cows.

b. For **beef** operations, percentage of cattle inventory lost by region and by size of operation:

Percent Inventory*										
Size of Operation (Number of Cows)										
Region	1-49		50-99		100-499		500 or More		All Operations	
	Pct.	Std. Error	Pct.	Std. Error	Pct.	Std. Error	Pct.	Std. Error	Pct.	Std. Error
North Central	1.8	(0.3)	1.2	(0.1)	1.1	(0.1)	1.1	(0.1)	1.2	(0.1)
Northeast	1.7	(0.1)	1.4	(0.1)	1.2	(0.1)	1.1	(0.1)	1.5	(0.1)
Northwest	1.6	(0.2)	1.2	(0.2)	0.9	(0.1)	0.8	(0.1)	1.0	(0.1)
South Central	3.1	(0.2)	2.3	(0.2)	2.0	(0.2)	1.3	(0.1)	2.3	(0.1)
Southeast	2.3	(0.1)	2.2	(0.2)	1.9	(0.2)	1.1	(0.1)	2.0	(0.1)
Southwest	2.2	(0.3)	1.7	(0.2)	1.2	(0.1)	1.3	(0.1)	1.4	(0.1)
All operations	2.4	(0.1)	1.8	(0.1)	1.5	(0.1)	1.1	(0.1)	1.7	(0.0)

*2005 death loss divided by end-of-year (January 1, 2006) inventory of cattle 500 lb or greater x 100.

Overall, 3.6 percent of cattle inventory on dairy operations was lost to nonpredator causes, ranging from 3.1 percent of inventory in the Northeast region to 6.1 percent in the South Central region. Across all regions, the percentage of inventory lost was higher on dairy operations than on beef operations (table b).

c. For **dairy** operations, percentage of cattle inventory lost by region and by size of operation:*

Percent Inventory														
Size of Operation (Number of Cows)														
Region	1-29		30-49		50-99		100-199		200-499		500 or More		All Operations	
	Pct.	Std. Error	Pct.	Std. Error	Pct.	Std. Error	Pct.	Std. Error	Pct.	Std. Error	Pct.	Std. Error	Pct.	Std. Error
North Central	3.8	(1.1)	3.7	(0.7)	3.3	(0.3)	3.6	(0.4)	4.1	(0.4)	9.0	(0.6)	4.5	(0.2)
Northeast	3.1	(0.5)	2.4	(0.2)	2.8	(0.1)	3.0	(0.2)	3.4	(0.2)	4.0	(0.3)	3.1	(0.1)
Northwest	2.0	(0.6)	4.4	(1.1)	3.9	(0.5)	3.3	(0.3)	3.7	(0.3)	4.0	(0.4)	3.9	(0.3)
South Central	2.2	(1.1)	7.2	(1.8)	6.0	(0.9)	4.9	(0.6)	5.6	(0.8)	7.1	(1.1)	6.1	(0.6)
Southeast	**		3.9	(0.3)	4.6	(0.5)	4.4	(0.3)	3.3	(0.2)	4.0	(0.3)	4.0	(0.2)
Southwest	**		**		3.0	(0.6)	3.9	(0.7)	3.8	(0.3)	3.0	(0.3)	3.2	(0.3)
All operations	3.9	(0.9)	2.7	(0.2)	3.1	(0.1)	3.4	(0.1)	3.7	(0.2)	3.8	(0.2)	3.6	(0.1)

*2005 death loss divided by end-of-year (January 1, 2006) inventory of cattle 500 lb or greater X 100.

**Too few operations to report.

On mixed operations, 2.7 percent of inventory was lost to nonpredator causes.

d. For *mixed* operations, percentage of cattle inventory lost by region and by size of operation:*

Percent Inventory								
Size of Operation (Number of Cows)								
	1-99		100-499		500 or More		All Operations	
Region	Pct.	Std. Error	Pct.	Std. Error	Pct.	Std. Error	Pct.	Std. Error
North Central	3.3	(0.5)	2.2	(0.4)	2.6	(0.9)	2.6	(0.3)
Northeast	2.8	(0.5)	2.3	(0.3)	3.2	(0.6)	2.6	(0.2)
Northwest	**		2.2	(0.5)	1.9	(0.3)	2.0	(0.3)
South Central	4.4	(1.3)	2.5	(0.4)	2.2	(0.8)	2.9	(0.4)
Southeast	2.6	(0.8)	2.7	(0.3)	2.6	(0.5)	2.7	(0.3)
Southwest	**		0.9	(0.2)	3.6	(0.3)	3.0	(0.6)
All operations	3.2	(0.3)	2.1	(0.2)	3.1	(0.3)	2.7	(0.2)

*2005 death loss divided by end-of-year (January 1, 2006) inventory of cattle 500 lb or greater X 100.
 ** Too few operations to report.

On “other” operations, 2.7 percent of inventory was lost to nonpredator causes.

e. For **other** operations, percentage of cattle inventory lost by region and by size of operation:*

Percent Inventory										
Size of Operation (Number of Cattle and Calves)										
Region	1-99		100-499`		500-999		1,000 or More		All Operations	
	Pct.	Std. Error	Pct.	Std. Error	Pct.	Std. Error	Pct.	Std. Error	Pct.	Std. Error
North Central	4.8	(0.9)	2.6	(0.3)	1.9	(0.2)	3.1	(0.2)	3.0	(0.1)
Northeast	2.7	(0.4)	2.0	(0.3)	1.8	(0.3)	1.4	(0.2)	2.1	(0.2)
Northwest	6.9	(1.7)	1.6	(0.5)	1.0	(0.2)	1.7	(0.2)	2.0	(0.2)
South Central	9.0	(2.6)	2.9	(0.6)	1.8	(0.4)	3.0	(0.5)	3.2	(0.4)
Southeast	5.1	(1.6)	1.5	(0.6)	3.2	(0.5)	2.3	(0.4)	3.5	(0.6)
Southwest	4.5	(1.5)	2.6	(0.8)	2.1	(0.4)	2.6	(0.2)	2.6	(0.2)
All operations	5.0	(0.6)	2.4	(0.2)	2.0	(0.2)	2.8	(0.3)	2.9	(0.2)

*2005 death loss divided by end-of-year (January 1, 2006) inventory of cattle 500 lb or greater X100.

2. Cause of loss—1991, 1995, 2000, and 2005

Since 1991, respiratory problems accounted for the highest percentage of nonpredator cattle losses (24.8 percent). Other common causes of loss in 2005 included unknown causes (13.3 percent) and calving problems (11.1 percent).

a. Percentage of cattle death losses, by cause and by year:


Percent Nonpredator Loss				
Nonpredator Cause	1991	1995	2000	2005
Digestive problems (bloat, scours, parasites, enterotoxemia, acidosis, etc.)	12.8	12.2	10.5	11.1
Respiratory problems (pneumonia, shipping fever, etc.)	26.1	26.0	27.1	24.8
Metabolic problems (milk fever, grass tetany, etc.)	N/A	N/A	4.0	3.6
Weather-related causes (chilling, drowning, lightning, etc.)	7.4	8.3	8.0	6.5
Theft (stolen)	1.0	0.5	1.2	0.6
Poisoning (nitrate poisoning, noxious feeds, noxious weeds, etc.)	3.2	2.0	1.6	1.4
Calving problems	16.2	15.3	12.4	11.1
Mastitis*	N/A	N/A	N/A	4.0
Lameness/injury**	N/A	N/A	N/A	5.3
Other diseases***	N/A	N/A	6.1	5.8
Other causes***	33.3	35.7	13.2	12.5
Unknown causes***	N/A	N/A	15.9	13.3
Total	100.0	100.0	100.0	100.0

*Mastitis included in "other causes" in 1991 and 1995 and in "other diseases" in 2000.

**Lameness/injury included in "other causes" in 1991, 1995, and 2000.

*** "Other diseases" and "unknown causes" included in "other causes" in 1991 and 1995.

Percentage of 2005 Cattle Death Losses, by Cause



*Mastitis included in "other causes" in 1991 and 1995 and in "other diseases" in 2000.

**Lameness/injury included in "other causes" in 1991, 1995, and 2000.

***"Other diseases" and "unknown causes" included in "other causes" in 1991 and 1995.

Respiratory problems accounted for just under one-fifth of cattle losses (18.7 percent) on beef operations. Not surprisingly, the percentage of losses due to mastitis was higher on dairy operations than on beef operations (12.8 and 0.5 percent, respectively.) Mastitis also accounted for 15.8 percent of losses on mixed operations, likely because almost 10 percent of the dairy cow inventory was on mixed operations (see table 2a, p. 18). Respiratory problems accounted for over half of nonpredator losses (54.2 percent) on “other” operations; many operations in the this category are feedlots, which commonly experience problems with bovine respiratory disease.

b. For **beef** operations, percentage of cattle death losses by cause and by year:

Percent Nonpredator Loss						
Nonpredator Cause	1991 ¹	1995 ¹	2000	Std. Error	2005	Std. Error
Digestive problems (bloat, scours, parasites, enterotoxemia, acidosis, etc.)	9.3	10.3	7.5	(0.4)	7.5	(0.5)
Respiratory problems (pneumonia, shipping fever, etc.)	12.8	14.2	16.9	(0.8)	18.7	(1.8)
Metabolic problems (milk fever, grass tetany, etc.)	N/A	N/A	1.7	(0.2)	1.2	(0.1)
Weather-related causes (chilling, drowning, lightning, etc.)	10.0	10.0	11.9	(0.7)	9.1	(0.6)
Theft (stolen)	2.3	0.6	2.2	(1.2)	0.9	(0.2)
Poisoning (nitrate poisoning, noxious feeds, noxious weeds, etc.)	6.1	3.2	2.6	(0.3)	2.2	(0.4)
Calving problems	18.9	19.0	15.5	(0.7)	14.2	(0.6)
Mastitis*	N/A	N/A	N/A	(--)	0.5	(0.1)
Lameness/injury**	N/A	N/A	N/A	(--)	3.4	(0.2)
Other diseases***	N/A	N/A	3.4	(0.4)	5.4	(0.5)
Other causes***	40.6	42.7	16.8	(0.7)	19.8	(0.9)
Unknown causes***	N/A	N/A	21.5	(0.9)	17.1	(0.8)
Total	100.0	100.0	100.0		100.0	

¹Standard errors not available for 1991 and 1995.

*Mastitis included in “other causes” in 1991 and 1995 and in “other diseases” in 2000.

**Lameness/injury included in “other causes” in 1991, 1995, and 2000.

*** “Other diseases” and “unknown causes” included in “other causes” in 1991 and 1995.

c. For *dairy* operations, percentage of cattle death losses by cause and by year:

Percent Nonpredator Loss						
Nonpredator Cause	1991 ¹	1995 ¹	2000	Std. Error	2005	Std. Error
Digestive problems (bloat, scours, parasites, enterotoxemia, acidosis, etc.)	14.3	15.2	12.2	(0.5)	12.3	(0.6)
Respiratory problems (pneumonia, shipping fever, etc.)	11.4	18.6	14.6	(0.6)	14.4	(0.8)
Metabolic problems (milk fever, grass tetany, etc.)	N/A	N/A	10.6	(0.5)	9.9	(0.5)
Weather-related causes (chilling, drowning, lightning, etc.)	4.4	8.2	4.3	(0.8)	2.0	(0.2)
Theft (stolen)	0.3	0.0	0.2	(0.1)	0.1	(0.1)
Poisoning (nitrate poisoning, noxious feeds, noxious weeds, etc.)	1.9	0.9	0.6	(0.2)	0.5	(0.1)
Calving problems	26.3	19.3	13.8	(0.5)	12.5	(0.6)
Mastitis*	N/A	N/A	N/A	(--)	12.8	(0.5)
Lameness/injury**	N/A	N/A	N/A	(--)	10.5	(0.5)
Other diseases***	N/A	N/A	15.4	(0.7)	8.3	(1.4)
Other causes***	41.4	37.8	15.3	(1.0)	6.2	(0.6)
Unknown causes***	N/A	N/A	13.0	(0.5)	10.5	(0.7)
Total	100.0	100.0	100.0		100.0	

¹Standard errors not available for 1991 and 1995.

*Mastitis included in "other causes" in 1991 and 1995 and in "other diseases" in 2000.

**Lameness/injury included in "other causes" in 1991, 1995, and 2000.

*** "Other diseases" and "unknown causes" included in "other causes" in 1991 and 1995.

d. For *mixed* operations, percentage of cattle death losses by cause and by year:

Percent Nonpredator Loss						
Nonpredator Cause	1991 ¹	1995 ¹	2000	Std. Error	2005	Std. Error
Digestive problems (bloat, scours, parasites, enterotoxemia, acidosis, etc.)	12.8	19.9	11.3	(1.3)	7.7	(1.7)
Respiratory problems (pneumonia, shipping fever, etc.)	9.4	22.5	16.8	(1.7)	15.2	(3.6)
Metabolic problems (milk fever, grass tetany, etc.)	N/A	N/A	10.1	(1.4)	11.9	(3.2)
Weather-related causes (chilling, drowning, lightning, etc.)	8.7	8.8	10.0	(1.5)	3.5	(1.0)
Theft (stolen)	0.4	0.01	1.7	(1.2)	0.1	(0.1)
Poisoning (nitrate poisoning, noxious feeds, noxious weeds, etc.)	3.1	0.01	0.4	(0.2)	0.8	(0.4)
Calving problems	28.3	15.0	16.0	(1.4)	12.7	(3.2)
Mastitis*	N/A	N/A	N/A	(--)	15.8	(6.4)
Lameness/injury**	N/A	N/A	N/A	(--)	7.4	(1.8)
Other diseases***	N/A	N/A	7.3	(0.9)	4.3	(1.3)
Other causes***	37.3	33.8	10.5	(1.5)	10.1	(2.6)
Unknown***	N/A	N/A	15.9	(1.5)	10.5	(3.5)
Total	100.0	100.0	100.0		100.0	

¹Standard errors not available for 1991 and 1995.

*Mastitis included in "other causes" in 1991 and 1995 and in "other diseases" in 2000.

**Lameness/injury included in "other causes" in 1991, 1995, and 2000.

*** "Other diseases" and "unknown causes" included in "other causes" in 1991 and 1995.

e. For **other** operations, percentage of cattle death losses by cause and by year:

Percent Nonpredator Loss						
Nonpredator Cause	1991 ¹	1995 ¹	2000	Std. Error	2005	Std. Error
Digestive problems (bloat, scours, parasites, enterotoxemia, acidosis, etc.)	15.6	12.1	14.9	(0.9)	18.8	(2.9)
Respiratory problems (pneumonia, shipping fever, etc.)	58.4	67.2	64.1	(2.5)	54.2	(2.7)
Metabolic problems (milk fever, grass tetany, etc.)	N/A	N/A	1.2	(0.1)	0.6	(0.1)
Weather-related causes (chilling, drowning, lightning, etc.)	6.9	3.3	3.5	(0.3)	6.0	(1.5)
Theft (stolen)	0.3	0.01	0.1	(0.0)	0.4	(0.2)
Poisoning (nitrate poisoning, noxious feeds, noxious weeds, etc.)	0.7	0.01	0.6	(0.3)	0.4	(0.2)
Calving problems	1.7	1.5	3.6	(2.0)	1.4	(0.6)
Mastitis*	N/A	N/A	N/A	(--)	0.1	(0.0)
Lameness/injury**	N/A	N/A	N/A	(--)	3.7	(1.0)
Other diseases***	N/A	N/A	1.6	(0.4)	3.9	(0.9)
Other causes***	16.4	15.9	3.5	(0.8)	2.4	(0.5)
Unknown causes***	N/A	N/A	6.9	(0.9)	8.1	(1.0)
Total	100.0	100.0	100.0		100.0	

¹Standard errors not available for 1991 and 1995.

*Mastitis included in "other causes" in 1991 and 1995 and in "other diseases" in 2000.

**Lameness/injury included in "other causes" in 1991, 1995, and 2000.

*** "Other diseases" and "unknown causes" included in "other causes" in 1991 and 1995.

3. Cause of loss by size of operation—2005

The percentage of losses due to calving problems decreased as the size of operation increased, ranging from 18.9 percent on operations with 1 to 49 cows to 10.7 percent on operations with 500 or more cows.

a. For **beef** operations, percentage of cattle death losses by cause and by size of operation:

Nonpredator Cause	Percent Nonpredator Loss							
	Size of Operation (Number of Cows)							
	1-49		50-99		100-499		500 or More	
	Std. Pct. Error	Std. Pct. Error	Std. Pct. Error	Std. Pct. Error	Std. Pct. Error	Std. Pct. Error	Std. Pct. Error	
Digestive problems (bloat, scours, parasites, enterotoxemia, acidosis, etc.)	8.2	(1.0)	7.4	(1.0)	7.1	(0.7)	6.8	(0.8)
Respiratory problems (pneumonia, shipping fever, etc.)	12.9	(1.3)	14.8	(1.7)	26.6	(4.5)	21.0	(1.9)
Metabolic problems (milk fever, grass tetany, etc.)	1.3	(0.3)	1.8	(0.4)	0.7	(0.1)	1.1	(0.2)
Weather-related causes (chilling, drowning, lightning, etc.)	8.0	(0.9)	10.4	(1.5)	10.0	(1.4)	8.0	(0.8)
Theft (stolen)	0.4	(0.1)	0.8	(0.3)	1.1	(0.4)	2.6	(1.1)
Poisoning (nitrate poisoning, noxious feeds, noxious weeds, etc.)	2.3	(0.9)	2.1	(0.7)	1.8	(0.3)	3.3	(0.4)
Calving problems	18.9	(1.2)	12.6	(1.0)	11.0	(1.1)	10.7	(0.9)
Mastitis	0.6	(0.2)	0.5	(0.2)	0.4	(0.1)	0.6	(0.2)
Lameness/injury	4.0	(0.5)	3.4	(0.5)	2.6	(0.4)	3.7	(0.5)
Other diseases	4.7	(0.7)	5.5	(0.9)	5.8	(1.0)	6.0	(0.9)
Other	22.4	(1.6)	23.7	(1.7)	16.5	(1.5)	14.0	(1.6)
Unknown	16.3	(1.2)	17.0	(1.5)	16.4	(1.6)	22.2	(2.1)
Total	100.0		100.0		100.0		100.0	

Respiratory problems accounted for a higher percentage of losses on operations with 1-29 cows (31.6 percent) than on all other operation sizes. Mastitis accounted for 15.0 of nonpredator losses on operations with 500 or more cows and 3.9 percent of losses on operations with 1 to 29 cows.

b. For *dairy* operations, percentage of cattle death losses by cause and by size of operation:

Nonpredator Cause	Percent Nonpredator Loss					
	Size of Operation (Number of Cows)					
	1-29	30-49	50-99	100-199	200-499	500 or More
	Std. Pct. Error	Std. Pct. Error	Std. Pct. Error	Std. Pct. Error	Std. Pct. Error	Std. Pct. Error
Digestive problems (bloat, scours, parasites, enterotoxemia, acidosis, etc.)	14.1 (5.0)	13.7 (3.1)	10.0 (1.1)	15.1 (2.9)	12.2 (1.0)	12.2 (0.8)
Respiratory problems (pneumonia, shipping fever, etc.)	31.6 (5.4)	8.3 (2.0)	8.9 (1.1)	13.3 (1.3)	11.6 (1.0)	17.0 (1.5)
Metabolic problems (milk fever, grass tetany, etc.)	9.2 (3.9)	10.3 (1.6)	10.7 (1.2)	8.9 (1.3)	11.2 (1.3)	9.5 (0.7)
Weather-related causes (chilling, drowning, lightning, etc.)	1.1 (0.7)	1.0 (0.5)	2.3 (0.7)	2.1 (0.4)	2.0 (0.4)	2.0 (0.4)
Theft (stolen)	0.0 (0.0)	0.0 (0.0)	0.3 (0.3)	0.0 (0.0)	0.0 (0.0)	0.1 (0.1)
Poisoning (nitrate poisoning, noxious feeds, noxious weeds, etc.)	0.3 (0.2)	0.5 (0.3)	1.5 (0.6)	0.4 (0.2)	0.4 (0.1)	0.2 (0.1)
Calving problems	9.0 (2.8)	19.2 (2.4)	17.0 (2.2)	14.0 (1.1)	13.3 (1.3)	9.9 (0.7)
Mastitis	3.9 (2.0)	9.6 (2.0)	10.6 (1.5)	9.9 (1.0)	13.0 (1.1)	15.0 (0.9)
Lameness/injury	8.7 (2.3)	11.2 (1.8)	13.2 (1.3)	10.2 (1.0)	13.0 (1.3)	9.0 (0.6)
Other diseases	7.8 (3.9)	9.9 (2.6)	8.1 (1.5)	7.0 (1.0)	9.3 (1.8)	8.4 (2.7)
Other	8.7 (3.0)	7.2 (1.4)	7.3 (1.2)	9.3 (1.4)	5.3 (1.2)	5.0 (1.0)
Unknown	5.6 (2.9)	9.1 (1.8)	10.1 (1.6)	9.8 (1.3)	8.7 (1.3)	11.7 (1.3)
Total	100.0	100.0	100.0	100.0	100.0	100.0

Almost one-fifth of losses on mixed operations with 500 or more cows (19.3 percent) were due to metabolic problems, while only 2.5 percent of losses on operations 1 to 99 cows were due to metabolic problems.

c. For *mixed* operations, percentage of cattle death losses by cause and by size of operation:

Nonpredator Cause	Percent Nonpredator Loss					
	Size of Operation (Number of Cows)					
	1-99		100-499		500 or More	
	Pct.	Std. Error	Pct.	Std. Error	Pct.	Std. Error
Digestive problems (bloat, scours, parasites, enterotoxemia, acidosis, etc.)	9.2	(2.8)	7.4	(1.4)	7.1	(3.2)
Respiratory problems (pneumonia, shipping fever, etc.)	14.4	(4.5)	16.3	(2.3)	14.8	(7.4)
Metabolic problems (milk fever, grass tetany, etc.)	2.5	(1.0)	8.2	(1.4)	19.3	(4.0)
Weather-related causes (chilling, drowning, lightning, etc.)	7.2	(2.5)	4.8	(1.2)	0.9	(0.5)
Theft (stolen)	0.0	(0.0)	0.0	(0.0)	0.2	(0.2)
Poisoning (nitrate poisoning, noxious feeds, noxious weeds, etc.)	1.5	(1.4)	1.3	(0.5)	0.0	(0.0)
Calving problems	21.6	(8.3)	16.1	(2.1)	5.9	(2.7)
Mastitis	3.4	(1.3)	8.6	(2.2)	27.2	(9.5)
Lameness/injury	8.0	(2.3)	10.8	(2.6)	4.5	(2.3)
Other diseases (mastitis, etc.)	5.2	(1.7)	6.9	(2.6)	1.9	(1.0)
Other	6.5	(2.0)	9.6	(2.0)	12.2	(4.8)
Unknown	20.5	(11.2)	10.0	(2.5)	6.0	(3.1)
Total	100.0		100.0		100.0	

Respiratory problems accounted for the majority of nonpredator losses on “other” operations. Operations with 1 to 99 head had a smaller percentage of losses due to respiratory problems compared to all other operation sizes.

d. For **other** operations, percentage of cattle death losses by cause and by size of operation:

Nonpredator Cause	Percent Nonpredator Loss							
	Size of Operation (Number of Cattle and Calves)							
	1-99		100-499		500-999		1,000 or More	
	Std. Pct.	Std. Error	Std. Pct.	Std. Error	Std. Pct.	Std. Error	Std. Pct.	Std. Error
Digestive problems (bloat, scours, parasites, enterotoxemia, acidosis, etc.)	9.3	(2.2)	5.5	(1.1)	10.6	(2.0)	24.0	(3.8)
Respiratory problems (pneumonia, shipping fever, etc.)	33.9	(5.3)	66.1	(4.3)	66.3	(3.6)	55.5	(3.5)
Metabolic problems (milk fever, grass tetany, etc.)	2.1	(0.8)	0.9	(0.5)	1.0	(0.4)	0.2	(0.1)
Weather-related causes (chilling, drowning, lightning, etc.)	12.5	(7.7)	2.5	(0.9)	5.4	(1.5)	5.1	(1.2)
Theft (stolen)	0.6	(0.6)	2.4	(1.7)	0.7	(0.6)	0.0	(0.0)
Poisoning (nitrate poisoning, noxious feeds, noxious weeds, etc.)	1.9	(1.1)	0.0	(0.0)	0.5	(0.3)	0.2	(0.1)
Calving problems	8.8	(3.6)	0.5	(0.3)	0.4	(0.3)	0.0	(0.0)
Mastitis	0.3	(0.1)	0.0	(0.0)	0.1	(0.0)	0.0	(0.0)
Lameness/injury	9.0	(5.2)	3.2	(0.9)	3.2	(0.9)	2.6	(0.7)
Other diseases	3.6	(1.3)	5.2	(3.2)	3.5	(1.7)	3.8	(1.1)
Other	9.8	(2.3)	4.8	(1.9)	2.4	(0.9)	0.4	(0.2)
Unknown	8.2	(2.0)	8.9	(2.7)	5.9	(1.8)	8.2	(1.4)
Total	100.0		100.0		100.0		100.0	

4. Cause of loss by region—2005

On beef operations, the percentage of losses due to respiratory problems ranged from 8.5 percent in the Southeast region to 29.1 in the North Central region. The percentage of losses due to weather-related causes ranged from 4.7 percent in the Northwest region to 16.7 percent in the Southeast region.

a. For **beef** operations, percentage of cattle death losses by cause and by region:

Nonpredator Cause	Percent Nonpredator Loss					
	Region					
	North Central	Northeast	Northwest	South Central	Southeast	Southwest
	Std. Pct. Error	Std. Pct. Error	Std. Pct. Error	Std. Pct. Error	Std. Pct. Error	Std. Pct. Error
Digestive problems (bloat, scours, parasites, entero- toxemia, acidosis, etc.)	9.4 (1.0)	9.9 (1.2)	9.2 (1.2)	6.0 (0.9)	7.0 (1.0)	9.6 (1.5)
Respiratory problems (pneumonia, shipping fever, etc.)	29.1 (2.1)	16.3 (2.0)	14.5 (1.6)	20.0 (3.6)	8.5 (1.1)	17.0 (2.0)
Metabolic problems (milk fever, grass tetany, etc.)	0.7 (0.3)	1.8 (0.4)	2.5 (1.0)	0.9 (0.2)	1.7 (0.4)	0.3 (0.1)
Weather-related causes (chilling, drowning, lightning, etc.)	11.2 (1.0)	6.9 (1.0)	4.7 (0.6)	7.5 (0.9)	16.7 (2.5)	5.8 (0.9)
Theft (stolen)	0.5 (0.2)	0.3 (0.1)	0.9 (0.4)	1.1 (0.4)	0.1 (0.0)	3.3 (1.0)
Poisoning (nitrate poisoning, noxious feeds, noxious weeds, etc.)	0.8 (0.3)	1.8 (0.6)	4.5 (0.6)	1.7 (0.7)	1.9 (0.5)	8.5 (1.8)
Calving problems	7.8 (1.7)	17.5 (1.5)	7.1 (0.8)	15.7 (1.2)	18.5 (1.3)	10.6 (1.7)
Mastitis	0.3 (0.1)	0.2 (0.1)	0.3 (0.2)	0.4 (0.1)	0.9 (0.3)	1.4 (0.6)
Lameness/injury	5.0 (0.7)	4.8 (0.6)	4.8 (0.7)	2.2 (0.3)	4.2 (0.8)	3.1 (0.6)
Other diseases	6.5 (1.4)	3.5 (0.5)	7.7 (2.3)	5.7 (0.8)	3.7 (0.8)	4.4 (0.8)
Other	14.4 (1.3)	16.8 (1.4)	17.8 (1.5)	22.4 (1.7)	21.8 (1.5)	15.7 (1.6)
Unknown	14.3 (1.6)	20.2 (1.7)	26.0 (3.0)	16.4 (1.4)	15.0 (1.3)	20.3 (2.0)
Total	100.0	100.0	100.0	100.0	100.0	100.0

On dairy operations, the percentage of losses due to mastitis ranged from 7.7 percent in the South Central region to 16.9 percent in the Southwest region.

b. For *dairy* operations, percentage of cattle death losses by cause and by region:

Nonpredator Cause	Percent Nonpredator Loss											
	Region											
	North Central		Northeast		Northwest		South Central		Southeast		Southwest	
	Std. Pct.	Std. Error	Std. Pct.	Std. Error	Std. Pct.	Std. Error	Std. Pct.	Std. Error	Std. Pct.	Std. Error	Std. Pct.	Std. Error
Digestive problems (bloat, scours, parasites, enterotoxemia, acidosis, etc.)	14.7	(2.0)	11.2	(0.8)	10.6	(1.7)	10.7	(1.8)	16.3	(6.9)	14.2	(1.0)
Respiratory problems (pneumonia, shipping fever, etc.)	12.0	(1.4)	11.9	(0.8)	19.0	(5.2)	14.6	(2.4)	10.8	(1.2)	18.7	(1.6)
Metabolic problems (milk fever, grass tetany, etc.)	8.8	(1.2)	11.3	(0.8)	10.5	(1.3)	5.6	(1.2)	7.2	(1.2)	10.0	(1.1)
Weather-related causes (chilling, drowning, lightning, etc.)	1.6	(0.4)	1.6	(0.3)	0.2	(0.1)	2.5	(0.7)	5.5	(0.9)	2.7	(0.8)
Theft (stolen)	0.0	(0.0)	0.1	(0.1)	0.2	(0.1)	0.5	(0.4)	0.0	(0.0)	0.0	(0.0)
Poisoning (nitrate poisoning, noxious feeds, noxious weeds, etc.)	0.3	(0.1)	0.7	(0.2)	0.0	(0.0)	1.2	(0.5)	0.1	(0.1)	0.1	(0.0)
Calving problems	12.7	(1.1)	12.9	(0.7)	8.9	(1.5)	16.2	(3.8)	14.6	(1.6)	11.2	(1.0)
Mastitis	11.8	(1.1)	13.1	(0.9)	9.5	(1.4)	7.7	(1.5)	12.3	(1.7)	16.9	(1.3)
Lameness/injury	11.6	(1.6)	12.9	(0.7)	9.1	(1.3)	6.2	(1.2)	10.8	(1.5)	8.2	(0.8)
Other diseases	6.3	(1.0)	8.0	(0.9)	5.9	(1.0)	20.5	(10.3)	4.5	(1.1)	6.1	(1.4)
Other	11.2	(1.8)	5.1	(0.5)	3.2	(0.6)	10.8	(2.7)	6.0	(1.1)	5.2	(1.9)
Unknown	9.0	(2.0)	11.2	(1.2)	22.9	(3.6)	3.5	(0.8)	11.9	(2.2)	6.7	(0.9)
Total	100.0		100.0		100.0		100.0		100.0		100.0	

On mixed operations, metabolic problems such as milk fever or grass tetany accounted for 22.5 percent of losses in the Southwest region and 2.9 percent of losses in the North Central region.

c. For *mixed* operations, percentage of cattle death losses by cause and by region:

Nonpredator Cause	Percent Nonpredator Loss					
	Region					
	North Central	Northeast	Northwest	South Central	Southeast	Southwest
	Std. Pct. Error	Std. Pct. Error	Std. Pct. Error	Std. Pct. Error	Std. Pct. Error	Std. Pct. Error
Digestive problems (bloat, scours, parasites, enterotoxemia, acidosis, etc.)	8.0 (3.2)	9.9 (2.1)	9.3 (1.8)	9.9 (2.7)	8.6 (1.9)	3.9 (2.7)
Respiratory problems (pneumonia, shipping fever, etc.)	29.8 (10.0)	13.9 (3.1)	16.8 (3.5)	21.1 (4.9)	17.8 (4.1)	4.4 (3.1)
Metabolic problems (milk fever, grass tetany, etc.)	2.9 (1.1)	8.0 (1.9)	12.4 (4.2)	10.4 (3.6)	4.6 (1.3)	22.5 (3.7)
Weather-related causes (chilling, drowning, lightning, etc.)	9.3 (3.1)	2.7 (0.9)	2.8 (0.9)	3.2 (1.7)	9.0 (2.8)	0.2 (0.2)
Theft (stolen)	0.0 (0.0)	0.0 (0.0)	0.0 (0.0)	0.0 (0.0)	0.0 (0.0)	0.2 (0.2)
Poisoning (nitrate poisoning, noxious feeds, noxious weeds, etc.)	0.4 (0.4)	0.9 (0.6)	0.4 (0.4)	2.4 (1.7)	0.7 (0.6)	0.1 (0.0)
Calving problems	9.2 (3.1)	13.5 (3.0)	11.3 (3.0)	24.4 (9.1)	22.0 (3.4)	4.7 (3.1)
Mastitis	4.4 (1.2)	8.5 (2.0)	7.9 (2.9)	8.5 (4.1)	7.6 (1.5)	35.3 (9.1)
Lameness/injury	8.2 (2.3)	13.1 (3.7)	11.2 (3.7)	4.5 (1.5)	7.9 (1.9)	3.3 (2.8)
Other diseases	9.4 (5.3)	4.0 (1.2)	4.2 (1.8)	3.4 (1.4)	2.6 (0.8)	2.8 (2.1)
Other	10.4 (3.9)	6.2 (1.6)	7.0 (2.3)	7.6 (2.7)	7.6 (2.2)	15.7 (4.9)
Unknown	8.0 (3.3)	19.3 (9.9)	16.7 (5.9)	4.6 (1.4)	11.6 (2.0)	6.9 (5.5)
Total	100.0	100.0	100.0	100.0	100.0	100.0

Across all regions, respiratory problems accounted for the highest percentage of nonpredator losses on “other” operations, ranging from 34.7 percent in the Southwest region to 62.3 percent in the North Central region.

d. For **other** operations, percentage of cattle death losses by cause and by region:

Nonpredator Cause	Percent Nonpredator Loss											
	Region											
	North Central		Northeast		Northwest		South Central		Southeast		Southwest	
	Std. Pct.	Error	Std. Pct.	Error	Std. Pct.	Error	Std. Pct.	Error	Std. Pct.	Error	Std. Pct.	Error
Digestive problems (bloat, scours, parasites, enterotoxemia, acidosis, etc.)	13.9	(1.4)	13.7	(2.4)	10.3	(1.7)	23.9	(5.2)	1.9	(0.8)	17.3	(2.3)
Respiratory problems (pneumonia, shipping fever, etc.)	62.3	(2.4)	58.6	(4.3)	45.5	(4.6)	56.3	(5.0)	53.2	(12.4)	34.7	(2.1)
Metabolic problems (milk fever, grass tetany, etc.)	0.6	(0.3)	2.0	(0.8)	0.1	(0.1)	0.4	(0.2)	4.2	(3.3)	0.1	(0.1)
Weather-related causes (chilling, drowning, lightning, etc.)	11.3	(2.2)	2.6	(0.8)	1.5	(0.4)	6.1	(2.9)	3.7	(2.2)	0.5	(0.2)
Theft (stolen)	0.0	(0.0)	0.0	(0.0)	0.1	(0.1)	0.6	(0.4)	3.6	(3.4)	0.0	(0.0)
Poisoning (nitrate poisoning, noxious feeds, noxious weeds, etc.)	0.4	(0.3)	1.0	(1.0)	0.9	(0.4)	0.2	(0.2)	4.8	(4.8)	0.0	(0.0)
Calving problems	0.4	(0.2)	0.8	(0.4)	0.9	(0.5)	1.9	(1.1)	10.0	(8.1)	0.1	(0.1)
Mastitis	0.1	(0.0)	0.5	(0.3)	0.0	(0.0)	0.0	(0.0)	0.0	(0.0)	0.0	(0.0)
Lameness/injury	3.2	(0.5)	4.0	(0.7)	6.6	(2.6)	2.9	(1.8)	0.7	(0.3)	6.4	(1.9)
Other diseases	1.9	(0.5)	6.1	(4.5)	5.5	(1.5)	1.2	(0.5)	0.1	(0.1)	15.3	(2.7)
Other	1.4	(0.4)	3.7	(1.4)	7.1	(3.9)	2.7	(0.8)	6.7	(2.5)	0.7	(0.3)
Unknown	4.5	(0.9)	7.0	(1.7)	21.5	(3.7)	3.8	(1.3)	11.1	(7.3)	24.9	(2.9)
Total	100.0		100.0		100.0		100.0		100.0		100.0	

C. Nonpredator Calf Death Loss

1. Losses as a percentage of calf crop—2005


Overall, 5.8 percent of the calf crop was lost to nonpredator causes. Losses were highest in the Northeast and Southwest regions (7.9 and 7.5 percent of calf crop, respectively).

a. Percentage of calf crop (born during 2005) lost,* by region:

Region	Percent Calf Crop*	
	Percent	Standard Error
North Central	5.8	(0.1)
Northeast	7.9	(0.4)
Northwest	4.6	(0.1)
South Central	5.1	(0.2)
Southeast	4.3	(0.1)
Southwest	7.5	(1.1)
All operations	5.8	(0.1)

*Calf deaths in 2005 / 2005 calf crop x 100.

Percentage of 2005 Calf Crop lost, by Region



On beef operations, 4.5 percent of the calf crop was lost to nonpredator causes in 2005. The lowest percentage of losses (3.0 percent) occurred on operations with 500 or more cows.

b. For **beef** operations, percentage of calf crop (born during 2005) lost* by region and by size of operation:

Percent Calf Crop										
Size of Operation (Number of Cows)										
Region	1-49		50-99		100-499		500 or More		All Operations	
	Pct.	Std. Error	Pct.	Std. Error	Pct.	Std. Error	Pct.	Std. Error	Pct.	Std. Error
North Central	7.3	(0.6)	5.7	(0.3)	4.0	(0.1)	3.2	(0.2)	4.7	(0.1)
Northeast	6.4	(0.3)	5.5	(0.4)	4.6	(0.3)	3.0	(0.2)	5.7	(0.2)
Northwest	6.2	(0.8)	4.5	(0.4)	3.5	(0.1)	3.6	(0.2)	3.9	(0.1)
South Central	5.4	(0.3)	4.5	(0.2)	4.6	(0.3)	2.8	(0.4)	4.6	(0.1)
Southeast	4.4	(0.2)	4.0	(0.2)	3.3	(0.2)	2.1	(0.1)	3.8	(0.1)
Southwest	8.1	(1.3)	5.1	(0.6)	3.9	(0.3)	2.9	(0.3)	4.3	(0.2)
All Operations	5.7	(0.2)	4.8	(0.1)	4.1	(0.1)	3.0	(0.1)	4.5	(0.1)

*Calf deaths in 2005 / 2005 calf crop x 100.

Overall, 7.1 percent of the calf crop on dairy operations was lost to nonpredator causes in 2005. The percentage of calf crop lost varied across regions, ranging from 4.5 percent in the Southwest region to 10.5 percent in the North Central region.

c. For **dairy** operations, percentage of calf crop (born during 2005) lost* by region and by size of operation:

Percent Calf Crop														
Size of Operation (Number of Cows)														
Region	1-29		30-49		50-99		100-199		200-499		500 or More		All Operations	
	Pct.	Std. Err.	Pct.	Std. Err.	Pct.	Std. Err.	Pct.	Std. Err.	Pct.	Std. Err.	Pct.	Std. Err.	Pct.	Std. Err.
North Central	10.1	(2.8)	11.1	(1.5)	11.8	(0.8)	12.0	(1.1)	8.9	(0.8)	7.7	(1.4)	10.5	(0.5)
Northeast	12.1	(2.4)	7.7	(0.5)	9.2	(0.4)	8.6	(0.5)	6.7	(0.4)	4.7	(0.3)	7.7	(0.2)
Northwest	5.9	(2.8)	11.4	(1.4)	8.8	(0.9)	8.1	(0.7)	6.5	(0.5)	5.5	(0.3)	5.9	(0.3)
South Central	3.0	(1.5)	8.3	(1.8)	10.9	(2.0)	9.8	(1.0)	9.6	(1.1)	7.6	(0.7)	8.4	(0.6)
Southeast	**		12.3	(2.3)	9.5	(1.1)	11.5	(2.6)	8.9	(0.7)	6.8	(0.5)	8.5	(0.7)
Southwest	**		**		11.0	(1.2)	8.6	(1.2)	6.1	(0.5)	4.3	(0.3)	4.5	(0.3)
All Operations	9.5	(1.5)	8.3	(0.5)	9.6	(0.4)	9.4	(0.4)	7.1	(0.3)	5.0	(0.2)	7.1	(0.1)

*Calf deaths in 2005 / 2005 calf crop x 100.

**Too few operations to report.

Overall, 8.7 percent of the calf crop on mixed operations was lost to nonpredator causes.

d. For *mixed* operations, percentage of calf crop (born during 2005) lost* by region and by size of operation:

Region	Percent Calf Crop							
	Size of Operation (Number of Cows)							
	1-99		100-499		500 or More		All Operations	
	Pct.	Std. Error	Pct.	Std. Error	Pct.	Std. Error	Pct.	Std. Error
North Central	14.7	(3.1)	8.8	(1.2)	8.7	(1.8)	10.6	(1.3)
Northeast	12.4	(3.3)	9.0	(1.1)	7.0	(1.3)	10.0	(1.4)
Northwest	**		7.2	(1.8)	4.9	(0.5)	6.1	(0.9)
South Central	8.7	(2.8)	7.9	(1.0)	6.8	(1.9)	7.6	(1.0)
Southeast	10.1	(3.5)	7.3	(0.8)	5.6	(0.6)	6.9	(0.6)
Southwest	**		8.0	(1.7)	8.0	(2.7)	8.0	(2.1)
All Operations	12.2	(2.0)	8.3	(0.5)	7.1	(1.3)	8.7	(0.6)

*Calf deaths in 2005 / 2005 calf crop x 100.

**Too few operations to report.

Note: Very few calves were born on “other” operations. Therefore, inventory of calves weighing less than 500 lb was used in place of calf crop.

Overall, 11.6 percent of calf inventory was lost to nonpredator causes on “other” operations. On operations with fewer than 100 head, 25.1 percent of inventory was lost, compared to 3.9 percent of inventory on operations with 500-999 head. Regionally, losses ranged from 8.1 percent of calf inventory in the South Central region to 17.2 percent of inventory in the North Central region.

e. For **other** operations, percentage of **calf inventory** lost* by region and by size of operation:

Percent Inventory										
Size of Operation (Number of Cattle and Calves)										
Region	1-99		100-499		500-999		1,000 or More		All Operations	
	Pct.	Std. Error	Pct.	Std. Error	Pct.	Std. Error	Pct.	Std. Error	Pct.	Std. Error
North Central	26.3	(6.2)	8.2	(2.4)	10.5	(3.6)	24.2	(14.9)	17.2	(3.2)
Northeast	34.9	(15.7)	8.4	(1.4)	3.5	(0.8)	1.9	(1.5)	9.1	(4.1)
Northwest	29.2	(11.0)	13.9	(4.4)	5.0	(3.3)	13.5	(2.1)	14.5	(2.0)
South Central	14.8	(4.1)	11.7	(4.5)	3.1	(1.0)	7.3	(1.6)	8.1	(1.3)
Southeast	14.8	(5.7)	7.8	(3.8)	2.1	(0.7)	8.9	(3.3)	10.6	(2.4)
Southwest	26.3	(13.4)	16.7	(8.0)	3.3	(1.7)	16.0	(4.1)	16.1	(3.8)
All Operations	25.1	(6.3)	9.6	(1.4)	3.9	(0.7)	10.6	(3.1)	11.6	(2.0)

*2005 death loss divided by end-of-year (January 1, 2006) inventory of calves 500 lbs or less X100.

2. Cause of loss—1991, 1995, 2000, and 2005

The highest percentages of nonpredator calf losses in 2005 were due to respiratory problems (31.8 percent) and digestive problems (21.2 percent).


a. Percentage of calf death losses, by cause and by year:

Percent Nonpredator Loss				
Nonpredator Cause	1991	1995	2000	2005
Digestive problems (bloat, scours, parasites, enterotoxemia, acidosis, etc.)	26.0	25.2	19.8	21.2
Respiratory problems (pneumonia, shipping fever, etc.)	35.2	29.6	30.0	31.8
Metabolic problems (milk fever, grass tetany, etc.)	N/A	N/A	0.8	0.8
Weather-related causes (chilling, drowning, lightning, etc.)	6.2	10.7	10.6	7.6
Theft (stolen)	0.6	0.4	0.4	0.5
Poisoning (nitrate poisoning, noxious feeds, noxious weeds, etc.)	0.8	0.6	0.5	0.7
Calving problems	13.2	15.1	16.7	17.7
Lameness/injury*	N/A	N/A	N/A	1.9
Other diseases	N/A	N/A	3.2	3.5
Other causes**	18.0	18.4	3.2	2.8
Unknown causes**	N/A	N/A	14.8	11.5
Total	100.0	100.0	100.0	100.0

*Lameness/injury included with other causes in 1991, 1995, and 2000.

**Other causes and unknown causes combined in 1991 and 1995.

Percentage of 2005 Calf Death Losses, by Cause



*Lameness/injury included with other causes in 1991, 1995, and 2000.

**Other causes and unknown causes combined in 1991 and 1995.

On beef operations in 2005, the two most common causes of calf losses were respiratory problems and calving problems (27.7 and 24.2 percent of losses, respectively).

b. For **beef** operations, percentage of calf death losses by cause and by year:

Percent Nonpredator Loss						
Nonpredator Cause	1991 ¹	1995 ¹	2000	Std. Error	2005	Std. Error
Digestive problems (bloat, scours, parasites, enterotoxemia, acidosis, etc.)	19.2	17.4	13.8	(0.5)	13.0	(0.5)
Respiratory problems (pneumonia, shipping fever, etc.)	25.6	23.5	23.2	(0.7)	27.7	(0.8)
Metabolic problems (milk fever, grass tetany, etc.)	N/A	N/A	0.7	(0.1)	0.6	(0.1)
Weather-related causes (chilling, drowning, lightning, etc.)	10.8	14.7	13.5	(0.5)	10.2	(0.5)
Theft (stolen)	1.0	0.4	0.6	(0.1)	0.7	(0.2)
Poisoning (nitrate poisoning, noxious feeds, noxious weeds, etc.)	1.2	0.8	0.7	(0.1)	0.9	(0.1)
Calving problems	20.4	20.4	23.5	(0.6)	24.2	(0.7)
Lameness/injury*	N/A	N/A	N/A	(--)	1.7	(0.2)
Other diseases**	N/A	N/A	2.1	(0.3)	3.5	(0.3)
Other causes**	21.8	22.8	3.5	(0.3)	3.5	(0.3)
Unknown causes**	N/A	N/A	18.4	(0.6)	14.0	(0.5)
Total	100.0	100.0	100.0		100.0	

¹Standard errors not available for 1991 and 1995.

*Lameness/injury included in "other causes" in 1991, 1995, and 2000.

** "Other diseases" and "unknown causes" included in "other causes" in 1991 and 1995.

On dairy operations in 2005, digestive and respiratory problems were the most common causes of calf loss (35.8 and 33.2 percent of losses, respectively).

c. For *dairy* operations, percentage of calf death losses by cause and by year:

Percent Nonpredator Loss						
Nonpredator Cause	1991 ¹	1995 ¹	2000	Std. Error	2005	Std. Error
Digestive problems (bloat, scours, parasites, enterotoxemia, acidosis, etc.)	36.6	40.5	35.4	(1.4)	35.8	(1.0)
Respiratory problems (pneumonia, shipping fever, etc.)	33.0	35.3	30.7	(1.3)	33.2	(0.9)
Metabolic problems (milk fever, grass tetany, etc.)	N/A	N/A	1.1	(0.1)	1.5	(0.4)
Weather-related causes (chilling, drowning, lightning, etc.)	2.5	3.9	6.8	(3.2)	2.2	(0.2)
Theft (stolen)	0.2	0.1	0.1	(0.0)	0.1	(0.0)
Poisoning (nitrate poisoning, noxious feeds, noxious weeds, etc.)	0.5	0.2	0.2	(0.0)	0.3	(0.2)
Calving problems	10.6	8.3	9.8	(0.6)	10.4	(0.6)
Lameness/injury*	N/A	N/A	N/A	(--)	1.5	(0.2)
Other diseases**	N/A	N/A	2.5	(0.3)	4.4	(0.7)
Other causes**	16.6	11.7	3.1	(0.9)	2.0	(0.7)
Unknown causes**	N/A	N/A	10.3	(0.6)	8.6	(0.6)
Total	100.0	100.0	100.0		100.0	

¹Standard errors not available for 1991 and 1995.

*Lameness/injury included in "other causes" in 1991, 1995, and 2000.

** "Other diseases" and "unknown causes" included in "other causes" in 1991 and 1995.

On mixed operations in 2005, digestive and respiratory problems were the most common causes of calf loss (28.9 and 30.0 percent of losses, respectively).

d. For *mixed* operations, percentage of calf death losses by cause and by year:

Percent Nonpredator Loss						
Nonpredator Cause	1991 ¹	1995 ¹	2000	Std. Error	2005	Std. Error
Digestive problems (bloat, scours, parasites, enterotoxemia, acidosis, etc.)	37.5	41.3	31.0	(2.1)	28.9	(4.6)
Respiratory problems (pneumonia, shipping fever, etc.)	34.9	41.5	34.1	(2.0)	30.0	(3.9)
Metabolic problems (milk fever, grass tetany, etc.)	N/A	N/A	1.5	(0.7)	1.2	(0.7)
Weather-related causes (chilling, drowning, lightning, etc.)	4.1	3.1	6.8	(0.9)	7.1	(4.6)
Theft (stolen)	0.4	0.0	0.0	(0.0)	0.1	(0.1)
Poisoning (nitrate poisoning, noxious feeds, noxious weeds, etc.)	0.2	0.0	0.0	(0.0)	0.0	(0.0)
Calving problems	11.3	5.3	10.7	(1.4)	11.8	(1.9)
Lameness/injury*	N/A	N/A	N/A	(--)	3.1	(1.8)
Other diseases**	N/A	N/A	1.3	(0.5)	3.4	(1.1)
Other causes**	N/A	N/A	1.8	(0.6)	2.3	(0.9)
Unknown causes**	11.6	8.8	12.8	(2.4)	12.1	(3.4)
Total	100.0	100.0	100.0		100.0	

¹Standard errors not available for 1991 and 1995.

*Lameness/injury included in "other causes" in 1991, 1995, and 2000.

** "Other diseases" and "unknown causes" included in "other causes" in 1991 and 1995.

On “other” operations, digestive and respiratory problems accounted for the highest percentage of calf losses (32.5 and 47.1 percent of losses, respectively). The percentage of calf losses due to respiratory causes was higher on other operations than on beef, dairy, or mixed operations.

e. For **other** operations, percentage of calf death losses by cause and by year:


Percent Nonpredator Loss						
Nonpredator Cause	1991 ¹	1995 ¹	2000	Std. Error	2005	Std. Error
Digestive problems (bloat, scours, parasites, enterotoxemia, acidosis, etc.)	16.0	27.7	15.1	(3.3)	32.5	(5.4)
Respiratory problems (pneumonia, shipping fever, etc.)	65.9	61.7	56.5	(9.2)	47.1	(3.0)
Metabolic problems (milk fever, grass tetany, etc.)	N/A	N/A	0.3	(0.2)	0.8	(0.4)
Weather-related causes (chilling, drowning, lightning, etc.)	3.1	2.5	5.6	(2.3)	4.5	(1.3)
Theft (stolen)	0.5	<0.01	0.0	(0.0)	0.2	(0.1)
Poisoning (nitrate poisoning, noxious feeds, noxious weeds, etc.)	0.4	<0.01	0.3	(0.2)	0.8	(0.5)
Calving problems	0.7	1.8	1.7	(0.5)	1.8	(0.4)
Lameness/injury*	N/A	N/A	N/A	(--)	3.4	(1.5)
Other diseases**	N/A	N/A	9.9	(5.3)	2.3	(0.7)
Other causes**	13.4	6.3	2.5	(1.2)	1.0	(0.4)
Unknown causes**	N/A	N/A	8.1	(2.9)	5.6	(1.3)
Total	100.0	100.0	100.0		100.0	

¹Standard errors not available for 1991 and 1995.

*Lameness/injury included in “other causes” in 1991, 1995, and 2000.

** “Other diseases” and “unknown causes” included in “other causes” in 1991 and 1995.

Percentage of 2005 Calf Death Losses, by Cause



3. Cause of loss by size of operation—2005

Calving problems accounted for 31.4 percent of losses on beef operations with 1 to 49 cows and 19.5 percent of losses on operations with 500 or more cows.

a. For **beef** operations, percentage of calf death losses by cause and by size of operation:

Nonpredator Cause	Percent Nonpredator Loss							
	Size of Operation (Number of Cows)							
	1-49		50-99		100-499		500 or More	
	Std. Pct.	Std. Error	Std. Pct.	Std. Error	Std. Pct.	Std. Error	Std. Pct.	Std. Error
Digestive problems (bloat, scours, parasites, enterotoxemia, acidosis, etc.)	11.2	(0.9)	14.7	(1.1)	13.1	(0.8)	16.0	(1.8)
Respiratory problems (pneumonia, shipping fever, etc.)	19.7	(1.2)	26.8	(1.4)	35.5	(1.6)	30.6	(1.8)
Metabolic problems (milk fever, grass tetany, etc.)	0.8	(0.2)	0.5	(0.2)	0.4	(0.1)	0.5	(0.2)
Weather-related causes (chilling, drowning, lightning, etc.)	8.7	(0.7)	10.8	(1.2)	11.0	(0.8)	10.8	(1.4)
Theft (stolen)	0.8	(0.3)	0.7	(0.3)	0.3	(0.1)	2.2	(1.0)
Poisoning (nitrate poisoning, noxious feeds, noxious weeds, etc.)	0.8	(0.2)	0.7	(0.2)	1.0	(0.2)	1.2	(0.3)
Calving problems	31.4	(1.4)	20.6	(1.2)	20.4	(0.9)	19.5	(2.6)
Lameness/injury	2.7	(0.4)	1.0	(0.3)	1.2	(0.2)	1.0	(0.2)
Other diseases	3.5	(0.5)	5.1	(1.2)	2.7	(0.4)	3.6	(0.8)
Other	5.1	(0.7)	3.8	(0.7)	2.1	(0.3)	2.1	(0.8)
Unknown	15.3	(0.9)	15.3	(1.2)	12.3	(0.9)	12.5	(1.3)
Total	100.0		100.0		100.0		100.0	

The percentage of calf losses due to respiratory problems was higher on operations with 500 or more cows than on operations with 1-29 cows.

b. For *dairy* operations, percentage of calf death losses by cause and by size of operation:

Percent Nonpredator Loss						
Size of Operation (Number of Cows)						
	1-29	30-49	50-99	100-199	200-499	500 or More
Cause	Std. Pct. Error	Std. Pct. Error	Std. Pct. Error	Std. Pct. Error	Std. Pct. Error	Std. Pct. Error
Digestive problems (bloat, scours, parasites, enterotoxemia, acidosis, etc.)	34.5 (8.1)	35.7 (3.5)	38.6 (2.1)	40.6 (3.0)	38.9 (2.0)	29.9 (1.5)
Respiratory problems (pneumonia, shipping fever, etc.)	20.2 (4.8)	31.9 (3.0)	31.7 (2.1)	33.1 (2.3)	35.1 (1.7)	35.3 (1.4)
Metabolic problems (milk fever, grass tetany, etc.)	0.8 (0.8)	0.0 (0.0)	1.9 (1.5)	0.6 (0.2)	0.9 (0.3)	2.6 (0.8)
Weather-related causes (chilling, drowning, lightning, etc.)	1.3 (0.6)	4.4 (1.4)	1.5 (0.5)	2.2 (0.5)	1.5 (0.3)	2.6 (0.4)
Theft (stolen)	0.0 (0.0)	0.0 (0.0)	0.1 (0.1)	0.0 (0.0)	0.0 (0.0)	0.1 (0.1)
Poisoning (nitrate poisoning, noxious feeds, noxious weeds, etc.)	0.7 (0.7)	0.1 (0.1)	0.2 (0.1)	1.2 (1.1)	0.3 (0.3)	0.0 (0.0)
Calving problems	10.5 (3.2)	13.3 (2.2)	14.2 (1.7)	9.0 (1.2)	9.0 (1.2)	8.4 (0.8)
Lameness/injury	1.4 (0.7)	2.6 (0.8)	0.9 (0.2)	1.0 (0.2)	1.6 (0.4)	1.8 (0.3)
Other diseases	13.3(10.5)	2.6 (1.0)	3.0 (0.7)	6.3 (2.4)	4.5 (1.0)	3.6 (0.4)
Other	13.6(11.5)	1.1 (0.6)	1.4 (0.6)	0.7 (0.3)	0.5 (0.2)	2.6 (1.3)
Unknown	3.7 (1.9)	8.3 (1.7)	6.5 (0.9)	5.3 (0.9)	7.7 (1.5)	13.1 (1.5)
Total	100.0	100.0	100.0	100.0	100.0	100.0

Across all sizes of mixed operations, at least one-quarter of calf losses were due to respiratory problems. Operations with 500 or more cows had a disproportionately high percentage of weather-related losses.

c. For *mixed* operations, percentage of calf death losses by cause and by size of operation:

Nonpredator Cause	Percent Nonpredator Loss					
	Size of Operation (Number of Cows)					
	1-99		100-499		500 or More	
	Pct.	Std. Error	Pct.	Std. Error	Pct.	Std. Error
Digestive problems (bloat, scours, parasites, enterotoxemia, acidosis, etc.)	35.0	(10.4)	32.4	(3.6)	17.1	(6.6)
Respiratory problems (pneumonia, shipping fever, etc.)	30.7	(6.4)	32.3	(3.0)	25.4	(10.2)
Metabolic problems (milk fever, grass tetany, etc.)	0.2	(0.2)	0.6	(0.2)	3.1	(2.7)
Weather-related causes (chilling, drowning, lightning, etc.)	1.4	(1.0)	2.2	(0.9)	20.9	(11.9)
Theft (stolen)	0.1	(0.1)	0.0	(0.0)	0.4	(0.3)
Poisoning (nitrate poisoning, noxious feeds, noxious weeds, etc.)	0.0	(0.0)	0.1	(0.1)	0.0	(0.0)
Calving problems	10.1	(3.3)	11.3	(2.1)	14.4	(4.1)
Lameness/injury	1.0	(0.5)	0.8	(0.4)	9.0	(4.5)
Other diseases	3.7	(2.8)	4.1	(1.6)	1.8	(0.8)
Other	1.9	(1.5)	3.4	(1.6)	0.7	(0.7)
Unknown	15.9	(9.1)	12.8	(4.1)	7.2	(2.9)
Total	100.0		100.0		100.0	

For other operations, the causes of calf loss were similar across operation sizes.

d. For **other** operations, percentage of calf death losses by cause and by size of operation:

Nonpredator Cause	Percent Nonpredator Loss							
	Size of Operation (Number of Cattle and Calves)							
	1-99		100-499		500-999		1,000 or More	
	Std. Pct.	Std. Error	Std. Pct.	Std. Error	Std. Pct.	Std. Error	Std. Pct.	Std. Error
Digestive problems (bloat, scours, parasites, enterotoxemia, acidosis, etc.)	21.0	(3.9)	38.0	(5.7)	20.3	(4.4)	38.6	(8.8)
Respiratory problems (pneumonia, shipping fever, etc.)	46.6	(3.9)	53.4	(5.2)	58.4	(6.2)	44.5	(4.7)
Metabolic problems (milk fever, grass tetany, etc.)	1.2	(0.9)	2.1	(2.1)	0.8	(0.5)	0.1	(0.0)
Weather-related causes (chilling, drowning, lightning, etc.)	6.7	(1.4)	1.2	(0.8)	2.0	(0.9)	4.5	(2.2)
Theft (stolen)	0.1	(0.1)	0.0	(0.0)	2.8	(2.6)	0.0	(0.0)
Poisoning (nitrate poisoning, noxious feeds, noxious weeds, etc.)	2.7	(1.3)	0.0	(0.0)	0.4	(0.4)	0.0	(0.0)
Calving problems	5.1	(1.6)	0.3	(0.2)	0.4	(0.3)	0.4	(0.2)
Lameness/injury	3.9	(1.9)	0.8	(0.5)	0.5	(0.3)	4.1	(2.7)
Other diseases	2.9	(0.8)	0.4	(0.3)	7.8	(3.8)	2.2	(1.3)
Other	1.8	(0.9)	1.0	(0.8)	0.0	(0.0)	0.6	(0.5)
Unknown	8.0	(2.6)	2.8	(1.1)	6.6	(2.5)	5.0	(2.1)
Total	100.0		100.0		100.0		100.0	

4. Cause of loss by region—2005

On beef operations, the percentage of calf losses due to unknown causes ranged from 7.8 percent in the North Central region to 20.4 percent in the Southeast region.

a. For **beef** operations, percentage of calf death losses by cause and by region:

Nonpredator Cause	Percent Nonpredator Loss											
	Region											
	North Central		Northeast		Northwest		South Central		Southeast		Southwest	
	Pct.	Std. Error	Pct.	Std. Error	Pct.	Std. Error	Pct.	Std. Error	Pct.	Std. Error	Pct.	Std. Error
Digestive problems (bloat, scours, parasites, enterotoxemia, acidosis, etc.)	18.1	(1.2)	15.3	(1.3)	18.1	(1.7)	8.9	(0.8)	8.6	(0.9)	15.7	(2.1)
Respiratory problems (pneumonia, shipping fever, etc.)	29.0	(1.3)	22.5	(1.8)	25.8	(1.6)	33.0	(1.8)	19.6	(1.5)	23.8	(2.2)
Metabolic problems (milk fever, grass tetany, etc.)	0.4	(0.2)	0.9	(0.2)	0.4	(0.1)	0.4	(0.1)	1.2	(0.4)	0.3	(0.1)
Weather-related causes (chilling, drowning, lightning, etc.)	12.4	(1.1)	10.8	(1.3)	13.8	(1.2)	7.3	(0.7)	11.2	(1.4)	10.4	(1.6)
Theft (stolen)	0.1	(0.1)	0.8	(0.6)	0.7	(0.2)	0.7	(0.2)	0.6	(0.3)	2.9	(1.4)
Poisoning (nitrate poisoning, noxious feeds, noxious weeds, etc.)	0.3	(0.1)	0.7	(0.3)	1.4	(0.3)	1.1	(0.2)	0.7	(0.3)	1.4	(0.4)
Calving problems	24.0	(1.3)	22.8	(2.0)	22.2	(1.5)	24.4	(1.3)	27.6	(1.5)	24.3	(3.1)
Lameness/injury	1.1	(0.2)	2.2	(0.5)	1.6	(0.4)	1.8	(0.3)	2.4	(0.5)	0.6	(0.2)
Other diseases	3.3	(0.8)	3.3	(0.6)	2.8	(0.8)	3.8	(0.7)	3.4	(0.6)	4.4	(1.3)
Other	3.5	(0.7)	2.7	(0.6)	2.7	(0.4)	3.8	(0.7)	4.3	(0.7)	2.7	(0.7)
Unknown	7.8	(0.7)	18.0	(1.6)	10.5	(1.0)	14.8	(1.1)	20.4	(1.3)	13.5	(1.7)
Total	100.0		100.0		100.0		100.0		100.0		100.0	

For dairy operations, digestive problems accounted for a higher percentage of calf losses in the North Central region (40.7 percent) than in the South Central and Southwest regions (27.2 and 26.8 percent, respectively).

b. For *dairy* operations, percentage of calf death losses by cause and by region:

Nonpredator Cause	Percent Nonpredator Loss											
	Region											
	North Central		Northeast		Northwest		South Central		Southeast		Southwest	
	Std. Pct.	Std. Error	Std. Pct.	Std. Error	Std. Pct.	Std. Error	Std. Pct.	Std. Error	Std. Pct.	Std. Error	Std. Pct.	Std. Error
Digestive problems (bloat, scours, parasites, enterotoxemia, acidosis, etc.)	40.7	(2.7)	39.0	(1.4)	31.9	(2.4)	27.2	(2.9)	37.1	(8.2)	26.8	(2.1)
Respiratory problems (pneumonia, shipping fever, etc.)	30.0	(2.1)	32.9	(1.3)	32.3	(2.0)	41.4	(4.3)	25.4	(4.1)	36.5	(1.9)
Metabolic problems (milk fever, grass tetany, etc.)	1.0	(0.5)	0.6	(0.2)	0.9	(0.3)	6.2	(4.7)	1.0	(0.2)	3.6	(1.5)
Weather-related causes (chilling, drowning, lightning, etc.)	3.5	(0.8)	1.4	(0.2)	0.7	(0.2)	1.7	(0.6)	4.5	(1.6)	3.8	(0.9)
Theft (stolen)	0.0	(0.0)	0.0	(0.0)	0.0	(0.0)	0.6	(0.3)	0.0	(0.0)	0.2	(0.1)
Poisoning (nitrate poisoning, noxious feeds, noxious weeds, etc.)	0.0	(0.0)	0.7	(0.4)	0.1	(0.1)	0.3	(0.2)	0.1	(0.1)	0.0	(0.0)
Calving problems	16.9	(1.9)	8.9	(0.9)	9.9	(1.4)	13.7	(2.2)	9.5	(1.7)	8.5	(1.3)
Lameness/injury	1.0	(0.3)	1.8	(0.3)	1.1	(0.2)	0.8	(0.4)	1.5	(0.4)	1.4	(0.4)
Other diseases	1.8	(0.7)	4.9	(1.0)	3.5	(0.7)	2.7	(0.8)	11.3	(7.3)	3.8	(0.7)
Other	1.3	(0.4)	1.8	(1.0)	0.5	(0.1)	3.1	(1.9)	0.6	(0.4)	4.1	(2.7)
Unknown	3.8	(0.9)	8.0	(0.8)	19.1	(3.5)	2.3	(0.6)	9.0	(1.9)	11.3	(1.6)
Total	100.0		100.0		100.0		100.0		100.0		100.0	

The percentage of losses due to calving problems ranged from 7.5 percent in the Northeast region to 18.5 percent in the Southwest region. The Southwest region had a disproportionately high percentage of weather-related losses.

c. For *mixed* operations, percentage of calf death losses by cause and by region:

Nonpredator Cause	Percent Nonpredator Loss											
	Region											
	North Central		Northeast		Northwest		South Central		Southeast		Southwest	
	Pct.	Std. Error	Pct.	Std. Error	Pct.	Std. Error	Pct.	Std. Error	Pct.	Std. Error	Pct.	Std. Error
Digestive problems (bloat, scours, parasites, enterotoxemia, acidosis, etc.)	31.6	(6.2)	37.8	(8.9)	27.7	(4.5)	33.1	(6.7)	22.3	(3.6)	7.2	(4.6)
Respiratory problems (pneumonia, shipping fever, etc.)	43.9	(6.0)	23.2	(4.6)	37.9	(4.1)	38.3	(5.8)	33.1	(4.9)	13.0	(8.0)
Metabolic problems (milk fever, grass tetany, etc.)	0.0	(0.0)	0.6	(0.3)	2.4	(1.2)	5.0	(4.1)	0.4	(0.3)	0.0	(0.0)
Weather-related causes (chilling, drowning, lightning, etc.)	1.0	(0.5)	1.0	(0.4)	2.2	(1.0)	3.5	(2.3)	7.3	(4.0)	32.5	(12.9)
Theft (stolen)	0.0	(0.0)	0.1	(0.1)	0.0	(0.0)	0.0	(0.0)	0.1	(0.0)	0.6	(0.6)
Poisoning (nitrate poisoning, noxious feeds, noxious weeds, etc.)	0.1	(0.1)	0.0	(0.0)	0.0	(0.0)	0.0	(0.0)	0.0	(0.0)	0.0	(0.0)
Calving problems	13.4	(4.0)	7.5	(2.5)	16.7	(4.1)	8.0	(2.6)	15.1	(3.5)	18.5	(4.1)
Lameness/injury	0.5	(0.3)	1.7	(0.7)	0.0	(0.0)	1.7	(1.4)	0.0	(0.0)	13.1	(5.1)
Other diseases	2.5	(1.6)	6.9	(3.1)	1.0	(0.4)	0.6	(0.3)	2.4	(0.9)	1.2	(1.0)
Other	3.1	(2.5)	1.0	(0.6)	2.5	(1.1)	4.4	(2.8)	0.0	(0.0)	2.5	(2.7)
Unknown	3.9	(1.5)	20.2	(8.5)	9.6	(4.0)	5.4	(1.9)	19.3	(4.5)	11.4	(9.5)
Total	100.0		100.0		100.0		100.0		100.0		100.0	

On “other” operations in all regions, digestive and respiratory problems combined accounted for at least 60 percent of calf losses.

d. For **other** operations, percentage of calf death losses by cause and by region:

Nonpredator Cause	Percent Nonpredator Loss											
	Region											
	North Central		Northeast		Northwest		South Central		Southeast		Southwest	
	Pct.	Std. Error	Pct.	Std. Error	Pct.	Std. Error	Pct.	Std. Error	Pct.	Std. Error	Pct.	Std. Error
Digestive problems (bloat, scours, parasites, enterotoxemia, acidosis, etc.)	32.8	(8.2)	25.8	(4.8)	47.5	(9.9)	15.8	(5.6)	18.3	(4.6)	43.8	(10.5)
Respiratory problems (pneumonia, shipping fever, etc.)	37.4	(6.8)	49.8	(3.3)	32.7	(7.1)	65.3	(5.7)	45.1	(7.3)	42.2	(5.2)
Metabolic problems (milk fever, grass tetany, etc.)	0.0	(0.0)	2.4	(1.6)	0.0	(0.0)	0.8	(0.8)	0.2	(0.0)	0.0	(0.0)
Weather-related causes (chilling, drowning, lightning, etc.)	16.0	(7.4)	3.7	(1.8)	6.1	(2.9)	1.9	(1.0)	9.9	(5.1)	2.3	(1.7)
Theft (stolen)	1.1	(1.0)	0.0	(0.0)	0.6	(0.5)	0.0	(0.0)	0.0	(0.0)	0.0	(0.0)
Poisoning (nitrate poisoning, noxious feeds, noxious weeds, etc.)	0.0	(0.0)	2.7	(1.5)	0.0	(0.0)	0.8	(0.7)	0.0	(0.0)	0.0	(0.0)
Calving problems	3.4	(1.3)	0.9	(0.4)	0.5	(0.3)	4.6	(1.7)	8.1	(3.1)	0.2	(0.2)
Lameness/injury	1.0	(0.4)	4.3	(2.1)	1.1	(0.6)	0.1	(0.1)	1.4	(0.8)	5.4	(3.6)
Other diseases	2.3	(1.4)	2.5	(0.7)	2.7	(1.0)	1.4	(0.7)	4.8	(2.9)	2.3	(1.7)
Other	2.8	(2.3)	0.6	(0.3)	2.9	(2.4)	1.8	(1.2)	2.0	(2.0)	0.1	(0.1)
Unknown	3.2	(1.3)	7.3	(2.6)	5.9	(2.0)	7.5	(3.1)	10.2	(3.2)	3.7	(2.4)
Total	100.0		100.0		100.0		100.0		100.0		100.0	

Section III: Methodology

A. Reliability of Cattle and Calf Death Loss Estimates

1. Survey procedures

A random sample of U.S. cattle producers was surveyed to provide data for these estimates. Survey procedures ensured that all producers (except those in Alaska) had an opportunity to participate in the survey, regardless of operation size. Large operations were sampled more heavily than small operations. Data were collected during the first half of January by mail, telephone, and face-to-face interviews from about 77,000 producers in 1992, 50,000 in 1996, 50,000 in 2001, and 50,000 in 2006. Regardless of when producers responded, they were asked to report death losses for cattle and calves for the 1991, 1995, 2000, and 2005 calendar years and the January 1, 1992, 1996, 2001, and 2006 cattle and calf inventories by class.

Based on the original data collected during January, NASS's Cattle and Calves Death Loss report was issued in May 1992, May 1996, and May 2006. NASS's Cattle Predator Loss report was published in May 2001. These reports contained data for death loss by cause, including predator losses and losses from other causes. These estimates were on State and regional levels. NASS's Cattle report was issued in February 1992, February 1996, January 2001, and January 2006 and contained January 1 inventory estimates.

Data in the reports mentioned above as well as additional summaries from the original data provided estimates for this report. The original data were resummarized to break the losses out by class and size group based on reported inventory. The classes included beef cattle, dairy cattle, mixed, and "other" cattle operations. These classes and size groups are further defined on p. 2 of this report (Terms Used in This Report).

2. Estimating procedures

Estimates of total death loss and death loss by cause were prepared by the Livestock, Dairy and Poultry Branch, NASS, and the National Animal Health Monitoring System, APHIS. Total cattle and calf death losses from all causes were published in NASS's 1991, 1995, 2000, and 2005 Meat Animal Production, Disposition, and Income reports, released in April 1992, 1996, 2001, and 2006. Breakdowns by cause of loss and value of loss were published in the NASS Cattle and Calves Death Loss reports issued in May 1992, 1996, and 2006. Only the NASS Cattle Predator Loss report was issued in May 2001.

3. Reliability

Since all cattle producers are not included in the sample, survey estimates are subject to sampling variability. Survey results also are subject to nonsampling errors such as omissions; duplications; and mistakes in reporting, recording, and processing data. The effects of these nonsampling errors cannot be measured directly. They are minimized through rigid quality controls in the data collection process and through a careful review of all reported data for consistency and reasonableness.

Appendix: U.S. Populations and Operations*

Number of Cattle and Calves on January 1, 2006, and Number of Operations in 2005

Region	State	Number cattle and calves (1,000 head)	Number Operations
North Central	Iowa	3,800	31,000
	Minnesota	2,350	26,000
	Nebraska	6,500	24,000
	North Dakota	1,720	11,700
	South Dakota	3,750	17,000
	Total	18,120	109,700
Northeast	Connecticut	52	1,100
	Delaware	23	420
	Illinois	1,330	19,800
	Indiana	900	19,000
	Kentucky	2,380	45,000
	Maine	92	1,700
	Maryland	230	4,000
	Massachusetts	47	1,100
	Michigan	1,030	14,400
	New Hampshire	39	850
	New Jersey	42	1,500
	New York	1,400	14,700
	Ohio	1,280	27,000
	Pennsylvania	1,590	28,000
	Rhode Island	5	220
	Vermont	275	2,500
	Virginia	1,660	25,000
	West Virginia	410	12,500
Wisconsin	3,400	36,000	
Total	16,185	254,790	
Northwest	Idaho	2,110	10,400
	Montana	2,400	12,600
	Oregon	1,420	15,300
	Washington	1,100	13,000
	Wyoming	1,430	5,800
	Total	8,460	57,100
South Central	Arkansas	1,710	30,000
	Kansas	6,650	32,000
	Missouri	4,500	64,000
	Oklahoma	5,450	56,000
	Texas	14,100	150,000
	Total	32,410	332,000
Southeast	Alabama	1,280	25,000
	Florida	1,680	18,800
	Georgia	1,180	21,000
	Louisiana	820	14,500
	Mississippi	990	21,000
	North Carolina	860	21,000
	South Carolina	410	10,000
	Tennessee	2,240	48,000
Total	9,460	179,300	
Southwest	Arizona	940	2,600
	California	5,450	17,000
	Colorado	2,650	12,900
	Hawaii	161	800
	Nevada	500	1,600
	New Mexico	1,550	7,600
	Utah	800	7,000
	Total	12,101	49,500
Total (49 States)	96,686	982,390	
Total U.S. (50 States)	96,701.5	982,510	

*Source: NASS Cattle 2007; NASS Farms, Land in Farms, and Livestock Operations, 2007