

Vaccine	Pathogen	Pig type	Label
3FLEX® (Boehringer Ingelheim)	M. Hyopneumoniae (helps with PCV2 and PRRS)	3 weeks of age or older	
Anthrax Spore Vaccine (Colorado Serum)	B. anthracis	not stated	
Argus® SC/ST with Diluent (Intervet/Merck Animal Health)	S.choleraesuis/typhimurium	3 weeks of age or older	
Argus® SC/ST without Diluent (Intervet/Merck Animal Health)	S.choleraesuis/typhimurium	3 weeks of age or older	
Bordetella Bronchiseptica Intranasal Vaccine (MVP Labs.)	B. bronchiseptica	Piglets within 24h of birth	
Circumvent® PCV (Intervet/Merck Animal Health)	PCV2	3 weeks of age or older	
Circumvent® PCV G2 (Intervet/Merck Animal Health)	PCV2	3 weeks of age or older	
Circumvent® PCV M (Intervet/Merck Animal Health)	PCV2 and M. Hyopneumoniae	3 weeks of age or older	
Circumvent® PCV-M G2 (Intervet/Merck Animal Health)	PCV2 and M. Hyopneumoniae	3 weeks of age or older	
Citadel® L5 (Boehringer Ingelheim)	Leptospirosis	Any age (Healthy swine- probably not piglets)	
Clostridium Perfringens Types C&D Toxoid (Professional Biological)	C. perfringens C and D	Any age (Healthy swine- probably not piglets)	
Cl. Perfringens Type A Toxoid (Swine) (Elanco (Farm Animal))	C. perfringens A	Any age (Healthy swine- probably not piglets)	
Edema Vac 100/500 (Arko)	E. coli	18 days of age or older	
Emulsibac® APP (Phibro/MVP Labs.)	Actinobacillus pleuropneumoniae	>4 weeks of age	
ENDOVAC-Porci® with ImmunePlus® (Immvac)	S.choleraesuis/typhimurium	Any age (Healthy swine- probably not piglets)	

Vaccine	Pathogen	Pig type	Label
Enterisol® Ileitis (Boehringer Ingelheim)	Lawsonia intracellularis (PPE, ileitis)	>3 weeks of age	
Enterisol® Ileitis FF (Boehringer Ingelheim)	Lawsonia intracellularis (PPE, ileitis)	>3 weeks of age	
Enterisol Salmonella T/C® (Boehringer Ingelheim)	Salmonellosis (systemic)	>=2 weeks of age	
Enterisol Salmonella T/C® FF (Boehringer Ingelheim)	Salmonellosis (systemic)	>=2 weeks of age	
Enterisol® SC-54 (Boehringer Ingelheim)	Salmonellosis (systemic)	>=1 day of age	
Enterisol® SC-54 FF (Boehringer Ingelheim)	Salmonellosis (systemic)	>3 weeks of age	
Entero Vac 100/500 (Arko)	E. coli	Weaning assuming a 17 day or older weaning age	
ER Bac®/L5 GOLD (Zoetis)	Erysipelas and Leptospirosis	1st dose and then 2nd 2-4 weeks prior to breeding	
ER Bac® Plus (Zoetis)	Erysipelas	>=3 weeks of age	
ER Bac® Plus/Leptoferm-5® (Zoetis)	Erysipelas and Leptospirosis	1st dose and then 2nd 2-4 weeks prior to breeding	
Ery Shield® (Elanco (Farm Animal))	Erysipelas	Any age	
Ery Vac 100/500 (Arko)	Erysipelas	>8 weeks of age	
Essential 3 (Colorado Serum)	Clostridium perfringens B, C & D	Any age (Healthy swine- probably not piglets)	
FarrowSure® GOLD (Zoetis)	Erysipelas, Leptospirosis and Parvovirus	1st dose and then 2nd 2-4 weeks prior to breeding	
FarrowSure® GOLD B (Zoetis)	Erysipelas, Leptospirosis and Parvovirus	1st dose and then 2nd 2-4 weeks prior to breeding	

Vaccine	Pathogen	Pig type	Label
FarrowSure® Plus (Zoetis)	Erysipelas, Leptospirosis and Parvovirus	1st dose and then 2nd 2-4 weeks prior to breeding	
FarrowSure® Plus B (Zoetis)	Erysipelas, Leptospirosis and Parvovirus	1st dose and then 2nd 2-4 weeks prior to breeding	
FLEXcombo® (Boehringer Ingelheim)	PCV2 and M. Hyopneumoniae	>=3 weeks of age	
FLEXMycoPRRS™ (Boehringer Ingelheim)	PRRS and M. Hyopneumoniae	>=3 weeks of age	
FluSure® Pandemic (Zoetis)	Swine influenza H1N1	>3 weeks of age	
FluSure XP® (Zoetis)	Swine influenza H1N1, H1N2 and H3N2	>=3 weeks of age	
FluSure XP®/ER Bac Plus® (Zoetis)	Erysipelas and Swine influenza H1N1, H1N2 and H3N2	>=3 weeks of age	
FluSure XP®/FarrowSure® GOLD (Zoetis)	Erysipelas, Leptospirosis, Parvovirus and Swine influenza H1N1, H1N2 and H3N2	1st dose and then 2nd 2-4 weeks prior to breeding	
FluSure XP®/FarrowSure® GOLD B (Zoetis)	Erysipelas, Leptospirosis, Parvovirus and Swine influenza H1N1, H1N2 and H3N2	1st dose and then 2nd 2-4 weeks prior to breeding	
FluSure XP®/RespiSure® (Zoetis)	M. Hyopneumoniae and Swine influenza H1N1, H1N2 and H3N2	>=3 weeks of age	
FluSure XP®/RespiSure-ONE® (Zoetis)	M. Hyopneumoniae and Swine influenza H1N1, H1N2 and H3N2	>=3 weeks of age	
FluSure XP®/RespiSure-ONE®/ER Bac Plus® (Zoetis)	Erysipelas, M. Hyopneumoniae and Swine influenza H1N1, H1N2 and H3N2	>=3 weeks of age	
Fostera® Gold PCV (Zoetis)	PCV2	>=3 weeks of age	
Fostera® Gold PCV MH (Zoetis)	PCV2 and M. Hyopneumoniae	>=3 weeks of age	

Vaccine	Pathogen	Pig type	Label
Fostera™ PCV (Zoetis)	PCV2	>=3 weeks of age	
Fostera™ PCV MetaStim® (Zoetis)	PCV3	>=3 weeks of age	
Fostera™ PCV MH (Zoetis)	PCV2 and M. Hyopneumoniae	>=3 weeks of age	
Fostera™ PRRS (Zoetis)	PRRS	>=1 day of age	
Haemo Shield® P (Elanco (Farm Animal))	Actinobacillus pleuropneumoniae and Pasteurella multocida	>=5 weeks of age	
Ingelvac CircoFLEX® (Boehringer Ingelheim)	PCV2	>=3 weeks of age	
Ingelvac® ERY-ALC (Boehringer Ingelheim)	Erysipelas	>8 weeks of age	
Ingelvac MycoFLEX® (Boehringer Ingelheim)	M. Hyopneumoniae	>=3 weeks of age	
Ingelvac MycoMAX™ (Boehringer Ingelheim)	Mycoplasma hyorhinis	>=3 weeks of age	
Ingelvac Provenza™ (Boehringer Ingelheim)	Swine influenza H1N2 and H3N2	>=1 day of age	
Ingelvac PRRS® ATP (Boehringer Ingelheim)	PRRS	Sows and gilts: 3-4 weeks prior to breeding and Piglets >=3 weeks of age	
Ingelvac PRRS® MLV (Boehringer Ingelheim)	PRRS	Sows and gilts: anytime during pregnancy and Piglets >=3 weeks of age	
Lepto-5 (Colorado Serum)	Leptospirosis	>3 weeks prior to breeding	
Leptoferm-5® (Zoetis)	Leptospirosis	Any age (Healthy swine- probably not piglets)	
Lepto Shield™ 5 (Elanco (Farm Animal))	Leptospirosis	not stated	

Vaccine	Pathogen	Pig type	Label
LitterGuard® (Zoetis)	E. coli	2 doses 3 weeks apart during last half of pregnancy	
LitterGuard® LT (Zoetis)	E. coli	3 doses 3 weeks apart during last half of pregnancy	
LitterGuard® LT-C (Zoetis)	E. coli and C. perfringens C	4 doses 3 weeks apart during last half of pregnancy	
Magestic® 7 with Spur® (Intervet/Merck Animal Health)	Erysipelas, Leptospirosis and Parvovirus	6 weeks before breeding	
Maxi/Guard® Nasal Vac (Addison)	B. bronchiseptica	1-3 days of age	
MaxiVac Excell® 5.0 (Intervet/Merck Animal Health)	Swine influenza H1N2 and H3N2	>=5 weeks of age	
M+Pac® (Intervet/Merck Animal Health)	M. Hyopneumoniae	7-10 days of age	
Myco Shield® (Elanco (Farm Animal))	M. Hyopneumoniae	>=2 weeks of age	
Myco Silencer® ONCE (Intervet/Merck Animal Health)	M. Hyopneumoniae	>3 weeks of age	
Nitro-Ery ALC (Arko)	Erysipelas	>=5 weeks of age	
Nitro-Sal (Arko)	Salmonellosis (systemic)	not stated	
Nitro-Sal F.D. (Arko)	Salmonellosis (systemic)	>=5 weeks of age	
Parapleuro Shield® P (Elanco (Farm Animal))	H. parasuis, Actinobacillus pleuropneumoniae and Pasteurella multocida	>=5 weeks of age	
Parapleuro Shield® P+BE (Elanco (Farm Animal))	Erysipelas, B. bronchiseptica and Pasteurella multocida	>=7 weeks of age	

Vaccine	Pathogen	Pig type	Label
ParaSail® (Newport Laboratories)	H. parasuis	>=3 weeks of age	
Para Shield® (Elanco (Farm Animal))	H. parasuis	3-4 weeks of age	
Parvo Shield® (Elanco (Farm Animal))	Parvovirus	3-4 weeks prior to breeding	
Parvo Shield® L5 (Elanco (Farm Animal))	Leptospirosis and Parvovirus	4-6 weeks prior to breeding	
Parvo Shield® L5E (Elanco (Farm Animal))	Erysipelas, Leptospirosis and Parvovirus	4-6 weeks prior to breeding	
PneumoSTAR® Myco (Elanco (Farm Animal))	M. Hyopneumoniae	>=3 weeks of age	
PneumoSTAR® SIV COMPLETE (Elanco (Farm Animal))	Swine influenza subtypes H1N1, H1N2, and H3N2	>=3 weeks of age	
Porcilis® Ileitis (Intervet/Merck Animal Health)	Lawsonia intracellularis (PPE, ileitis)	>=3 weeks of age	
Porcine Epidemic Diarrhea Vaccine (Zoetis)	Porcine Epidemic Diarrhea	5 weeks prior to farrowing	
Porcine Epidemic Diarrhea Vaccine, RNA (Harrisvaccines)	Porcine Epidemic Diarrhea	>=3 weeks of age	
Porcine Pili Shield® (Elanco (Farm Animal))	E. coli	5 and 2 weeks prior to farrowing	
Porcine Pili Shield®+C (Elanco (Farm Animal))	E. coli and C. perfringens C	5 and 2 weeks prior to farrowing	
Prefarrow Shield® 9 (Elanco (Farm Animal))	Erysipelas, B. bronchiseptica, Pasteurella multocida, E. coli and C. perfringens C	5 and 2 weeks prior to farrowing	
Prefarrow Strep Shield® (Elanco (Farm Animal))	Streptococcus suis	5 and 2 weeks prior to farrowing	

Vaccine	Pathogen	Pig type	Label
Prime Pac™ PRRS+ (Intervet/Merck Animal Health)	PRRS	Sows and gilts: 3-6 weeks prior to breeding and Piglets >=3-4 weeks of age	
Prime Pac™ PRRS RS (Intervet/Merck Animal Health)	PRRS	>=3 weeks of age	
ProSystem® CE (Intervet/Merck Animal Health)	E. coli (enteritis and systemic) and C. perfringens C	5 and 2 weeks prior to farrowing	
ProSystem® RCE (Intervet/Merck Animal Health)	E. coli, C. perfringens C and Rotavirus	5 and 2 weeks prior to first farrowing and subsequent farrowings: 2 weeks before farrowing	
ProSystem® Rota (Intervet/Merck Animal Health)	Rotavirus	7-10 days pre-weaning	
ProSystem® TGE/Rota (Intervet/Merck Animal Health)	Rotavirus and Transmissible gastroenteritis (TGE)	5, 3 and 1 week prior to farrowing	
ProSystem® TREC (Intervet/Merck Animal Health)	E. coli, C. perfringens C, Rotavirus and Transmissible gastroenteritis (TGE)	5 and 2 weeks prior to farrowing	
PRV/Marker Gold® (Intervet/Merck Animal Health)	Pseudorabies	Piglets from non-immune dams >=3 days of age, piglets from immune dams after maternal antibody wanes	
RespiSure® (Zoetis)	M. Hyopneumoniae	Sows and gilts: 6 and 2 weeks prior to farrowing and Piglets ~1 week of age	
RespiSure 1 ONE® (Zoetis)	M. Hyopneumoniae	>=1 week of age	
RespiSure 1 ONE®/ER Bac Plus® (Zoetis)	Erysipelas and M. Hyopneumoniae	>3 weeks of age	
RespiSure®/ER Bac Plus® (Zoetis)	Erysipelas and M. Hyopneumoniae	>3 weeks of age	
Rhini Shield® TX4 (Elanco (Farm Animal))	Erysipelas, B. bronchiseptica and Pasteurella multocida	Sows and gilts: 5 and 2 weeks prior to farrowing and Piglets 7-10 days of age	

Vaccine	Pathogen	Pig type	Label
Salmo Shield® Live (Elanco (Farm Animal))	Salmonellosis (systemic)	Orally >= 1 day of age or I.M. at >=3 weeks of age	
Scourmune ®-C (Intervet/Merck Animal	E. coli and C. perfringens C	6-7 weeks prior to first farrowing and subsequent farrowings: 2-3 weeks before farrowing	
Suvaxyn® E (Zoetis)	Erysipelas	Sows and gilts: time of selection and Piglets 7-10 days of age	
Suvaxyn® E-oral (Zoetis)	Erysipelas	>6 weeks of age	
Suvaxyn® MH-one (Zoetis)	M. Hyopneumoniae	>=3 weeks of age	
Suvaxyn® P (Zoetis)	Parvovirus	Gilts: >6 months of age and at least 2 weeks before breeding, Sows: 8-2 weeks before breeding, Boars: > 8 months of age.	
Suvaxyn® RespiFend® APP (Zoetis)	Actinobacillus pleuropneumoniae	Piglets:>=4 weeks of age or older, Feeder pigs: upon arrival, Breeding stock: >=6 months of age or prior to introduction.	
Suvaxyn® RespiFend® HPS (Zoetis)	H. parasuis	Piglets: 7-10 days of age or older, Feeder pigs: upon arrival, Breeding stock: prior to introduction to herd.	
Suvaxyn® RespiFend® MH (Zoetis)	M. Hyopneumoniae	Piglets: 7-10 days of age or prior to weaning, Feeder pigs: upon arrival, Breeding stock: prior to introduction to herd.	
Suvaxyn® RespiFend® MH/HPS (Zoetis)	M. Hyopneumoniae and H. parasuis	Piglets: 7-10 days of age or older, Feeder pigs: upon arrival, Breeding stock: prior to introduction to herd.	
Swine Influenza Vaccine, RNA (Harrisvaccines)	Swine Influenza RNA	>=3 weeks of age	