

United States Department of Agriculture  
Center for Veterinary Biologics  
Testing Protocol

SAM 630

Supplemental Assay Method for Potency Testing of Fowl Cholera (*Pasteurella multocida*) Bacterins, Type 4

Date: May 27, 2016  
Number: SAM 630.04  
Supersedes: SAM 630.03, December 29, 2009  
Standard Requirement: 9 CFR 113.116  
Contact Person: Janet M. Wilson, (515) 337-7245

Approvals: /s/Larry R. Ludemann Date: 06Jun16  
Larry R. Ludemann, Section Leader  
Bacteriology  
  
/s/Paul J. Hauer Date: 20Jun16  
Paul J. Hauer, Director  
Policy, Evaluation, and Licensing  
Center for Veterinary Biologics

Animal and Plant Health Inspection Service  
P. O. Box 844  
Ames, IA 50010

Mention of trademark or proprietary product does not constitute a guarantee or warranty of the product by USDA and does not imply its approval to the exclusion of other products that may be suitable.

Entered into CVB Quality Management System by: <u>/s/Linda S. Snavelly</u> <u>23Jun16</u> Linda S. Snavelly Date Quality Management Program Assistant
--

UNCONTROLLED COPY

Supplemental Assay Method for Potency Testing Fowl Cholera (*Pasteurella multocida*) Bacterins, Type 4

Table of Contents

1. Introduction
2. Materials
  - 2.1 Equipment/instrumentation
  - 2.2 Reagents/supplies
  - 2.3 Animals
3. Preparation for the Test
  - 3.1 Personnel qualifications/training
  - 3.2 Selection and handling of test birds
  - 3.3 Preparation of supplies/equipment
  - 3.4 Preparation of reagents
4. Performance of the Test
  - 4.1 Vaccination of test animals
  - 4.2 Preparation of challenge in a biological safety cabinet
  - 4.3 Timing and administration of challenge
  - 4.4 Postinoculation plate count
  - 4.5 Observation of turkeys after challenge
5. Interpretation of Test Results
6. Report of Test Results
7. References
8. Summary of Revisions

**UNCONTROLLED COPY**

**Supplemental Assay Method for Potency Testing Fowl Cholera (*Pasteurella multocida*) Bacterins, Type 4**

**1. Introduction**

This Supplemental Assay Method (SAM) describes procedures for potency testing biological products containing avian *Pasteurella multocida*, type 4, as prescribed in title 9, *Code of Federal Regulations* (9 CFR), part 113.116. Turkeys are vaccinated twice, 21 days apart, and challenged with a standard dose of virulent *P. multocida*, type 4, 14 days after the second vaccination. This is a 2-stage test in which the second stage is applied when 7 or 8 vaccinated turkeys die in the first stage.

**2. Materials**

**2.1 Equipment/instrumentation**

Equivalent equipment or instrumentation may be substituted for any brand name listed below.

- 2.1.1 Spectrophotometer, Spectronic 20D+ (Spectronic Instruments)
- 2.1.2 Sterile inoculating loop
- 2.1.3 Bunsen burner or Bacti-Cinerator<sup>®</sup> (if non-sterile wire loop is used)
- 2.1.4 Incubator, 35°- 37°C
- 2.1.5 Micropipettors, 20- to 1000-µL
- 2.1.6 Crimper for aluminum seals on serum vials
- 2.1.7 Test tube mixer, vortex-type
- 2.1.8 Biological safety cabinet

**2.2 Reagents/supplies**

Equivalent reagents or supplies may be substituted for any brand name listed below.

- 2.2.1 *P. multocida*, type 4, strain P-1662. This culture must be obtained from the United States Department of Agriculture, Veterinary Services, Center for Veterinary Biologics (CVB). Refer to the current reagent data sheet for details.
- 2.2.2 Test bacterin(s) containing *P. multocida*, type 4

**UNCONTROLLED COPY**

**Supplemental Assay Method for Potency Testing Fowl Cholera (*Pasteurella multocida*) Bacterins, Type 4**

- 2.2.3 Syringes, Luer-lock, 3-mL or 5-mL
- 2.2.4 Needles, 18-gauge x 1 1/2-inch
- 2.2.5 Glass serum bottle, 20-mL
- 2.2.6 Rubber stopper, 13 x 20-mm, and aluminum cap for serum bottle
- 2.2.7 Screw-top glass tubes, 13 x 100-mm, with caps
- 2.2.8 Pipettes, 5-mL, 10-mL, 25-mL
- 2.2.9 Micropipette tips, up to 1000- $\mu$ L capacity
- 2.2.10 Bovine blood agar plates
- 2.2.11 Tryptose broth
- 2.2.12 Sterile cotton swabs
- 2.2.13 Poultry leg bands (size 11 or 14) or livestock spray paint, 1 color per treatment group, for animal identification

**2.3 Animals**

Turkeys, broad-breasted white, at least 6 weeks of age. Twenty turkeys are required for each serial to be tested. Ten additional turkeys are required as controls. All birds must be from the same source and hatch. The birds must be from flocks with no history of fowl cholera. Birds must not be previously vaccinated with any products containing *P. multocida*.

**3. Preparation for the Test**

**3.1 Personnel qualifications/training**

Technical personnel need working knowledge of the use of general laboratory chemicals, equipment, and glassware and must have specific training and experience in sterile technique, the handling of live bacterial cultures, and the handling of poultry.

**3.2 Selection and handling of test birds**

- 3.2.1 Turkeys of either sex may be used.

**UNCONTROLLED COPY**

**Supplemental Assay Method for Potency Testing Fowl Cholera (*Pasteurella multocida*) Bacterins, Type 4**

**3.2.2** House and feed all turkeys in a similar manner.

**3.2.3** It is permissible to house vaccinates and controls in the same enclosure, provided that space allocation is sufficient to meet requirements set forth by the CVB/National Veterinary Services Laboratories (NVSL) Animal Care and Use Committee.

**3.2.4** Positively identify each bird by treatment group. Identification may be by means of leg bands or livestock body paint.

1. If leg bands are used, band each leg in case 1 band is lost.
2. If body paint is used, it should be freshened at least every 3 weeks.

**3.2.5** If any turkeys die after vaccination for suspected vaccine related causes, but prior to challenge with live *P. multocida*, these birds shall be necropsied. If cause of death is unrelated to vaccination, the pathologist's report is filed with the test records and no additional action is taken. If death is attributable to the test bacterin, the death must be reported immediately to the CVB-Inspection and Compliance, which may request further safety testing of the bacterin.

**3.2.6** When the test is concluded, instruct the animal caretakers to euthanize and incinerate the birds and to sanitize contaminated rooms.

**3.3 Preparation of supplies/equipment**

**3.3.1** Sterilize all glassware before use.

**3.3.2** Use only sterile bacteriological supplies (pipettes, syringes, needles, rubber stoppers, saline, etc.).

**3.3.3** All equipment must be operated according to manufacturers' recommendations and applicable standard operating procedures.

**3.4 Preparation of reagents**

**3.4.1** *P. multocida*, type 4 (Lyon and Little classification), strain P-1662 challenge culture. Refer to the current reagent data sheet for details on storage and preparation.

**Supplemental Assay Method for Potency Testing Fowl Cholera (*Pasteurella multocida*) Bacterins, Type 4**

**3.4.2** Tryptose broth – National Centers for Animal Health (NCAH) Media #10404

Tryptose broth powder	26 g
Deionized water	q.s. 1 L

Autoclave 15 minutes at  $\geq 121^{\circ}\text{C}$ . Cool before using. Store at  $20^{\circ}$ -  $25^{\circ}\text{C}$  for no more than 6 months.

**3.4.3** Bovine blood agar – NCAH Media #10006

Blood agar base powder	40 g
Deionized water	q.s. 950 mL

Autoclave 20 minutes at  $\geq 121^{\circ}\text{C}$ . Cool to  $45^{\circ}$ -  $47^{\circ}\text{C}$ .

Add:

Defibrinated bovine blood	50 mL
---------------------------	-------

Pour into sterile petri dishes. Allow to cool to  $20^{\circ}$ -  $25^{\circ}\text{C}$ . Store at  $2^{\circ}$ -  $7^{\circ}\text{C}$  for no more than 6 months.

**4. Performance of the Test**

**4.1 Vaccination of test animals**

**4.1.1** Check the label on each product and/or Section VI of the current Outline of Production to confirm identity, recommended field dose, and route of injection.

**4.1.2** Thoroughly mix product by inverting end-to-end at least 10 times before the syringes are filled. Use 3- or 5-mL syringes, fitted with 18-gauge x 1 1/2-inch needles.

**4.1.3** Vaccinate separate groups of no more than 21 turkeys with each of the test bacterins. Use the dose volume and injection route recommended on the product label and/or Section IV of the current Outline of Production for each bacterin. Unless otherwise specified on the product label, subcutaneous injections must be given in the unfeathered, loose skin on the back of the lower neck.

**4.1.4** Revaccinate the turkeys in a similar manner 21 days after the first vaccination.

**UNCONTROLLED COPY**

**Supplemental Assay Method for Potency Testing Fowl Cholera (*Pasteurella multocida*) Bacterins, Type 4**

**4.1.5** Retain no more than 11 turkeys as non-vaccinated controls.

**4.2 Preparation of challenge in a biological safety cabinet**

**4.2.1** Reconstitute a vial of challenge in 1 mL tryptose broth.

**4.2.2** Inoculate 2 blood agar plates with 100 µL of reconstituted culture and streak for isolation.

**4.2.3** Incubate the inoculated blood agar plates at 35°- 37°C for 16 to 19 hours.

**4.2.4** Use plates that have pure growth by visual inspection to prepare the challenge inoculum.

**4.2.5** Scrape several bacterial colonies from the surface of the blood agar plates using a sterile cotton swab and suspend in tryptose broth in a 13 x 100-mm tube. Add bacterial growth until the suspension measures 73-78%T at 630 nm using a Spectronic 20D+ spectrophotometer, or equivalent. Use sterile tryptose broth in a 13 x 100-mm tube as a blank for the spectrophotometer.

**4.2.6** Prepare a 10<sup>-5</sup> dilution of the standardized culture in tryptose broth. **This is the inoculum used to challenge the turkeys.** Dispense challenge liquid into a serum vial and seal with a rubber stopper and aluminum ring.

**4.2.7** Prepare 10<sup>-6</sup>, 10<sup>-7</sup>, and 10<sup>-8</sup> dilutions for postinoculation plate counts or alternatively, save an aliquot of the challenge dilution in a separate vial and prepare these additional dilutions later (see **Section 4.4**).

**4.2.8** Place vial(s) of challenge inoculum and additional dilution tubes on ice. Keep on ice through challenge procedure and until added to plates for postinoculation plate count.

**4.3 Timing and administration of challenge**

**4.3.1** Challenge 20 vaccinates per serial of product 14 to 18 days after the second vaccination. Euthanize any additional birds at this time.

**4.3.2** Challenge non-vaccinated controls at the same time as the vaccinates. Euthanize any additional control birds at this time.

**4.3.3** Inoculate each turkey with 0.5 mL of challenge inoculum (10<sup>-5</sup> dilution of standardized culture, see **Section 4.2.6**) intramuscularly in the breast muscle, using a 3-mL syringe and 18-gauge x 1 1/2-inch needle.

**UNCONTROLLED COPY**

**Supplemental Assay Method for Potency Testing Fowl Cholera (*Pasteurella multocida*) Bacterins, Type 4**

**4.4 Postinoculation plate count**

**4.4.1** After birds are challenged, prepare  $10^{-6}$ ,  $10^{-7}$ , and  $10^{-8}$  dilutions using tryptose broth as the diluent.

**4.4.2** All bacterial suspensions must be mixed well prior to placing an aliquot on an agar plate. Plate each dilution in triplicate using 0.1 mL on bovine blood agar. Inoculum must be spread evenly on the surface of the agar plates and not allowed to pool around the edges. Complete all plate inoculations within 1 hour of challenge.

**4.4.3** Incubate the plates aerobically at 35°- 37°C for 18 to 30 hours.

**4.4.4** Using the dilution yielding 30-300 colonies per plate, calculate the colony forming units (CFU)/challenge dose according to the following formula:

$$\frac{\text{Colony count sum}}{\text{Number of plates}} \times \frac{1}{\text{Dilution factor plated}} \times \frac{1}{\text{Plated volume (mL)}} \times \frac{\text{Challenge dilution}}{1} \times \frac{\text{Challenge vol. (mL)}}{\text{Dose}} = \frac{\text{CFU}}{\text{Dose}}$$

**4.5 Observation of turkeys after challenge**

**4.5.1** Observe the turkeys up to twice daily for 14 days after challenge. Record deaths and euthanize any moribund birds as recommended by the Institutional Animal Care and Use Committee.

**4.5.2** If deaths occurring after challenge are suspected to be due to causes other than fowl cholera, such turkeys are necropsied to determine cause of death. If cause of death is unrelated to vaccination and/or challenge, the deaths are not included in the total deaths for the test.

**5. Interpretation of Test Results**

The test is interpreted as prescribed in the 9 CFR 113.116.

**5.1** For a valid test, at least 8 of 10 control turkeys shall die during the 14-day postchallenge period.



**Supplemental Assay Method for Potency Testing Fowl Cholera (*Pasteurella multocida*) Bacterins, Type 4**

Stage	Number of vaccinates	Cumulative number of vaccinates	Cumulative number of dead vaccinates for...	
			Satisfactory serial	Unsatisfactory serial
1	20	20	6 or less	9 or more
2	20	40	15 or less	16 or more

**5.2** The second stage may be conducted when 7 or 8 vaccinates die in the first stage of a valid test. The serial is Unsatisfactory if the test is not repeated. The second stage test is performed in a manner identical to the first stage test and evaluated according to the Table in **Section 5.1**.

**5.3** If fewer than 8 of 10 control turkeys die during the postchallenge period, the test is considered invalid due to insufficient challenge and is reported as Inconclusive. The test may be repeated without prejudice, and the repeat test is considered to be a first-stage test.

**5.4** The plate count (CFU/Dose) of the challenge is recorded on the test result form for informational purposes to track trends and to troubleshoot problem tests, but the 9 CFR does not specify a minimum or maximum CFU/Dose for this test.

## **6. Report of Test Results**

Report results of the test(s) as described by standard operating procedures.

## **7. References**

Title 9, *Code of Federal Regulations*, part 113.116, U.S. Government Printing Office, Washington, DC.

## **8. Summary of Revisions**

### **Version .04**

- The Section Leader and Director information has been updated.

### **Version .03**

- The Contact information has been updated.

**UNCONTROLLED COPY**

**Supplemental Assay Method for Potency Testing Fowl Cholera (*Pasteurella multocida*) Bacterins, Type 4**

- **2.1.3:** This section has been updated to reflect current practices.
- **2.2.12/4.2.5:** Sterile cotton swabs have been added.
- **4.1.3/4.1.5:** Bird group numbers have been updated to reflect 9 CFR 113.116.
- **4.3.1/4.3.2:** Instructions on euthanizing extra birds have been added.
- **4.5.1:** This section has been updated to reflect current practices.

**Version .02**

This document was revised to clarify practices currently in use at the Center for Veterinary Biologics and to provide additional detail. While no significant changes were made that impact the outcome of the test, the following changes were made to the document:

- **2.1.8** Biological safety cabinet has been added for equipment and references to its use have been added throughout the document.
- **3.2.5** Additional information regarding necropsy has been added.
- **4.1.3** Additional details on the location of SC vaccination have been added.
- **4.2.6** The option of preparation of dilutions for use in the plate count at 2 separate places in the protocol has been added.
- Information regarding the current spectrophotometer in use has been added.
- Plate count dilution details have been added.
- Details regarding references supplied by CVB have been replaced with references to the current reagent data sheets throughout the document.
- The size of the needle currently in use has been indicated throughout the document.
- The size of poultry leg bands have been indicated throughout the document.
- The contact person has been changed to Janet Wilson.

**UNCONTROLLED COPY**