

Breadcrumb

1. [Home](#)
2. Print
3. Pdf
4. Node
5. Entity Print

# Detections of Highly Pathogenic Avian Influenza in Wild Birds

Last Modified:

Wild birds can be infected with highly pathogenic avian influenza (HPAI) and show no signs of illness. They can carry the disease to new areas when migrating, potentially exposing domestic poultry to the virus.

APHIS' wild bird surveillance program provides an early warning system for the introduction and distribution of avian influenza viruses of concern in the United States, allowing APHIS and the poultry industry to take timely and rapid action to reduce the risk of spread to our poultry industry and other populations of concern.

Users may need to refresh the page to see the latest table data. To refresh the page, hold down the SHIFT key and click the Reload Page button in your browser.

## Confirmed HPAI Detections

Captive wild birds, defined by the [World Organisation for Animal Health \(WOAH\)](#) as a wild animal that is captive or otherwise lives under or requires human supervision or control, are included in the numbers reported on this page. Captive wild birds, including sick wild birds that may have died after being found and taken to a rehabilitator or sanctuary, will have the designation of "captive wild bird" in the

WOAH Classification column. To remain consistent with current reporting, information will continue to be broken down to the county level. To protect privacy, private or business names will not be released. This is consistent with current reporting of wild birds as well as commercial and backyard flocks.



APHIS is continuing to process samples collected over the past year from State and partner agencies and will continue to post this information on this page. The detection date is the date of [National Veterinary Services Laboratories](#) confirmatory testing. APHIS will provide updates below when there are significant differences between sample collection and detection dates.

For submitting agencies who notice data errors or omissions, please send an email with supporting documentation (laboratory report/accession number/data collection) to [wslabresults@usda.gov](mailto:wslabresults@usda.gov).

The two maps below represent:

1. A comprehensive view of all HPAI detections in wild birds since the beginning of the U.S. outbreak in January 2022, and
2. The number of HPAI detections in wild birds from the past 30 days.

Both maps represent samples collected by APHIS [Wildlife Services](#) as well as morbidity/mortality samples submitted by State agencies and private facilities. The data presented visually in these maps is also available in the table below.



See More See Less

Loading...

Loading...

Loading...

## Table Notes

**Date Detected:** Specimens detected by the NAHLN H5 assay were further tested by a developmental real-time RT PCR targeting the Eurasian lineage goose/Guangdong H5 clade 2.3.4.4b. "Date Detected" indicates the date when a positive detection was obtained by the developmental RRT PCR targeting the Eurasian lineage goose/Guangdong H5 clade 2.3.4.4b.

### HPAI Strain:

EA = Eurasian; AM = North American; the EA H5 (2.3.4.4) viruses are highly pathogenic to poultry.

EA/AM: reassortant of H5 goose/Guangdong and North American wild bird lineage

Virus lineage, subtype, and pathotype per cleavage site analysis are determined from sequence data direct from the sample or virus isolate. An incomplete subtype indicates either (1) the specimen is pending virus isolation and/or sequencing results, or (2) the specimen was detected by the developmental H5 RRT PCR targeting the Eurasian lineage goose/Guangdong H5 clade 2.3.4.4b but could not be further characterized, often due to a low level of virus or viral RNA present in a given sample.

### Sampling Sources:

AK DFG = Alaska Department of Fish and Game  
CA DFW = California Department of Fish and Wildlife  
DOI = Department of the Interior  
FL FWCC = Florida Fish and Wildlife Conservation Commission  
GA DNR = Georgia Department of Natural Resources  
IA DNR = Iowa Department of Natural Resources  
ID Fish and Game = Idaho Fish & Game  
KS DWP = Kansas Department of Wildlife and Parks  
KY DFW = Kentucky Department of Fish and Wildlife  
LA DWF = Louisiana Department of Wildlife and Fisheries  
MA DFW = Massachusetts Department of Fish and Wildlife  
MD DNR = Maryland Department of Natural Resources  
ME DIFW = Maine Department of Inland Fisheries and Wildlife  
MN DNR = Minnesota Department of Natural Resources  
MO DOC = Missouri Department of Conservation  
MT FWP = Montana Fish, Wildlife, and parks  
NC WRC = North Carolina Wildlife Resources Commission  
ND Game and Fish = North Dakota Game and Fish

NE Dept of Ag = Nebraska Department of Agriculture  
NH FGD = New Hampshire Fish and Game Department  
NPS = DOI National Park Service  
NV DOW = Nevada Division of Wildlife  
NWDP = USDA Wildlife Services National Wildlife Disease Program  
NY DEC = New York State Department of Environmental Conservation  
NY State Dept of Ag and Markets = New York State Department of Agriculture and Markets  
OH DNR = Ohio Department of Natural Resources  
PA Game Commission = Pennsylvania Game Commission  
SC DNR = South Carolina Department of Natural Resources  
SCWDS = Southeastern Cooperative Wildlife Disease Study  
SD GFP = South Dakota Game, Fish, and Parks  
USACE = U.S. Army Corps of Engineers  
USFWS = DOI U.S. Fish and Wildlife Service  
USGS = U.S. Geological Survey National Wildlife Health Center  
VA DWR = Virginia Department of Wildlife Resources  
WA DFW = Washington Department of Fish and Wildlife  
WI DNR = Wisconsin Department of Natural Resources

[See More](#) [See Less](#)

## About Avian Influenza

Avian influenza is caused by influenza Type A virus (influenza A). Avian-origin influenza viruses are broadly categorized based on a combination of two groups of proteins on the surface of the influenza A virus: hemagglutinin or “H” proteins, of which there are 16 (H1–H16), and neuraminidase or “N” proteins, of which there are 9 (N1–N9). Many different combinations of “H” and “N” proteins are possible. Each combination is considered a different subtype, and related viruses within a subtype may be referred to as a lineage. Avian influenza viruses are classified as either “low pathogenic” or “highly pathogenic” based on their genetic features and the severity of the disease they cause in poultry. Most viruses are of low pathogenicity, meaning they cause no signs or only minor clinical signs of infection in poultry.

---

## Related Links

- [Avian Influenza](#)
- [Confirmations of HPAI in Commercial and Backyard Flocks](#)
- [Detections of HPAI in Mammals](#)

[Print](#)