

TERRESTRIAL ANIMAL HEALTH STANDARDS COMMISSION  
FEBRUARY 2013 REPORT

CHAPTER 14.5.

**INFECTION WITH CHLAMYDOPHILA ABORTUS INFECTION**  
**( ENZOOTIC ABORTION OF EWES , OVINE**  
**CHLAMYDIOSIS )**

Article 14.5.1.

**General provisions**

For the purposes of the *Terrestrial Code*, enzootic abortion of ewes (EAE), also known as ovine chlamydiosis or ovine enzootic abortion, is an *infection* of domestic sheep and goats by the bacterium *Chlamydomphila abortus*.

Susceptible *animals* become infected through ingestion of infectious materials. In lambs and non-pregnant ewes, the *infection* remains latent until conception. Ewes exposed to *infection* late in pregnancy may not exhibit signs of *infection* until the subsequent pregnancy. Countries should take account of these risk factors.

Standards for diagnostic tests are described in the *Terrestrial Manual*.

Article 14.5.2.

**Recommendations for the importation of sheep and/or goats for breeding**

*Veterinary Authorities of importing countries* should require the presentation of an *international veterinary certificate* attesting that the *animals*:

- 1) have remained since birth, or for the previous two years, in *establishments* where no EAE has been diagnosed during the past two years;
- 2) showed no clinical sign of EAE on the day of shipment;
- 3) were subjected to a diagnostic test for EAE with negative results within the 30 days prior to shipment.

Article 14.5.3.

**Sheep flocks and/or goat herds free from EAE infection**

To qualify as free from EAE *infection*, a sheep *flock* or goat *herd* shall satisfy the following requirements:

- 1) it is under official veterinary *surveillance*;
- 2) all sheep and goats showed no clinical evidence of EAE *infection* during the past two years;
- 3) a statistically valid number of sheep and goats over six months of age were subjected to a diagnostic test for EAE with negative results within the past six months;
- 4) all sheep or goats are permanently identified;
- 5) no sheep or goat has been added to the *flock* or *herd* since 30 days prior to the *flock* or *herd* test referred to in point 3 above unless:
  - a) either the additions were isolated from other members of the *flock* or *herd* in the *establishment* of origin for a minimum period of 30 days and then were subjected to a diagnostic test for EAE with negative results, before entry into the new *flock* or *herd*; or

- b) they originated from an *establishment* of equal health status.

Article 14.5.4.

**Recommendations for the importation of semen of sheep or goats**

*Veterinary Authorities of importing countries* should require the presentation of an *international veterinary certificate* attesting that

4. the donor *animals* showed no clinical signs on the day of semen collection; and:
  - 1a) have been kept in *establishments* or *artificial insemination centres* free from EAE according to Article 14.5.3. during for the past two years prior to collection, and have not been in contact with *animals* of a lower health status; or
  - 2b) have remained since birth, or for the previous two years prior to collection, in establishments where no EAE has been diagnosed during the past two years and were subjected to a diagnostic test for EAE with negative results two to three weeks after collection of the semen;.
2. an aliquot of the semen to be exported was shown to be free of *Chlamydomphila abortus*.

Article 14.5.5.

**Recommendations for the importation of embryos of sheep or goats**

*Veterinary Authorities of importing countries* should require the presentation of an *international veterinary certificate* attesting that the donor *animals* showed no clinical signs on the day of embryo collection and:

- 1) have been kept in *establishments* free from EAE according to Article 14.5.3. during for the past two years prior to collection and have not been in contact with *animals* of a lower health status; or
- 2) have remained since birth, or for the previous two years prior to collection, in establishments where no EAE has been diagnosed during the past two years and were subjected to a diagnostic test for EAE with negative results two to three weeks after collection of the embryos.

The embryos should be collected, processed and stored in accordance with Chapter 4.7.