



Pest-Free Areas

The following table lists countries and associated areas that meet the APHIS requirements for designated pest-free areas in accordance with 7 CFR 319.56-5.

Country	Pest(s)	Pest-free Area
Argentina	Mediterranean fruit fly (Medfly) <i>Ceratitis capitata</i> and <i>Anastrepha</i> spp. fruit flies	The Patagonia Provinces of Neuguen, Rio Negro, Chabut, Santa Cruz, and Tierra del Fuego. This includes areas along the valleys of the Rio Colorado and Rio Negro rivers and areas of the southern part of the Mendoza Province, south of the following coordinates: lat 33° 13' 40.98" S, length 69° 54' 36.86" W; lat 33° 13' 40.98" S, length 69° 04' 18.24" W; lat 33° 29' 29" S, length 68° 59' 20" W; lat 34° 02' 47" S, length 67° 57' 17" W; lat 34° 02' 47" S, length 66° 44' 06.05" W
Australia	Mediterranean fruit fly (Medfly) (<i>Ceratitis capitata</i>), the Queensland fruit fly (<i>Bactrocera tryoni</i> or QFF) and other fruit flies destructive of citrus	The Riverland district of South Australia, defined as the county of Hamley; the geographical subdivisions, called hundreds, of Bookpurnong, Cadell, Eba, Fisher, Forster, Gordon, Hay, Holder, Katarapko, Loveday, Markaranka, Morook, Murbko, Murtho, Nildottie, Paisley, Parcoola, Paringa, Pooginook, Pyap, Ridley, Skurray, Stuart, and Waikerie; and the Parish of Onley of the Shire of Mildura, Victoria
	Mediterranean fruit fly (Medfly) (<i>Ceratitis capitata</i>) and the Queensland fruit fly (<i>Bactrocera tryoni</i>)	Tasmania
	Mediterranean fruit fly (Medfly) (<i>Ceratitis capitata</i>)	Eastern Australia, defined as the Northern Territory, Queensland, South Australia, New South Wales, and Victoria
	Queensland fruit fly (<i>Bactrocera tryoni</i>)	Western Australia
Brazil	South American cucurbit fruit fly (<i>Anastrepha grandis</i>)	The portion of Brazil bounded on the north by the Atlantic Ocean; on the east by the River Assu (Acu) from the Atlantic Ocean to the city of Assu; on the south by Highway BR 304 from the city of Assu (Acu) to Mossoro, and by Farm Road RN-015 from Mossoro to the Ceara State line; and on the west by the Ceara State line to the Atlantic Ocean The Municipalities of Aracati, Icapuí, Itaiçaba, Jaguaruana, Limoeiro do Norte, Quixerê, and Russas in the State of Ceará, and the municipalities of Açú, Afonso Bezerra, Alto do Rodrigues, Areia Branca, Baraúna, Camaubais, Grossos, Ipanguaçú, Mossoró, Porto do Mangue, Serra do Mel, Tibau, Upanema in the State of Rio Grande do Norte

Country	Pest(s)	Pest-free Area
Ecuador	South American cucurbit fruit fly (<i>Anastrepha grandis</i>)	The area within 5 kilometers of either side of the following roads: (i) Beginning in Guayaquil, the road north through Nobol, Palestina, and Balzar to Velasco-Ibarra (Empalme); (ii) Beginning in Guayaquil, the road south through E1 26, Puerto Inca, Naranjal, and Camilo Ponce to Enriquez; (iii) Beginning in Guayaquil, the road east through Palestina to Vinces; (iv) Beginning in Guayaquil, the road west through Piedrahita (Novol) to Pedro Carbo; or (v) Beginning in Guayaquil, the road west through Progreso, Engngunga, Tugaduaja, and Zapotal to El Azucar
Guatemala	Mediterranean fruit fly (Medfly) (<i>Ceratitis capitata</i>)	Department of Peten
Mexico	Mediterranean Fruit fly (Medfly) (<i>Ceratitis capitata</i>), Mexican Fruit fly (<i>Anastrepha ludens</i>), Sapote fruit fly (<i>A. serpentine</i>), West Indian fruit fly (<i>A. oblique</i>), and South American fruit fly (<i>A. fraterculus</i>)	The Municipalities of Comondu, La Paz, Loreto, Los Cabos, and Mulegé in the State of Baja California Sur; the Municipalities of Bachiniva, Casas Grandes, Cuahutemoc, Guerrero, Namiquipa, and Nuevo Casas Grandes in the State of Chihuahua; the Municipalities of Ahome, Choix, El Fuerte, Guasave, and Sinaloa de Leyva in the State of Sinaloa; and the Municipalities of Altar, Atil, Bacum, Benito Juarez, Caborca, Cajeme, Carbo, Empalme, Etchojoa, Guaymas, Hermosillo, Huatabampo, Navojoa, Pitiquito, Plutarco Elias Calles, Puerto Penasco, San Luis Rio Colorado, San Miguel, and San Ignacio Rio Muerto in the State of Sonora
Peru	South American cucurbit fly (<i>Anastrepha grandis</i>), gray pineapple mealybug (<i>Dysmicoccus neobrevipes</i>)	The Departments of Lima, Ica, Arequipa, Moquegua, and Tacna
Republic of South Africa	Citrus black spot (CBS) (<i>Guignardia citricarpa</i> Kiely)	Magisterial districts of Bellville, Bredasdorp, Caledon, Cape, Ceres, Clanwilliam, Goodwood, Grabouw, Heidelberg, Hermanus, Hopefield, Kuilsriver, Ladismith, Malmesbury, Mitchell's Plain, Montagu, Moorreesburg, Paarl, Piketberg, Robertson, Somerset-West, Stellenbosch, Strand, Swellendam, Tulbagh, Villiersdrop, Vredenburg, Wellington, Worcester, and Wynberg of the Western Cape Province; Magisterial districts of Barkly-west/west, Gordonia, Hartswater, Hay, Herbert, Hopetown, Kenhardt, Kimberley, Namakwaland, Prieska, and Warrenton of the Northern Cape Province; Magisterial districts of Boshof, Fauresmith, Jacobsdal, Koffiefontein, and Philippolis of the Free State Province; Magisterial districts of Christiania and Taung of the North West Province

Country	Pest(s)	Pest-free Area
Venezuela	South American cucurbit fly (<i>Anastrepha grandis</i>)	The Paraguana Peninsula, located in the State of Falcon, bounded on the north and east by the Caribbean Ocean, on the south by the Gulf of Coro and an imaginary line dividing the autonomous districts of Falcon and Miranda, and on the west by the Gulf of Venezuela