



United States
Department of
Agriculture

Marketing and
Regulatory
Programs

Animal and Plant
Health Inspection
Service

Mexican Fruit Fly Cooperative Eradication Program

Zapata County, Texas

Environmental Assessment July 2016

Mexican Fruit Fly Cooperative Eradication Program

Zapata County, Texas

**Environmental Assessment,
July 2016**

Agency Contact:

John C. Stewart
National Fruit Fly Policy Manager
Center for Plant Health Science and Technology Plant
Protection and Quarantine
Animal and Plant Health Inspection Service
U.S. Department of Agriculture
1730 Varsity Drive, Suite 400
Raleigh, NC 27606

Non-Discrimination Policy

The U.S. Department of Agriculture (USDA) prohibits discrimination against its customers, employees, and applicants for employment on the bases of race, color, national origin, age, disability, sex, gender identity, religion, reprisal, and where applicable, political beliefs, marital status, familial or parental status, sexual orientation, or all or part of an individual's income is derived from any public assistance program, or protected genetic information in employment or in any program or activity conducted or funded by the Department. (Not all prohibited bases will apply to all programs and/or employment activities.)

To File an Employment Complaint

If you wish to file an employment complaint, you must contact your agency's EEO Counselor (PDF) within 45 days of the date of the alleged discriminatory act, event, or in the case of a personnel action. Additional information can be found online at http://www.ascr.usda.gov/complaint_filing_file.html.

To File a Program Complaint

If you wish to file a Civil Rights program complaint of discrimination, complete the USDA Program Discrimination Complaint Form (PDF), found online at http://www.ascr.usda.gov/complaint_filing_cust.html, or at any USDA office, or call (866) 632-9992 to request the form. You may also write a letter containing all of the information requested in the form. Send your completed complaint form or letter to us by mail at U.S. Department of Agriculture, Director, Office of Adjudication, 1400 Independence Avenue, S.W., Washington, D.C. 20250-9410, by fax (202) 690-7442 or email at program.intake@usda.gov.

Persons With Disabilities

Individuals who are deaf, hard of hearing, or have speech disabilities and you wish to file either an EEO or program complaint please contact USDA through the Federal Relay Service at (800) 877-8339 or (800) 845-6136 (in Spanish).

Persons with disabilities who wish to file a program complaint, please see information above on how to contact us by mail directly or by email. If you require alternative means of communication for program information (e.g., Braille, large print, audiotope, etc.) please contact USDA's TARGET Center at (202) 720-2600 (voice and TDD).

Mention of companies or commercial products in this report does not imply recommendation or endorsement by USDA over others not mentioned. USDA neither guarantees nor warrants the standard of any product mentioned. Product names are mentioned to report factually on available data and to provide specific information.

This publication reports research involving pesticides. All uses of pesticides must be registered by appropriate State and/or Federal agencies before they can be recommended.

CAUTION: Pesticides can be injurious to humans, domestic animals, desirable plants, and fish and other wildlife—if they are not handled or applied properly. Use all pesticides selectively and carefully. Follow recommended label practices for the use and disposal of pesticides and pesticide containers.

Table of Contents

I. Need for Proposal.....	1
II. Alternatives	3
A. No Action	3
B. Quarantine and Commodity Certification	3
C. Eradication Using an IBM Approach (Preferred Approach)	4
1. Delimitation	5
2. Treatments.....	5
III. Potential Environmental Consequences.....	7
A. No Action	7
B. Quarantine and Commodity Certification	7
C. Eradication Using and IPM Approach (Preferred Alternative)	7
1. Affected Environment	8
2. Human Health	12
3. Other Aspects of the Human Environment	15
4. Nontarget Species	18
5. Environmental Quality.....	21
6. Cumulative Impacts	23
IV. Agencies Consulted.....	27
V. References Cited	28
 Appendix A. San Ygnacio Mexfly Program Area, Texas—as of June 30, 2016	
 Appendix B. Active Texas Mexfly Quarantines—as of June 30, 2016	
 Appendix C. Outside-APHIS Spatial Data Resources Used to Prepare this Document	

I. Need for the Proposal

The Mexican fruit fly (Mexfly), *Anastrepha ludens* (Loew), is native to central Mexico, and is a major pest capable of devastating crops throughout many parts of the Western Hemisphere. Mexfly has been introduced into the United States repeatedly since its first detection in Texas in 1927 (NAPIS, n.d.). Successful eradication programs have prevented Mexfly from becoming an established pest in the conterminous United States.

Adult Mexflies are long lived (up to 11 months), highly fertile, strong fliers, and highly mobile (UFL, 2012). Because Mexfly can damage over 50 species of fruits, a wide range of commercial crops and dooryard production in the United States would suffer if Mexfly populations became established. Fruit infested by Mexfly is unfit to eat because the larvae tunnel through the fleshy part of the fruit, damaging it and subjecting it to decay from bacteria and fungi (CDFA, n.d.). Mexfly outbreaks have occurred repeatedly in southern Texas due to the proximity of infested areas in Mexico.

On June 29, 2016, a larva collected from a citrus host in the town of San Ygnacio was confirmed as Mexfly (APHIS, 2016a). As a result of the larval detection, the San Ygnacio Quarantined Area¹ was established: 29-square miles that contain residential suburban and waterfront property, with parcels of commercial and undeveloped land (see map in appendix A). Delimitation surveys continue around the Mexfly detection site; fruit stripping and spinosad applications were begun after confirmation of the finds; and, releases of sterile Mexflies (sterile insect technique, or SIT) are planned (APHIS, 2016a).

San Ygnacio is about 14 miles from an active Mexfly quarantine and eradication program in Zapata, Texas (APHIS, 2016b). The San Ygnacio and Zapata Mexfly program areas do not overlap; their current quarantine boundaries are about 1.5 miles apart at the closest point (see map in appendix B) (Smith, pers. comm., 07/05/2016). There are no acres of commercial citrus production inside the current San Ygnacio quarantine; the nearest commercial Mexfly-host production is in eastern Hidalgo County, about 120 miles away (APHIS, 2016a). Commercial citrus harvest in the Lower Rio Grande Valley (LRGV) region of Texas is complete for 2016 (Nash, pers. comm., 06/17/2016). However, many Mexfly-host plant species are cultivated across south Texas, increasing the potential environmental impact of the San Ygnacio infestation.

The Texas Department of Agriculture (TDA) and APHIS and are proposing a cooperative program to eradicate the Mexfly infestation and prevent the

¹ For the purposes of this document, and unless specified otherwise in the text, the terms "Quarantined Area" and "program area" signify the same place. A core area is where program chemical treatments may be applied.

spread of Mexfly to noninfested areas of the United States. APHIS' authority for cooperation in the program is the Plant Protection Act (Title 4 of the Agricultural Risk Protection Act of 2000), which authorizes the Secretary of Agriculture to carry out operations to eradicate insect pests, and to use emergency measures to prevent the dissemination of plant pests new to, or not widely distributed throughout, the United States.

Working cooperatively with States and territories, APHIS identifies and eradicates Mexfly infestations. APHIS has cooperated with the California, Florida, Puerto Rico, and Texas Departments of Agriculture on fruit fly eradication programs since 1984. To date, every fruit fly population targeted by APHIS' cooperative control programs was successfully eradicated.

Monitoring for Mexfly is ongoing in susceptible agricultural regions of Texas. State officials initiate *Anastrepha* spp. delimitation and eradication programs in locations where the types and number of detections are not yet triggering quarantine regulatory actions. Delimitation and eradication programs try to eliminate fruit fly infestations before reaching a quarantine threshold and imposing regulatory quarantines. Following Texas program protocols for Mexfly eradication, the use of sterile insect technique (SIT) continues year-round at a rate upwards of 500 flies per acre in designated at-risk counties which includes the commercial citrus production areas in the Lower Rio Grande Valley. Program officials have approved SIT for all current Mexfly quarantines, including the San Ygnacio program area (APHIS, 2016a and 2016b; Stewart, pers. comm., 7/6/16). (See appendix B for Texas Mexfly quarantine locations.)

APHIS and its cooperating partners have discussed and comprehensively analyzed alternatives for Mexfly eradication since 1984. APHIS first evaluated the environmental impacts of fruit fly control technologies in the *Fruit Fly Cooperative Control Program, Final Environmental Impact Statement—2001* (EIS1) (APHIS, 2001). APHIS reexamined its findings and introduced an additional tool for eradication in the *Use of Genetically Engineered Fruit Fly and Pink Bollworm in APHIS Plant Pest Control Programs, Final Environmental Impact Statement—2008* (EIS2) (APHIS, 2008). Both EIS1 and EIS2 consider fruit fly risks and mitigations at the programmatic level. This case-specific environmental analysis (EA) incorporates the findings of EIS1 and EIS2 by reference.

This environmental assessment (EA) analyzes the environmental consequences of alternatives considered for Mexfly eradication, and analyzes, from a site-specific perspective, environmental issues relevant to this particular program. The eradication measures being considered for this program were discussed and comprehensively analyzed within APHIS' fruit fly chemical risk assessments (APHIS, 2014, 2003, 1999, 1998a, and 1998b). These documents are also incorporated by reference and summarized within this EA. Environmental documentation for APHIS'

fruit fly control programs may be viewed online via the following links: [APHIS fruit fly control program environmental documentation](#) and [APHIS GE control applications for plant health](#).

II. Alternatives

Alternatives considered for this proposed program include: (A) no Federal action, (B) quarantine and commodity certification, and (C) the preferred alternative, eradication using an integrated pest management (IPM) approach. Component techniques of alternative C may include the use of regulatory controls, high density trapping, host larval survey, and chemical and biological control (SIT) to facilitate the timely elimination of the current Mexfly infestation. These alternatives and their component techniques were discussed and comprehensively analyzed within EIS1 and EIS2 (APHIS, 2001 and 2008), and are incorporated by reference and summarized within this EA.

A. No Action

Under the no action alternative, there would be no Federal efforts to eradicate Mexfly or restrict expansion of the Mexfly population from the infested area. In the absence of a Federal effort, quarantine and control would be left to State and local government, grower groups, and individuals. Expansion of the infestation would be influenced by any controls exerted over it, by the proximity of host plants, and by climatic conditions.

“No treatment” might be the only reasonable alternative for some sensitive sites. In such cases, lack of treatment could lead to a continuing and expanding infestation. An expansion of the infestation would likely result in substantial economic losses to growers in the United States, as well as the loss of U.S. export agricultural markets.

Under the no action alternative, APHIS would continue cooperative practices to support the TDA detection trapping program and research. SIT Mexfly releases would continue over commercial citrus groves and high-risk areas in the LRGV of Texas and Mexico (APHIS, 2010). (For details about the Texas State program to control Mexfly, please use the following link: [Texas Mexfly program information](#).)

B. Quarantine and Commodity Certification

The Quarantine and Commodity Certification alternative combines a Federal quarantine with commodity treatment and certification, as stipulated under Title 7 of the Code of Federal Regulations (CFR) § 301.32. Regulated commodities harvested within the quarantine area would not be allowed to move unless treated with prescribed applications, and certified

for movement outside the area. For a large infestation, intensive quarantine enforcement activities could be necessary, including safeguarding local fruit stands, mandatory baggage inspection at airports, and judicious use of road patrols and regulatory checks. The quarantine actions of this alternative are designed to reduce Mexfly movement outside treated areas, and reduce human-mediated transport of Mexfly in host plant materials to areas outside the quarantined area; however, the infestation could remain established within the quarantine boundaries. Any Mexfly eradication efforts would be managed by, and wholly under the control of, TDA.

The interstate movement of regulated commodities would require the issuance of a certificate, or limited permit, contingent upon commodity treatment or the grower or shipper complying with specific conditions designed to minimize pest risk and prevent the spread of Mexfly. Eradication methods that may be used in this alternative include (1) regulatory chemicals, (2) cold treatment, (3) vapor heat treatment, and (4) irradiation treatment. Regulatory chemical treatments may include fumigation with methyl bromide (MB), and bait spray with a mixture of protein hydrolysate (a food bait) and spinosad. (Refer to EIS1 (APHIS, 2001) for more detailed information about these chemicals and their uses.) Cold treatment, vapor heat treatment, or irradiation treatment of certain produce, as a requirement for certification and shipping, must be done in facilities that are inspected and approved by APHIS.

C. Eradication Using an IPM Approach (Preferred Alternative)

APHIS' preferred alternative for the San Ygnacio Mexfly program is eradication using an IPM approach. This alternative combines quarantine and commodity certification with eradication treatments, and is designed to be biologically effective while minimizing impacts to the environment, public intrusiveness, and program operating costs (TDA, 2016a). Successful eradication of a Mexfly infestation in the LRGV, using a similar IPM strategy, was declared in December 2015 (APHIS, 2015).

For many species of exotic fruit flies, effective nonchemical control or eradication techniques do not exist (APHIS, 2001). APHIS' and TDA's Mexfly eradication program rely primarily on surveillance, bait sprays, and SIT (TDA, 2016b). Eradication efforts for the San Ygnacio program may include any or all of the following:

- no action,
- regulatory quarantine treatment and movement control of host materials and regulated articles,
- host survey for evidence of breeding Mexfly populations,
- host removal,
- eradication chemical applications,

- mass trapping to delimit the infestation and monitor post-treatment Mexfly populations, and
- SIT.

Program areas for Mexfly infestations are centered on Mexfly detection sites (see map of proposed program area in appendix A). Program surveillance, quarantine, and treatment boundaries may be expanded to include other properties if additional adult flies or life stages are found.

APHIS' cooperative programs to eradicate Mexfly infestations in Texas use established procedures and treatments (including delimitation and treatment) designed with the species' life stages in mind,.

1. Delimitation

The use of McPhail traps delimits the infestation and helps determine the efficacy of treatments. To do this, the traps are baited with a female attractant called biolure or natural food and preservative formulation (e.g., torula yeast and borax) and placed in varying densities inside core and buffer areas surrounding Mexfly detection sites (USDA, 2015; APHIS, 2016a). All monitoring traps will be serviced for a period equal to three Mexfly life cycles beyond the date of the last fly detection. Fruit of host plants will be sampled for the presence of eggs and larvae in a 200-meter radius around each detection site.

2. Treatment

Confirmation of a breeding Mexfly population leads to the application of a targeted, ground-based foliar bait treatment to host trees and plants within a 500-meter radius of the find site. Treatments are highly localized sprays consisting of an organic formulation of spinosad (pesticide) and protein hydrolysate (a food bait) (APHIS, 2016a). Spinosad is relatively nontoxic to mammals and beneficial arthropods; it is certified organic and has approved uses for the control of certain pests of agriculture, livestock, pets, and humans (DeAngelis, 2004). Spinosad applications in Texas occur at 7-10-day intervals for three life cycles (APHIS, 2010).

Malathion (an alternative to spinosad) has not been used for APHIS fruit fly programs in Texas for over 10 years; its use is not planned for the San Ygnacio program. APHIS allows the use of malathion in a regulatory quarantine treatment by growers in order to permit movement of their produce. Any activity or application that allows growers to apply a control measure to permit the movement of their produce must be analyzed in the EA of the program. Even though the proposed Mexfly program avoids using malathion, private growers may opt to use that chemical on their individual properties.

Protein hydrolysate is a common attractant used in fruit fly treatments, increasing the efficacy of chemical applications, and reducing the area of pesticide treatments needed for control (Prokopy et al., 1992). Pest fruit

flies are attracted to the protein hydrolysate (which can be derived from plants or yeast) where they then receive a lethal dose of the pesticide that is mixed with the attractant.

Evidence of a breeding population (immature life stages, mated female Mexfly, or multiple adult captures within a certain distance and timeframe) will result in removal of host fruit from each detection site, and from all properties within a 100-meter radius of each detection site.

SIT will be used to eradicate, as well as limit expansion of the Mexfly infestation—the eradication area receives a periodic release of sterilized male Mexflies in order to disrupt the reproduction cycle and control the wild population. The release area covers a 3.5-mile radius around each Mexfly detection site. Releases are repeated twice a week to achieve a minimum weekly release rate of 250,000 sterile Mexflies per square mile, and continue for two life cycles beyond the last Mexfly detection date (typically 4 to 6 months, dependant on temperature).

A quarantine boundary will be established to ensure any host material that leaves the program area is free of Mexfly. Host material may be treated in enclosed areas or containers by cold treatment, vapor heat treatment, irradiation, or fumigation with MB. Should the Mexfly quarantine spread to federally protected historical sites, wilderness, or tribal lands, program treatments will be restricted to those approved for the type of site in question.

Growers will be able to move their harvested fruit out of the quarantined area, under a limited permit, to enclosed facilities for processing into juice or for packing, after the fruit receives APHIS-approved MB treatment in the field or at the packing shed. Growers of host fruits may also treat their production areas using approved program treatments (field and/or premise treatment) and, under compliance agreement, have crops certified for movement to packing sheds.

Before taking action, program officials are to inform the public and impacted industry via press releases, meetings, and other forms of communication appropriate for the recipients. Residents whose property will be treated, or whose fruit will be removed, are to be notified at least 48 hours in advance. Notification letters will be sent to trading partners as they are identified. Given the potential impacts to commercial production, grove owners, packing sheds, nurseries, vendors, and other industry operations handling Mexfly-host material will be notified of the Mexfly quarantine location and treatment schedule.

For more detailed information regarding the alternatives considered for Mexfly control and their component methods, refer to the previously mentioned fruit fly risk assessments (APHIS, 2014, 2003, 1999, 1998a, and 1998b).

III. Potential Environmental Consequences

This EA analyzes the potential environmental consequences of alternatives considered for Mexfly control. The site-specific characteristics of the Mexfly program area were considered with respect to the preferred alternative's potential to affect human health, nontarget species (including threatened and endangered species), and environmental quality. Potentially sensitive sites were identified, considered, and accommodated through special selection of eradication methods and use of specific mitigation measures. APHIS will conduct any necessary additional environmental analyses if Mexfly detections lead to an expansion of the program boundary.

A. No Action

Lack of Federal action would place the burden of eradication on the State of Texas. It is reasonable to expect that Mexfly populations would continue to expand in number and area, leading to increased quarantine efforts. Any failure of those efforts could lead to the establishment of this pest within the conterminous United States. If eradication attempts are unsuccessful, APHIS expects substantial economic losses to growers in the United States. Crop loss is likely to lead to commodity scarcity, higher costs for U.S. consumers, and the temporary or permanent loss of valuable U.S. export markets.

B. Quarantine and Commodity Certification

The quarantine actions of this alternative are designed to reduce the human-mediated movement of Mexfly by preventing the transportation of host-plant materials beyond the quarantine boundary. A resident Mexfly population would be expected to remain within the quarantine boundary. Any failure in quarantine actions could lead to Mexfly establishment outside the program area. The commodity certification requirement would create a necessary but new layer of ongoing governmental presence in the marketplace. This situation could create inspection jobs; however, it would restrict trade until the produce was inspected and certified for sale. Host plants would likely cease being grown for domestic use as landowners shifted to non-Mexfly host plants.

C. Eradication Using an IPM Approach (Preferred Alternative)

This section considers the extent to which implementation of the preferred alternative might affect the human environment. It begins with a brief description of the physical aspects of the region and its residents, both

within and near the proposed program area. The preferred alternative, eradication using an IPM approach, may employ any or a combination of the following measures:

- no action,
- regulatory quarantine treatment and movement control of host materials and regulated articles,
- host survey for evidence of breeding Mexfly populations,
- host removal,
- eradication chemical applications (foliar bait spray),
- mass trapping using pheromone lures or food bait as an attractant, and
- SIT.

Female Mexican fruit fly attractants present little or no risk to human health or to the general environment, based on their low toxicity in animal testing, high target specificity, and low exposure to humans and the environment (Reilly, 2003). Review of the treatment protocols by APHIS indicates the chemical formulations used as female lures in Mexfly program traps are unlikely to result in adverse environmental or human health risks (APHIS, 2014, 2003, 1999, 1998a, and 1998b). Also, APHIS considers SIT to be a routine measure that is categorically excluded under the agency's NEPA implementing procedures (7 CFR § 372.5.i.C). Mexfly program use of SIT is not expected to result in adverse impacts to the human environment. Therefore, the discussion in chapter III will focus on the other eradication measures of the preferred alternative.

1. Affected Environment

a. Land Characteristics and Demographics

Mexfly infestions in Texas that resulted in cooperative eradication programs in 2015 and 2016 include portions of Cameron, Hidalgo, Webb, Willacy, and Zapata Counties (TDA, 2016b). San Ygnacio (Ignacio), the oldest town in Zapata County, is at the intersection of U.S. Highway 83 and Texas Farm-to-Market Road 3169, 30 miles south of Laredo and 14 miles northwest of Zapata (Heller Jr., 2015). Of the active Mexfly cooperative eradication efforts, the Laredo and Zapata Mexfly programs are nearest to the San Ygnacio Mexfly program.

Zapata County is located in South Texas in the Rio Grande Valley and is bounded by Webb County on the north, Jim Hogg County on the east, Starr County on the south, and the Rio Grande and Mexico on the west (see figure 1). Ninety-eight percent of the county is used as rangeland; urban areas and irrigated lands are concentrated near the Rio Grande (TWDB, 1996). Less than 1 percent of the county is considered prime farmland. The county's largest town and county seat is Zapata, located on the Rio Grande at the junction of U.S. Highway 83 and State Highway 16. Zapata County covers about 999 square miles, with elevations from 200 to 700 feet above sea level. The county generally has light-colored loamy

soils over reddish or mottled clayey subsoils; limestone lies in places within 40 inches of the surface. The flora includes thorny shrubs, grasses, mesquite, and cacti. Natural resources include caliche, clay, lignite coal, sand, gravel, oil, and gas. Zapata County's climate is subtropical-subhumid. The average annual temperature is 74 °F. Rainfall averages 19 inches a year, and the growing season lasts 295 days (Garza and Long, 2016).




Figure 1. Counties in southern Texas. (Source: <http://www.digital-topo-maps.com/county-map/texas-county-map.gif>)

In 2015 the U.S. Census estimated 14,374 people lived in Zapata County. In 1990 San Ygnacio had a population of 895; in 2010 it reported a population of 667. San Ygnacio is a Census-Designated Place that occupies about 1.5 square miles of land along the Rio Grande (Heller Jr., 2015; USCB, 2016a and 2016b).

Agriculture, forestry, fishing, and hunting (23%), mining, quarrying, and oil and gas extraction (18%), and construction (16%) are the primary industries employing San Ygnacio residents (City-data.com, 2016). There are 41 colonias² listed in Zapata County; 4 of these colonias contain land within the San Ygnacio Mexfly program area (see appendix C for data sources).

Recreational facilities in Zapata County include the International Falcon Reservoir, Falcon State Park, the San Ygnacio Historic District, Corralitos

² The term "colonia" in Spanish means a community or neighborhood. The Texas Secretary of State defines a "colonia" as a residential area along the Texas-Mexico border that may lack some of the most basic living necessities, such as potable water and sewer systems, electricity, paved roads, and safe and sanitary housing (Cascos, n.d.).

Ranch, and San Francisco Ranch. The Texas Tropical Trail, which links the counties of the lower Valley, runs through the area. There are extensive hunting and fishing opportunities throughout the year (Garza and Long, 2016).

The current Mexfly infestation centers on private property where fruit is grown near houses. Local land use is mainly residential, with scattered urban and light industrial districts. Mexfly host plants in the local communities are cultivated primarily for non-commercial purposes, but may represent subsistence agriculture in certain neighborhoods. (For land site information see table 1.)

Table 1. Distance from Center of Detection to Certain Land Sites.*

Designated Land Use	Distance Rounded Off to Nearest Tenth of a Mile
Nearest International Airports	<ul style="list-style-type: none"> • Laredo International Airport, 47.3 • Alice International Airport, 87.8 • McAllen International Airport, 93.4 • Kingsville Naval Air Station, 94.3
Mexico	To border, 0.2
Nearest Seaports	<ul style="list-style-type: none"> • Port Isabel: 171.0 • Port Aransas: 193.0
Colonias	<p>Within the proposed treatment area</p> <ul style="list-style-type: none"> • Valle Verde • San Ygnacio <p>Within the proposed quarantine</p> <ul style="list-style-type: none"> • San Ygnacio Viejo Unit 2 • Ramireno
City, State and Federal Lands	<p>Within the proposed treatment area</p> <ul style="list-style-type: none"> • Plaza Blas Maria Uribe • Uribe Cemetery • San Ygnacio Baseball Park
Federally Registered Historic Sites	<p>Within the proposed treatment area</p> <ul style="list-style-type: none"> • Trevino-Uribe Rancho, 0.3 • San Ygnacio Historic District, 0.3 <p>Within the proposed quarantine</p> <ul style="list-style-type: none"> • Corralitos Ranch, 5.3
Nearest Native American Reservation	<ul style="list-style-type: none"> • Kickapoo Reservation, about 120 from quarantine boundary • Ceded lands of the Comanche and Kiowa are located inside the treatment area and quarantine boundary
Organic Production and Certified Farmers Markets	None registered within 50 miles
Schools and Academic Institutions	<ul style="list-style-type: none"> • 1 inside proposed treatment area

* See appendix C for data sources.

b. Water Resources

Texas remained free of drought in June 2016, following a much-wetter-than-normal May. However, recent 100 °F heat and short-term dryness have raised concerns over the potential for a return to “flash” drought (i.e., rapidly occurring drought caused by a combination of dryness, high heat, and strong winds) (Luebehusen, 2016).

The Rio Grande is a natural river system that defines much of the international border between Mexico and the United States, forms Zapata County’s western border, and is the county’s main source of potable and irrigation water. All of the water provided by public water suppliers in the county is surface water pumped from the Rio Grande. Eight counties in the Rio Grande river basin form a State water planning area known as Region M (see figure 2). The water-use category with the largest demand in Region M is irrigation, followed by municipal (TWDB, 2016a and 2016b).




Figure 2. Regional water planning area M. (Source: TWDB, 2016a)

San Ygnacio is located within two Falcon Reservoir watersheds—El Grullo Creek (HUC 12 ID: 1308000030502) and Las Palomas Creek (HUC 12 ID: 1308000030504). Both watersheds are also within the proposed Mexfly quarantine; the El Grullo watershed is within the proposed treatment area. There are no impaired waters listed for the San Ygnacio program area. The U.S. Environmental Protection Agency defines impaired waters as “waters with chronic or recurring monitored violations of the applicable numeric or narrative water quality standards” (EPA, 2015). (Water resource data

sources provided in appendix C. See table 2 for information on other water resources as they relate to the proposed program area.)

In southern Texas, the spread of invasive aquatic weeds, international treaty issues, and increased demand are also threatening long-term water availability (LRGVDC, 2009). The vast majority of the Rio Grande water available to the region comes from supplies stored in the international Amistad and Falcon Reservoir System, owned by the United States and Mexico, and administered by the International Water and Boundary Commission (RGRWPG, 2016).

Table 2. Distance from Center of Detections to Certain Water Resources.*

Type of Resource	Distance Rounded Off to Nearest Tenth of a Mile
U.S. Wetlands within 5 miles	<ul style="list-style-type: none"> • Riverine • Lake • Fresh water, emergent • Fresh water, forested/shrub • Fresh water, pond
Water Bodies within 6 miles	<p>Within the proposed treatment area</p> <ul style="list-style-type: none"> • Falcon International Reservoir, 0.2 • El Grullo Creek, 0.3 • Rio Grande, 0.3 <p>Within the proposed quarantine</p> <ul style="list-style-type: none"> • Arroyo San Francisco, 2.7 • Las Palomas Creek, 3.2 • Rio Grande, 3.3 • Refugio Ramirez Tank, 5.1 • Marcial Tank, 5.6
Impaired Waters within 15 miles	<ul style="list-style-type: none"> • None listed

* See appendix C for data sources.

2. Human Health

The principal concerns for human health are related to potential program use of chemical pesticides, including spinosad or malathion bait, and MB (as a fumigant). Factors that influence the human health risk are associated with pesticide use, and include pesticide toxicity and exposure to humans. These factors are influenced by the use pattern and environmental fate for a particular pesticide.

Spinosad is toxic to specific invertebrate species but has low toxicity to humans and other mammals (APHIS, 2014). Limited data exists regarding the toxicity of the protein hydrolysate bait used in the spinosad formulation, however, the available data suggests low acute toxicity to human health.

Malathion is an organophosphate insecticide that targets the nervous system and acts by inhibiting acetylcholinesterase. In humans and other mammals, malathion is metabolized to its oxon (malaaxon), a more potent cholinesterase inhibitor than malathion. Carboxylesterases and other

metabolic processes detoxify malathion and malaoxon to polar, and water-soluble compounds that are excreted. Mammals are less sensitive to the effects of malathion than insects due to increased carboxylesterase compared resulting in less accumulation of malaoxon. Malathion has low acute toxicity via oral, dermal, and inhalation exposure routes. EPA classifies malathion as having “suggestive evidence of carcinogenicity, but not sufficient to assess human carcinogenic potential” (EPA, 2006). This indicates that any carcinogenic potential of malathion is so low that it cannot be quantified based upon the weight of evidence. At high doses, human health effects from malathion may include headache, nausea, vomiting, abdominal cramps, diarrhea, blurred vision, muscle weakness and twitching (ATSDR, 2003). If malathion is used, high dose exposures are not expected from program applications of malathion. Malathion may have synergistic effects when used with other organophosphate or carbamate pesticides (USFS, 2008). However, other organophosphate or carbamate pesticides are not proposed to be used in the Mexfly eradication program.

Exposure to program pesticides can vary, depending upon the pesticide and the use pattern. The Mexfly eradication program will employ ground-based targeted applications of spinosad (or malathion, if it is used) combined with protein bait. Workers who mix, load, and apply pesticides, and members of the public who live or visit the Mexfly eradication program area are the potentially exposed human populations. Proper use of personal protective equipment and engineering controls limit the exposure of program workers.

Exposure to the general public is not expected based on the targeted foliar applications and program mitigation measures. Commercial applications, should they become necessary, will be applied to properties owned by commercial growers and producers, making exposure to the general public unlikely. Residential neighborhoods and other areas of public traffic within the Mexfly eradication program receive only targeted foliar applications. Another mitigation measure designed to minimize exposure of humans to program pesticides is the requirement for public notification. Information concerning the Mexfly eradication project will be shared via press releases and media announcements to the general public. Either the county agricultural commissioner or public information officer will serve as the primary contact to the media. Any resident with property to be treated will be contacted directly or notified in writing at least 48 hours prior to treatment. Following the treatment, notices will be left with homeowners detailing precautions to take, and safe intervals of time that should elapse before harvesting fruit on the property.

Spinosad bait applications in Texas occur at 7-to 10-day intervals for three life cycles. Applications of malathion bait sprays in commercial or production orchards in Texas are conducted by a private contractor hired by

industry, and occur at 10- to 14 day intervals for three life cycles (APHIS, 2010). The potential for exposure to the general public after pesticide application is low because spinosad and malathion are not persistent in the environment with a half-life of 2.0 to 11.7 days (spinosad), and 1 to 6 days (malathion) on foliage. An additional summary of the environmental fate of the pesticides is discussed in the in the environmental quality portion of this document (chapter III, section 5).

If spinosad bait and malathion bait applications are restricted to target surfaces and made in accordance with EPA label instructions, effects to human health and the environment are expected to be incrementally negligible. The use of protein hydrolysate as an attractant in the Mexfly program is also expected to present a low risk to human health. The attractant has low toxicity and its ground-based, targeted method of application results in a low probability of exposure and risk to workers and the general public.

Should treatment by MB fumigation be indicated, adherence to EPA label restrictions and application in enclosed areas or containers will protect applicators and the general public from risk of exposure to the fumigant (APHIS, 2007 and 2002).

The analyses and data of EIS1 and EIS2, and the associated human health risk assessments indicate exposures to pesticides from normal program operations are not likely to result in substantial adverse human health effects. (Refer to EIS1 and EIS2 (APHIS, 2001 and 2008) and the human health risk assessments (APHIS, 2014, 1999 and 1998a) for more detailed information relative to human health risk.).

APHIS recognizes a small portion of the population may have greater than usual sensitivity to certain chemicals, and program treatments may pose a higher risk for these individuals. Special communication strategies to mitigate this risk are discussed in detail in appendix C of EIS1 (APHIS, 2001).

Trap placement and chemical applications may be rescheduled if rainfall or strong winds are forecast for the program area. Site inspections will continue to ensure existing program treatments are not likely to affect humans. The destruction or relocation of traps and treatments due to weather events is unlikely to result in adverse impacts to the human environment because the potential pesticide toxicity is reduced by dilution during the storm's water and air movement.

Of the three alternatives considered, a well-coordinated eradication program using IPM technologies results in the least use of chemical pesticides and minimizes their potential to adversely affect human health. The no action alternative and quarantine/commodity certification alternative are not expected to eliminate Mexfly as readily or as effectively as the preferred alternative. Over a protracted period of no action, there would likely be broader and more

widespread use of pesticides by homeowners and commercial growers, with correspondingly greater potential for adverse impacts to human health.

3. Other Aspects of the Human Environment

APHIS summarizes its findings on potential environmental impacts of implementing the action alternatives on historic sites, minority and/or low-income communities, and tribal interactions in the proposed quarantine program area in this section.

The National Historic Preservation Act of 1966, as amended (16 U.S. Code (U.S.C.) § 470 et seq.) requires Federal agencies to consider the impact on properties included in, or eligible for inclusion in, the National Register of Historic Places (36 CFR §§ 63 and 800). APHIS identified six locations listed on the National Register within Zapata County. The historic places include three ranches (Corralitos Ranch, San Francisco Ranch, and Trevino-Uribe Rancho), and a historic district where buildings may have surrounding landscaping that includes Mexfly host plants (Anon., 2016). Only one of the 19 cemeteries in Zapata County is listed as a Texas Historical Cemetery (Anon., 2016). The structures would not be affected by activities conducted under any of the alternatives analyzed in this assessment because APHIS does not anticipate any disturbance of the ground or the facility. The surrounding landscape plants may have fruit removed by hand-picking. If treatments are needed, then, where possible, bait stations would not be placed on the historic site's property. If any pesticide applications are deemed essential, then ground-based targeted bait applications or backpack sprayers would be used.

APHIS initiated consultation with the State Historic Preservation Office (SHPO) for Mexfly activities in Zapata County, Texas, in April 2016. The SHPO evaluated the Zapata County program area, and determined that historic properties would not be affected by the proposed action. To ensure historic properties will not be adversely affected by the proposed action, APHIS will not conduct aerial chemical applications at locations with historical or archeological importance. Instead, surveillance trapping and fruit stripping by hand may occur. Hand spraying with a backpack sprayer may be permitted after consultation with the SHPO. If needed, modifications of normal program activities would be designed to reduce pesticide release at these locations.

In general, APHIS' fruit fly eradication programs are compatible with the preservation of historic sites because control activities are inconspicuously integrated into the site, do not disturb the ground, and do not affect human-made structures. APHIS restricts program treatments and activities to an as-needed basis, and also can modify normal program activities at historically significant locations to reduce pesticide release, if necessary. APHIS will not conduct aerial chemical applications; spraying will be ground-based, directly targeting foliage. This may include hand spraying with a backpack sprayer. In this program, APHIS intends to use surveillance trapping or bait treatments and, when necessary, fruit stripping by hand. In addition, SIT will be used to limit expansion of the Mexfly infestation. If APHIS discovers any archaeological resources, the appropriate individuals will be notified.

Federal agencies identify and address disproportionately high and adverse human health or environmental effects of its proposed activities, as described in Executive Order 12898, *Federal Actions to Address Environmental Justice in Minority Populations and Low-Income Populations*. APHIS considered the potential environmental impacts of implementing the action alternatives on minority and/or low-income communities, tribal interactions, and historical and culturally sensitive sites in the program area. A lack of Federal action could result in adverse economic and health impacts on affected producers and consumers, such as decreased harvests, higher consumer prices, loss of local employment, reduced nutritional options, loss of market share, compromised mental and physical health, loss of property, and so on. These indirect impacts are expected to occur to a lesser extent under the quarantine and commodity certification alternative. Adverse effects are not anticipated as a result of carrying out the preferred alternative's surveillance activities, trapping, SIT, or program chemical applications.

“Colonia” is a term used in the Southwestern States to describe a subdivision where developers divide the land into small lots, and offer affordable housing to low-income families. These lots are often purchased through a contract for a deed with a low down payment and low monthly payments. The title for the house is not issued until the final payment is made by the homeowner (Anon., 2013). Housing in these locations is built by residents over time as they can afford materials. Consequently, many residences lack connections to sewers or running water, and residents may not be able to access water lines because their homes do not meet county building codes (Anon., 2013).

Residents of Zapata County reported a median household income of less than \$31,000 per year, and the median value of owner-occupied houses was estimated as below \$55,000. The percentage of residents below the poverty level from 2009–2013 in the county exceeded 32 percent (USCB, 2016b). Based on this data, APHIS finds Zapata County is a low-income area. The ability of APHIS program delivery systems to consistently minimize the potential for pesticide exposure ensures this low-income population does not disproportionately experience adverse impacts.

In Zapata County, 93.9 percent of the recorded population identifies itself as Hispanic or Latino (USCB, 2016b). To meet the linguistic needs of this group, advance notice of program activities and potential exposure hazards will be provided in Spanish to members of colonias and other non-English-speaking populations or people in areas that generally lack access to news media.

Federal agencies comply with Executive Order 13045, *Protection of Children from Environmental Health Risks and Safety Risks*. The preferred alternative does not pose any disproportionate adverse effects to children because maintenance of traps and any pesticide applications would not occur when children are present in the immediate area. The intermittent presence of children at shelters, playgrounds, parks and picnic areas, religious centers, public/private campgrounds and trailer parks, athletic fields, bus depots, and

outdoor community facilities means they are likely to frequent locations where fruit fly baits are in use; however, the placement of bait traps is likely to be far above their reach.

There are several schools within the quarantine area, and some may occur within the treatment zone (e.g., schools within the Zapata County Independent School District). Any exposure of children to applied products is negligible based on the program's application methods and the product formulations. APHIS will meet the following program requirements for every school that may require treatments:

- Maintenance of traps and any pesticide applications will occur when children are not present in the immediate area.
- The surrounding landscape plants may have fruit removed by hand-picking.
- Baits will not be applied on school property.
- A bait or backpack sprayer would be used if any pesticide applications are deemed essential.

The proposed program is not expected to result in any disproportionate adverse effects to children, minority, or low-income populations because these individuals are unlikely to be present when APHIS applies treatments or otherwise become exposed to the applied products.

Executive Order 13175, *Consultation and Coordination with Indian Tribal Governments*, calls for agency communication and collaboration with tribal officials when proposed Federal actions have potential tribal implications. The Archaeological Resources Protection Act of 1979 (16 U.S.C. §§ 470aa-mm), secures the protection of archaeological resources and sites on public and tribal lands. Using the Native American Graves Protection and Repatriation Act Online Databases (NPS, 2016; 25 U.S.C. §§ 3001 et. seq.), APHIS finds the nearest reservation is more than 250 miles away (Kickapoo Traditional Tribe of Texas). Based on the best available information, several Indian tribes ceded land in the area in treaties dating from 1865–1874 while retaining hunting interests. This suggests it is prudent for APHIS to contact the present day Comanche Nation of Oklahoma, and the Kiowa Indian Tribe of Oklahoma, in addition to the Kickapoo Tribe to ascertain their interests. APHIS met with the Kickapoo Tribe on February 4, 2013, to review the tribe's needs, interests, and concerns. During the discussion, the Tribal Administrator reiterated that the Kickapoo Tribe does not have any land holdings within the Lower Rio Grande Valley area (Duhaime, pers. comm., 01/20/2014). APHIS does not expect program activities to affect any tribes or tribal lands. The proposed action will not disturb the ground, so it is unlikely to affect Native American sites or artifacts. Should APHIS discover any archaeological resources, it will notify the appropriate individuals. APHIS is in the process of notifying the tribal

governments regarding the program activities, and will consult with tribal entities, as needed.

Individual tribal members living within the quarantine zone will not be disproportionately affected in comparison to other individuals in the area because eradication treatments are applied to host foliage above the unassisted reach of most humans, and foliar canopy spraying is highly targeted. The proposed action will not disturb the ground, so it is unlikely to affect Native American sites or artifacts. For these reasons, APHIS does not expect any tribal members to be directly affected by program activities.

4. Nontarget Species

Potential environmental impacts of alternative A (the no action alternative) or alternative B (quarantine and commodity certification) on nontarget species could include loss of animal and plant life, and habitat from unregulated pesticide use by the public, or from Mexfly host damage. Under the preferred alternative, the principal concerns for nontarget species, including threatened and endangered species, relate to potential harm from the use of program pesticides. Paralleling human health risk, the risk to nontarget species is related to the pesticides' fate in the environment, their toxicity to the nontarget species, and their exposure to nontarget species.

Current eradication activities in Zapata County will be limited to ground-based foliar applications of spinosad, combined with protein hydrolysate bait that is applied to host plants, and the use of SIT to control invasive Mexfly populations. Malathion may be used as an alternative to spinosad, although it has not been used by the Texas program for at least 10 years. The malathion and spinosad bait treatments target Mexfly life stages on host plants in a manner that minimizes potential exposure and associated risks to nontarget species.

Protein hydrolysate is a common attractant used in fruit fly treatments, increasing the efficacy of chemical applications and reducing the area of pesticide treatments needed for control (Prokopy et al., 1992). Mexflies are attracted to the protein hydrolysate where they then receive a lethal dose of the pesticide (spinosad or malathion) that is mixed with the attractant. Protein hydrolysate alone is expected to have minimal impacts to environmental quality and nontarget species because of its low toxicity.

Malathion is an organophosphate pesticide whose mode of toxic action is primarily through acetylcholinesterase inhibition (Klaassen et al., 1996; Smith, 1987). The toxicity of malathion is very slight to moderate for mammals and birds. For fish, the acute toxicity of malathion varies from moderately toxic to some species of fish, to very highly toxic to other species (Beyers and Sikoski, 1994; Mayer and Ellersieck, 1986; USFS, 2008). Malathion is moderately to very highly toxic to most aquatic invertebrates on an acute basis, depending on the sensitivity of the species.

Spinosad has low to moderate toxicity to wild mammals and birds. Spinosad's toxicity to fish is moderate, while aquatic invertebrates are more sensitive in acute and chronic exposures. Toxicity to terrestrial invertebrates is variable; however, spinosad is considered highly toxic to honey bees. Risks to nontarget species from the use of either malathion or spinosad baits are anticipated to be negligible because the proposed use pattern (targeted, hand application of the bait) results in a low potential for exposure to most taxa. The bait applications attract only a small number of invertebrate species other than Mexfly. (Refer to EIS1 and EIS2 (APHIS, 2001, 2008) and the supporting nontarget risk assessments (APHIS, 2014, 2003, 1998) for more information on risks to all classes of nontarget species.)

Conservation areas in Zapata County provide important habitat for a wide variety of wildlife. For example, Zapata County contains a portion of the Lower Rio Grande Valley National Wildlife Refuge. Falcon State Park and the Falcon International Reservoir provide habitat for wildlife including birds and fish in the county. Seed eater Sanctuary is a Zapata County park where white-collared seed eaters breed in the cane beds along the river.

APHIS' Mexfly programs are designed to prevent the introduction of program chemicals into non-targeted areas. Sites near the program area that might require special consideration, should the program area expand, include irrigation canals or wetlands of potential ecological importance. No program chemical applications will be permitted at these sites or other protected areas. However, surveillance trapping will continue, and fruit stripping by hand and SIT will be undertaken if Mexfly detections occur at such locations. Pesticide applications would only occur in national wildlife refuges with the approval of the U.S. Fish and Wildlife Service (FWS) and coordination with the refuge manager.

a. Migratory Birds

Unless permitted by regulation, the Migratory Bird Treaty Act of 1918 (16 U.S.C. §§ 703–712) provides that it is unlawful to pursue, hunt, take, capture, kill, attempt to take, capture or kill, possess, offer for sale, sell, offer to purchase, purchase, deliver for shipment, ship, cause to be shipped, deliver for transportation, transport, cause to be transported, carry, or cause to be carried by any means whatever, receive for shipment, transportation or carriage, or export, at any time, or in any manner, any migratory bird or any part, nest, or egg of any such bird.

Executive Order 13186, Responsibilities of Federal Agencies to Protect Migratory Birds, directs Federal agencies taking actions with a measurable negative effect on migratory bird populations to develop and implement a Memorandum of Understanding (MOU) with FWS that promotes the conservation of migratory bird populations. On August 2, 2012, an MOU

between FWS and APHIS was signed to facilitate the implementation of this Executive order.

Approximately 290 bird species are found in Zapata County, including the red-billed pigeon, Audubon's oriole, Cassin's sparrow, and the varied bunting, green jay, great kiskadee, vermilion flycatcher, hooded oriole, and black phoebe (ZCC, 2012). APHIS evaluated the proposed Mexfly program in terms of potential impact on migratory avian species. Implementation of the preferred alternative is not expected to have any adverse effect on migratory birds or their flight corridors.

Malathion is only slightly to moderately toxic to birds, and spinosad's acute and chronic toxicity to birds is low (APHIS, 2014). The targeted application of the insecticide baits to Mexfly host plants within 500 meters of Mexfly detections (usually in residential areas) and the short half-life of malathion on vegetation would result in limited to no exposure of birds to malathion. The localized and direct application of malathion and spinosad baits to host plants would not result in any impacts to food of birds. Birds would not be exposed to methyl bromide treatments. SIT would provide an additional food source to migratory birds.

FWS recommends that any activities requiring vegetation removal or disturbance should avoid the peak nesting period of March through August to avoid destruction of individual birds, nests, or eggs (FWS, 2015). If project activities must be conducted during this time, FWS recommends surveying for nests prior to commencing work. If a nest is found, if possible, FWS recommends a buffer of vegetation (≥ 50 feet) remain around the nest until young have fledged or the nest is abandoned.

b. Endangered Species Act

Section 7 of the Endangered Species Act (ESA) and ESA's implementing regulations require Federal agencies to consult with FWS and/or the National Marine Fisheries Service to ensure that their actions are not likely to jeopardize the continued existence of threatened or endangered species or result in the destruction or adverse modification of critical habitat.

There are 5 federally listed species in Zapata County: ocelot (*Leopardus pardalis*), Gulf Coast jaguarundi (*Felis yagouaroundi*), least tern (*Sterna antillarum*), ashy dogweed (*Thymophylla tephroleuca*), and Zapata bladderpod (*Lesquerella thamnophila*). APHIS contacted the FWS, Ecological Services Field Offices in Corpus Christi and Alamo, Texas, regarding the proposed treatment. The FWS reviewed a map of the quarantined area, and indicated that their only concern is the critical habitat of the Zapata bladderpod (E. Reyes, U.S. Fish and Wildlife Service, pers. comm., July 15, 2016). Critical habitat for the Zapata bladderpod does not occur within the currently quarantined area, and only occurs on national wildlife refuge property in Zapata County. The treatment areas where

spinosad or malathion will be applied are on private property in dooryard fruit. Local land use is mainly residential, with scattered urban and light industrial districts, and is not habitat for listed species. Therefore, APHIS has determined that the proposed program will have no effect on listed species or critical habitat in the current treatment areas. Should the program area expand or further outbreaks be detected that are not considered herein, APHIS will consult with FWS and other appropriate agencies, as necessary. In the meantime, APHIS has submitted a programmatic biological assessment requesting consultation on threatened and endangered species and their critical habitats in Zapata and Webb Counties to ensure that protection measures are in place should program activities co-occur with listed species in those counties. APHIS has had a programmatic consultation in place since 2009 for Mexfly program activities for Willacy, Hidalgo, and Cameron Counties.

5. Environmental Quality

The principal environmental quality concerns are for the protection of air quality, water quality, and the minimization of the potential for environmental contamination. In relation to preserving environmental quality, program pesticides remain the major concern for the public and the program. Although program pesticide use is limited, especially in comparison to other agricultural pesticide use, the proposed action would result in a controlled release of chemicals into the environment. The fate of those chemicals varies with respect to the environmental component (air, water, or other substrate) and its characteristics (temperature, pH, dilution, etc.). The environmental fates of spinosad, malathion, and MB are outlined below. (Refer to EIS1 and EIS2 (APHIS, 2001 and 2008) and the risk assessments (APHIS, 2014, 2003, 1999, 1998a and 1998b) for a more detailed consideration of program pesticides' environmental fates.)

Attractant ingredients (e.g., yeast and protein hydrolysate baits) have minimal affect on environmental quality, based on EPA-approved use patterns and the ingredients' rapid degradation. Use of these attractants in Mexfly program treatments is not expected to result in impacts to environmental quality beyond those described for the below chemicals.

- **Spinosad** is not considered mobile in soil as it adsorbs strongly to soil particles and is unlikely to leach to great depths. Dissipation half-lives for spinosad in the field may last 0.3 to 0.5 day. It is photodegraded quickly on soil exposed to sunlight. Spinosad is quickly metabolized by soil micro-organisms under aerobic conditions, and has a half-life of 9.4 to 17.3 days. Spinosad is not sensitive to hydrolysis, but aqueous photolysis is rapid in natural sunlight (half-life of less than 1.0 to 1.6 days), and is the primary route of degradation in aquatic systems exposed to sunlight. Under anaerobic conditions, the degradation rate is slower, between 161 and 250 days. Spinosad has a half-life of 2.0 to 11.7 days on plant surfaces. After initial photodegradation, residues are available for metabolism by plant biochemical processes. Effects from residues

of individual treatments are no longer detectable in environmental substrates within a few weeks of application (APHIS, 2014; Kollman, 2003).

- **Malathion** is toxic to many nontarget species; it is used less widely than spinosad, and primarily by commercial growers on private property. Malathion is considered lower in toxicity and less persistent (1 to 25 days in soil) than other organophosphorus pesticides. In water, malathion has a half-life of approximately 1 week, and is more stable in acidic aquatic conditions. Malathion is soluble in water, and can be highly mobile in soil. Generally, degradation occurs rapidly (a half-life of less than 1 to nearly 9 days) (Gervais et al., 2009); application to foliage allows for exposure of residues to degradation from processes (e.g., photolysis), resulting in a reduced potential for significant movement to ground water. Malaoxon is an oxygen analogue of malathion, and it can be found either as an impurity in malathion products, or can be generated during the oxidation of malathion in air or soil. Malathion and malaoxon can be transported in air over large distances and elevations (Newhart, 2006).
- **Methyl bromide (MB)** fumigation will not be used as an eradication treatment, but may be employed as a regulatory treatment. MB volatilizes into air from soil and water, and is known to contribute to stratospheric ozone depletion. The volatilization half-life for MB from surface water ranges from 3.1 hours to 5 days. The degradation half-life of MB in water ranges from 20 to 38 days, depending on temperature and pH. Volatilization of MB from surface soil is rapid, with a half-life ranging from 0.2 to 0.5 day. The degradation half-life of MB in soil ranges from 31 to 55 days. MB has a low affinity to bind to soils, however, is not considered a major contaminant of ground water (NPIC, 2000). The small quantities used to treat for Mexfly disperse when fumigation chambers are vented. (See section 6 of chapter 3 regarding MB's potential cumulative impacts to the environment.)

Urban and agricultural runoff may flow directly into local waters, picking up trash, dirt, chemicals, and other contaminants along the way. If treatment is indicated in close proximity to a body of water where pesticides might be directly discharged into the water, TDA will analyze the environmental setting, and establish and follow site-specific best management practices. The prescribed method of spray application directly to host plants is designed to minimize drift and runoff. Mitigation measures will be applied to protect marine and freshwater resources. Personnel will maintain a minimum distance of 98 feet (30 meters) from surface water. Personnel applying pesticides will adhere to label directions, State and Federal laws, and recommendations of the environmental

compliance staff associated with the program. Water body contact is not anticipated due to the targeted application measures and the environmental fate of the pesticides used in *Anastrepha* spp. cooperative eradication programs.

The alternatives (A) no Federal action, (B) quarantine and commodity certification, and (C) the preferred alternative, eradication using an IPM approach, were compared with respect to their potential to affect environmental quality. Risk to environmental quality is considered minimal. Again, a well-coordinated eradication program using IPM technologies would result in the least use of chemical pesticides overall, with minimal adverse impacts on environmental quality. The no action alternative and the quarantine and commodity certification alternative would likely result in broader and more widespread use of pesticides by homeowners and commercial growers, with correspondingly greater potential for adverse impacts.

The proposed program area was examined to identify characteristics that would tend to influence the effects of program operations. Potentially sensitive areas were identified, considered, and accommodated, as necessary, through special selection of control methods and use of specific mitigation measures. Allowances were made for the special site-specific characteristics that would require a departure from the standard operating procedures. The approaches used to mitigate for adverse impacts to bodies of water are described in EIS1 (APHIS, 2001).

6. Cumulative Impacts

This section considers the potential of the alternatives to cause cumulative impacts on the human environment. Not taking Federal action is expected to result in the cumulative impacts that arise from tolerating uncontrolled Mexfly infestations in the United States. Federal quarantine restrictions and commodity certification requirements would place the burden of control efforts and expense on producers already engaged in complying with other quarantine and commodity certification requirements. Either of these alternatives may increase the time it takes for commodities to reach their intended markets, or may prevent them from reaching consumers at all, which may contribute to consumer shortages and negative public perception of the affected industry.

APHIS considered implementation of the preferred alternative in the context of, and in conjunction with, other pest insect eradication and quarantine projects in the Zapata program area (e.g., pink bollworm and cattle fever tick eradication efforts). These programs use pesticides with different chemistries. They target different pests, and are applied at different times. The combination of these different pesticide chemistries, targets for application, and application timings suggest limited interacting or multiple exposures that are not likely to create significant cumulative impacts in the human environment.

Current and future in-State Mexfly programs could potentially merge into one larger program area. When Mexfly eradication programs are combined with trapping and eradication actions across Texas counties, APHIS expects a beneficial cumulative impact on the environment from reduced Mexfly populations causing damage to fruit combined with overall fewer chemical treatments. Trapping and surveys for Mexfly continue under the Texas fruit fly detection and monitoring program, and sterile Mexflies continue to be released over high-risk regions as a preventive measure.

The San Ygnacio Mexfly program activities and treatments are designed not to overlap (in time or space) with similar activities and treatments. Pesticide use in a Mexfly control program that overlaps with another *Anastrepha* spp. program is monitored and adjusted, where necessary, to minimize environmental impacts. During 2014, for example, infestations of West Indian fruit fly (*Anastrepha obliqua* (Macquart)) were detected in areas being treated for Mexfly. The eradication program protocols for West Indian fruit fly being the same as that for Mexfly, no additional chemical treatments were considered necessary. Due to the passage of time and the prevailing weather conditions in southern Texas during 2015 and early 2016, no chemical residues are believed to remain from previous Mexfly programs that could result in additive or synergistic chemical effects with previous program chemical applications.

The Mexfly programs for Zapata County, Webb County, and the LRGV were examined for potential synergistic and cumulative environmental impacts. APHIS considered the potential for pesticide exposure from the use of pesticides on adjacent properties and in residential and community gardens, as well as urban communities in close proximity to commercial cultivation of Mexfly-host plant species (Belson et al., 2003; Donnelly and Cizmas, 2007).

Malathion is a pesticide approved for use against Mexfly but is unlikely to be employed in program treatments. Malathion is also a regulatory treatment for the Texas cotton boll weevil eradication program. Uses of malathion by a Mexfly program within the Texas boll weevil quarantine (currently active in the counties of Brooks, Cameron, Hidalgo, Jim Hogg, Kenedy, Maverick, Starr, Webb, Willacy, and Zapata) should, therefore, be monitored and adjusted, where necessary, in order to minimize environmental impacts (TBWEF, 2016). Other treatments for potentially overlapping eradication programs in southern Texas target different arthropod species, and do not affect the same nontarget organisms (TDA, 2016c).

No significant environmental impacts are expected to result from proper implementation of this Mexfly eradication and control program. The differences in pesticide chemistries, targets for application, affected species and resources, and application timing between the Mexfly program and other pest control programs in Texas are not likely to create significant

cumulative impacts in the human environment. No synergistic, additive, or cumulative impacts from pesticide applications in the San Ygnacio Quarantined Area are expected with the following active control programs (TDA, 2016c)—

- **Asian citrus psyllid:** Quarantine over the entire State; pesticide applications in the citrus-growing areas of eight counties, including Zapata.
- **Citrus greening:** Quarantine over six counties, not including Zapata.
- **Exotic fruit fly species:** Mexfly programs in Cameron, Hidalgo, Webb, Willacy, and Zapata Counties.

Care should be taken, however, when multiple pest species in the same area are targeted for treatment using the same chemical. Spinosad, for example, has other labeled food and non-food uses, and is currently used in a variety of pest control efforts, including the control of termites and European grapevine moth (APHIS, 2014). Implementation of a Mexfly eradication program could lead to an increase in spinosad use, and the possible overlap of APHIS and non-APHIS program treatments. The Mexfly treatment schedule will be adjusted in locations where another CDFA or APHIS program may have scheduled similar treatments to avoid additive chemical impacts.

It is uncertain how pesticides may be used by private entities in the Mexfly program area. In terms of Federal and Texas State program activity, there are no significant cumulative impacts anticipated as a consequence of implementing the preferred alternative or its component treatment measures. The preferred alternative is designed for pesticide applications to avoid overlapping treatment areas, and to prevent nontarget exposure until pesticide residues are degraded.

APHIS determined uses of MB for fruit fly quarantine treatments pose negligible potential for cumulative impacts to the environment. For information on potential depletion of the ozone layer related to MB released into the atmosphere, see the *Rule for the Importation of Unmanufactured Wood Articles from Mexico, with Consideration for Cumulative Impact of Methyl Bromide Use, Final Environmental Impact Statement* (APHIS, 2002) and subsequent analyses, such as the *Importation of Solid Wood Packing Material, Supplement to the Final Environmental Impact Statement* (APHIS, 2007).

There were no residual impacts from previous Federal and non-Federal actions targeting fruit fly infestations in the State of Texas, and there are no reasonably foreseeable future actions that could result in incremental increases in environmental effects. Based on APHIS' review of the context and intensity of the existing, ongoing, and potential future treatments, there

will be no cumulative impacts to the human environment resulting from this Mexfly cooperative eradication program.

As discussed previously, additional actions may be implemented in this program, including additional quarantines and regulatory treatments. The anticipated use of these treatments is considered to pose minimal risk to the human environment, as determined in EIS1 and EIS2 (APHIS, 2001 and 2008), and the nontarget species and human health risk assessments (APHIS, 2014, 2003, 1999, 1998a, and 1998b).

IV. Agencies Consulted

State Historic Preservation Officer
Texas Historical Commission
108 W. 16th Street
Austin, TX 78701

U.S. Department of Agriculture
Animal and Plant Health Inspection Service
Plant Protection and Quarantine
Center for Plant Health Science and Technology
1730 Varsity Drive, Suite 400
Raleigh, NC 27606

U.S. Department of Agriculture
Animal and Plant Health Inspection Service
Policy and Program Development
Environmental Risk and Analysis Services
4700 River Road, Unit 149
Riverdale, MD 20737

U.S. Fish and Wildlife Service
Texas Coastal Ecological Services
P.O. Box 81468
Corpus Christi, TX 78468

V. References Cited

Anon.—See Anonymous

Anonymous, 2016. Texas Historic Sites Atlas. Texas Historical Commission. [Online]. Available: <http://atlas.thc.state.tx.us/> [2016, July 8].

Anonymous, 2013. Office of the Texas Secretary of State. Colonias FAQs. [Online]. Available: <http://www.sos.state.tx.us/border/colonias/faqs.shtml> [2016, July 8].

APHIS—See U.S. Department of Agriculture, Animal and Plant Health Inspection Service

ATSDR—See U.S. Department of Health and Human Services, Public Health Service, Agency for Toxic Substances and Disease Registry

Beyers, D.W., and Sikoski, P.J., 1994. Acetylcholinesterase inhibition in federally endangered Colorado squawfish exposed to carbaryl and malathion. *Environ. Toxicol. and Chem.* 13(6): 935–939.

Belson, M., Kieszak, S., Watson, W. Blindauer, K.M., Phan, K., Backer, L., and Rubin, C., 2003. Childhood Pesticide Exposures on the Texas-Mexico Border: Clinical Manifestations and Poison Center Use. *American Journal of Public Health* 93(8): 1310-1315. 6 pp.

California Department of Food and Agriculture, n.d. Mexican Fruit Fly Pest Profile. [Online]. Available: https://www.cdffa.ca.gov/plant/pdep/target_pest_disease_profiles/mexican_ff_profile.html [2016, July 5].

Cascos, C. H., n.d. What is a Colonia? [Online]. Available: http://www.sos.state.tx.us/border/colonias/what_colonia.shtml [2016, July 8].

CDFA—See California Department of Food and Agriculture

City-data.com, 2016. San Ygnacio, Texas. [Online]. Available: <http://www.city-data.com/city/San-Ygnacio-Texas.html> [2016, July 5].

DeAngelis, J., 2004. Spinosad is a natural insecticide. [Online]. Available: <http://www.livingwithbugs.com/spinosad.html> [2016, July 5].

Donnelly, K.C., and Cizmas, L., 2007. Texas A&M University - Border Health Environmental Coordination Program – August, 2007. U.S. EPA Archive Document. 5 pp.

EIS1—See U.S. Department of Agriculture, Animal and Plant Health Inspection Service, 2001

EIS2—See U.S. Department of Agriculture, Animal and Plant Health Inspection Service, 2008

EPA—See U.S. Environmental Protection Agency

FWS—See U.S. Department of the Interior, Fish and Wildlife Service

Garza, A.A., and Long, C., 2016. Handbook of Texas Online, "Zapata County." Uploaded on June 15, 2010. Modified on February 22, 2016. Published by the Texas State Historical Association [Online]. Available: <http://www.tshaonline.org/handbook/online/articles/hcz01> [2016, July 5].

Gervais, J.A., Luukinen, B., Buhl, K., Stone, D., 2009. Malathion Technical Fact Sheet. National Pesticide Information Center, Oregon State University Extension Services. [Online]. Available: <http://npic.orst.edu/factsheets/archive/malatech.html> [2016, July 5].

Heller Jr., D.D., 2015. Handbook of Texas Online, "San Ygnacio, TX." Uploaded on June 15, 2010. Modified on July 24, 2015. Published by the Texas State Historical Association. [Online]. Available: <http://www.tshaonline.org/handbook/online/articles/hls15> [2016, July 5].

Klaassen, C.D., Amdur, M.O., and Doull, J., 1986. Casarett and Doull's Toxicology, the basic science of poisons. 5th ed., Macmillan Publishing Co., New York.

Kollman, W.S., 2003. Environmental Fate of Spinosad. California Department of Pesticide Regulation, Environmental Monitoring Branch. [Online]. Available: http://www.cdpr.ca.gov/docs/emon/pubs/fatememo/spinosad_fate.pdf [2016, July 5].

Lower Rio Grande Valley Development Council, 2009. Water Resources. [Online]. Available: <http://www.lrgvdc.org/water.html> [2016, July 5].

LRGV—Lower Rio Grande Valley

LRGVDC—See Lower Rio Grande Valley Development Council

Luebehusen, E., 2016. U.S. Drought Monitor: June 28, 2016. (Released Thursday June 30, 2016) Valid 8 a.m. EDT. [Online]. Available: <http://droughtmonitor.unl.edu/Home.aspx> [2016, July 5].

Mayer, F.L., Jr., and Ellersieck, M.C., 1986. Manual of acute toxicity: interpretation and data base for 410 chemicals and 66 species of freshwater animals. Resour. Publ. 160. U.S. Department of the Interior, Fish and Wildlife Service, Washington, DC.

NAPIS—See National Agricultural Pest Information System

National Agricultural Pest Information System, n.d. Pest Tracker. Mexican Fruit Fly (Mesfly), *Anastrepha ludens*. [Online]. Available: <http://pest.ceris.purdue.edu/pest.php?code=IOBMABA> [2016, July 5].

National Pesticide Information Center, 2000. Methyl Bromide. Technical fact sheet. [Online]. Available: <http://npic.orst.edu/factsheets/MBgen.pdf> [2016, July 5].

Newhart, K., 2006. Environmental fate of malathion. California Environmental Protection Agency, Department of Pesticide Regulation, Environmental Monitoring Branch. Oct. 11, 2006. [Online]. Available: http://www.cdpr.ca.gov/docs/emon/pubs/fatememo/efate_malathion.pdf [2016, July 5].

NPIC—See National Pesticide Information Center

NPS—See U.S. National Park Service

Prokopy, R.J., Papaj, D.R., Hendrichs, J., and Wong, T.T.Y., 1992. Behavioral responses of *Ceratitidis capitata* flies to bait spray droplets and natural food. *Entomologia Experimentalis et Applicata* 64: 247–257.

Reilly, S.K., 2003. Pheromones: risk assessment and decision making. U.S. Environmental Protection Agency presentation to the NAFTA Technical Working Group on Pesticides. 15 MS PowerPoint slides.

RGRWPG—See Rio Grande Regional Water Planning Group

Rio Grande Regional Water Planning Group, 2016. 2016 Rio Grande Regional Water Plan: Volume I. December 1, 2015. 553 pp.

Smith, G.J., 1987. Pesticide use and toxicology in relation to wildlife: organophosphate and carbamate compounds. Resource Publ. 170. U.S. Department of the Interior, Fish and Wildlife Service, Washington, DC.

TBWEF—See Texas Boll Weevil Eradication Foundation

TDA—See Texas Department of Agriculture

Texas Boll Weevil Eradication Foundation, 2016. Lower Rio Grande Valley Zone. [Online]. Available: <http://txbollweevil.org/Zones/lrgv.html> [2016, July 5].

Texas Department of Agriculture, 2016a. School Integrated Pest Management (IPM). [Online]. Available: <http://www.texasagriculture.gov/regulatoryprograms/pesticides/structuralpestcontrolservice/schoolintegratedpestmanagement.aspx> [2016, July 5].

Texas Department of Agriculture, 2016b. Mexican Fruit Fly. Mexican Fruit Fly Quarantine. [Online]. Available: <http://www.texasagriculture.gov/RegulatoryPrograms/PlantQuality/PestandDiseaseAlerts/MexicanFruitFly.aspx> [2016, July 5].

- Texas Department of Agriculture, 2016c. Regulatory programs: Plant quality. Pest and disease alerts. [Online]. Available: <http://www.texasagriculture.gov/RegulatoryPrograms/PlantQuality/PestandDiseaseAlerts.aspx> [2016, July 5].
- Texas Water Development Board, 2016a. Rio Grande Regional Water Planning Group: 2016 Region M Water Plan. Chapter 1: Description of Regional Water Planning Area. 30 pp.
- Texas Water Development Board, 2016b. 2017 State Water Plan: Water for Texas. Transitional Documentation of Final Plan as Adopted by the TWDB on May 19, 2016. 164 pp.
- TWDB—See Texas Water Development Board
- UFL—See University of Florida
- University of Florida, 2012. Mexican fruit fly—*Anastrepha ludens* (Loew) (Insecta: Diptera: Tephritidae). Publication EENY-201. Reviewed August 2015. [Online]. Available: http://entnemdept.ufl.edu/creatures/fruit/tropical/mexican_fruit_fly.htm [2016, July 5].
- U.S. Census Bureau, 2016a. Community Facts. San Ygnacio CDP, Texas. [Online]. Available: http://factfinder.census.gov/faces/nav/jsf/pages/community_facts.xhtml [2016, July 5].
- U.S. Census Bureau, 2016b. QuickFacts: Zapata County/Zapata CDP/Laredo City, Texas. [Online]. Available: <https://www.census.gov/quickfacts/table/PST045215/48505,4880716,4841464> [2016, July 5].
- U.S. Department of Agriculture, 2015. National Exotic Fruit Fly Detection Trapping Guidelines. Prepared by employees of State and Federal plant protection and regulatory agencies. First edition: June 2015. 126 pp.
- U.S. Department of Agriculture, Animal and Plant Health Inspection Service, 2016a. Plant Protection and Quarantine. San Ygnacio, Zapata County, Texas MXFF Fruit Fly Outbreak Information Needed Within 96 Hours 06/30/2016. 2 pp.
- U.S. Department of Agriculture, Animal and Plant Health Inspection Service, 2016b. Mexican Fruit Fly Cooperative Eradication Program, Zapata County, Texas. Environmental Assessment and FONSI—April 2016. Riverdale, MD.
- U.S. Department of Agriculture, Animal and Plant Health Inspection Service, 2015. Plant Protection and Quarantine. SPRO letter. Subject: Removal of Mexican Fruit Fly (*Anastrepha ludens*) Quarantine Areas in Rangerville Cameron County and McAllen, Mission and La Villa, Hidalgo County, Texas. For information and action. DA-2015-60. December 10, 2015.

- U.S. Department of Agriculture, Animal and Plant Health Inspection Service, 2014. Human health and ecological risk assessment for STATIC™ spinosad ME Bait Applications. March 2014. Riverdale, MD.
- U.S. Department of Agriculture, Animal and Plant Health Inspection Service, 2010. United States and Mexico Lower Rio Grande Valley Mexican Fruit Fly Program Review: Final Report. Based on August 2009 site visits by expert review panel. Coordinator: Edward F. Gersabeck, Riverdale, MD. 130 pp.
- U.S. Department of Agriculture, Animal and Plant Health Inspection Service, 2008. Use of Genetically Engineered Fruit Fly and Pink Bollworm in APHIS Plant Pest Control Programs. Final Environmental Impact Statement—October 2008. Riverdale, MD.
- U.S. Department of Agriculture, Animal and Plant Health Inspection Service, 2007. Importation of Solid Wood Packing Material. Supplement to the Final Environmental Impact Statement—October 2007. Riverdale, MD.
- U.S. Department of Agriculture, Animal and Plant Health Inspection Service, 2003. Spinosad Bait Spray Applications. Nontarget Risk Assessment. October 2003. USDA–APHIS, Riverdale, MD.
- U.S. Department of Agriculture, Animal and Plant Health Inspection Service, 2002. Rule for the Importation of Unmanufactured Wood Articles From Mexico, with Consideration for Cumulative Impact of Methyl Bromide Use. Final Environmental Impact Statement—September 2002. Riverdale, MD.
- U.S. Department of Agriculture, Animal and Plant Health Inspection Service, 2001. Fruit Fly Cooperative Control Program, Final Environmental Impact Statement—2001. USDA–APHIS, Riverdale, MD.
- U.S. Department of Agriculture, Animal and Plant Health Inspection Service, 1999. Spinosad Bait Spray Applications. Human Health Risk Assessment. March 1999. USDA–APHIS, Riverdale, MD.
- U.S. Department of Agriculture, Animal and Plant Health Inspection Service, 1998a. Human Health Risk Assessment for Fruit Fly Cooperative Control Programs. USDA–APHIS, Riverdale, MD.
- U.S. Department of Agriculture, Animal and Plant Health Inspection Service, 1998b. Nontarget Species Risk Assessment for Fruit Fly Cooperative Control Programs. USDA–APHIS, Riverdale, MD.
- U.S. Department of Agriculture, Forest Service, 2008. Malathion – human health and ecological risk assessment – final report. Submitted by Patrick R. Durkin of Syracuse Environmental Research Associates, Inc. on May 12, 2008. SERA TR–052–02–02c. 328 pp.

U.S. Department of Health and Human Services, Public Health Service, Agency for Toxic Substances and Disease Registry, 2003. Toxicological Profile for Malathion. September 2003. 327 pp.

U.S. Environmental Protection Agency, 2015. Terms and Acronyms: search results. Last updated September 1, 2015. [Online]. Available: https://iaspub.epa.gov/sor_internet/registry/termreg/searchandretrieve/termsandacronyms/search.do [2016, Apr. 14].

U.S. Environmental Protection Agency, 2006. Appendix J. July 31, 2006 memorandum. Subject: Malathion: Revised Human Health Risk Assessment for the Reregistration Eligibility Decision Document (RED). PC Code: 057701. Case No. 0248. DP Barcode: D330680. 171 pp.

USCB—See U.S. Census Bureau


USDA—See U.S. Department of Agriculture

USFS—See U.S. Department of Agriculture, Forest Service

U.S. National Park Service, 2016. National NAGPRA online databases. [Online]. Available: <http://www.nps.gov/nagpra/onlinedb/index.htm> [2016, July 8].

Zapata Chamber of Commerce. 2012. Zapata is for the birds! [Online]. Available: <http://www.zapatachamber.com/index.php/attractions/birding> [2016, July 18].

Appendix A. San Ygnacio Mexfly Program Area, Texas—as of June 30, 2016



Source: USDA-APHIS-PPQ

Appendix C. Outside-APHIS Spatial Data Resources Used to Prepare this Document

The following resources were used by USDA-APHIS-PPD, 1 July 2016.

- **Crop data:** <http://nassgeodata.gmu.edu/CropScape/>
- **Historic sites:** <http://www.nps.gov/nr/research/>
- **Migratory birds, wetlands:** <http://ecos.fws.gov/ipac/>
- **National wildlife refuges, Native American areas:**
<http://viewer.nationalmap.gov/>
- **Organic farms:** <http://www.ams.usda.gov/AMSV1.0/nop>
- **US FWS critical habitat:** <http://ecos.fws.gov/crithab/>
- **Waters, superfund sites, historic sites, schools, demographics:**
<http://nepassisttool.epa.gov/nepassist/entry.aspx>