

DRAFT - Environmental Assessment Rangeland Grasshopper and Mormon Cricket Suppression Program

ARIZONA
EA Number: AZ-21-01

Prepared by:

Animal and Plant Health Inspection Service
3640 East Wier Ave.
Phoenix, Arizona 85040

Site-Specific

Graham and Gila County portion within the San Carlos Apache Reservation.

January 25, 2021

Non-Discrimination Policy

The U.S. Department of Agriculture (USDA) prohibits discrimination against its customers, employees, and applicants for employment on the bases of race, color, national origin, age, disability, sex, gender identity, religion, reprisal, and where applicable, political beliefs, marital status, familial or parental status, sexual orientation, or all or part of an individual's income is derived from any public assistance program, or protected genetic information in employment or in any program or activity conducted or funded by the Department. (Not all prohibited bases will apply to all programs and/or employment activities.)

To File an Employment Complaint

If you wish to file an employment complaint, you must contact your agency's EEO Counselor (PDF) within 45 days of the date of the alleged discriminatory act, event, or in the case of a personnel action. Additional information can be found online at http://www.ascr.usda.gov/complaint_filing_file.html.

To File a Program Complaint

If you wish to file a Civil Rights program complaint of discrimination, complete the USDA Program Discrimination Complaint Form (PDF), found online at http://www.ascr.usda.gov/complaint_filing_cust.html, or at any USDA office, or call (866) 632-9992 to request the form. You may also write a letter containing all of the information requested in the form. Send your completed complaint form or letter to us by mail at U.S. Department of Agriculture, Director, Office of Adjudication, 1400 Independence Avenue, S.W., Washington, D.C. 20250-9410, by fax (202) 690-7442 or email at program.intake@usda.gov.

Persons With Disabilities

Individuals who are deaf, hard of hearing, or have speech disabilities and you wish to file either an EEO or program complaint please contact USDA through the Federal Relay Service at (800) 877-8339 or (800) 845-6136 (in Spanish).

Persons with disabilities who wish to file a program complaint, please see information above on how to contact us by mail directly or by email. If you require alternative means of communication for program information (e.g., Braille, large print, audiotape, etc.) please contact USDA's TARGET Center at (202) 720-2600 (voice and TDD).

Mention of companies or commercial products in this report does not imply recommendation or endorsement by USDA over others not mentioned. USDA neither guarantees nor warrants the standard of any product mentioned. Product names are mentioned to report factually on available data and to provide specific information.

This publication reports research involving pesticides. All uses of pesticides must be registered by appropriate State and/or Federal agencies before they can be recommended.

CAUTION: Pesticides can be injurious to humans, domestic animals, desirable plants, and fish and other wildlife—if they are not handled or applied properly. Use all pesticides selectively and carefully. Follow recommended label practices for the use and disposal of pesticides and pesticide containers

Table of Contents

I.	Need for Proposed Action.....	1
A.	Purpose and Need Statement	1
B.	Background Discussion	2
C.	About This Process	5
II.	Alternatives	6
A.	No Suppression Program Alternative	7
B.	Insecticide Applications at Conventional Rates or Reduced Agent Area Treatments with Adaptive Management Strategy (Preferred Alternative)	7
III.	Affected Environment.....	12
A.	Description of Affected Environment.....	12
B.	Site-Specific Considerations.....	13
1.	Human Health	13
2.	Nontarget Species	14
3.	Socioeconomic Issues	17
4.	Cultural Resources and Events	18
5.	Special Considerations for Certain Populations	18
IV.	Environmental Consequences.....	19
A.	Environmental Consequences of the Alternatives	19
1.	No Suppression Program Alternative	19
2.	Insecticide Applications at Conventional Rates or Reduced Agent Area Treatments with Adaptive Management Strategy.....	20
B.	Other Environmental Considerations.....	28
1.	Cumulative Impacts	28
2.	Executive Order No. 12898, Federal Actions to Address Environmental Justice in Minority Populations and Low-Income Populations	29
3.	Executive Order No. 13045, Protection of Children from Environmental Health Risks and Safety Risks	29
4.	Tribal Consultation	30
5.	Executive Order 13186, Responsibilities of Federal Agencies to Protect Migratory Birds	30
6.	Endangered Species Act	31
7.	Bald and Golden Eagle Protection Act	32
8.	Additional Species of Concern	32
9.	Fires and Human Health Hazards	33
10.	Cultural and Historical Resources	33
V.	Literature Cited	33
VI.	Listing of Agencies and Persons Consulted.....	41
	Appendix A : APHIS Rangeland Grasshopper and Mormon Cricket Suppression Program	43
	Appendix B: Hazard Map of the Affected Environment.....	47
	Appendix C: Map of Proposed Action Areas Covered by the Two Environmental Assessments for the State of Arizona. (EA# AZ-21-01 and AZ-21-02).....	48
	Appendix D: Map of the San Carlos Apache Tribal Proposed Action Area.	479
	Appendix E FWS/NMFS Correspondence	50
	Appendix F: Public Comments and APHIS Response	51

Acronyms and Abbreviations

ac	acre
a.i.	active ingredient
AChE	acetylcholinesterase
APHIS	Animal and Plant Health Inspection Service
BCF	bioconcentration factor
BLM	Bureau of Land Management
CEQ	Council of Environmental Quality
CFR	Code of Federal Regulations
EA	environmental assessment
e.g.	example given (Latin, <i>exempli gratia</i> , “for the sake of example”)
EIS	environmental impact statement
E.O.	Executive Order
FONSI	finding of no significant impact
FR	Federal Register
FS	Forest Service
g	gram
ha	hectare
HHERA	human health and ecological risk assessments
i.e.	in explanation (Latin, <i>id est</i> “in other words.”)
IPM	integrated pest management
lb	pound
MBTA	Migratory Bird Treaty Act
MOU	memorandum of understanding
NEPA	National Environmental Policy Act
NHPA	National Historic Preservation Act
NIH	National Institute of Health
ppm	parts per million
PPE	personal protective equipment
PPQ	Plant Protection and Quarantine
RAATs	reduced agent area treatments
S&T	Science and Technology
ULV	ultra-low volume
U.S.C.	United States Code
USDA	United States Department of Agriculture
USEPA	United States Environmental Protection Agency
USFWS	United States Fish and Wildlife Services

[DRAFT] Site-Specific Environmental Assessment

Rangeland Grasshopper and Mormon Cricket Suppression Program

ARIZONA

EA Number: AZ-21-01

I. Need for Proposed Action

A. Purpose and Need Statement

An infestation of grasshoppers or Mormon crickets may occur on rangeland in Graham and Gila County, San Carlos Apache Reservation. The Animal and Plant Health Inspection Service (APHIS) and may, upon request by land managers or State departments of agriculture, conduct treatments to suppress grasshopper infestations as part of the Rangeland Grasshopper and Mormon Cricket Suppression Program (program). The term “grasshopper” used in this environmental assessment (EA) refers to both grasshoppers and Mormon crickets, unless differentiation is necessary.

Populations of grasshoppers that trigger the need for a suppression program are normally considered on a case-by-case basis. Participation is based on potential damage such as; grasshoppers which defoliate grasses by direct feeding on leaf and stem tissue and by cutting off leaves or stems and heads while feeding. High populations of grasshoppers on rangeland can damage plant crowns so severely that many grass plants will not recover. Some grasshopper species not only reduces grass forage by consuming it but also by cutting it down. The cut grass may become litter on the ground where it may also be used for food by grasshoppers or becomes wasted biomass. Potential areas where large populations may occur can be found in the 2021 Grasshopper Hazard Map in appendix B. The benefits of treatments include the suppressing of over abundant grasshopper populations to lower adverse impacts to range plants and adjacent crops. Treatment would also decrease the economic impact to local agricultural operations and permit normal range plant utilization by wildlife and livestock.

The goal of the proposed suppression program analyzed in this EA is to reduce grasshopper populations below economical infestation levels in order to protect rangeland ecosystems or cropland adjacent to rangeland.

This EA analyzes potential effects of the proposed action and its alternatives. This EA applies to a proposed suppression program that would take place from 04/01/21 to 09/30/21 on rangeland in Graham and Gila County, San Carlos Apache Reservation (Appendix D).

This EA is prepared in accordance with the requirements under the National Environmental Policy Act of 1969 (NEPA) (42 United States Code § 4321 *et. seq.*) and the NEPA procedural requirements promulgated by the Council on Environmental Quality, United States Department of Agriculture (USDA), and APHIS. A decision will be made by APHIS based on the analysis presented in this EA, the results of public involvement, and consultation with other agencies and individuals. A selection of one of the program alternatives will be made by APHIS for the 2021 Control Program for infested rangeland in Graham and Gila County, San Carlos Apache Reservation.

B. Background Discussion

Rangelands provide many goods and services, including food, fiber, recreational opportunities, and grazing land for cattle (Havstad et al., 2007; Follett and Reed, 2010). Grasshoppers and Mormon crickets are part of rangeland ecosystems, serving as food for wildlife and playing an important role in nutrient cycling. However, grasshoppers and Mormon crickets have the potential to occur at high population levels (Belovsky et al., 1996) that result in competition with livestock and other herbivores for rangeland forage and can result in damage to rangeland plant species.

In rangeland ecosystem areas of the United States, grasshopper populations can build up to economic infestation levels¹ despite even the best land management and other efforts to prevent outbreaks. At such a time, a rapid and effective response may be requested and needed to reduce the destruction of rangeland vegetation. In some cases, a response is needed to prevent grasshopper migration to cropland adjacent to rangeland. In most circumstances, APHIS is not able to accurately predict specific treatment areas and treatment strategies months or even weeks before grasshopper populations reach economic infestation levels. The need for rapid and effective response when an outbreak occurs limits the options available to APHIS to inform the public other than those stakeholders who could be directly affected by the actual application. The emergency response aspect is why site-specific treatment details cannot be known, analyzed, and published in advance.

The site-specific data used to make treatment decisions in real time is gathered during spring nymphal surveys. The general site-specific data include: grasshopper densities, species complex, dominant species, dominant life stage, grazing allotment terrain, soil types, range conditions, local weather patterns (wind, temp., precipitation), slope and aspect for hatching beds, animal unit months (AUM's) present in grazing allotment, forage damage estimates, number of potential AUM's consumed by grasshopper population, potential AUM's managed for allotment and value of the AUM, estimated cost of replacement feed for livestock, rotational time frame for grazing allotments, number of livestock in grazing allotment. These are all factors that are considered when determining the economic infestation level.

APHIS surveys grasshopper populations on rangeland in the Western United States, provides technical assistance on grasshopper management to land owners and managers, and may cooperatively suppress grasshoppers when direct intervention is requested by a Federal land management agency or a State agriculture department (on behalf of a State or local government, or a private group or individual). APHIS' enabling legislation provides, in relevant part, that 'on request of the administering agency or the agriculture department

¹ The "economic infestation level" is a measurement of the economic losses caused by a particular population level of grasshoppers to the infested rangeland. This value is determined on a case-by-case basis with knowledge of many factors including, but not limited to, the following: economic use of available forage or crops; grasshopper species, age, and density present; rangeland productivity and composition; accessibility and cost of alternative forage; and weather patterns. In decision making, the level of economic infestation is balanced against the cost of treating to determine an "economic threshold" below which there would not be an overall benefit for the treatment. Short-term economic benefits accrue during the years of treatments, but additional long-term benefit may accrue and be considered in deciding the total value gained by treatment. Additional losses to rangeland habitat and cultural and personal values (e.g., aesthetics and cultural resources), although a part of decision making, are not part of the economic values in determining the necessity of treatment.

of an affected State, the Secretary, to protect rangeland, shall immediately treat Federal, State, or private lands that are infested with grasshoppers or Mormon crickets' ... (7 U.S.C. § 7717(c)(1)). The need for rapid and effective response when an outbreak occurs limits the options available to APHIS. The application of an insecticide within all or part of the outbreak area is the response available to APHIS to rapidly suppress or reduce grasshopper populations and effectively protect rangeland.

In June 2002, APHIS completed an environmental impact statement (EIS) document concerning suppression of grasshopper populations in 17 Western States (Rangeland Grasshopper and Mormon Cricket Suppression Program, Environmental Impact Statement, June 21, 2002). The EIS described the actions available to APHIS to reduce the damage caused by grasshopper populations in Arizona, California, Colorado, Idaho, Kansas, Montana, Nebraska, Nevada, New Mexico, North Dakota, Oklahoma, Oregon, South Dakota, Texas, Utah, Washington, and Wyoming. During November 2019, APHIS published an updated EIS to incorporate the available data and analyze the environmental risk of new program tools. The risk analysis in the 2019 EIS is incorporated by reference.

APHIS has authority under the Plant Protection Act of 2000 (PPA) (7 United States Code (U.S.C.) § 7701) to take actions to control and minimize the economic, ecological, and human health impacts that harmful plant pests can cause. APHIS uses this authority to protect U.S. agriculture, forests, and other natural resources from harmful pest species.

Section 417 of the PPA (7 U.S.C. § 7717) authorizes APHIS' efforts to minimize the economic impacts of grasshoppers. Section 417(a) states that subject to the availability of funds, the Secretary "shall carry out a program to control grasshoppers and Mormon crickets on all Federal lands to protect rangeland." Section 417(c) (1) states that "Subject to the availability of funds pursuant to this section, on request of the administering agency or the agriculture department of an affected State, the Secretary, to protect rangeland, shall immediately treat Federal, State, or private lands that are infested with grasshoppers or Mormon crickets at levels of economic infestation, unless the Secretary determines that delaying treatment will not cause greater economic damage to adjacent owners of rangeland." Section 417(c)(2) states, "In carrying out this section, the Secretary shall work in conjunction with other Federal, State, and private prevention, control, or suppression efforts to protect rangeland."

APHIS has the authority to implement Section 417 of the PPA through the Rangeland Grasshopper and Mormon Cricket Suppression Program. The priorities of the APHIS program are:

- to conduct surveys for grasshopper and Mormon cricket populations on rangelands in the western United States,
- to provide technical assistance on grasshopper management to landowners/managers, and
- subject to the availability of funds, to suppress grasshoppers and Mormon crickets on rangeland when direct intervention is requested by the landowner/manager.

Additional information regarding technical assistance and other aspects of the program can be obtained from the USDA Agricultural Research Service site at <http://www.sidney.ars.usda.gov/grasshopper/index.htm>.

On September 16, 2016, APHIS and the Bureau of Indian Affairs (BIA) signed a Memorandum of Understanding (MOU) detailing cooperative efforts between the two groups on suppression of grasshoppers on BIA managed lands. This MOU clarifies that APHIS will prepare and issue to the public, site-specific environmental documents that evaluate potential impacts associated with the proposed measures to suppress economically damaging grasshopper populations. The MOU also states that these documents will be prepared under the APHIS NEPA implementing procedures with cooperation and input from the BIA.

The MOU further states that the responsible BIA official will request in writing the inclusion of appropriate lands in the APHIS suppression project when treatment on BIA land is necessary. The BIA must also approve a Pesticide Use Proposal for APHIS to treat infestations. According to the provisions of the MOU, APHIS can begin treatments after APHIS issues an appropriate decision document and BIA approves the Pesticide Use Proposal.

On November 6, 2019, APHIS and the Forest Service (FS) signed a Memorandum of Understanding (MOU) detailing cooperative efforts between the two groups on suppression of grasshoppers on FS managed lands (Document #19-8100-0573-MU, November 6, 2019). This MOU clarifies that APHIS will prepare and issue to the public, site-specific environmental documents that evaluate potential impacts associated with the proposed measures to suppress economically damaging grasshopper populations. The MOU also states that these documents will be prepared under the APHIS NEPA implementing procedures with cooperation and input from the FS.

The MOU further states that the responsible FS official will request in writing the inclusion of appropriate lands in the APHIS suppression project when treatment on FS land is necessary. The FS must also approve a Pesticide Use Proposal for APHIS to treat infestations. According to the provisions of the MOU, APHIS can begin treatments after APHIS issues an appropriate decision document and FS approves the Pesticide Use Proposal.

On October 15, 2015, APHIS and the Bureau of Land Management (BLM) signed a Memorandum of Understanding (MOU) detailing cooperative efforts between the two groups on suppression of grasshoppers on BLM managed lands (Document #15-8100-0870-MU, October 15, 2015). This MOU clarifies that APHIS will prepare and issue to the public, site-specific environmental documents that evaluate potential impacts associated with the proposed measures to suppress economically damaging grasshopper populations. The MOU also states that these documents will be prepared under the APHIS NEPA implementing procedures with cooperation and input from the BLM. The MOU further states that the responsible BLM official will request in writing the inclusion of appropriate lands in the APHIS suppression project when treatment on BLM land is necessary. The BLM must also approve a Pesticide Use Proposal for APHIS to treat infestations.

According to the provisions of the MOU, APHIS can begin treatments after APHIS issues an appropriate decision document and BLM approves the Pesticide Use Proposal.

APHIS supports the use of Integrated Pest Management (IPM) principles in the management of grasshoppers and Mormon Crickets. APHIS provides technical assistance to Federal, Tribal, State and private land managers including the use of IPM. However, implementation of on-the-ground IPM activities is limited to land management agencies and Tribes, as well as private landowners. In addition, APHIS' authority under the Plant Protection Act is to treat Federal, State and private lands for grasshoppers and Mormon cricket populations. APHIS' technical assistance occurs under each of the three alternatives proposed in the EIS.

In addition to providing technical assistance, APHIS completed the Grasshopper Integrated Pest Management (GIPM) project. One of the goals of the GIPM is to develop new methods of suppressing grasshopper and Mormon cricket populations that will reduce non-target effects. RAATs are one of the methods that has been developed to reduce the amount of pesticide used in suppression activities and is a component of IPM. APHIS continues to evaluate new suppression tools and methods for grasshopper and Mormon cricket populations, including biological control, and as stated in the EIS, will implement those methods once proven effective and approved for use in the United States.

C. About This Process

The NEPA process for grasshopper management is complicated by the fact that there is very little time between requests for treatment and the need for APHIS to act swiftly with respect to those requests. Surveys help to determine general areas, among the millions of acres where harmful grasshopper infestations may occur in the spring of the following year. Survey data provides the best estimate of future grasshopper populations, while short-term climate or environmental factors change where the specific treatments will be needed. Therefore, examining specific treatment areas for environmental risk analysis under NEPA is typically not possible. At the same time, the program strives to alert the public in a timely manner to its more concrete treatment plans and avoid or minimize harm to the environment in implementing those plans.

Intergovernmental agreements between APHIS and cooperators with Tribal Nations may preclude disclosure of Tribal site-specific information to the public without the consent of the Tribal Administrator. Individuals may request information on the specific treatment areas on Tribal Lands from the individual Tribal Nations.

Public involvement under the CEQ Regulations for Implementing the Procedural Provisions of NEPA distinguishes federal actions with effects of national concern from those with effects primarily of local concern (40 CFR 1506.6). The grasshopper and Mormon cricket suppression program EIS was published in the Federal Register (APHIS-2016-0045), and met all applicable notice and comment requirements for a federal action with effects of national concern. This process provided individuals and national groups the ability to participate in the development of alternatives and provide comment. Our subsequent state-

based actions have the potential for effects of local concern, and we publish them according to the provisions that apply to federal actions with effects primarily of local concern. This includes the USDA APHIS NEPA Implementation Procedures, which allows for EAs and findings of no significant impact (FONSIs) where the effects of an action are primarily of regional or local concern, to normally provide notice of publication in a local or area newspaper of general circulation (7 CFR 372.7(b)(3)). These notices provide potentially locally affected individuals an additional opportunity to provide input into the decision-making process. Some states also provide additional opportunities for local public involvement, such as public meetings. In addition, when an interested party asks to be informed APHIS ensures their contact information is added to the list of interested stakeholders.

APHIS uses the scoping process to enlist land managers and the public to identify alternatives and issues to be considered during the development of a grasshopper or Mormon cricket suppression program. Scoping was helpful in the preparation of the draft EAs. The process can occur formally and informally through meetings, conversations, or written comments from individuals and groups.

The current EIS provides a solid analytical foundation; however, it may not be enough to satisfy NEPA completely for actual treatment proposals. The program typically prepares a Draft EA tiered to the current EIS for each of the 17 Western States, or portion of a state, that may receive a request for treatment. The Draft EA analyzes aspects of environmental quality that could be affected by treatments in the area where grasshopper outbreaks are anticipated. The Draft EA will be made available to the public for a 30-day comment period. **The comment period will begin February 1st and end March 2nd, 2021.** Comments can be sent to USDA, APHIS, 3640 East Wier Ave. Phoenix, Arizona 85040, or contacting local USDA, APHIS Arizona State Office. When the program receives a treatment request and determines that treatment is necessary, the specific site within the state will be evaluated to determine if environmental factors were thoroughly evaluated in the Draft EA. If all environmental issues were accounted for in the Draft EA, the program will prepare a Final EA and FONSI. Once the FONSI has been finalized copies of those documents will be sent to any parties that submitted comments on the Draft EA, and to other appropriate stakeholders. To allow the program to respond to comments in a timely manner, the Final EA and FONSI will be posted to the APHIS website. The program will also publish a notice of availability in the same manner used to advertise the availability of the Draft EA.

II. Alternatives

To engage in comprehensive NEPA risk analysis APHIS must frame potential agency decisions into distinct alternative actions. These program alternatives are then evaluated to determine the significance of environmental effects. The 2002 EIS presented three alternatives: (A) No Action; (B) Insecticide Applications at Conventional Rates and Complete Area Coverage; and (C) Reduced Agent Area Treatments (RAATs), and their potential impacts were described and analyzed in detail. The 2019 EIS was tiered to and updated the 2002 EIS. Therefore the 2019 EIS considered the environmental background or 'No Action' alternative of maintaining the program that was described in the 2002 EIS and Record of Decision. The 2019 EIS also considered an alternative where APHIS would not

fund or participate in grasshopper suppression programs. The preferred alternative of the 2019 EIS allowed APHIS to update the program with new information and technologies that not were analyzed in the 2002 EIS. Copies of the complete 2002 and 2019 EIS documents are available for review at USDA, APHIS, 3640 East Wier Ave. Phoenix, Arizona 85040. These documents are also available at the Rangeland Grasshopper and Mormon Cricket Program web site, <http://www.aphis.usda.gov/plant-health/grasshopper>.

All insecticides used by APHIS for grasshopper suppression are used in accordance with applicable product label instructions and restrictions. Representative product specimen labels can be accessed at the Crop Data Management Systems, Incorporated web site at www.cdms.net/manuf/manuf.asp. Labels for actual products used in suppression programs will vary, depending on supply issues. All insecticide treatments conducted by APHIS will be implemented in accordance with APHIS' treatment guidelines and operational procedures, included as Appendix A to this Draft EA.

This Draft EA analyzes the significance of environmental effects that could result from the alternatives described below. These alternatives differ from those described in the 2019 EIS because grasshopper treatments are not likely to occur in most of the rangeland in Arizona and therefore the environmental baseline should describe a no treatment scenario in those rangeland areas.

A. No Suppression Program Alternative

Under Alternative A, the No Action alternative, APHIS would not conduct a program to suppress grasshopper infestations within Arizona. Under this alternative, APHIS may opt to provide limited technical assistance, but any suppression program would be implemented by a Federal land management agency, a State agriculture department, a local government, or a private group or individual.

B. Insecticide Applications at Conventional Rates or Reduced Agent Area Treatments with Adaptive Management Strategy (Preferred Alternative)

Under Alternative B, the Preferred Alternative, APHIS would manage a grasshopper treatment program using techniques and tools discussed hereafter to suppress outbreaks. The insecticides available for use by APHIS include the U.S. Environmental Protection Agency (USEPA) registered chemicals carbaryl and diflubenzuron. These chemicals have varied modes of action. Carbaryl work by inhibiting acetylcholinesterase (enzymes involved in nerve impulses) and diflubenzuron inhibits the formation of chitin by insects. APHIS would make a single application per year to a treatment area and could apply insecticide at an APHIS rate conventionally used for grasshopper suppression treatments, or more typically as reduced agent area treatments (RAATs). APHIS selects which insecticides and rates are appropriate for suppression of a grasshopper outbreak based on several biological, logistical, environmental, and economical criteria. The identification of grasshopper species and their life stage largely determines the choice of insecticides used among those available to the program. RAATs are the most common application method for all program insecticides, and only rarely do rangeland pest conditions warrant full coverage and higher rates.

Typically, the decision to use diflubenzuron, the pesticide most commonly used by the program, is determined by the life stage of the dominant species within the outbreak population. Diflubenzuron can produce 90 to 97% grasshopper mortality in nascent populations with a greater percentage of early instars. If the window for the use of diflubenzuron closes, because of treatment delays, then carbaryl or rarely malathion are the remaining control options. Certain species are more susceptible to carbaryl bait, and sometimes that pesticide is the best control option.

The RAATs strategy is effective for grasshopper suppression because the insecticide controls grasshoppers within treated swaths while conserving grasshopper predators and parasites in swaths not directly treated. RAATs can decrease the rate of insecticide applied by either using lower insecticide concentrations or decreasing the deposition of insecticide applied by alternating one or more treatment swaths. Both options are most often incorporated simultaneously into RAATs. Either carbaryl and diflubenzuron would be considered under this alternative, typically at the following application rates:

- 8.0 fluid ounces (0.25 lb a.i.) of carbaryl ULV spray per acre.
- 10.0 pounds (0.20 lb a.i.) of 2 percent carbaryl bait per acre.
- 0.75 or 1.0 fluid ounce (0.012 lb a.i.) of diflubenzuron per acre; or

The width of the area not directly treated (the untreated swath) under the RAATs approach is not standardized. The proportion of land treated in a RAATs approach is a complex function of the rate of grasshopper movement, which is a function of developmental stage, population density, and weather (Narisu et al., 1999, 2000), as well as the properties of the insecticide (insecticides with longer residuals allow wider spacing between treated swaths). Foster et al. (2000) left 20 to 50% of their study plots untreated, while Lockwood et al. (2000) left 20 to 67% of their treatment areas untreated. Currently the grasshopper program typically leaves 50% of a spray block untreated for ground applications where the swath width is between 20 and 45 feet. For aerial applications, the skipped swath width is typically no more than 100 feet for carbaryl and 200 feet for diflubenzuron. The selection of insecticide and the use of an associated swath widths is site dependent. Rather than suppress grasshopper populations to the greatest extent possible, the goal of this method is to suppress grasshopper populations to less than the economic infestation level.

Applicators use of Trimble GPS Navigation equipment is used to navigate and capture shapefiles of the treatment areas. All sensitive sites are buffered out of the treatment area using flagging which is highly visible to the aerial applicator. All sensitive sites are reviewed in the daily briefing with APHIS personnel including the applicator working on the treatment site. Treatments are conducted to suppress large grasshopper populations to protect rangeland vegetation. Treatments are conducted using the Reduced Agent Area Treatment (RAAT's) method. This method of skipping swaths (fig.1) decreases the amount



Figure 1. Reduced Agent Area Treatment (RAAT's)

of chemical and acreage treated still maintaining an effective kill rate. Swath widths usually range from 40-45 feet depending on ground equipment used. In Arizona, only ground equipment is used, no aerial treatments are conducted. Grasshoppers in untreated areas will tend to move to treated areas, thus becoming exposed to the insecticide. For example, if the area in figure 1 was 100 acres, with 50% RAAT's the acreage actually treated would be 50 acres. Protection would include the entire 100 acres, only exposing half the area with half the chemical amount compared to a conventional blanket treatment covering the entire 100 acres and the label rate of application.

Insecticide applications at conventional rates and complete area coverage, is an approach that APHIS has used in the past but is currently uncommon. Under this alternative, carbaryl and diflubenzuron would cover all treatable sites within the designated treatment block per label directions. The application rates under this alternative are typically at the following application rates:

- 16.0 fluid ounces (0.50 lb a.i.) of carbaryl spray per acre.
- 10.0 pounds (0.50 lb a.i.) of 5 percent carbaryl bait per acre.
- 1.0 fluid ounce (0.016 lb a.i.) of diflubenzuron per acre; or

The potential generalized environmental effects of the application of carbaryl and diflubenzuron under this alternative are discussed in detail in the 2019 EIS. A description of anticipated site-specific impacts from this alternative may be found in Part IV of this document.

C. Experimental Treatments Alternative

APHIS-PPQ continues to refine its methods of grasshopper and Mormon cricket management in order to improve the abilities of the Rangeland Grasshopper and Mormon Cricket Suppression Program (herein referred to as the Program) to make it more economically feasible, and environmentally acceptable. These refinements can include reduced rates of currently used pesticides, improved formulations, development of more target-specific baits, development of biological pesticide suppression alternatives, and improvements to aerial (e.g., incorporating the use of Unmanned Aircraft Systems (UAS)) and ground application equipment. A division of APHIS-PPQ, Science and Technology's (S&T) Phoenix Lab is in Arizona and its Rangeland Grasshopper and Mormon Cricket Management Team (Rangeland Unit) conducts methods development and evaluations on behalf of the Program. The Rangeland Unit's primary mission is to comply with Section 7717 of the Plant Protection Act and protect the health of rangelands (wildlife habitats and where domestic livestock graze) against economically damaging cyclical outbreaks of grasshoppers and Mormon crickets. The Rangeland Unit tests and develops more effective, economical, and less environmentally harmful management methods for the Program and its federal, state, tribal, and private stakeholders.

To achieve this mission, experimental plots ranging in area from less than one foot to 640 acres are used and often replicated. The primary purpose of these experiments is to test and

develop improved methods of management for grasshoppers and Mormon crickets. This often includes testing and refining pesticide and biopesticide formulations that may be incorporated into the Program. These investigations often occur in the summer (May-August) and the locations typically vary annually. The plots often include “no treatment” (or control) areas that are monitored to compare with treated areas. Some of these plots may be monitored for additional years to gather information on the effects of utilized pesticides on non-target arthropods. Note that an [Experimental Use Permit](#) is not needed when testing non-labeled experimental pesticides if the use is limited to laboratory or greenhouse tests, or limited replicated field Trials involving 10 acres or less per pest for terrestrial tests.

Studies and experimental plots are typically located on large acreages of rangelands and the Rangeland Unit often works on private land with the permission of landowners. Locations of experimental trials will be made available to the appropriate agencies in order to ensure these activities are not conducted near sensitive species or habitats. Due to the small size of the experimental plots, no adverse effects to the environment, including protected species and their critical habitats, are expected, and great care is taken to avoid sensitive areas of concern prior to initiating studies.

Methods Development Studies

Methods development studies may use planes and all-terrain vehicles (ATVs) to apply labeled pesticides using conventional applications and the Reduced Agent Area Treatments (RAATs) methodology. The experiments may include the use of an ultra-low volume sprayer system for applying biopesticides (such as native fungal pathogens). Mixtures of native pathogens and low doses of pesticides may be conducted to determine if these multiple stressor combinations enhance mortality. Aircraft will be operated by Federal Aviation Administration-licensed pilots with an aerial pesticide applicator’s permit.

Rangeland Unit often uses one square foot micro plots covered by various types of cages depending on the study type and species used. These types of study plots are preferred for Mormon cricket treatments and those involving non-labeled experimental pesticides or biopesticides. Our most common application method for micro plots is simulating aerial applications via the Field Aerial Application Spray Simulation Tower Technique (FAASSTT). This system consists of a large tube enclosed on all sides except for the bottom, so micro plot treatments can be accurately applied to only the intended treatment target. Treatments are applied with the FAASSTT in micro doses via a syringe and airbrush apparatus mounted in the top.

Rangeland Unit is also investigating the potential use of Unmanned Aerial Systems (UAS) for a number of purposes related to grasshopper and Mormon cricket detection and treatment. UAS will be operated by FAA-licensed pilots with an aerial pesticide applicator’s permit.

Pesticides and Biopesticides Used in Studies

Pesticides likely to be involved in studies currently include:

- 1) Liquids: diflubenzuron (Dimilin 2L and generics: currently Unforgiven and Cavalier 2L) and chlorantraniliprole (Prevathon). Program standard application rates are diflubenzuron -

1.0 fl. oz./acre in a total volume of 31 fl. oz./acre; chlorantraniliprole - 2.0 fl. oz./acre (RAATs) or 4.0 fl. oz./acre (conventional coverage), both in a total volume of 32 fl. oz./acre. Experimental rates often vary, but the doses are lower than standard Program rates unless otherwise noted.

2) Baits: carbaryl. Program standard application rates: 2% bait at 10 lbs. /acre (2 lbs. AI/acre) or 5% bait at 4 lbs. /acre (2 lbs. AI/acre).

3) LinOilEx (Formulation 103), a proprietary combination of easily available natural oils and some commonly encountered household products, created by Manfred Hartbauer, University of Graz, Austria. Note that LinOilEx (Formulation 103) is experimental; for more information, see “Potential Impacts of LinOilEx Applications” in the section “Information on Experimental Treatments.”

Biopesticides likely to be involved in studies currently include:

1) *Metarhizium robertsii* (isolate DWR2009), a native fungal pathogen. Note that *Metarhizium robertsii* (isolate DWR2009) is experimental; for more information, see “Potential Impacts of *Metarhizium robertsii* Applications” in the section “Information on Experimental Treatments.”

2) *Beauveria bassiana* GHA, a native fungal pathogen sold commercially and registered for use across the U.S.

At this time, we are unsure where in the 17 states we will be doing most of the following proposed experimental field studies. The final location decision is dependent upon grasshopper and/or Mormon cricket population densities, and availability of suitable sites, but we plan to most likely work in Arizona, Idaho, New Mexico, Oregon, Montana, or Washington.

Study 1: Evaluate efficacy of a UAS-mounted bait spreader applying 2% carbaryl bait at 5 lbs/acre. This study plans to use replicated 40-acre plots (320 acres total) on Colville Confederated Tribes land in Washington sometime in May/June, but is contingent upon a population of sufficient size. Mortality will be then be observed for a duration of time to determine efficacy.

Study 2: Evaluate persistence of the experimental biopesticide DWR2009 in bait form by coating wheat bran with the pathogen. A species of local abundance will be placed into replicated microplot cages and fed the baits by hand. Mortality and sporulation will be then be observed for a duration of time to determine persistence in both the field and lab.

Study 3: Evaluate efficacy of the experimental biopesticide DWR2009 in bait form by coating wheat bran with the pathogen. A species of local abundance will be placed into replicated microplot cages and fed the baits by hand. Mortality and sporulation will be then be observed for a duration of time to determine efficacy in both the field and lab.

Study 4: A stressor study to evaluate efficacy of the experimental biopesticide DWR2009 in liquid form when combined with Dimilin 2L. The FAASSTT will be utilized to apply varying dose levels of Dimilin 2L (below label rates) in order to compare efficacy, starting

at the rate of 1.0 fl. oz. /acre. Replicated microplots will be treated and then a species of local abundance will be placed into each cage. Mortality will be then be observed for a duration of time to determine efficacy.

Study 5: Evaluate efficacy of the experimental biopesticide DWR2009 in liquid and bait form (by coating wheat bran with the pathogen) using ultra-ultra low volume RAATs (involves a timing device and ULV nozzles) and a 10-acre plot. ATV-mounted liquid and bait spreaders will be utilized to apply DWR2009. Specimens will be periodically collected to observe mortality and sporulation for a duration of time to determine efficacy.

Study 6: Evaluate efficacy of the experimental, non-traditional pesticide LinOilEx (Formulation 103). A micro-FAASSTT (airbrush system mounted on a 5-gal bucket) will be utilized to apply varying dose levels in order to compare efficacy, starting at the base rate of 6.64 ml/cage. A species of local abundance will be placed into replicated microplot cages and sprayed directly. Mortality will be then be observed for a duration of time to determine efficacy.

III. Affected Environment

A. *Description of Affected Environment*

The Site-Specific Graham and Gila County portion within the San Carlos Apache Reservation proposed suppression program area in the EA encompasses 332,120 acres. This is the total estimated acres within the proposed action area (Appendix D map). Acres treated will be from somewhere within this total. Actual acres treated will be far less than this amount. For example, 2020 season only 2,875 actual acres were treated from within this proposed action area. The vegetative communities are semiarid grasslands; Plains & Great Basin Grasslands; Great Basin Conifer woodland; Interior Chaparral covered in this area. Soil types include basalt and basalt flows, weakly consolidated sandstone and siltstone, unconsolidated alluvial sand, silt, and some gravel. All rangeland covered in this EA is managed by the San Carlos Apache Tribe and BIA.

Elevations range from approximately 3,500 to over 6,000 feet. Potential treatment sites are within watersheds which drain into tributaries of the Bonita Creek, Hackberry Creek, Hackberry Draw, Cottonwood Canyon Salt Creek, and San Carlos River. There are stock tanks in the potential treatment area. All potential treatment areas fall within the Arizona Interior Chaparral biome (Brown, 1994), grassland representative species of this biome include:

Plants: Emory oak (*Quercus emoryi*), alligator bark juniper (*Juniperus deppeana*), pinyon pine (*Pinus edulis*), gray oak (*Quercus grisea*), canyon live oak (*Quercus chrysolepis*), Arizona oak (*Quercus arizonica*), western chokecherry (*Prunus virginiana*), shrub live-oak (*Quercus turbinella*), ceanothus (*Ceanothus greggii*), crucifixion thorn (*Canotia holocantha*), penstemon (*Penstemon spp.*), desert verbena (*Verbena wrightii*), Wright buckwheat (*Eriogonum wrightii*), narrowleaf yerbasanta (*Eriodictyon angustifolium*), sideoats grama (*Bouteloua curtipendula*), cane bluestem (*Bothriochloa barbinodis*), plains lovegrass (*Eragrostis intermedia*), Black grama (*Bouteloua eriopoda*), Blue grama, (*Bouteloua gracilis*) Hairy grama, (*Bouteloua hirsuta*) Rothrock's grama, (*Bouteloua*

rothrockii), Fendler three-awn (*Aristida spp.*), agave (*Agave parryi*), beargrass (*Nolina microcarpa*), sotol (*Dasyilirion wheeleri*), banana yucca (*Yucca baccata*), , squirreltail, (*Elymus elymoides*), Arizona cottontop, (*Digitaria californica*), Green sprangletop (*Leptochloa dubia*), Junegrass, (*Koeleria spp.*), Western wheatgrass (*Pascopyrum smithii*), Tobosagrass, (*Pleuraphis mutica*), Vine Mesquite, (*Panicum obtusum*), curly-mesquite (*Hilaria belangeri*), Cholla (*Opuntia spp.*), Prickly Pear (*Opuntia spp.*),

Mammals: cliff chipmunk (*Eutamias dorsalis*), white-throated woodrat (*Neotoma albigula*), mule deer (*Odocoileus hemionus*), brush mouse (*Peromyscus boylei*), rock mouse (*P. difficilis*), white- footed mouse (*P. leucopus*), eastern cottontail (*Sylvilagus floridanus holzeri*), pronghorn antelope (*Antilocapra americana*), elk (*Cervus elaphus*) javalina (*Pecari tajacu*), jackrabbit (*Lepus spp.*), coyote (*Canis latran*), White-tailed deer (*Odocoileus virginianus*).

Birds: rufous-crowned sparrow (*Aimophila ruficeps*), scrub jay (*Aphelocoma coerulescens*), canyon wren (*Catherpes mexicanus*), rufous-sided towhee (*Pipilo erythrophthalmus*), brown towhee (*P. fuscus*), bushtit (*Psaltiriparus minimus*), black-chinned sparrow (*Spizella atrogularis*), crissal thrasher (*Toxostoma dorsale*), burrowing owl (*Athene cunicularia*).

Amphibians and reptiles: glossy snake (*Arizona elegans*), Arizona alligator lizard (*Gerrhonotus kingi*), night snake (*Hypsiglena torquata*), Sonoran mountain kingsnake (*Lampropeltis pyromelana*), southwestern blind snake (*Leptotyphlops humilis*), Sonora whipsnake (*Masticophis bilineatus*), desert striped whipsnake (*M. taeniatus*), western fence lizard (*Sceloporus occidentalis*), eastern fence lizard (*S. undulates*), western blackhead snake (*Tantilla planiceps*), Sonoran lyre snake (*Trimorphodon biscutatus lambda*), Texas lyre snake (*T. b. vilkinsoni*), side-blotched lizard (*Uta stansburiana*), Arizona night lizard (*Zantusia arizonae*), Western Diamond-backed Rattlesnake (*Crotalus atrox*), Black-tailed Rattlesnake (*Crotalus molossus*), Arizona Black Rattlesnake (*Crotalus cerberus*)

B. Site-Specific Considerations

1. Human Health

The 2019 EIS contains detailed hazard, exposure, and risk analyses for the chemicals available to APHIS. APHIS has incorporated by reference the analysis from the EIS and the associated risk assessments of pesticides which are mentioned this EA. These documents are titled, The Final Human Health and Ecological Risk Assessments (USDA, APHIS 2018a, 2018b, 2018c, 2018d) for program pesticides which are available at the following website, <http://www.aphis.usda.gov/plant-health/grasshopper>.

Impacts to workers and the general public were analyzed for all possible routes of exposure (dermal, oral, inhalation) under a range of conditions designed to overestimate risk. The operational procedures and spraying conditions examined in those analyses conform to those expected for operations. The following discussion summarizes the hazards, potential

exposure, and risk to workers and the general public for operations within these potential proposed treatment areas detailed in this EA. The operational procedures identified in Appendix 1 would be required in all cases and further mitigation measures are identified in this section, as appropriate.

The suppression program would be conducted on federally managed rangelands. No treatments will occur over congested or residential areas, recreation areas, and schools. The nearest residential or populated area to potential treatment areas are at least 6 miles away. Refer to the Operational Procedures, Specific Procedures for Aerial and Ground Applications in Appendix A for further information.

Groundwater wells are a major source of domestic water supplies. Groundwater and surface water are the major rural and livestock water sources. No impact is anticipated. Strict adherence to label requirements and the USDA treatment guidelines (appendix A) will be followed in regard to treatments bordering open surface waters.

2. Nontarget Species

Threatened & Endangered Species and Sensitive Species of Concern

The area assessed by this EA includes a variety of organisms i.e., terrestrial vertebrates and invertebrates, migratory birds, biocontrol agents, pollinators, aquatic organisms, plants (both native and introduced), etc. APHIS will employ measures, such as buffer zones, to protect these species and their habitat. APHIS will also consult with local agency officials to determine appropriate protective measures.

Federally Listed Threatened and Endangered Species Covered by this EA:

MAMMALS

Endangered Mexican gray wolf, *Canis lupus baileyi* - Endangered

BIRDS

Mexican spotted owl, *Strix occidentalis lucida* - Threatened

Southwestern willow flycatcher, *Empidonax traillii extimus* - Endangered

Western yellow-billed cuckoo, *Coccyzus americanus* – Threatened

FISH

Desert pupfish, *Cyprinodon macularius* - Endangered

Gila chub, *Gila intermedia* - Endangered

Razorback sucker, *Xyrauchen texanus* – Endangered

AMPHIBIANS

Chiricahua leopard frog, *Rana chiricahuensis* - Threatened

PLANTS

Arizona cliffrose, *Purshia subintegra* - Endangered

REPTILES

Northern Mexican gartersnake, *Thamnophis eques megalops* - Threatened

Sensitive Species of Concern:

Sonoran Desert tortoise, *Gopherus morafkai* – Candidate

Headwater chub, *Gila nigra* - Candidate

Roundtail chub, *Gila robusta* – Candidate

Bald and Golden Eagle Protection Act (BGEPA)

The Eagle Act (16 U.S.C. 668-668c), enacted in 1940, and amended several times since then, prohibits anyone, without a permit issued by the Secretary of the Interior, from “taking” bald eagles, including their parts, nests, or eggs. The Act provides criminal and civil penalties for persons who “take, possess, sell, purchase, barter, offer to sell, purchase or barter, transport, export or import, at any time or any manner, any bald eagle ... [or any golden eagle], alive or dead, or any part, nest, or egg thereof.” The Act defines “take” as “pursue, shoot, shoot at, poison, wound, kill, capture, trap, collect, molest or disturb.” “Disturb” means: "Disturb means to agitate or bother a bald or golden eagle to a degree that causes, or is likely to cause, based on the best scientific information available, 1) injury to an eagle, 2) a decrease in its productivity, by substantially interfering with normal breeding, feeding, or sheltering behavior, or 3) nest abandonment, by substantially interfering with normal breeding, feeding, or sheltering behavior." In addition to immediate impacts, this definition also covers impacts that result from human- induced alterations initiated around a previously used nest site during a time when eagles are not present, if, upon the eagles return, such alterations agitate or bother an eagle to a degree that injures an eagle or substantially interferes with normal breeding, feeding, or sheltering habits and causes, or is likely to cause, a loss of productivity or nest abandonment.

As listed in the National Bald Eagle Management Guidelines (USFWS, May 2007) and adapting recommendations from (Driscoll et al. 2006) the following mitigation measures will be followed.

Category G Helicopters and fixed-wing aircraft. Except for authorized biologists trained in survey techniques, avoid operating aircraft within 2,000 feet of the nest during the breeding season, except where eagles have demonstrated tolerance for such activity. In addition, Category A (Agriculture) and Category D (Off Road Vehicle Use) both provide the same

guidance for use of ATV's or trucks: No buffer is necessary around nest sites outside the breeding season. During the breeding season, do not operate off-road vehicles within 1,000 feet of the nest. In open areas, where there is increased visibility and exposure to noise, this distance should be extended to 1,000 feet.

Representative wildlife and plant spp.

See Table 1 for list of representative wildlife, and plant spp.

Under the no action alternative, destruction of grasses and forbs by grasshoppers could cause localized disruption of food and cover for a number of wildlife species. Under chemical control there is a possibility of indirect effects on local wildlife populations, particularly insectivorous birds that depend on a readily available supply of insects, including grasshoppers, for their own food supply and for their young. We have found no valid data which suggests that (absent a spill) any species other than certain mice would be subjected to a dosage in excess of 1/5 of the LD50 for carbaryl (Pg. B-37 GH EIS.) Therefore, it is not apparent that any fatalities would be likely to occur as a result of carbaryl intoxication.

Malathion and carbaryl have been shown to reduce brain cholinesterase (ChE) (an enzyme important in nerve cell transmissions) levels in birds. Effects of ChE inhibition are not fully understood but could cause inability to gather food, escape predation, or care for young.

In any given treatment season, only a fraction (**less than 1 percent**) of the total rangeland in a region is likely to be sprayed for grasshopper control. For species that are wide spread and numerous lowered survival and lowered reproductive success in a small portion of their habitat would not constitute a significant threat to the population.

The wildlife risk assessment in APHIS FEIS 2002 estimated wildlife doses of Malathion and carbaryl to representative rangeland species and compared them with toxicity reference levels. No dose of Malathion will approach or exceed the reference species LD50. Some individual animals may be at risk of fatality or behavioral alterations that make them more susceptible to predation resulting from ChE level changes in Malathion spraying for grasshopper control. However, most individual animals would not be seriously affected.

Carbaryl also poses a low risk to wildlife, with few fatalities likely to occur and a low risk of behavioral anomalies caused by cholinesterase depression. There is some chance of adverse effects on bird reproduction through the use of any of these chemicals or diesel oil through direct toxicity to developing embryos in birds' eggs.

Some species of herbivorous mammals and birds may consume wheat bran bait after it has been applied to grasshopper-infested areas. Carbaryl is moderately toxic to mammals and slightly toxic to birds. We have found no valid data which suggests that (absent a spill) any species other than certain mice would be subjected to a dosage in excess of 1/5 of the LD50 for carbaryl (Pg. B-37 GH EIS.) Therefore, it is not apparent that any fatalities would be likely to occur as a result of carbaryl intoxication. Additionally, we note that carbaryl 5% bait is labeled at 3 lbs. /1000ft² in poultry houses when poultry are present. (<http://www.cdms.net/manuf/>)

Chitin or chitin-like substances are not as important to terrestrial mammals, birds, and other vertebrates as chitin is to insects; therefore, the chitin inhibiting properties of diflubenzuron applications under the conditions of Alternative 2 such as reductions in the food base for insectivorous wildlife species, especially birds. As stated above, diflubenzuron is practically nontoxic to birds, including those birds that ingest moribund grasshoppers resulting from diflubenzuron applications, as described in Alternative 2.

While immature grasshoppers and other immature insects can be reduced up to 98 percent in area covered with diflubenzuron, some grasshoppers and other insects remain in the treatment area. Although the density of grasshoppers and other insects may be low, it is most likely sufficient to sustain birds and other insectivores until insect populations recover. Those rangeland birds that feed primarily on grasshoppers may switch to other diet items. However, in some areas the reduced number of invertebrates necessary for bird survival and development may result in birds having less available food. In these cases, birds will either have less than optimal diets or travel to untreated areas for suitable prey items, causing a greater foraging effort and a possible increased susceptibility to predation. It also should be noted that suppressing grasshopper populations conserves rangeland vegetation that often is important habitat to rangeland wildlife. Habitat loss is frequently the most important factor leading to the decline of a species and reducing grasshopper densities can be an aid in reducing habitat loss.

APHIS is the lead agency in Arizona regarding biological control for invasive weeds. All biocontrol programs are coordinated between APHIS and Federal, Tribal, State agencies and Weed Management Districts and City Municipalities. APHIS has GIS data for all Biological Control programs throughout Arizona. There has been no overlap between biocontrol programs and grasshopper treatments. If this does become the case in the future, the grasshopper program would eliminate questionable acreage from the treatment area.

3. Socioeconomic Issues

Livestock grazing and hunting are the main uses of the potential treatment area. These grasslands provide forage for cattle and wildlife. Farming, forestry occupations, agriculture, fishing and hunting, and mining provide 10.6% of the employment on San Carlos Apache Reservation (U.S. Bureau of the Census, Census 2000).

As of August 2014, the San Carlos Apache tribe had an enrollment of 15,393 tribal members. Currently there is approximately 9,945-10,945 living on the Reservation according to the My Tribal Data. US Census. Retrieved 20 July 2020.

The San Carlos Reservation's annual median household income of approximately \$27,542, according to the US Census. About 49.2 percent of the people live under the poverty line, and 36.7 percent of the active labor force is unemployed. Replacement feed for damage rangeland would be almost impossible to afford under these circumstances. It is critical that APHIS provide the Rangeland Grasshopper Suppression Program to assist the Tribal Ranches management of resources.

The principal economic activities are tourism, cattle ranching, and arts and crafts. The San Carlos is rich in hunting, fishing. The tribe sells guided big-game hunting permits for desert bighorn sheep, trophy elk, antelope, and mountain lion.

The possible treatment areas are subject to reoccurring drought. A combination of drought and grasshopper damage causes economic stress to landowners and permittees. The control of grasshoppers in this area would have beneficial economic impacts to local landowners. The forage not utilized by grasshoppers will be available for livestock consumption and harvesting. This will allow greater livestock grazing, decreased needs for supplemental feed, and increased monetary returns.

4. Cultural Resources and Events

To ensure that historical or cultural sites, monuments, buildings or artifacts of special concern are not adversely affected by program treatments, APHIS will confer with Tribal Officials, BIA, or other appropriate land management agencies on a local level to protect these areas of special concern. APHIS will also confer with the appropriate Tribal Authority and with the BIA office at a local level to ensure that the timing and location of planned program treatments do not coincide or conflict with cultural events or observances, on Tribal and/or allotted lands.

5. Special Considerations for Certain Populations

a) Executive Order No. 12898, Federal Actions to Address Environmental Justice in Minority Populations and Low-Income Populations

Executive Order (E.O.) 12898, Federal Actions to Address Environmental Justice in Minority Populations and Low-Income Populations, was signed by President Clinton on February 11, 1994 (59 *Federal Register* (FR) 7269). This E.O. requires each Federal agency to make achieving environmental justice part of its mission by identifying and addressing, as appropriate, disproportionately high and adverse human health or environmental effects of its programs, policies, and activities on minority populations and low-income populations. Consistent with this E.O., APHIS will consider the potential for disproportionately high and adverse human health or environmental effects on minority populations and low-income populations for any of its actions related to grasshopper suppression programs.

The San Carlos Reservation's annual median household income of approximately \$31,696 according to the US Census and BUREAU OF WOMEN'S AND CHILDREN'S HEALTH, Arizona Department of Health Services. About 49.2 percent of the people live under the poverty line, and 36.7 percent of the active labor force is unemployed.

b) Executive Order No. 13045, Protection of Children from Environmental Health Risks and Safety Risks

The increased scientific knowledge about the environmental health risks and safety risks associated with hazardous substance exposures to children and recognition of these issues in Congress and Federal agencies brought about legislation and other requirements to

protect the health and safety of children. On April 21, 1997, President Clinton signed E.O. 13045, Protection of Children from Environmental Health Risks and Safety Risks (62 FR 19885). This E.O. requires each Federal agency, consistent with its mission, to identify and assess environmental health risks and safety risks that may disproportionately affect children and to ensure that its policies, programs, activities, and standards address those risks. APHIS has developed agency guidance for its programs to follow to ensure the protection of children (USDA, APHIS, 1999).

According to the BUREAU OF WOMEN'S AND CHILDREN'S HEALTH, Arizona Department of Health Services, there are 3,235 children between the ages of 0-14. There is approximately 1,151 youth from the ages of 15-19 according to the Arizona Department of Health Services. The risk for children to be exposed to treatment pesticides is very low due to the remote nature of the Tribal Rangeland. The nearest communities are approximately 50 miles from the Tribal rangeland areas. There will be no aerial treatments conducted in Arizona only by ground based equipment.

IV. Environmental Consequences

Each alternative described in this EA potentially has adverse environmental effects. The general environmental impacts of each alternative are discussed in detail in the 2002 and 2019 EIS. The specific impacts of the alternatives are highly dependent upon the particular action and location of infestation. The principal concerns associated with the alternatives are: (1) the potential effects of insecticides on human health (including subpopulations that might be at increased risk); and (2) impacts of insecticides on nontarget organisms (including threatened and endangered species).

APHIS has written human health and ecological risk assessments (HHERAs) to assess the insecticides and use patterns that are specific to the program. The risk assessments provide an in-depth technical analysis of the potential impacts of each insecticide to human health, and non-target fish and wildlife along with its environmental fate in soil, air, and water. The assessments rely on data required by the USEPA for pesticide product registrations, as well as peer-reviewed and other published literature. The HHERAs are heavily referenced in the EIS and this EA. These Environmental Documents can be found at the following website: <http://www.aphis.usda.gov/plant-health/grasshopper>.

A. Environmental Consequences of the Alternatives

Site-specific environmental consequences of the alternatives are discussed in this section.

1. No Suppression Program Alternative

Under this alternative, APHIS would not conduct a program to suppress grasshoppers. If APHIS does not participate in any grasshopper suppression program, Federal land management agencies, State agriculture departments, local governments, private groups or individuals, may not effectively combat outbreaks in a coordinated effort. Without the technical assistance and coordination that APHIS provides during grasshopper outbreaks, the uncoordinated programs could use insecticides that APHIS considers too environmentally harsh. Multiple treatments and excessive amount of insecticide could be applied in efforts to suppress or even locally eradicate grasshopper populations. There are approximately 100 pesticide products registered by USEPA for use on rangelands and

against grasshoppers (Purdue University, 2018). It is not possible to accurately predict the environmental consequences of the No Action alternative because the type and amount of insecticides that could be used in this scenario are unknown. However, the environmental impacts could be much greater than under the APHIS led suppression program alternative due to lack of treatment knowledge or coordination among the groups.

The potential environmental impacts from the No Action alternative, where other agencies and land managers do not control outbreaks, stem primarily from grasshoppers consuming vast amounts of vegetation in rangelands and surrounding areas. Grasshoppers are generalist feeders, eating grasses and forbs first and often moving to cultivated crops. High grasshopper density of one or several species and the resulting defoliation may reach an economic threshold where the damage caused by grasshoppers exceeds the cost of controlling the grasshoppers. Researchers determined that during typical grasshopper infestation years, approximately 20% of forage rangeland is removed, valued at a dollar adjusted amount of \$900 million. This value represents 32 to 63% of the total value of rangeland across the western states (Rashford et al., 2012). Other market and non-market values such as carbon sequestration, general ecosystem services, and recreational use may also be impacted by pest outbreaks in rangeland.

Vegetation damage during serious grasshopper outbreaks may be so severe that all grasses and forbs are destroyed; thus, plant growth is impaired for several years. Rare plants may be consumed during critical times of development such as during seed production, and loss of important plant species, or seed production may lead to reduced biological diversity of the rangeland habitats, potentially creating opportunities for the expansion of invasive and exotic weeds (Lockwood and Latchininsky, 2000). When grasshoppers consume plant cover, soil is more susceptible to the drying effects of the sun, making plant roots less capable of holding soil in place. Soil damage results in erosion and disruption of nutrient cycling, water infiltration, seed germination, and other ecological processes which are important components of rangeland ecosystems (Latchininsky et al., 2011).

When the density of grasshoppers reaches economic infestation levels, grasshoppers begin to compete with livestock for food by reducing available forage (Wakeland and Shull, 1936; Belovsky, 2000; Pfadt, 2002; Branson et al., 2006; Bradshaw et al., 2018). Ranchers could offset some of the costs by leasing rangeland in another area and relocating their livestock, finding other means to feed their animals by purchasing hay or grain, or selling their livestock. Ranchers could also incur economic losses from personal attempts to control grasshopper damage to rangeland. Local communities could see adverse economic impacts to the entire area. Grasshoppers that infest rangeland could move to surrounding croplands. Farmers could incur economic losses from attempts to chemically control grasshopper populations or due to the loss of their crops. The general public could see an increase in the cost of meat, crops, and their byproducts.

2. Insecticide Applications at Conventional Rates or Reduced Agent Area Treatments with Adaptive Management Strategy

Under Alternative 2, APHIS would participate in grasshopper programs with the option of using one of the insecticides carbaryl or diflubenzuron depending upon the various factors related to the grasshopper outbreak and the site-specific characteristics. The use of an

insecticide would typically occur at half the conventional application rates following the RAATs strategy. APHIS would apply a single treatment to affected rangeland areas to suppress grasshopper outbreak populations by a range of 35 to 98 percent, depending upon the insecticide used.

a) Carbaryl

Carbaryl is a member of the N-methyl carbamate class of insecticides, which affect the nervous system via cholinesterase inhibition. Inhibiting the enzyme acetylcholinesterase (AChE) causes nervous system signals to persist longer than normal. While these effects are desired in controlling insects, they can have undesirable impacts to non-target organisms that are exposed. The APHIS HHERA assessed available laboratory studies regarding the toxicity of carbaryl on fish and wildlife. In summary, the document indicates the chemical is highly toxic to insects, including native bees, honeybees, and aquatic insects; slightly to highly toxic to fish; highly to very highly toxic to most aquatic crustaceans, moderately toxic to mammals, minimally toxic to birds; moderately to highly toxic to several terrestrial arthropod predators; and slightly to highly toxic to larval amphibians (USDA APHIS, 2018a). However, adherence to label requirements and additional program measures designed to prevent carbaryl from reaching sensitive habitats or mitigate exposure of non-target organisms will reduce environmental effects of treatments.

The offsite movement and deposition of carbaryl after treatments is unlikely because it does not significantly vaporize from the soil, water, or treated surfaces (Dobroski et al., 1985). Temperature, pH, light, oxygen, and the presence of microorganisms and organic material are factors that contribute to how quickly carbaryl will degrade in water. Hydrolysis, the breaking of a chemical bond with water, is the primary degradation pathway for carbaryl at pH 7 and above. In natural water, carbaryl is expected to degrade faster than in laboratory settings due to the presence of microorganisms. The half-lives of carbaryl in natural waters varied between 0.3 to 4.7 days (Stanley and Trial, 1980; Bonderenko et al., 2004). Degradation in the latter study was temperature dependent with shorter half-lives at higher temperatures. Aerobic aquatic metabolism of carbaryl reported half-life ranged of 4.9 to 8.3 days compared to anaerobic (without oxygen) aquatic metabolism range of 15.3 to 72 days (Thomson and Strachan, 1981; USEPA, 2003). Carbaryl is not persistent in soil due to multiple degradation pathways including hydrolysis, photolysis, and microbial metabolism. Little transport of carbaryl through runoff or leaching to groundwater is expected due to the low water solubility, moderate sorption, and rapid degradation in soils. There are no reports of carbaryl detection in groundwater, and less than 1% of granule carbaryl applied to a sloping plot was detected in runoff (Caro et al., 1974).

Acute and chronic risks to mammals are expected to be low to moderate based on the available toxicity data and conservative assumptions that were used to evaluate risk. There is the potential for impacts to small mammal populations that rely on terrestrial invertebrates for food. However, based on the toxicity data for terrestrial plants, minimal risks of indirect effects are expected to mammals that rely on plant material for food. Carbaryl has a reported half-life on vegetation of three to ten days, suggesting mammal exposure would be short-term. Direct risks to mammals from carbaryl bait applications is expected to be minimal based on oral, dermal, and inhalation studies (USDA APHIS, 2018a).

A number of studies have reported no effects on bird populations in areas treated with carbaryl (Buckner et al., 1973; Richmond et al., 1979; McEwen et al., 1996). Some applications of formulated carbaryl were found to cause depressed AChE levels (Zinkl et al., 1977; Gramlich, 1979); however, the doses were twice those proposed for the full coverage application in the grasshopper program.

While sublethal effects have been noted in fish with depressed AChE, as well as some impacts to amphibians (i.e. days to metamorphosis) and aquatic invertebrates in the field due to carbaryl, the application rates and measured aquatic residues observed in these studies are well above values that would be expected from current program operations. Indirect risks to amphibian and fish species can occur through the loss of habitat or reduction in prey, yet data suggests that carbaryl risk to aquatic plants that may serve as habitat, or food, for fish and aquatic invertebrates is very low.

Product use restrictions appear on the USEPA-approved label and attempt to keep carbaryl out of waterways. Carbaryl must not be applied directly to water, or to areas where surface water is present (USEPA, 2012c). The USEPA-approved use rates and patterns and the additional mitigations imposed by the grasshopper program, such as using RAATs and application buffers, where applicable, further minimize aquatic exposure and risk.

The majority of rangeland plants require insect-mediated pollination. Native, solitary bee species are important pollinators on western rangeland (Tepedino, 1979). Potential negative effects of insecticides on pollinators are of concern because a decrease in their numbers has been associated with a decline in fruit and seed production of plants. Laboratory studies have indicated that bees are sensitive to carbaryl applications, but the studies were at rates above those proposed in the program. The reduced rates of carbaryl used in the program and the implementation of application buffers should significantly reduce exposure of carbaryl applications to pollinators. In areas of direct application where impacts may occur, alternating swaths and reduced rates (i.e., RAATs) would reduce risk. Potential negative effects of grasshopper program insecticides on bee populations may also be mitigated by the more common use of carbaryl baits than the ULV spray formulation. Studies with carbaryl bran bait have found no sublethal effects on adults or larvae bees (Peach et al., 1994, 1995).

Carbaryl can cause cholinesterase inhibition (i.e., overstimulate the nervous system) in humans resulting in nausea, headaches, dizziness, anxiety, and mental confusion, as well as convulsions, coma, and respiratory depression at high levels of exposure (NIH, 2009a; Beauvais, 2014). USEPA classifies carbaryl as “likely to be carcinogenic to humans” based on vascular tumors in mice (USEPA, 2007, 2015a, 2017a).

USEPA regulates the amount of pesticide residues that can remain in or on food or feed commodities as the result of a pesticide application. The agency does this by setting a tolerance, which is the maximum residue level of a pesticide, usually measured in parts per million (ppm), that can legally be present in food or feed. USEPA-registered carbaryl products used by the grasshopper program are labeled with rates and treatment intervals that are meant to protect livestock and keep chemical residues in cattle at acceptable levels (thereby protecting human health). While livestock and horses may graze on rangeland the same day that the land is sprayed, in order to keep tolerances to acceptable levels, carbaryl

spray applications on rangeland are limited to half a pound active ingredient per acre per year (USEPA, 2012c). The grasshopper program would treat at or below use rates that appear on the label, as well as follow all appropriate label mitigations, which would ensure residues are below the tolerance levels.

Adverse human health effects from the proposed program ULV applications of the carbaryl spray (Sevin[®] XLR Plus) and bait applications of the carbaryl 5% and 2% baits formulations to control grasshoppers are not expected based on low potential for human exposure to carbaryl and the favorable environmental fate and effects data. Technical grade (approximately 100% of the insecticide product is composed of the active ingredient) carbaryl exhibits moderate acute oral toxicity in rats, low acute dermal toxicity in rabbits, and very low acute inhalation toxicity in rats. Technical carbaryl is not a primary eye or skin irritant in rabbits and is not a dermal sensitization in guinea pig (USEPA, 2007). This data can be extrapolated and applied to humans revealing low health risks associated with carbaryl.

The Sevin[®] XLR Plus formulation, which contains a lower percent of the active ingredient than the technical grade formulation, is less toxic via the oral route, but is a mild irritant to eyes and skin. The proposed use of carbaryl as a ULV spray or a bait, use of RAATs, and adherence to label requirements, substantially reduces the potential for exposure to humans. Program workers are the most likely human population to be exposed. APHIS does not expect adverse health risks to workers based on low potential for exposure to carbaryl when applied according to label directions and use of personal protective equipment (PPE) (e.g., long-sleeved shirt and long pants, shoes plus socks, chemical-resistant gloves, and chemical-resistant apron) (USEPA, 2012c) during loading and applications. APHIS quantified the potential health risks associated with accidental worker exposure to carbaryl during mixing, loading, and applications. The quantitative risk evaluation results indicate no concerns for adverse health risk for program workers (<http://www.aphis.usda.gov/plant-health/grasshopper>).

Adherence to label requirements and additional program measures designed to reduce exposure to workers and the public (e.g., mitigations to protect water sources, mitigations to limit spray drift, and restricted-entry intervals) result in low health risk to all human population segments.

b) Diflubenzuron

Diflubenzuron is a restricted use pesticide (only certified applicators or persons under their direct supervision may make applications) registered with USEPA as an insect growth regulator. It specifically interferes with chitin synthesis, the formation of the insect's exoskeleton. Larvae of affected insects are unable to molt properly. While this effect is desirable in controlling certain insects, it can have undesirable impacts to non-target organisms that are exposed.

USEPA considers diflubenzuron relatively non-persistent and immobile under normal use conditions and stable to hydrolysis and photolysis. The chemical is considered unlikely to contaminate ground water or surface water (USEPA, 1997). The vapor pressure of diflubenzuron is relatively low, as is the Henry's Law Constant value, suggesting the chemical will not volatilize readily into the atmosphere from soil, plants or water.

Therefore, exposure from volatilization is expected to be minimal. Due to its low solubility (0.2 mg/L) and preferential binding to organic matter, diflubenzuron seldom persists more than a few days in water (Schaefer and Dupras, 1977; Schaefer et al., 1980). Mobility and leachability of diflubenzuron in soils is low, and residues are usually not detectable after seven days (Eisler, 2000). Aerobic aquatic half-life data in water and sediment was reported as 26.0 days (USEPA, 1997). Diflubenzuron applied to foliage remains adsorbed to leaf surfaces for several weeks with little or no absorption or translocation from plant surfaces (Eisler, 1992, 2000). Field dissipation studies in California citrus and Oregon apple orchards reported half-life values of 68.2 to 78 days (USEPA, 2018). Diflubenzuron persistence varies depending on site conditions and rangeland persistence is unfortunately not available. Diflubenzuron degradation is microbially mediated with soil aerobic half-lives much less than dissipation half-lives. Diflubenzuron treatments are expected to have minimal effects on terrestrial plants. Both laboratory and field studies demonstrate no effects using diflubenzuron over a range of application rates, and the direct risk to terrestrial plants is expected to be minimal (USDA APHIS, 2018c).

Dimilin[®] 2L is labeled with rates and treatment intervals that are meant to protect livestock and keep residues in cattle at acceptable levels (thereby, protecting human health). Tolerances are set for the amount of diflubenzuron that is allowed in cattle fat (0.05 ppm) and meat (0.05 ppm) (40 CFR Parts 180.377). The grasshopper program would treat at application rates indicated on product labels or lower, which should ensure approved residues levels.

APHIS' literature review found that on an acute basis, diflubenzuron is considered toxic to some aquatic invertebrates and practically non-toxic to adult honeybees. However, diflubenzuron is toxic to larval honeybees (USEPA, 2018). It is slightly nontoxic to practically nontoxic to fish and birds and has very slight acute oral toxicity to mammals, with the most sensitive endpoint from exposure being the occurrence of methemoglobinemia (a condition that impairs the ability of the blood to carry oxygen). Minimal direct risk to amphibians and reptiles is expected, although there is some uncertainty due to lack of information (USDA APHIS, 2018c; USEPA, 2018).

Risk is low for most non-target species based on laboratory toxicity data, USEPA approved use rates and patterns, and additional mitigations such as the use of lower rates and RAATs that further reduces risk. Risk is greatest for sensitive terrestrial and aquatic invertebrates that may be exposed to diflubenzuron residues.

In a review of mammalian field studies, Dimilin[®] applications at a rate of 60 to 280 g a.i./ha had no effects on the abundance and reproduction in voles, field mice, and shrews (USDA FS, 2004). These rates are approximately three to 16 times greater than the highest application rate proposed in the program. Potential indirect impacts from application of diflubenzuron on small mammals includes loss of habitat or food items. Mice on treated plots consumed fewer lepidopteran (order of insects that includes butterflies and moths) larvae compared to controls; however, the total amount of food consumed did not differ between treated and untreated plots. Body measurements, weight, and fat content in mice collected from treated and non-treated areas did not differ.

Poisoning of insectivorous birds by diflubenzuron after spraying in orchards at labeled rates is unlikely due to low toxicity (Muzzarelli, 1986). The primary concern for bird species is related to an indirect effect on insectivorous species from a decrease in insect prey. At the proposed application rates, grasshoppers have the highest risk of being impacted while other taxa have a much reduced risk because the lack of effects seen in multiple field studies on other taxa of invertebrates at use rates much higher than those proposed for the program. Shifting diets in insectivorous birds in response to prey densities is not uncommon in undisturbed areas (Rosenberg et al., 1982; Cooper et al., 1990; Sample et al., 1993).

Indirect risk to fish species can be defined as a loss of habitat or prey base that provides food and shelter for fish populations, however these impacts are not expected based on the available fish and invertebrate toxicity data (USDA APHIS, 2018c). A review of several aquatic field studies demonstrated that when effects were observed it was at diflubenzuron levels not expected from program activities (Fischer and Hall, 1992; USEPA, 1997; Eisler, 2000; USDA FS, 2004).

Diflubenzuron applications have the potential to affect chitin production in various other beneficial terrestrial invertebrates. Multiple field studies in a variety of application settings, including grasshopper control, have been conducted regarding the impacts of diflubenzuron to terrestrial invertebrates. Based on the available data, sensitivity of terrestrial invertebrates to diflubenzuron is highly variable depending on which group of insects and which life stages are being exposed. Immature grasshoppers, beetle larvae, lepidopteran larvae, and chewing herbivorous insects appear to be more susceptible to diflubenzuron than other invertebrates. Within this group, however, grasshoppers appear to be more sensitive to the proposed use rates for the program. Honeybees, parasitic wasps, predatory insects, and sucking insects show greater tolerance to diflubenzuron exposure (Murphy et al., 1994; Eisler, 2000; USDA FS, 2004).

Diflubenzuron is moderately toxic to spiders and mites (USDA APHIS, 2018c). Deakle and Bradley (1982) measured the effects of four diflubenzuron applications on predators of *Heliothis* spp. at a rate of 0.06 lb a.i./ac and found no effects on several predator groups. This supported earlier studies by Keever et al. (1977) that demonstrated no effects on the arthropod predator community after multiple applications of diflubenzuron in cotton fields. Grasshopper integrated pest management (IPM) field studies have shown diflubenzuron to have a minimal impact on ants, spiders, predatory beetles, and scavenger beetles. There was no significant reduction in populations of these species from seven to 76 days after treatment. Although ant populations exhibited declines of up to 50 percent, these reductions were temporary, and population recovery was described as immediate (Catangui et al., 1996).

Due to its mode of action, diflubenzuron has greater activity on immature stages of terrestrial invertebrates. Based on standardized laboratory testing diflubenzuron is considered practically non-toxic to adult honeybees. The contact LD50 value for the honeybee, *Apis mellifera*, is reported at greater than 114.8 µg a.i./bee while the oral LD50 value was reported at greater than 30 µg a.i./bee. USEPA (2018) reports diflubenzuron toxicity values to adult honeybees are typically greater than the highest test concentration using the end-use product or technical active ingredient. The lack of toxicity to honeybees, as well as other bees, in laboratory studies has been confirmed in additional studies (Nation

et al., 1986; Chandel and Gupta, 1992; Mommaerts et al., 2006). Mommaerts et al. (2006) and Thompson et al. (2005) documented sublethal effects on reproduction-related endpoints for the bumble bee, *Bombus terrestris* and *A. mellifera*, respectively, testing a formulation of diflubenzuron. However, these effects were observed at much higher use rates relative to those used in the program.

Insecticide applications to rangelands have the potential to impact pollinators, and in turn, vegetation and various rangeland species that depend on pollinated vegetation. Based on the review of laboratory and field toxicity data for terrestrial invertebrates, applications of diflubenzuron are expected to have minimal risk to pollinators of terrestrial plants. The use of RAATs provide additional benefits by using reduced rates and creating untreated swaths within the spray block that will further reduce the potential risk to pollinators.

APHIS reduces the risk to native bees and pollinators through monitoring grasshopper and Mormon cricket populations and making pesticide applications in a manner that reduces the risk to this group of nontarget invertebrates. Monitoring grasshopper and Mormon cricket populations allows APHIS to determine if populations require treatment and to make treatments in a timely manner reducing pesticide use and emphasizing the use of Program insecticides that are not broad spectrum. Historical use of Program insecticides demonstrate that diflubenzuron is the preferred insecticide for use. Over 90% of the acreage treated by the Program has been with diflubenzuron. Diflubenzuron poses a reduced risk to native bees and pollinators compared to liquid carbaryl and malathion applications.

Adverse human health effects from ground or aerial ULV applications of diflubenzuron to control grasshoppers are not expected based on the low acute toxicity of diflubenzuron and low potential for human exposure. The adverse health effects of diflubenzuron to mammals and humans involves damage to hemoglobin in blood and the transport of oxygen. Diflubenzuron causes the formation of methemoglobin. Methemoglobin is a form of hemoglobin that is not able to transport oxygen (USDA FS, 2004). USEPA classifies diflubenzuron as non-carcinogenic to humans (USEPA, 2015b).

Program workers adverse health risks are not likely when diflubenzuron is applied according to label directions that reduce or eliminate exposures. Adverse health risk to the general public in treatment areas is not expected due to the low potential for exposure resulting from low population density in the treatment areas, adherence to label requirements, program measures designed to reduce exposure to the public, and low toxicity to mammals.

c) Reduced Area Agent Treatments (RAATs)

The use of RAATS is the most common application method for all program insecticides and would continue to be so, except in rare pest conditions that warrant full coverage and higher rates. The goal of the RAATs strategy is to suppress grasshopper populations to a desired level, rather than to reduce those populations to the greatest possible extent. This strategy has both economic and environmental benefits. APHIS would apply a single application of insecticide per year, typically using a RAATs strategy that decreases the rate of insecticide applied by either using lower insecticide spray concentrations, or by alternating one or more treatment swaths. Usually RAATs applications use both lower concentrations and skip treatment swaths. The RAATs strategy suppresses grasshoppers

within treated swaths, while conserving grasshopper predators and parasites in swaths that are not treated.

The concept of reducing the treatment area of insecticides while also applying less insecticide per treated acre was developed in 1995, with the first field tests of RAATs in Wyoming (Lockwood and Schell, 1997). Applications can be made either aerially or with ground-based equipment (Deneke and Keyser, 2011). Studies using the RAATs strategy have shown good control (up to 85% of that achieved with a total area insecticide application) at a significantly lower cost and less insecticide, and with a markedly higher abundance of non-target organisms following application (Lockwood et al., 2000; Deneke and Keyser, 2011). Levels of control may also depend on variables such as body size of targeted grasshoppers, growth rate of forage, and the amount of coverage obtained by the spray applications (Deneke and Keyser, 2011). Control rates may also be augmented by the necrophilic and necrophagic behavior of grasshoppers, in which grasshoppers are attracted to volatile fatty acids emanating from cadavers of dead grasshoppers and move into treated swaths to cannibalize cadavers (Lockwood et al., 2002; Smith and Lockwood, 2003). Under optimal conditions, RAATs decrease control costs, as well as host plant losses and environmental effects (Lockwood et al., 2000; Lockwood et al., 2002).

The efficacy of a RAATs strategy in reducing grasshoppers is, therefore, less than conventional treatments and more variable. Foster et al. (2000) reported that grasshopper mortality using RAATs was reduced 2 to 15% from conventional treatments, depending on the insecticide, while Lockwood et al. (2000) reported 0 to 26% difference in mortality between conventional and RAATs methods. APHIS will consider the effects of not suppressing grasshoppers to the greatest extent possible as part of the treatment planning process.

RAATs reduces treatment costs and conserves non-target biological resources in untreated areas. The potential economic advantages of RAATs was proposed by Larsen and Foster (1996), and empirically demonstrated by Lockwood and Schell (1997). Widespread efforts to communicate the advantages of RAATs across the Western States were undertaken in 1998 and have continued on an annual basis. The viability of RAATs at an operational scale was initially demonstrated by Lockwood et al. (2000), and subsequently confirmed by Foster et al. (2000). The first government agencies to adopt RAATs in their grasshopper suppression programs were the Platte and Goshen County Weed and Pest Districts in Wyoming; they also funded research at the University of Wyoming to support the initial studies in 1995. This method is now commonly used by government agencies and private landowners in States where grasshopper control is required.

Reduced rates should prove beneficial for the environment. All APHIS grasshopper treatments using carbaryl, diflubenzuron, or malathion are conducted in adherence with USEPA-approved label directions. Labeled application rates for grasshopper control tend to be lower than rates used against other pests. In addition, use rates proposed for grasshopper control by APHIS are lower than rates used by private landowners.

B. Other Environmental Considerations

1. Cumulative Impacts

Cumulative impact, as defined in the Council on Environmental Quality (CEQ) NEPA implementing regulations (40 CFR § 1508.7) “is the impact on the environment which results from the incremental impact of the action when added to the past, present, and reasonably foreseeable future actions regardless of what agency (Federal or non-Federal) or person undertakes such actions. Cumulative impacts can result from individually minor but collectively significant actions taking place over a period of time.”

Potential cumulative impacts associated with the No Action alternative where APHIS would not take part in any grasshopper suppression program include the continued increase in grasshopper populations and potential expansion of populations into neighboring range and cropland. In addition, State and private land managers could apply insecticides to manage grasshopper populations however, land managers may opt not to use RAATs, which would increase insecticides applied to the rangeland. Increased insecticide applications from the lack of coordination or foregoing RAATs methods could increase the exposure risk to non-target species. In addition, land managers may not employ the extra program measures designed to reduce exposure to the public and the environment to insecticides.

Potential cumulative impacts associated with the Preferred Alternative are not expected to be significant because the program applies an insecticide application once during a treatment. The program may treat an area with different insecticides but does not overlap the treatments. The program does not mix or combine insecticides. Based on historical outbreaks in the United States, the probability of an outbreak occurring in the same area where treatment occurred in the previous year is unlikely; however, given time, populations eventually will reach economically damaging thresholds and require treatment

The insecticide application reduces the insect population down to levels that cause an acceptable level of economic damage. The duration of treatment activity, which is relatively short since it is a one-time application, and the lack of repeated treatments in the same area in the same year reduce the possibility of significant cumulative impacts.

Potential cumulative impacts resulting from the use of insecticides include insect pest resistance, synergistic chemical effects, chemical persistence and bioaccumulation in the environment. The program use of reduced insecticide application rates (i.e. ULV and RAATs) are expected to mitigate the development of insect resistance to the insecticides. Grasshopper outbreaks in the United States occur cyclically so applications do not occur to the same population over time further eliminating the selection pressure increasing the chances of insecticide resistance.

The insecticides proposed for use in the program have a variety of agricultural and non-agricultural uses. There may be an increased use of these insecticides in an area under suppression when private, State, or Federal entities make applications to control other pests. However, the vast majority of the land where program treatments occur is uncultivated rangeland and additional treatments by landowners or managers are very uncommon making possible cumulative or synergistic chemical effects extremely unlikely.

The insecticides proposed for use in the grasshopper program are not anticipated to persist in the environment or bioaccumulate. Therefore, a grasshopper outbreak that occurs in an area previously treated for grasshoppers is unlikely to cause an accumulation of insecticides from previous program treatments.

Herbicide treatments for invasive species have been conducted on the San Carlos Reservation by Tribal management agencies. These areas are not located in the 2021 Action area for the rangeland grasshopper suppression program. Therefore, there would be no synergistic effect from an overlap of pesticide and herbicide treatments.

2. Executive Order No. 12898, Federal Actions to Address Environmental Justice in Minority Populations and Low-Income Populations

Federal agencies identify and address the disproportionately high and adverse human health or environmental effects of their proposed activities, as described in E.O. 12898, “Federal Actions to Address Environmental Justice in Minority Populations and Low-Income Populations.”

APHIS has evaluated the proposed grasshopper program and has determined that there is no disproportionately high and adverse human health or environmental effects on minority populations or low-income populations.

3. Executive Order No. 13045, Protection of Children from Environmental Health Risks and Safety Risks

Federal agencies consider a proposed action’s potential effects on children to comply with E.O. 13045, “Protection of Children from Environmental Health Risks and Safety Risks.” This E.O. requires each Federal agency, consistent with its mission, to identify and assess environmental health and safety risks that may disproportionately affect children and to ensure its policies, programs, activities, and standards address disproportionate risks to children that result from environmental health risks or safety risks. APHIS has developed agency guidance for its programs to follow to ensure the protection of children (USDA APHIS, 1999).

APHIS’ HHERAs evaluated the potential exposure to each insecticide used in the program and risks associated with these insecticides to residents, including children. The HHERAs for the proposed program insecticides, located at <http://www.aphis.usda.gov/plant-health/grasshopper>, suggest that no disproportionate risks to children, as part of the general public, are anticipated.

According to the BUREAU OF WOMEN'S AND CHILDREN'S HEALTH, Arizona Department of Health Services, there are 3,235 children between the ages of 0-14. There is approximately 1,151 youth from the ages of 15-19 according to the Arizona Department of Health Services. The risk for children to be exposed to treatment pesticides is very low due to the remote nature of the Tribal Rangeland. The nearest communities are approximately 50 miles from the Tribal rangeland areas. There will be no aerial treatments conducted in Arizona only by ground-based equipment.

APHIS grasshopper insecticide treatments are conducted in rural rangeland areas, where agriculture is a primary industry. The areas consist of widely scattered, single, rural dwellings in ranching communities with low population density. The program notifies residents within treatment areas, or their designated representatives, prior to proposed operations to reduce the potential for incidental exposure to residents including children. Treatments are conducted primarily on open rangelands where children would not be expected to be present during treatment or to enter should there be any restricted entry period after treatment. The program also implements mitigation measures beyond label requirements to ensure that no treatments occur within the required buffer zones from structures, such as a 500-foot treatment buffer zone from schools and recreational areas. Program insecticides are not applied while school buses are operating in the treatment area.

4. Tribal Consultation

Executive Order 13175 "Consultation and Coordination with Indian Tribal Governments," calls for agency communication and collaboration with tribal officials when proposed Federal actions have potential tribal implications. The Archaeological Resources Protection Act of 1979 (16 U.S.C. §§ 470aa-mm), secures the protection of archaeological resources and sites on public and tribal lands.

Prior to the treatment season, program personnel notify Tribal land managers of the potential for grasshopper and Mormon cricket outbreaks on their lands. Consultation with local Tribal representatives takes place prior to treatment programs to inform fully the Tribes of possible actions APHIS may take on Tribal lands. Treatments typically do not occur at cultural sites, and drift from a program treatment at such locations is not expected to adversely affect natural surfaces, such as rock formations and carvings. APHIS would also confer with the appropriate Tribal authority to ensure that the timing and location of a planned program treatment does not coincide or conflict with cultural events or observances on Tribal lands.

5. Executive Order 13186, Responsibilities of Federal Agencies to Protect Migratory Birds

The Migratory Bird Treaty Act (MBTA) of 1918 (16 U.S.C. 703–712) established a Federal prohibition, unless permitted by regulations, to pursue, hunt, take, capture, kill, attempt to take, capture or kill, possess, offer for sale, sell, offer to purchase, purchase, deliver for shipment, ship, cause to be shipped, deliver for transportation, transport, cause to be transported, carry, or cause to be carried by any means whatever, receive for shipment, transportation or carriage, or export, at any time, or in any manner, any migratory bird or any part, nest, or egg of any such bird.

APHIS will support the conservation intent of the migratory bird conventions by integrating bird conservation principles, measures, and practices into agency activities and by avoiding or reducing, to the extent practicable, adverse impacts on migratory bird resources when conducting agency actions. Impacts are minimized as a result of buffers to water, habitat, nesting areas, riparian areas, and the use of RAATs. For any given treatment, only a portion of the environment will be treated, therefore minimizing potential impacts to migratory bird populations.

6. Endangered Species Act

Section 7 of the Endangered Species Act (ESA) and its implementing regulations require Federal agencies to ensure their actions are not likely to jeopardize the continued existence of listed threatened or endangered species or result in the destruction or adverse modification of critical habitat. Numerous federally listed species and areas of designated critical habitat occur within the 17-State program area, although not all occur within or near potential grasshopper suppression areas or within the area under consideration by through this EA.

APHIS considers whether listed species, species proposed for listing, experimental populations, or critical habitat are present in the proposed suppression area. Before treatments are conducted, APHIS contacts the U.S Fish and Wildlife Service (USFWS) or the National Marine Fisheries Service (NMFS) (where applicable) to determine if listed species are present in the suppression area, and whether mitigations or protection measures must be implemented to protect listed species or critical habitat.

APHIS completed a programmatic Section 7 consultation with NMFS for use of carbaryl, malathion, and diflubenzuron to suppress grasshoppers in the 17-state program area because of the listed salmonid (*Oncorhynchus* spp.) and critical habitat. To minimize the possibility of insecticides from reaching salmonid habitat, APHIS implements the following protection measures:

- RAATs are used in all areas adjacent to salmonid habitat
- ULV sprays are used, which are between 50% and 66% of the USEPA recommended rate
- Insecticides are not aerially applied in a 3,500-foot buffer zones for carbaryl or malathion, or applied within a 1,500-foot buffer zones for diflubenzuron along stream corridors
- Insecticides will not be applied when wind speeds exceed 10 miles per hour. APHIS will attempt to avoid insecticide application if the wind is blowing towards salmonid habitat
- Insecticide applications are avoided when precipitation is likely or during temperature inversions

APHIS determined that with the implementation of these measures, the grasshopper suppression program may affect, but is not likely to adversely affect listed salmonids or designated critical habitat in the program area. NMFS concurred with this determination in a letter dated April 12, 2010.

APHIS submitted a programmatic biological assessment for grasshopper suppression in the 17-state program area and requested consultation with USFWS on March 9, 2015. With the incorporation and use of application buffers and other operational procedures APHIS anticipates that any impacts associated with the use and fate of program insecticides will be insignificant and discountable to listed species and their habitats. Based on an assessment of the potential exposure, response, and subsequent risk characterization of program operations, APHIS concludes the proposed action is not likely to adversely affect listed species or critical habitat in the program area. APHIS has requested concurrence from the USFWS on these determinations. Until this programmatic Section 7 consultation with

USFWS is completed, APHIS will conduct consultations with USFWS field offices at the local level.

APHIS considers the role of pollinators in any consultations conducted with the FWS to protect federally listed plants. Mitigation measures, such as no treatment buffers are applied with consideration of the protection of pollinators that are important to a listed plant species. Correspondence from FWS is in appendix E of this EA. There are no species in Arizona regulated by NMFS. No consultation or concurrence from NMFS is needed.

7. Bald and Golden Eagle Protection Act

The Bald and Golden Eagle Protection Act (16 U.S.C. 668–668c) prohibits anyone, without a permit issued by the Secretary of the Interior, from “taking” bald eagles, including their parts, nests, or eggs. During the breeding season, bald eagles are sensitive to a variety of human activities. Grasshopper management activities could cause disturbance of nesting eagles, depending on the duration, noise levels, extent of the area affected by the activity, prior experiences that eagles have with humans, and tolerance of the individual nesting pair. Also, disruptive activities in or near eagle foraging areas can interfere with bald eagle feeding, reducing chances of survival. USFWS has provided recommendations for avoiding disturbance at foraging areas and communal roost sites that are applicable to grasshopper management programs (USFWS, 2007).

No toxic effects are anticipated on eagles as a direct consequence of insecticide treatments. Toxic effects on the principle food source, fish, are not expected because insecticide treatments will not be conducted over rivers or lakes. Buffers protective of aquatic biota are applied to their habitats to ensure that there are no indirect effects from loss of prey.

8. Additional Species of Concern

There may be species that are of special concern to land management agencies, the public, or other groups and individuals in proposed treatment areas. For example, the sage grouse populations have declined throughout most of their entire range, with habitat loss being a major factor in their decline.

Grasshopper suppression programs reduce grasshoppers and at least some other insects in the treatment area that can be a food item for sage grouse chicks. As indicated in previous sections on impacts to birds, there is low potential that the program insecticides would be toxic to sage grouse, either by direct exposure to the insecticides or indirectly through immature sage grouse eating moribund grasshoppers.

Because grasshopper numbers are so high in an outbreak year, treatments would not likely reduce the number of grasshoppers below levels present in a normal year which would usually range from 3-7 gh/yd². Should grasshoppers be unavailable in small, localized areas, sage grouse chicks may consume other insects, which sage grouse chicks likely do in years when grasshopper numbers are naturally low. By suppressing grasshoppers, rangeland vegetation is available for use by other species, including sage grouse, and rangeland areas are less susceptible to invasive plants that may be undesirable for sage grouse habitat.

9. Fires and Human Health Hazards

Various compounds are released in smoke during wildland fires, including carbon monoxide (CO), carbon dioxide, nitrous oxides, sulfur dioxide, hydrogen chloride, aerosols, polynuclear aromatic hydrocarbons contained within fine particulate matter (a byproduct of the combustion of organic matter such as wood), aldehydes, and most notably formaldehyde produced from the incomplete combustion of burning biomass (Reisen and Brown, 2009; Burling et al., 2010; Broyles, 2013). Particulate matter, CO, benzene, acrolein, and formaldehyde have been identified as compounds of particular concern in wildland fire smoke (Reinhardt and Ottmar, 2004).

Many of the naturally occurring products associated with combustion from wildfires may also be present as a result of combustion of program insecticides that are applied to rangeland. These combustion byproducts will be at lower quantities due to the short half-lives of most of the program insecticides and their low use rates. Other minor combustion products specific to each insecticide may also be present as a result of combustion from a rangeland fire but these are typically less toxic based on available human health data (<http://www.aphis.usda.gov/plant-health/grasshopper>).

The safety data sheet for each insecticide identifies these combustion products for each insecticide as well as recommendations for PPE. The PPE is similar to what typically is used in fighting wildfires. Material applied in the field will be at a much lower concentration than what would occur in a fire involving a concentrated formulation. Therefore, the PPE worn by rangeland firefighters would also be protective of any additional exposure resulting from the burning of residual insecticides.

10. Cultural and Historical Resources

Federal actions must seek to avoid, minimize, and mitigate potential negative impacts to cultural and historic resources as part of compliance with the National Historic Preservation Act (NHPA), the Archaeological Resources Protection Act of 1979, and NEPA. Section 106 of the NHPA requires Federal agencies to provide the Advisory Council on Historic Preservation with an opportunity to comment on their findings. There are no known historic resources and National Trails within the proposed action area.

V. Literature Cited

- Beauvais, S. 2014. Human exposure assessment document for carbaryl. Page 136. California Environmental Protection Agency, Department of Pesticide Regulation.
- Belovsky, G. E., A. Joern, and J. Lockwood. 1996. VII.16 Grasshoppers—Plus and Minus: The Grasshopper Problem on a Regional Basis and a Look at Beneficial Effects of Grasshoppers. Pages 1-5 in G. L. Cunningham and M. W. Sampson, editors. Grasshopper Integrated Pest Management User Handbook, Technical Bulletin No. 1809. U.S. Department of Agriculture, Animal and Plant Health Inspection Service, Washington, DC.
- Belovsky, G. E. 2000. Part 1. Grasshoppers as integral elements of grasslands. 1. Do grasshoppers diminish grassland productivity? A new perspective for control based

- on conservation. Pages 7-29 in J. A. Lockwood et al, editor. Grasshoppers and Grassland Health. Kluwer Academic Publishers, Netherlands.
- Bonderenko, S., J. Gan, D. L. Haver, and J. N. Kabashima. 2004. Persistence of selected organophosphate and carbamate insecticides in waters from coastal watershed. *Env. Toxicol. Chem.* 23:2649-2654.
- Bradshaw, J. D., K. H. Jenkins, and S. D. Whipple. 2018. Impact of grasshopper control on forage quality and availability in western Nebraska. *Rangelands* 40:71-76.
- Branson, D., A. Joern, and G. Sword. 2006. Sustainable management of insect herbivores in grassland ecosystems: new perspectives in grasshopper control. *BioScience* 56:743-755.
- Broyles, G. 2013. Wildland firefighter smoke exposure. Page 26. U.S. Department of Agriculture, Forest Service.
- Buckner, C. H., P. D. Kingsbury, B. B. McLeod, K. L. Mortensen, and D. G. H. Ray. 1973. The effects of pesticides on small forest vertebrates of the spruce woods provincial forest, Manitoba. *The Manitoba Entomologist* 7:37-45.
- Burling, I., R. Yokelson, D. Griffith, T. Johson, P. Veres, J. Roberts, C. Warneke, S. Urbanski, J. Reardon, D. Weise, W. Hao, and J. de Gouw. 2010. Laboratory measures of trace gas emissions from biomass burning of fuel types from the southeastern and southwestern United States. *Atmospheric Chemistry and Physics* 10:11115-111130.
- Caro, J. H., H. P. Freeman, and B. C. Turner. 1974. Persistence in soil and losses in runoff of soil-incorporated carbaryl in a small watershed. *J. Agric. Food Chem.* 22:860-863.
- Catangui, M.A., Fuller, B.W., and Walz, A.W., 1996. Impact of Dimilin® on nontarget arthropods and its efficacy against rangeland grasshoppers. *In* U.S. Department of Agriculture, Animal and Plant Health Inspection Service, 1996. Grasshopper Integrated Pest Management User Handbook, Tech. Bul. No. 1809. Sec. VII.3. Washington, DC.
- Chandel, R.S., and P.R Gupta. 1992. Toxicity of diflubenzuron and penfluron to immature stages of *Apis cerana indica* and *Apis mellifera*. *Apidologie* 23:465-473.
- Cooper, R. J., K. M. Dodge, P. J. Marinat, S. B. Donahoe, and R. C. Whitmore. 1990. Effect of diflubenzuron application on eastern deciduous forest birds. *J. Wildl. Mgmt.* 54:486-493.
- Cordova, D., E. Benner, M. D. Sacher, J. J. Rauh, J. S. Sopa, G. Lahm, T. Selby, T. Stevenson, L. Flexner, S. Gutteridge, D. F. Rhoades, L. Wu, R. M. Smith, and Y. Tao. 2006. Anthranilic diamides: a new class of insecticides with a novel mode of action, ryanodine receptor activation. *Pesticide Biochemistry and Physiology* 84:196-214.
- Deakle, J. P. and J. R. Bradley, Jr. 1982. Effects of early season applications of diflubenzuron and azinphosmethyl on populations levels of certain arthropods in cotton fields. *J. Georgia Entomol. Soc.* 17:189-200.
- Deneke, D. and J. Keyser. 2011. Integrated Pest Management Strategies for Grasshopper Management in South Dakota. South Dakota State University Extension.
- Dinkins, M. F., A. L. Zimmermann, J. A. Dechant, B. D. Parkins, D. H. Johnson, L. D. Igl, C. M. Goldade, and B. R. Euliss. 2002. Effects of Management Practices on Grassland Birds: Horned Lark Northern Prairie Wildlife Research Center. Page 34. Northern Prairie Wildlife Research Center, Jamestown, ND.

- Dobroski, C. J., E. J. O'Neill, J. M. Donohue, and W. H. Curley. 1985. Carbaryl: a profile of its behaviors in the environment. Roy F. Weston, Inc. and V.J. Ciccone and Assoc., Inc., West Chester, PA; Woodbridge, VA.
- Eisler, R. 1992. Diflubenzuron Hazards to Fish, Wildlife, and Invertebrate: A Synoptic Review. U.S. Department of Interior, Fish and Wildlife Service, Washington, D.C.
- Eisler, R., 2000. Handbook of chemical risk assessment: health hazards to humans, plants, and animals. Lewis Publishers, New York.
- El-Refai, A. and T. L. Hopkins. 1972. Malathion adsorption, translocation, and conversion to malaoxon in bean plants. *J. Assoc. Official Analytical Chemists* 55:526-531.
- Fischer, S. A. and L. W. Hall, Jr. 1992. Environmental concentrations and aquatic toxicity data on diflubenzuron (Dimilin). *Critical Rev. in Toxicol.* 22:45-79.
- Follett, R. F. and D. A. Reed. 2010. Soil carbon sequestration in grazing lands: societal benefits and policy implications. *Rangeland Ecology & Management* 63:4-15.
- Foster, R. N., K. C. Reuter, K. Fridley, D. Kurtenback, R. Flakus, R. Bohls, B. Radsick, J. B. Helbig, A. Wagner, and L. Jeck. 2000. Field and Economic Evaluation of Operational Scale Reduced Agent and Reduced Area Treatments (RAATs) for Management of Grasshoppers in South Dakota Rangeland. U.S. Department of Agriculture, Animal and Plant Health Inspection Service, Plant Protection and Quarantine, Phoenix, AZ.
- George, T. L., L. C. McEwen, and B. E. Peterson. 1995. Effects of grasshopper control programs on rangeland breeding bird populations. *J. Range Manage.* 48:336-342.
- Gramlich, F. J. 1979. Effects of Sevin on songbird cholinesterase. Environmental Monitoring of Cooperative Spruce Budworm Control Projects. Maine Department of Conservation, Bureau of Forestry, Augusta, ME.
- Guerrant, G. O., L. E. Fetzer, Jr., and J. W. Miles. 1970. Pesticide residues in Hale County, Texas, before and after ultra-low-volume aerial applications of Malathion. *Pesticide Monitoring J.* 4:14-20.
- Havstad, K. M., D. P. Peters, R. Skaggs, J. Brown, B. Bestelmeyer, E. Fredrickson, J. Herrick, and J. Wright. 2007. Ecological services to and from rangelands of the United States. *Ecological Economics* 64:261-268.
- Howe, F. P. 1993. Effects of Grasshopper Insecticide Application on Diet, Food Delivery Rates, Growth, and Survival of Shrubsteppe Passarine. Page 108 PhD dissertation. Colorado State University, Fort Collins, CO.
- Howe, F. P., R. L. Knight, L. C. McEwen, and T. L. George. 1996. Direct and indirect effects of insecticide applications on growth and survival of nestling passerines. *Ecol. Appl.* 6:1314-1324.
- Kar, A., K. Mandal, and B. Singh. 2012. Environmental fate of chlorantraniliprole residues on cauliflower using QuEChERS technique. *Environ. Monit. Assess* 85:1255-1263.
- Keever, D. W., J. R. Bradley, Jr, and M. C. Ganyard. 1977. Effects of diflubenzuron (Dimilin) on selected beneficial arthropods in cotton fields. *J. Econ. Entomol.* 6:832-836.
- LaFleur, K. S. 1979. Sorption of pesticides by model soils and agronomic soils: rates and equilibria. *Soil Sci.* 127:94-101.
- Larsen, J. and R. N. Foster. 1996. Using Hopper to Adapt Treatments and Costs to Needs and Resources. U.S. Department of Agriculture, Animal and Plant Health Inspection Service Grasshopper Integrated Pest Management User Handbook, Washington, D.C.

- Larson, J. L., C. T. Redmond, and D. A. Potter. 2012. Comparative impact of an anthranilic diamide and other insecticidal chemistries on beneficial invertebrates and ecosystem services in turfgrass. *Pest Management Science* 68:740-748.
- Latchininsky, A., G. Sword, M. Sergeev, M. Cigiliano, and M. Lecoq. 2011. Locusts and grasshoppers: behavior, ecology, and biogeography. *Psyche* 2011:1-4.
- Lockwood, J. A. and S. P. Schell. 1997. Decreasing economic and environmental costs through reduced area and agent insecticide treatments (RAATs) for the control of rangeland grasshoppers: empirical results and their implications for pest management. *J. Orthoptera Res.* 6:19-32.
- Lockwood, J., S. Schell, R. Foster, C. Reuter, and T. Rahadi. 2000. Reduced agent-area treatments (RAAT) for management of rangeland grasshoppers: efficacy and economics under operational conditions. *International Journal of Pest Management* 46:29-42.
- Lockwood, J. A. and A. Latchininsky. 2000. The Risks of Grasshoppers and Pest Management to Grassland Agroecosystems: An International Perspective on Human Well-Being and Environmental Health. Pages 193-215 in A. Latchininsky and M. Sergeev, editors. *Grasshoppers and Grassland Health*. Kluwer Academic Publishers.
- Lockwood, J., R. Anderson-Sprecher, and S. Schell. 2002. When less is more: optimization of reduced agent-area treatments (RAATs) for management of rangeland grasshoppers. *Crop Protection* 21:551-562.
- Matsumara, F. 1985. *Toxicology of insecticides*. Plenum Press, New York.
- McEwen, L.C., Althouse, C.M., and Peterson, B.E., 1996. Direct and indirect effects of grasshopper integrated pest management (GHIPM) chemicals and biologicals on nontarget animal life. *In* U.S. Department of Agriculture, Animal and Plant Health Inspection Service, 1996. *Grasshopper Integrated Pest Management User Handbook*, Tech. Bul. No. 1809. Sec. III.2. Washington, DC.
- Miles, C. J. and S. Takashima. 1991. Fate of malathion and O.O.S. trimethyl phosphorothioate byproduct in Hawaiian soil and water. *Arch. Environ. Contam. Toxicol* 20:325-329.
- Mommaerts, V., Sterk, G., and G. Smagghe. 2006. Hazards and uptake of chitin synthesis inhibitors in bumblebees *Bombus terrestris*. *Pest Mgt. Science* 62:752-758.
- Murphy, C. F., P. C. Jepson, and B. A. Croft. 1994. Database analysis of the toxicity of antilocus pesticides to non-target, beneficial invertebrates. *Crop Protection* 13:413-420.
- Muzzarelli, R. 1986. Chitin synthesis inhibitors: effects on insects and on nontarget organisms. *CRC Critical Review of Environmental Control* 16:141-146.
- Narisu, J., A. Lockwood, and S. P. Schell. 1999. A novel mark-capture technique and its application to monitoring the direction and distance of local movements of rangeland grasshoppers (Orthoptera: Acridade) in context of pest management. *J. Appl. Ecol.* 36:604-617.
- Narisu, J., A. Lockwood, and S. P. Schell. 2000. Rangeland grasshopper movement as a function of wind and topography: implications for pest movement. *J. Appl. Ecol.* 36:604-617.
- Nation, J.L., Robinson, F.A., Yu, S.J., and A.B. Bolten. 1986. Influence upon honeybees of chronic exposure to very low levels of selected insecticides in their diet. *J. Apic. Res.* 25:170-177.
- Neary, D. G. 1985. Fate of pesticides in Florida's forests: an overview of potential impacts of water quality. Pages 18-24 in *Procs. Soil and Crop Sci. Soc. of FL.*

- Nigg, H. N., R. D. Cannizzaro, and J. H. Stamper. 1986. Diflubenzuron surface residues in Florida citrus. *Bul. Environ. Contam. Toxicol.* 36:833-838.
- NIH. 2009a. Carbaryl, CASRN: 63-25-2. National Institutes of Health, U.S. National Library of Medicine, Toxnet, Hazardous Substances Database.
- NIH. 2009b. National Institutes of Health, U.S. National Library of Medicine, Hazardous Substances Database.
- Norelius, E. E. and J. A. Lockwood. 1999. The effects of reduced agent-area insecticide treatments for rangeland grasshopper (Orthoptera: Acrididae) control on bird densities. *Archives of Environmental Contamination and Toxicology* 37:519-528.
- Pascual, J. A. 1994. No effects of a forest spraying of malathion on breeding blue tits (*Parus caeruleus*). *Environ. Toxicol. Chem.* 13:1127-1131.
- Peach, M. P., D. G. Alston, and V. J. Tepedino. 1994. Bees and bran bait: is carbaryl bran bait lethal to alfalfa leafcutting bee (Hymenoptera: Megachilidae) adults or larvae? *J. Econ. Entomol.* 87:311-317.
- Peach, M. P., D. G. Alston, and V. J. Tepedino. 1995. Sublethal effects of carbaryl bran bait on nesting performance, parental investment, and offspring size and sex ratio of the alfalfa leafcutting bee (Hymenoptera: Megachilidae). *Environ. Entomol.* 24:34-39.
- Pfadt, R. E. 2002. *Field Guide to Common Western Grasshoppers*, Third Edition. Wyoming Agricultural Experiment Station Bulletin 912. Laramie, Wyoming.
- Purdue University. 2018. National Pesticide Information Retrieval System. West Lafayette, IN.
- Quinn, M. A., R. L. Kepner, D. D. Walgenbach, R. N. Foster, R. A. Bohls, P. D. Pooler, K. C. Reuter, and J. L. Swain. 1991. Effect of habitat and perturbation on populations and community structure of darkling beetles (Coleoptera: tenebrionidae) on mixed grass rangeland. *Environ. Entomol.* 19:1746-1755.
- Rashford, B. S., A. V. Latchininsky, and J. P. Ritten. 2012. *An Economic Analysis of the Comprehensive Uses of Western Rangelands*. U.S. Department of Agriculture, Animal and Plant Health Inspection Service.
- Reinhardt, T. and R. Ottmar. 2004. Baseline measurements of smoke exposure among wildland firefighters. *Journal of Occupational and Environmental Hygiene* 1:593-606.
- Reisen, F. and S. Brown. 2009. Australian firefighters' exposure to air toxics during bushfire burns of autumn 2005 and 2006. *Environment International* 35:342-353.
- Richmond, M. L., C. J. Henny, R. L. Floyd, R. W. Mannan, D. W. Finch, and L. R. DeWeese. 1979. Effects of Sevin 4-oil, Dimilin, and Orthene on Forest Birds in Northeastern Oregon. USDA, Pacific SW Forest and Range Experiment Station.
- Rosenberg, K. V., R. D. Ohmart, and B. W. Anderson. 1982. Community organization of riparian breeding birds: response to an annual resource peak. *The Auk* 99:260-274.
- Sample, B. E., R. J. Cooper, and R. C. Whitmore. 1993. Dietary shifts among songbirds from a diflubenzuron-treated forest. *The Condor* 95:616-624.
- Schaefer, C. H., A. E. Colwell, and E. F. Dupras, Jr. 1980. The occurrence of p-chloroaniline and p-c hlorophenylurea from the degradation of pesticide in water and fish. *Proceedings of the 48th Ann. Meeting Mosquito Vector Cont. Assoc.*:84-89.
- Schaefer, C. H. and E. F. Dupras, Jr. 1977. Residues of diflubenzuron [1-(4-chlorophenyl)-3(2,6-difluorobenzoyl) urea] in pasture soil, vegetation, and water following aerial applications. *J. Agric. Food Chem.* 25:1026-1030.

- Smith, D. and J. Lockwood. 2003. Horizontal and trophic transfer of diflubenzuron and fipronil among grasshoppers and between grasshoppers and darkling beetles (Tenebrionidae). *Archives of Environmental Contamination and Toxicology* 44:377-382.
- Smith, D. I., J. A. Lockwood, A. V. Latchininsky, and D. E. Legg. 2006. Changes in non-target populations following applications of liquid bait formulations of insecticides for control of rangeland grasshoppers. *Internat. J. Pest Mgt.* 52:125-139.
- Stanley, J. G. and J. G. Trial. 1980. Disappearance constants of carbaryl from streams contaminated by forest spraying. *Bul. Environ. Contam. Toxicol.* 25:771-776.
- Swain, J. L. 1986. Effect of Chemical Grasshopper Controls on Non-Target Arthropods of Rangeland in Chaves County, New Mexico. New Mexico State University.
- Tepedino, V. J. 1979. The importance of bees and other insect planetaries in maintaining floral species composition. *Great Basin Naturalist Memoirs* 3:139-150.
- Thompson, H.M, Wilkins, S. Battersby, A.H., Waite, R.J., and D. Wilkinson. 2005. The effects of four insect growth-regulating (IGR) insecticides on honeybee (*Apis mellifera* L.) colony development, queen rearing and drone sperm production. *Ecotoxicology* 14:757-769.
- Thomson, D. L. K. and W. M. J. Strachan. 1981. Biodegradation of carbaryl in simulated aquatic environment. *Bul. Environ. Contam. Toxicol.* 27:412-417.
- USDA APHIS— see U.S. Department of Agriculture, Animal and Plant Health Inspection Service
- U.S. Department of Agriculture, Animal and Plant Health Inspection Service, 1999. APHIS Directive 5600.3, Evaluating APHIS programs and activities for ensuring protection of children from environmental health risks and safety risks. September 3, 1999. U.S. Department of Agriculture, Animal and Plant Health Inspection Service, Riverdale, MD. [online] available: <http://www.aphis.usda.gov/library/directives>.
- U.S. Department of Agriculture, Animal and Plant Health Inspection Service. 2015. Biological Assessment for the APHIS Rangeland Grasshopper and Mormon Cricket Suppression Program. Page 162. U.S. Department of Agriculture, Animal and Plant Health Inspection Service.
- U.S. Department of Agriculture, Animal and Plant Health Inspection Service. 2018a. Human Health and Ecological Risk Assessment for Carbaryl Rangeland Grasshopper and Mormon Cricket Suppression Applications. United States Department of Agriculture, Animal and Plant Health Inspection Service.
- U.S. Department of Agriculture, Animal and Plant Health Inspection Service. 2018c. Human Health and Ecological Risk Assessment for Diflubenzuron Rangeland Grasshopper and Mormon Cricket Suppression Applications. United States Department of Agriculture, Animal and Plant Health Inspection Service.
- U.S. Department of Agriculture, Animal and Plant Health Inspection Service. 2018d. Human Health and Ecological Risk Assessment for Malathion Rangeland Grasshopper and Mormon Cricket Suppression Applications. United States Department of Agriculture, Animal and Plant Health Inspection Service.
- U.S. Department of Agriculture, Animal and Plant Health Inspection Service. 2019. Rangeland Grasshopper and Mormon Cricket Suppression Program Final Environmental Impact Statement. United States Department of Agriculture, Animal and Plant Health Inspection Service.

- USDA FS. 2004. Control/eradication agents for the gypsy moth—human health and ecological risk assessment for diflubenzuron (final report). United States Department of Agriculture, Forest Service
- USDA FS. 2008. Malathion- Human Health and Ecological Risk Assessment. U.S. Department of Agriculture, Forest Service.
- USEPA – See U.S. Environmental Protection Agency
- U.S. Environmental Protection Agency. 1997. Reregistration Eligibility Decision (RED): Diflubenzuron. U.S. Environmental Protection Agency.
- U.S. Environmental Protection Agency. 2000a. Malathion Reregistration Eligibility Document Environmental Fate and Effects. Page 146. U.S. Environmental Protection Agency, Office of Prevention, Pesticides and Toxic Substances.
- U.S. Environmental Protection Agency. 2000b. Reregistration Eligibility Decision (RED) for Malathion. U.S. Environmental Protection Agency.
- USEPA. 2003. Environmental Fate and Ecological Risk Assessment for Re-Registration of Carbaryl. U.S. Environmental Protection Agency.
- U.S. Environmental Protection Agency. 2006. Malathion Reregistration Eligibility Document. Page 147. U.S. Environmental Protection Agency, Office of Pesticide Programs.
- U.S. Environmental Protection Agency. 2007. Reregistration Eligibility Decision (RED) for Carbaryl. Page 47. U.S. Environmental Protection Agency, Prevention, Pesticides and Toxic Substances.
- U.S. Environmental Protection Agency. 2012a. Fyfanon ULV AG. U.S. Environmental Protection Agency.
- U.S. Environmental Protection Agency. 2012c. Sevin XLR Plus Label. Pages 1-40 Pesticide Product and Label System. U.S. Environmental Protection Agency.
- U.S. Environmental Protection Agency. 2015a. Annual Cancer Report 2015, Chemicals Evaluated for Carcinogenic Potential Page 34. U.S. Environmental Protection Agency, Office of Pesticide Programs.
- U.S. Environmental Protection Agency. 2015b. Memorandum - Diflubenzuron: human health risk assessment for an amended Section 3 registration for carrot, peach subgroup 12-12B, plum subgroup 12-12C, pepper/eggplant subgroup 8010B, cottonseed subgroup 20C, alfalfa (regional restrictions) and R175 Crop Group Conversion for tree nut group 14-12. Page 71 U.S. Environmental Protection Agency, Office of Pesticide Programs.
- U.S. Environmental Protection Agency. 2016a. Appendix 3-1: Environmental transport and fate data analysis for malathion. In: Biological Evaluation Chapters for Malathion ESA Assessment.
- U.S. Environmental Protection Agency. 2016b. Chapter 2: Malathion Effects Characterization for ESA Assessment. In: Biological Evaluation Chapters for Malathion ESA Assessment.
- U.S. Environmental Protection Agency. 2016c. Malathion: Human Health Draft Risk Assessment for Registration Review. Page 258. U.S. Environmental Protection Agency.
- U.S. Environmental Protection Agency. 2017a. Memorandum - Carbaryl: Draft Human Health Risk Assessment in Support of Registration Review. Page 113 U.S. Environmental Protection Agency.
- U.S. Environmental Protection Agency. 2017b. Prevathon Label. U.S. Environmental Protection Agency.

- U.S. Environmental Protection Agency. 2018. Preliminary Risk Assessment to Support the Registration Review of Diflubenzuron.
- USFWS. 2007. National Bald Eagle Management Guidelines. Page 23 pp. U.S. Fish and Wildlife Service.
- Wakeland, C. and W. E. Shull. 1936. The Mormon cricket with suggestions for its control, Extension Bulletin No. 100. University of Idaho, College of Agriculture, Idaho Agricultural Extension.
- Zinkl, J. G., C. J. Henny, and L. R. DeWeese. 1977. Brain cholinesterase activities of birds from forests sprayed with trichlorfon (Dylox) and carbaryl (Sevin 4-oil). *Bul. Environ. Contam. Toxicol.* 17:379-386.

VI. Listing of Agencies and Persons Consulted

John Nystedt, Fish and Wildlife Biologist, U.S. Fish and Wildlife Service. Flagstaff Suboffice
323 N. Leroux Street, Suite 201
Flagstaff, Arizona 86001

Brenda Smith, Fish and Wildlife Biologist, U.S. Fish and Wildlife Service Flagstaff Suboffice
323 N. Leroux Street, Suite 201
Flagstaff, Arizona 86001

Shaula Hedwall, Fish and Wildlife Biologist, U.S. Fish and Wildlife Service Flagstaff Suboffice
323 N. Leroux Street, Suite 201
Flagstaff, Arizona 86001

Brian J. Wooldridge, Fish and Wildlife Biologist, U. S. Fish and Wildlife Service Flagstaff Suboffice
323 N. Leroux St., Suite 201
Flagstaff, AZ 86001

Terry Rambler, Tribal Chairman, San Carlos Apache Tribe
P.O. Box 0
#3 San Carlos Avenue
San Carlos, Arizona 85550

Clark Richins, Land Operation Coordinator, San Carlos Agency, Bureau of Indian Affairs #3 San Carlos Avenue
San Carlos, Arizona 85550

Teresa Goseyun, Rangeland Management Specialist, San Carlos Apache Tribe
#3 San Carlos Avenue
San Carlos, Arizona 85550

Daniel Juan, Biologist, San Carlos Apache Tribe PO Box 97
San Carlos, Arizona 85550

Paul Buck, Soil Scientist, San Carlos Apache Tribe PO Box 0
San Carlos, Arizona 85550

Michael J. Herder, Arizona Strip District Manager, BLM
345 East Riverside Drive
St. George, Utah 84790

Ben Otte, Rangeland Management Specialist, Arizona Strip District Manager, BLM
345 East Riverside Drive
St. George, Utah 84790

Justin Reeves, Rangeland Management Specialist, Arizona Strip District Manager, BLM
345 East Riverside Drive
St. George, Utah 84790

Travis Largent, Wildlife Biologist
Kaibab National Forest, Williams and Tusayan Ranger Districts
742 S Clover Rd.
Williams, AZ 86046

Kelly Hetzler, Forester, Inventory and GIS, San
Carlos Apache Tribe PO Box 0
San Carlos, Arizona 85550

Victoria Wesley, San Carlos
Apache Tribe, Forestry PO Box 0
San Carlos, Arizona 85550

Daniel Juan, Wildlife Technician, San
Carlos Apache Tribe PO Box 0
San Carlos, Arizona 85550

Jeff McFadden, Wildlife Technician, San
Carlos Apache Tribe PO Box 0
San Carlos, Arizona 85550

Sabrina Tuttle, Ph.D., Assistant Extension Agent, Gila County
Cooperative Extension, U of A, San Carlos Apache Reservation
PO Box 850
San Carlos, Arizona 85550

Jennifer Cordova, Wildlife Specialist II,
Arizona Game and Fish Dept.
PO Box 397
Seligman, AZ 86337

**Appendix A - APHIS Rangeland Grasshopper and Mormon Cricket
Suppression Program
FY-2021 Treatment Guidelines
Version DD/MM/YYYY**

The objectives of the APHIS Rangeland Grasshopper and Mormon Cricket Suppression Program are to 1) conduct surveys in the Western States; 2) provide technical assistance to land managers and private landowners; and 3) when funds permit, suppress economically damaging grasshopper and Mormon cricket outbreaks on Federal, Tribal, State, and/or private rangeland. The Plant Protection Act of 2000 provides APHIS the authority to take these actions.

General Guidelines for Grasshopper / Mormon Cricket Treatments

- 1) All treatments must be in accordance with:
 - a) the Plant Protection Act of 2000;
 - b) applicable environmental laws and policies such as: the National Environmental Policy Act, the Endangered Species Act, the Federal Insecticide, Fungicide, and Rodenticide Act, and the Clean Water Act (including National Pollutant Discharge Elimination System requirements – if applicable);
 - c) applicable state laws;
 - d) APHIS Directives pertaining to the proposed action;
 - e) Memoranda of Understanding with other Federal agencies.
- 2) Subject to the availability of funds, upon request of the administering agency, the agriculture department of an affected State, or private landowners, APHIS, to protect rangeland, shall immediately treat Federal, Tribal, State, or private lands that are infested with grasshoppers or Mormon crickets at levels of economic infestation, unless APHIS determines that delaying treatment will not cause greater economic damage to adjacent owners of rangeland. In carrying out this section, APHIS shall work in conjunction with other Federal, State, Tribal, and private prevention, control, or suppression efforts to protect rangeland.
- 3) Prior to the treatment season, conduct meetings or provide guidance that allows for public participation in the decision making process. In addition, notify Federal, State and Tribal land managers and private landowners of the potential for grasshopper and Mormon cricket outbreaks on their lands. Request that the land manager / land owner advise APHIS of any sensitive sites that may exist in the proposed treatment areas.
- 4) Consultation with local Tribal representatives will take place prior to treatment programs to fully inform the Tribes of possible actions APHIS may take on Tribal lands.
- 5) On APHIS run suppression programs, the Federal government will bear the cost of treatment up to 100 percent on Federal and Tribal Trust land, 50 percent of the cost on State land, and 33 percent of cost on private land. There is an additional 16.15% charge, however, on any funds received by APHIS for federal involvement with suppression treatments.
- 6) Land managers are responsible for the overall management of rangeland under their control to prevent or reduce the severity of grasshopper and Mormon cricket outbreaks. Land

managers are encouraged to have implemented integrated pest management systems prior to requesting a treatment. In the absence of available funding or in the place of APHIS funding, the Federal land management agency, Tribal authority or other party/ies may opt to reimburse APHIS for suppression treatments. Interagency agreements or reimbursement agreements must be completed prior to the start of treatments which will be charged thereto.

- 7) There are situations where APHIS may be requested to treat rangeland that also includes small areas where crops are being grown (typically less than 10 percent of the treatment area). In those situations, the crop owner pays the entire treatment costs on the croplands.

NOTE: The insecticide being considered must be labeled for the included crop as well as rangeland and current Worker Protection Standards must be followed by the applicator and private landowner.

- 8) In some cases, rangeland treatments may be conducted by other federal agencies (e.g., Forest Service, Bureau of Land Management, or Bureau of Indian Affairs) or by non-federal entities (e.g., Grazing Association or County Pest District). APHIS may choose to assist these groups in a variety of ways, such as:
 - a) loaning equipment (an agreement may be required);
 - b) contributing in-kind services such as surveys to determine insect species, instars, and infestation levels;
 - c) monitoring for effectiveness of the treatment;
 - d) providing technical guidance.
- 9) In areas considered for treatment, State-registered beekeepers and organic producers shall be notified in advance of proposed treatments. If necessary, non-treated buffer zones can be established.

Operational Procedures

GENERAL PROCEDURES FOR ALL AERIAL AND GROUND APPLICATIONS

- 1) Follow all applicable Federal, Tribal, State and local laws and regulations in conducting grasshopper and Mormon cricket suppression treatments.
- 2) Notify residents within treatment areas, or their designated representatives, prior to proposed operations. Advise them of the control method to be used, proposed method of application, and precautions to be taken.
- 3) One of the following insecticides that are labeled for rangeland use can be used for a suppression treatment of grasshoppers and Mormon crickets:
 - a) Carbaryl
 - i) solid bait
 - ii) ultra-low volume (ULV) spray
 - b) Diflubenzuron ULV spray
 - c) Malathion ULV spray
- 4) Do not apply insecticides directly to water bodies (defined herein as reservoirs, lakes, ponds, pools left by seasonal streams, springs, wetlands, and perennial streams and rivers).

Furthermore, provide the following buffers for water bodies:

- 500-foot buffer with aerial liquid insecticide.
 - 200-foot buffer with ground liquid insecticide.
 - 200-foot buffer with aerial bait.
 - 50-foot buffer with ground bait.
- 5) Instruct program personnel in the safe use of equipment, materials and procedures; supervise to ensure safety procedures are properly followed.
 - 6) Conduct mixing, loading, and unloading in an approved area where an accidental spill would not contaminate a water body.
 - 7) Each aerial suppression program will have a Contracting Officer's Representative (COR) OR a Treatment Manager on site. Each State will have at least one COR available to assist the Contracting Officer (CO) in GH/MC aerial suppression programs.

NOTE: A Treatment Manager is an individual that the COR has delegated authority to oversee the actual suppression treatment; someone who is on the treatment site and overseeing / coordinating the treatment and communicating with the COR. No specific training is required, but knowledge of the Aerial Application Manual and treatment experience is critical; attendance to the Aerial Applicators Workshop is very beneficial.

- 8) Each suppression program will conduct environmental monitoring as outlined in the current year's Environmental Monitoring Plan.

APHIS will assess and monitor rangeland treatments for the efficacy of the treatment, to verify that a suppression treatment program has properly been implemented, and to assure that any environmentally sensitive sites are protected.


- 9) APHIS reporting requirements associated with grasshopper / Mormon cricket suppression treatments can be found in the APHIS Grasshopper Program Guidebook:
http://www.aphis.usda.gov/import_export/plants/manuals/domestic/downloads/grasshopper.pdf

SPECIFIC PROCEDURES FOR AERIAL APPLICATIONS


- 1) APHIS Aerial treatment contracts will adhere to the current year's Statement of Work (SOW).
- 2) Minimize the potential for drift and volatilization by not using ULV sprays when the following conditions exist in the spray area:
 - a) Wind velocity exceeds 10 miles per hour (unless state law requires lower wind speed);
 - b) Rain is falling or is imminent;
 - c) Dew is present over large areas within the treatment block;
 - d) There is air turbulence that could affect the spray deposition;

- e) Temperature inversions (ground temperature higher than air temperature) develop and deposition onto the ground is affected.
- 3) Weather conditions will be monitored and documented during application and treatment will be suspended when conditions could jeopardize the correct spray placement or pilot safety.
- 4) Application aircraft will fly at a median altitude of 1 to 1.5 times the wingspan of the aircraft whenever possible or as specified by the COR or the Treatment Manager.
- 5) Whenever possible, plan aerial ferrying and turnaround routes to avoid flights over congested areas, water bodies, and other sensitive areas that are not to be treated.

Appendix B: Grasshopper Hazard Map of the Affected Environment



Appendix C: Map of Proposed Action Areas



Appendix D: Map of the San Carlos Apache Tribal Proposed Action Area



Appendix E: FWS/NMFS Correspondence (PENDING)

Appendix F: Public Comments and APHIS Response