



Pale Cyst Nematode (*Globodera pallida*) Eradication Program- Idaho Falls, Idaho

January 2010 Report

Background

Pale cyst nematodes (PCN), *Globodera G. pallida*, are soil-borne organisms that do not infest potato tubers. The pests infest feeder roots, where the females attach, feed, and become sedentary. Nematodes reproduce sexually. Females form cysts containing 200 to 600 eggs, which can stay dormant for up to 30 years while the eggs inside remain viable. On host plants, large numbers of PCN can cause wilting, stunted growth, poor root development, and early plant death. If left uncontrolled, PCN can reduce yields up to 80 percent in potato fields. Even with only minor symptoms showing on the foliage, PCN can significantly reduce tuber size. PCN spread primarily by the transport of cysts in soil. This may occur with the movement of soil on farming, construction, and other equipment; infested soil adhering to seed potatoes and other regulated crops; and any other items or means of transport such as water.

On April 19, 2006, officials of USDA's Animal and Plant Health Inspection Service (APHIS) and the Idaho State Department of Agriculture (ISDA) announced the detection of PCN, a major pest of potato crops. This was the first detection of the pest in the United States. The nematode cysts were detected during a routine survey of tare soil at an ISDA grading facility in eastern Idaho. Subsequent 2006 surveying to determine the possible origin and distribution of the pest in Idaho confirmed seven PCN-infested fields totaling 911 acres, all within a one mile radius in Bingham and Bonneville Counties, Idaho. The PCN-infested fields and an area surrounding the fields were placed under a Federal Domestic Quarantine Order and parallel State Rule in August 2006, establishing restrictions on movement of certain regulated articles from Idaho in order to prevent the spread of PCN.

As a result of continued intensive soil sampling in 2007 and 2008, an additional two PCN-infested fields were found in Bingham County, Idaho. The nine PCN-infested fields all continue to be within a one mile radius and the fields associated with them through shared tenancy, farming practices, equipment, and/or shared borders have been extensively surveyed and regulated. Since program inception, a total of 30,753 acres have been regulated due to their infestation or association with an infested field. Non-infested, associated fields have been eligible for federal deregulation following a sequence of soil surveys with no PCN detections. To date 29,048 acres have been released from federal regulation. Currently, 1,635 acres remain regulated, of which 1,100 are PCN-infested.

Eradication treatments of PCN-infested fields have been ongoing since the spring of 2007. Eradication treatments have included methyl bromide fumigation, Telone II fumigation, and biofumigant plantings. Testing of the top 3 inches of soil in infested fields indicate the average viability of eggs within the PCN cysts have declined by more than 90% since eradication treatments began.

A description of the current PCN regulated area can be found at:

http://www.aphis.usda.gov/plant_health/plant_pest_info/potato/pcn-maps.shtml.

The current Federal PCN rule, within docket # APHIS-2006-0143, is available at:

<http://regulations.gov>.

Survey Information

Type of survey	Idaho soil samples collected	
	Jan 2010	Since program inception
Detection	0	82,106
Delimiting	0	127,380
Eradication	0	48,264
Total	0	257,750

There were no soil survey activities in Idaho during January, 2010. Survey activities are set to resume when weather permits, likely around the middle of March.

Identification and Diagnostics

Type of survey	Samples processed by the Idaho PCN Laboratory			
	Jan 2010	Since program inception	Jan 2010 Results	Results since program inception
Detection	3,682	14,125	Negative	Negative
Delimiting	1,833	115,713	Negative	Negative ¹
Eradication	0	46,314	N/A	N/A
National	1,972	18,548	Negative	Negative
Total	7,487	194,700		

¹Except for samples confirmed for the nine infested fields

Type of survey	Samples processed by the Idaho Food Quality Assurance Laboratory	
	Since program inception	Results
Detection	35,524	Negative
Delimiting	31,097	Negative ²
Total	66,621	

²Except for samples confirmed for the nine infested fields

Program Research

None to Report

Eradication Activities

There were no eradication activities in Idaho during January, 2010. Methyl bromide application to the PCN-infested fields is scheduled to occur in spring, 2010. Methyl bromide was applied to the PCN-infested fields in the spring of 2007, 2008, and 2009.

Telone II application to the PCN-infested fields is scheduled for late summer, 2010. Telone II was applied in the late summer of 2007 and 2008. There was no Telone II application in 2009 due to a world-wide shortage of this chemical.

Biofumigants with nematicidal activity were planted in the infested fields in the summers of 2007 (oil radish) and 2009 (arugula).

Regulatory Actions

The PCN-regulated area was amended twice in January 2010. On January 8th, 1,562 acres were deregulated, and on January 15th, an additional 622 acres were deregulated. To date 29,048 acres have been released from federal regulation.

Regulatory Treatments

In January, two snow machines that entered PCN-infested fields were washed and steam sanitized. Since program inception, more than 7,600 pieces of equipment have been sanitized.

Regulatory Documentation

In January 2010, two certificates (PPQ form 540) and one limited permit (PPQ form 530) were issued to document the movement and treatment of snow machines that entered PCN-infested fields. Since program inception, more than 6,100 certificates and 1,100 limited permits have been issued for the sanitation and movement of regulated articles.

One compliance agreement was issued to an external stakeholder in January 2010. To date 129 compliance agreements have been issued.

Impacts on Commerce

In response to the initial PCN detection in 2006, Canada, Mexico and Korea shut off importation of potatoes from Idaho, while Japan cut off importation of potatoes from the entire U.S. The Mexican and Canadian export markets have both been re-opened with the exception of potatoes from PCN-regulated areas. Both require PCN soil surveys from origin fields. The Korean and Japanese markets remain closed to Idaho potatoes but negotiations are actively underway to re-gain market access. Because of extensive field surveys conducted throughout production areas in Idaho, all of which have been negative beyond the nine infested fields, the general opinion by our trading partners is that potatoes produced outside regulated areas do not pose the biological risk for introduction of PCN.

Communication and Outreach

In January, a meeting with infested field landowners and operators was held to discuss calendar year 2010 eradication planning. Topics discussed included methyl bromide fumigation logistics, summer biosecurity cover planting, and bioassay options in PCN-infested fields. The next eradication planning meeting with landowners and operators is scheduled for mid-March.

In January a PCN stakeholder update was issued by Legislative and Public Affairs. Stakeholder updates are available at:

http://www.aphis.usda.gov/plant_health/plant_pest_info/potato/pcn_stakeholder.shtml