ENVIRONMENTAL ASSESSMENT

MAMMAL DAMAGE MANAGEMENT IN TENNESSEE

UNITED STATES DEPARTMENT OF AGRICULTURE (USDA)
ANIMAL AND PLANT HEALTH INSPECTION SERVICE (APHIS)
WILDLIFE SERVICES (WS)

In cooperation with:

TENNESSEE VALLEY AUTHORITY (TVA)

September 2012

TABLE OF CONTENTS

ACRONYMS ..... *iii*

CHAPTER 1: PURPOSE AND NEED FOR ACTION
1.1 PURPOSE 1
1.2 NEED FOR ACTION 2
1.3 SCOPE OF THIS ENVIRONMENTAL ASSESSMENT 22
1.4 RELATIONSHIP OF THIS EA TO OTHER ENVIRONMENTAL DOCUMENTS 25
1.5 AUTHORITY OF FEDERAL AND STATE AGENCIES 26
1.6 COMPLIANCE WITH LAWS AND STATUTES 27
1.7 DECISIONS TO BE MADE 31

CHAPTER 2: AFFECTED ENVIRONMENT AND ISSUES
2.1 AFFECTED ENVIRONMENT 32
2.2 ISSUES ASSOCIATED WITH MAMMAL DAMAGE MANAGEMENT ACTIVITIES 34
2.3 ISSUES CONSIDERED BUT NOT IN DETAIL WITH RATIONALE 41

CHAPTER 3: ALTERNATIVES
3.1 DESCRIPTION OF THE ALTERNATIVES 48
3.2 ALTERNATIVES CONSIDERED BUT NOT ANALYZED IN DETAIL 56
3.3 STANDARD OPERATING PROCEDURES FOR MAMMAL DAMAGE MANAGEMENT .. 60
3.4 ADDITIONAL STANDARD OPERATING PROCEDURES SPECIFIC TO THE ISSUES 61

CHAPTER 4: ENVIRONMENTAL CONSEQUENCES
4.1 ENVIRONMENTAL CONSEQUENCES FOR ISSUES ANALYZED IN DETAIL 64
4.2 CUMULATIVE IMPACTS OF THE PROPOSED ACTION BY ISSUE 158

CHAPTER 5: LIST OF PREPARERS AND PERSONS CONSULTED
5.1 LIST OF PREPARERS 165
5.2 LIST OF PERSONS CONSULTED 165

LIST OF APPENDICES:

APPENDIX A – LITERATURE CITED A-1
APPENDIX B – METHODS AVAILABLE FOR RESOLVING OR PREVENTING MAMMAL DAMAGE IN TENNESSEE B-1
APPENDIX C – FEDERAL LISTED THREATENED AND ENDANGERED SPECIES C-1
APPENDIX D – STATE LISTED THREATENED AND ENDANGERED SPECIES D-1
APPENDIX E – CRITERIA FOR BEAVER DAM BREACHING/REMOVAL E-1
APPENDIX F – CRITERIA FOR BEAVER DAM BREACHING/REMOVAL WHERE THREATENED OR ENDANGERED SPECIES MAY BE PRESENT F-1

ACRONYMS

ABC	American Bird Conservancy
AMDUCA	Animal Medicinal Drug Use Clarification Act
APHIS	Animal and Plant Health Inspection Service
AVMA	American Veterinary Medical Association
CDC	Centers for Disease Control and Prevention
CEQ	Council on Environmental Quality
CFR	Code of Federal Regulations
CO_2	Carbon Dioxide
CWA	Clean Water Act
CWCS	Comprehensive Wildlife Conservation Strategy
DEA	Drug Enforcement Administration
EA	Environmental Assessment
EIS	Environmental Impact Statement
EPA	U.S. Environmental Protection Agency
ESA	Endangered Species Act
FAA	Federal Aviation Administration
FAR	Federal Aviation Regulation
FDA	Food and Drug Administration
FEIS	Final Environmental Impact Statement
FIFRA	Federal Insecticide, Fungicide, and Rodenticide Act
FLIR	Forward Looking Infrared
FR	Federal Register
FY	Fiscal Year
GAT	Georgia Alabama Tennessee (Oral Rabies Vaccine Bait Zone)
GnRH	Gonadotropin-releasing Hormone
GSMNP	Great Smoky Mountains National Park
IV	Intravenous
IC	Intracardiac
MOU	Memorandum of Understanding
MRLC	Multi-Resolution Land Characteristics
NASS	National Agricultural Statistics Service
NEPA	National Environmental Policy Act
NHPA	National Historic Preservation Act
NLCD	National Land Cover Dataset
NPS	National Park Service
NRCS	Natural Resources Conservation Service
NRP	Natural Resource Plan
NWP	Nationwide Permit
NWRC	National Wildlife Research Center
ORV	Oral Rabies Vaccination
PEP	Post - Exposure Prophylaxis
PL	Public Law
SOP	Standard Operating Procedure
T&E	Threatened and Endangered
TCA	Tennessee Code Annotated
TDA	Tennessee Department of Agriculture
TDEC	Tennessee Department of Environment and Conservation
TDH	Tennessee Department of Health
TNHP	Tennessee Natural Heritage Program
TNR	Trap, Neuter, Release Program

ACRONYMS (continued)

TVA	Tennessee Valley Authority
TWRA	Tennessee Wildlife Resources Agency
USC	United States Code
US	United States
USDA	U.S. Department of Agriculture
USDI	U.S. Department of the Interior
USFS	U.S. Forest Service
USFWS	U.S. Fish and Wildlife Services
USGS	U.S. Geological Survey
WS	Wildlife Services

CHAPTER 1: PURPOSE AND NEED FOR ACTION

1.1 PURPOSE

The United States Department of Agriculture (USDA), Animal and Plant Health Inspection Service (APHIS), Wildlife Services (WS)[1] program in Tennessee continues to receive requests for assistance or anticipates receiving requests for assistance to resolve or prevent damage occurring to agricultural resources, natural resources, and property, including threats to human safety, associated with beaver (*Castor canadensis*), muskrat (*Ondatra zibethicus*), nutria (*Myocastor coypus*), woodchuck (*Marmota monax*), gray squirrel (*Sciurus carolinensus*), fox squirrel (*Sciurus niger*), eastern chipmunk (*Tamia striatus*), pine vole (*Microtus pinetorum*), prairie vole (*Microtus ochrogaster*), meadow vole (*Microtus pennsylvanicus*), house mouse (*Mus musculus*), roof rat (*Rattus rattus*), Norway rat (*Rattus norvegicus*), eastern harvest mouse (*Reithrodontomys humulis*), deer mouse (*Peromyscus maniculatus*), white-footed mouse (*Peromyscus leucopus*), cotton mouse (*Peromyscus gossypinus*), rice rat (*Oryzomys palustris*), hispid cotton rat (*Sigmodon hispidus*), cottontail rabbit (*Sylvilagus floridanus*), black bear (*Ursus americanus*), raccoon (*Procyon lotor*), river otter (*Lutra canadensis*), long-tailed weasel *(Mustela frenata*), mink (*Mustela vison*), striped skunk (*Mephitis mephitis*), spotted skunk (*Spilogale putoris*), coyote (*Canis latrans*), gray fox (*Urocyon cinereoargenteus*), red fox (*Vulpes vulpes*), bobcat (*Felis rufus*), feral cat (*Felis domesticus*), feral dog (*Canis familiaris*), nine-banded armadillo (*Dasypus novemcinctus*), Virginia opossum (*Didelphis virginiana*), little brown myotis (*Myotis lucifugus*), silver-haired bat (*Lasionycteris noctivagans*), eastern pipistrelle (*Pipistrellus subflavus*), big brown bat (*Eptesicus fuscus*), evening bat (*Nycticeius humeralis*), eastern big-eared bat (*Plecotus rafinesquei*), eastern mole (*Scalopus aquaticus*), hairy-tailed mole (*Parascalops breweri*), star-nosed mole (*Condylura cristata*), feral swine (*Sus scrofa*), white-tailed deer (*Odocoileus virginianus*), and elk (*Cervus elaphus*). Normally, individual wildlife damage management projects conducted by the WS program could be categorically excluded from further analysis under the National Environmental Policy Act (NEPA), in accordance with APHIS implementing regulations for the NEPA (7 CFR 372.5(c), 60 FR 6000-6003). The Tennessee Valley Authority (TVA) also continues to experience damage and threats of damage associated with mammals at facilities or properties they own or manage in Tennessee. The TVA could request the assistance of WS to manage damage or threats of damage at those facilities and properties.

The purpose of this Environmental Assessment (EA) is to evaluate cumulatively the individual projects conducted by WS to manage damage and threats to agricultural resources, property, natural resources, and threats to humans caused by those mammal species identified previously. This EA will assist in determining if the proposed cumulative management of mammal damage could have a significant impact on the environment based on previous activities conducted by WS and based on the anticipation of receiving additional requests for assistance. Because the goal of WS is to conduct a coordinated program to alleviate mammal damage in accordance with plans, goals, and objectives developed to reduce damage, and because the program's goals and directives[2] are to provide services when requested, within the constraints of available funding and workforce, it is conceivable that additional damage management efforts could occur. Thus, this EA anticipates those additional efforts and the analyses would be intended to apply to actions that may occur in any locale and at any time within Tennessee as part of a coordinated program. This EA analyzes the potential effects of mammal damage management when requested, as coordinated between WS, TVA, and the Tennessee Wildlife Resources Agency (TWRA).

WS and the TVA are preparing this EA to: 1) facilitate planning, 2) promote interagency coordination, 3) streamline program management, 4) clearly communicate to the public the analysis of individual and

[1]The WS program is authorized to protect agriculture and other resources from damage caused by wildlife through the Act of March 2, 1931 (46 Stat. 1468; 7 USC 426-426b) as amended, and the Act of December 22, 1987 (101 Stat. 1329-331, 7 USC 426c).
[2]At the time of preparation, WS' Directives could be found at the following web address: http://www.aphis.usda.gov/wildlife_damage/ws_directives.shtml.

cumulative impacts of proposed activities; and 5) evaluate and determine if there would be any potentially significant or cumulative adverse effects from the proposed program. The analyses contained in this EA are based on information derived from WS' Management Information System, published documents (see Appendix A), interagency consultations, public involvement, and the EA previously prepared by WS and the TVA to address mammal damage in the State (USDA 2005*a*).

The EA evaluates the need for action to manage damage associated with mammals in the State, the potential issues associated with mammal damage management, and the environmental consequences of conducting different alternatives to meet the need for action while addressing the identified issues. The issues and alternatives associated with mammal damage management were initially developed by WS in cooperation with the TVA, and in consultation with the TWRA. The TWRA has regulatory authority to manage populations of mammal species in the State. To assist with the identification of additional issues and alternatives to managing damage associated with mammals in Tennessee this EA will be made available to the public for review and comment prior to the issuance a Decision[3].

WS previously developed an EA that addressed WS' activities to manage damage associated with mammals in the State, including areas managed and owned by the TVA (USDA 2005*a*). Based on the analyses in that EA, a Decision and Finding of No Significant Impact was signed selecting the proposed action alternative. The proposed action alternative implemented a damage management program using a variety of methods in an integrated approach (USDA 2005*a*). Changes in the need for action and the affected environment have prompted WS and the TVA to initiate this new analysis to address mammal damage in the State. This EA will address more recently identified changes and will assess the potential environmental impacts of program alternatives based on a new need for action, primarily a need to address damage and threats of damage associated with several additional species of mammals. In addition, this EA will: (1) assist in determining if the proposed management of damage associated with mammals could have a significant impact on the environment for both humans and other organisms, (2) analyze several alternatives to address the need for action and the identified issues, (3) coordinate efforts between WS, the TVA, and other entities, (4) inform the public, and (5) document the analyses of the environmental consequences of the alternatives to comply with the NEPA. Since activities conducted under the previous EA will be re-evaluated under this EA to address the new need for action and the associated affected environment, the previous EA that addresses mammal damage management will be superseded by this analysis and the outcome of the Decision issued for this EA.

1.2 NEED FOR ACTION

Some species of wildlife have adapted to and have thrived in human altered habitats. Those species, in particular, are often responsible for the majority of conflicts between people and wildlife. Those conflicts often lead people to request assistance with reducing damage to resources and to reduce threats to human safety. WS' programmatic Final Environmental Impact Statement (FEIS) summarizes the relationship of wildlife values and wildlife damage in this way (USDA 1997):

"*Wildlife has either positive or negative values, depending on varying human perspectives and circumstances...Wildlife is generally regarded as providing economic, recreational and aesthetic benefits...and the mere knowledge that wildlife exists is a positive benefit to many people. However... the activities of some wildlife may result in economic losses to agriculture and damage to property...Sensitivity to varying perspectives and values are required to manage the balance between human and wildlife needs. In addressing conflicts, wildlife managers must consider not only the needs of*

[3]After the development of the EA by WS and consulting agencies and after public involvement in identifying new issues and alternatives, WS will issue a Decision. Based on the analyses in the EA after public involvement, a decision will be made to either publish a Notice of Intent to prepare an Environmental Impact Statement or a Finding of No Significant Impact will be noticed to the public in accordance to NEPA and the Council of Environmental Quality regulations.

those directly affected by wildlife damage but a range of environmental, sociocultural and economic considerations as well."

Both sociological and biological carrying capacities must be considered when resolving wildlife damage problems. The wildlife acceptance capacity, or cultural carrying capacity, is the limit of human tolerance for wildlife or the maximum number of a given species that can coexist compatibly with local human populations. Biological carrying capacity is the land or habitat's ability to support healthy populations of wildlife without degradation to the species' health or their environment during an extended period of time (Decker and Purdy 1988). Those phenomena are especially important because they define the sensitivity of a person or community to a wildlife species. For any given damage situation, there are varying thresholds of tolerance exhibited by those people directly and indirectly affected by the species and any associated damage. This damage threshold determines the wildlife acceptance capacity. While the habitat may have a biological carrying capacity to support higher populations of wildlife, in many cases the wildlife acceptance capacity is lower or has been met. Once the wildlife acceptance capacity is met or exceeded, people begin to implement population or damage management to alleviate damage or address threats to human health and safety.

The alleviation of damage or other problems caused by or related to the behavior of wildlife is termed wildlife damage management and is recognized as an integral component of wildlife management (The Wildlife Society 1990, Berryman 1991). The imminent threat of damage or loss of resources is often sufficient for individual actions to be initiated and the need for damage management is derived from the specific threats to resources. Those species have no intent to do harm. They utilize habitats (*e.g.*, reproduce, walk, forage) where they can find a niche. If their activities result in lost economic value of resources or threaten human safety, people characterize this as damage. When damage exceeds or threatens to exceed an economic threshold and/or pose a threat to human safety, people seek assistance with resolving damage or reducing threats to human safety. The threshold triggering a request for assistance is often unique to the individual person requesting assistance and can be based on many factors (*e.g.*, economic, social, aesthetics). Therefore, what constitutes damage is often unique to the individual person and damage occurring to one individual may not be considered damage by another individual. However, the use of the term "*damage*" is consistently used to describe situations where the individual person has determined the losses associated with wildlife is actual damage requiring assistance (*i.e.*, has reached an individual threshold). The term "*damage*" is most often defined as economic losses to resources or threats to human safety; however, "*damage*" could also be defined as a loss in the aesthetic value of property and other situations where the behavior of wildlife was no longer tolerable to an individual person.

The need for action to manage damage and threats associated with mammals in Tennessee arises from requests for assistance[4] received by WS (USDA 2005*a*). WS receives requests to reduce and prevent damage from occurring to four major categories: agricultural resources, natural resources, property, and threats to human safety (USDA 2005*a*). In addition, the TVA often experiences damage and threats of damage to property and natural resources, electric system operational reliability, as well as threats to human safety at their facilities. WS and the TVA have identified those mammal species most likely to be responsible for causing damage to those four categories in the State based on previous requests for assistance (USDA 2005*a*). Table 1.1 lists WS' technical assistance projects involving mammal damage or threats of damage to those four major resource types in Tennessee from the federal fiscal year[5] (FY) 2005 through FY 2011.

[4] WS would only conduct mammal damage management after receiving a request for assistance. Before initiating damage management activities, a Memorandum of Understanding, cooperative service agreement, or other comparable document would be signed between WS and the cooperating entity that would list all the methods the property owner or manager would allow WS to use on property they own and/or manage.
[5] The federal fiscal year begins on October 1 and ends on September 30 the following year.

Table 1.1 – Technical assistance projects conducted by WS from FY 2005 through FY 2011.

Species	Projects	Species	Projects
Nine-banded Armadillo	6	**Red Fox**	48
Virginia Opossum	37	**Bobcat**	4
Bats (All species)	59	**Woodchuck**	33
Black Bear	5	**Gray Squirrel**	19
Raccoon	1,041	**Fox Squirrel**	10
River Otter	11	**Feral Swine**	123
Striped Skunk	127	**White-tailed Deer**	23
Coyote	115	**Feral Cat**	19
Gray Fox	26	**Feral Dog**	12
Beaver	1,238	**Eastern Chipmunk**	4
Muskrat	37	**Cottontail Rabbit**	8
Moles	38	**Voles**	9
Mice	6	**Rats**	2
Nutria	2		

Technical assistance has been provided by WS to those persons requesting assistance with resolving damage or the threat of damage. Technical assistance provides information and recommendations on activities to alleviate mammal damage that can be conducted by the requestor without WS' direct involvement in managing or preventing the damage. WS' technical assistance activities will be discussed further in Chapter 3 of this EA. Table 1.1 does not include direct operational assistance projects conducted by WS where WS was requested to provide assistance through the direct application of methods.

The technical assistance projects conducted by WS are representative of the mammal species that cause damage and threats in Tennessee. As shown in Table 1.1, WS has conducted 3,062 technical assistance projects in Tennessee that addressed damage and threats associated with those mammal species addressed in this assessment from FY 2005 through FY 2011. Over 74% of the requests for assistance were associated with raccoons and beaver. Raccoons can cause damage to a variety of resource types and can pose threats to human, pet, and livestock safety. For example, raccoons can cause damage to agricultural resources by consuming livestock feed. Raccoons can also pose disease threats to people, pets, and livestock.

Beaver cause damage by gnawing, girdling, and felling trees, as well as impounding water from the dams they build and by burrowing into earthen embankments. The gnawing, girdling, and felling of trees can be aesthetically displeasing to property owners since trees often die and can result lost economic value of property and timber resources. The felling of trees by beaver can cause damage to nearby structures, can block roads, and can pose a safety hazard to human safety.

Table 1.2 lists those mammal species addressed in this EA and the resource types that those mammal species can cause damage to in Tennessee. Many of the mammal species can cause damage to or pose threats to a variety of resources. In Tennessee, most requests for assistance received by WS are related to threats associated with those mammal species causing damage or posing threats of damage to property, agriculture, and human safety. Nearly all those species are known to cause damage to property, including posing strike risks at airport and airbases in Tennessee or posing as attractants for other species that are strike risks.

Table 1.2 – Mammal species that WS routinely receives requests for assistance and the resource type damage by those species

Species	Resource				Species	Resource			
	A	N	P	H		A	N	P	H
Beaver	X	X	X	X	**Striped Skunk**	X	X	X	X
Muskrat	X	X	X	X	**Spotted Skunk**	X	X	X	X
Nutria	X	X	X		**Coyote**	X	X	X	X
Woodchuck	X		X	X	**Gray Fox**	X	X	X	X
Gray Squirrel			X		**Red Fox**	X	X	X	X
Fox Squirrel			X		**Bobcat**	X	X	X	X
Eastern Chipmunk			X		**Feral Cat**	X	X	X	X
Voles			X	X	**Feral Dog**	X	X	X	X
Mice	X	X	X	X	**Nine-banded Armadillo**	X		X	
Rats	X	X	X	X	**Virginia Opossum**	X	X	X	X
Cottontail Rabbit			X	X	**Bats**			X	X
Black Bear	X		X	X	**Moles**			X	
Raccoon	X	X	X	X	**Feral Swine**	X	X	X	X
River Otter	X	X			**White-tailed Deer**	X	X	X	X
Long-tailed Weasel	X	X	X	X	**Elk**	X		X	X
Mink	X	X	X	X					

[a]A=Agriculture, N =Natural Resources, P=Property, H=Human Safety

More specific information regarding mammal damage to those main categories, including damages or threats that could occur on properties owned or managed by the TVA, are discussed in the following subsections of the EA:

Need for Mammal Damage Management on TVA Properties and Facilities

The TVA owns and manages over 293,000 acres in the Tennessee River system. All of these lands support TVA's goals of power generation and transmission, flood control, and economic development of the Tennessee River Valley. The TVA operates hydroelectric dams, coal-fired power plants, nuclear power plants, solar facilities, wind turbine facilities, and combustion turbine sites in Tennessee. The TVA also owns or maintains electrical power substations, switching stations, and the associated transmission lines and rights-of-way easements in Tennessee. In addition, the TVA operates public recreation areas throughout the Tennessee Valley region, including campgrounds, day-use areas, and boat launching ramps.

Mammal damage and threats of damage occurring at facilities and properties owned or managed by the TVA have occurred primarily to property, human safety, and potential electric system operational reliability impacts. Beaver, muskrats, woodchucks, and voles burrowing into earthen levees and dikes used to impound water can compromise the integrity of the structures and threaten the safety of humans downstream from these impoundments. Beaver build dams that impound water causing standing timber and crops to drown, limit the flow of water through irrigation and drainage ditches, and place excessive hydrostatic pressures on roadways and bridges causing millions of dollars in repairs.

Raccoons, opossums, foxes, coyotes, bobcats, feral cats, and feral dogs all reside on TVA lands. These animals frequently become overpopulated or lose their fear of humans, causing zoonotic disease transmission and aggressive behavior toward humans. Many of these lands are considered public or recreational lands and those individuals using these lands expect the TVA to manage mammal populations and reduce the possibilities of disease transmission and attack by wildlife. Mammals

frequently enter substations and power generation facilities and threaten the interruption of power by chewing on various plastic components or climbing into areas of electric current and shorting out electrical circuits. In one instance, numerous raccoons preyed upon tree swallow boxes eating both the eggs and adult birds, threatening research on bioaccumulation of toxins in these species from eating insects exposed to the Kingston Fly Ash Spill. In another instance, several species of mammals were seen hunting and killing nesting endangered least terns on TVA-managed lands. As a federal entity, the TVA is required to support the Endangered Species Act (ESA) and protect these nesting birds against depredation.

All of these damage issues and others occur throughout TVA owned and managed properties. The TVA has requested assistance from WS to address these in the past and may request assistance with additional mammal damage issues in the future. For several years, this cooperative relationship has allowed WS to address TVA's requests for assistance by conducting operational control of these species at 71 different Tennessee sites. Table 1.3 summarizes the mammal species and resource damage that have been addressed by WS on TVA lands during the six-year period, FY 2006 through FY 2011. As many of these species populations grow and thrive in these areas, both WS and TVA expect increases in the need for mammal damage management in the future.

Table 1.3 – Mammal species that WS has addressed on TVA properties and the resource type damage by those species, FY 2006 through FY 2011

Species	Resource				Species	Resource			
	A	N	P	H		A	N	P	H
Beaver	X		X	X	**Virginia Opossum**			X	X
Bobcat		X	X	X	**Raccoon**		X	X	X
Coyote		X	X	X	**Red Fox**		X	X	X
Feral Cat			X	X	**Striped Skunk**			X	X
Feral Dog				X	**Voles**			X	X
Muskrat			X		**Woodchuck**			X	X

[a]A=Agriculture, N =Natural Resources, P=Property, H=Human Safety

Need for Mammal Damage Management to Protect Human Health and Safety

Zoonoses (*i.e.*, wildlife diseases transmissible to people) are a major concern of cooperators when requesting assistance with managing threats from mammals. Disease transmission could occur from direct interactions between humans and mammals or from interactions with pets and livestock that have direct contact with wild mammals. Pets and livestock often encounter and interact with wild mammals, which can increase the opportunity of transmission of disease to humans. Table 1.4 shows common diseases affecting humans that can be transmitted by wild mammals in addition to diseases that affect other animals, including domestic species. These include viral, bacterial, mycotic (fungal), protozoal, and rickettsial diseases.

Table 1.4 - Wildlife diseases in the Eastern United States that pose potential health risks through transmission to humans (Beran 1994, Davidson 2006)[†]

Disease	Causative Agent	Hosts[‡]	Human Exposure
Anthrax	*Bacillus antracis*	cats, dogs	inhalation, ingestion
Tetanus	*Clostridium tetani*	mammals	direct contact
Dermatophilosis	*Dermatophilus congolensis*	mammals	direct contact
Leprosy	*Mycobacterium leprae*	armadillo	inhalation, direct contact
Pasteurellaceae	*Haemophilus influenzae*	mammals	bite or scratch
Salmonellosis	*Salmonella* spp.	mammals	ingestion

Disease	Causative Agent	Hosts[‡]	Human Exposure
Yersinosis	*Yersinia* spp.	cats	ingestion
Chlamydioses	*Chlamydophilia felis*	cats	inhalation, direct contact
Typhus	*Rickettsia prowazekii*	opossums	inhalation, ticks, fleas
Sarcoptic mange	*Sarcoptes scabiei*	red fox, coyotes, dogs	direct contact
Trichinosis	*Trichinella spiralis*	raccoons, fox	ingestion, direct contact
Rabies	*Lyssavirus* spp.	mammals	direct contact
Visceral larval	*Baylisascaris procyonis*	raccoons, skunks	ingestion, direct contact
Leptospirosis	*Leptospira interrogans*	mammals	ingestion, direct contact
Echinococcus	*Echinococcus multilocularis*	fox, coyotes	ingestion, direct contact
Toxoplasmosis	*Toxoplasma ondii*	cats, mammals	ingestion, direct contact
Spirometra	*Spirometra mansonoides*	bobcats, raccoons, fox	ingestion, direct contact
Giardiasis	*Giardia lamblia, G. duodenalis*	mammals	ingestion, direct contact

[†]Table 1.3 is not considered an exhaustive list of wildlife diseases that are considered infectious to humans that are carried by wildlife species. The zoonoses provided are the more common infectious diseases for the species addressed in this EA and are only a representation of the approximately 100 to 3,000 zoonoses known to exist.

[‡] The host species provided for each zoonosis includes only those mammalian species addressed in this EA unless the zoonoses listed potentially infects a broad range of mammalian wildlife. Zoonoses infecting a broad range of mammals are denoted by the general term "*mammals*" as the host species. The diseases listed do not necessarily infect only those mammalian species covered under this EA but likely infect several species of mammals or groups of mammals. For a complete discussion of the more prevalent diseases in free-ranging mammals, please refer to Beran (1994) and Davidson (2006).

Individuals or property owners that request assistance with mammals frequently are concerned about potential disease risks but are unaware of the types of diseases that can be transmitted by those animals. In those types of situations, assistance is requested because of a perceived risk to human health or safety associated with wild animals living in close association with humans, from animals acting out of character by roving in human-inhabited areas during daylight, or from animals showing no fear when humans are present. Under the proposed action, WS could assist in resolving those types of requests for assistance.

In many circumstances when human health concerns are the primary reason for requesting WS' assistance there may have been no actual cases of transmission of disease to humans by mammals. Thus, the risk of disease transmission would be the primary reason for requesting assistance from WS. Situations in Tennessee where the threat of disease associated with wild or feral mammal populations may include:

- Exposure of residents to the threat of rabies due to high densities of raccoons or from companion animals encountering infected raccoons.
- Exposure of humans to threats of rabies posed by skunks that den under buildings or from companion animals interacting with infected skunks.
- Threats of parasitic infections to humans from *Giardia* spp. resulting from high feral cat populations in a park or recreation area.
- Concern about the threat of histoplasmosis from the disturbance of a large deposit of guano in an attic where a large colony of bats routinely roosts or raise young.
- Accumulated droppings from denning or foraging raccoons and the subsequent exposure of the public to raccoon roundworm in fecal deposits.
- Exposure of domestic livestock to the bacterium, *Brucella suis*, by feral swine. *B. suis* causes swine brucellosis.

The most common disease concern expressed by individuals requesting assistance is the threat of rabies transmission to humans, pets, and companion animals. Rabies is an acute, fatal viral disease of mammals most often transmitted through the bite of a rabid animal that poses an indirect and direct threat to humans. Indirect threats to humans occur from exposure from pets or livestock that have been infected

from bites of a rabid animal. Direct threats can occur from handling infected wildlife or from aggressive animal behavior caused by rabies. The disease can be effectively prevented in humans when exposure is identified early and treated and domestic animals and pets can be vaccinated for rabies. However, the abundant and widely distributed reservoir among wild mammals complicates rabies control. The vast majority of rabies cases reported to the Centers for Disease Control and Prevention (CDC) each year occur in raccoons, skunks (primarily *Mephitis mephitis*), and bats (Order Chiroptera) (CDC 2011).

Over the last 100 years, the vector of rabies in the United States has changed dramatically. About 90% or greater of all animal cases reported annually to CDC now occur in wildlife (Krebs et al. 2000, CDC 2011). Before 1960, the majority of cases were reported in domestic animals. The principal rabies hosts today are wild carnivores and bats. The number of rabies-related human deaths in the United States has declined from more than 100 annually in the early 1900s to an average of one or two people per year in the 1990s. Modern day prophylaxis, which is the series of vaccine injections given to people who have been potentially or actually exposed, has proven nearly 100% successful in preventing mortality when administered promptly (CDC 2011). In the United States, human fatalities associated with rabies occur in people who fail to seek timely medical assistance, usually because they were unaware of their exposure to rabies. Although human rabies deaths are rare, the estimated public health costs associated with disease detection, prevention, and control have risen, exceeding $300 million annually. Those costs include the vaccination of companion animals, maintenance of rabies laboratories, medical costs such as those incurred for exposure case investigations, rabies post-exposure prophylaxis (PEP), and animal control programs (CDC 2011).

Accurate estimates of the aforementioned expenditures are not available. Although the number of PEPs given in the United States each year is unknown, it has been estimated to be as high as 40,000. When rabies becomes epizootic (*i.e.*, affecting a large number of animals over a large area) or enzootic (*i.e.*, present in an area over time but with a low case frequency) in a region, the number of PEPs in that area increases. Although the cost varies, a course of rabies immunoglobulin and five doses of vaccine given over a 4-week period typically exceeds $1,000 (CDC 2011) and has been reported to be as high as $3,000 or more (Meltzer 1996). As epizootics spread in wildlife populations, the risk of "*mass*" human exposures requiring treatment of large numbers of people that contact individual rabid domestic animals infected by wild rabid animals increases. One case in Massachusetts involving contact with, or drinking milk from, a single rabid cow required PEPs for 71 persons (CDC 1999). The total cost of this single incident exceeded $160,000 based on a median cost of $2,376 per PEP in Massachusetts. Likely, the most expensive single mass exposure case on record in the United States occurred in 1994 when a kitten from a pet store in Concord, New Hampshire tested positive for rabies after a brief illness. Because of potential exposure to the kitten or to other potentially rabid animals in the store, at least 665 persons received post-exposure rabies vaccinations at a total cost of more than $1.1 million (Noah et al. 1995). The American Veterinary Medical Association (AVMA) estimated the total cost for this specific incident, including investigation, laboratory testing, and rabies immunoglobulin and vaccines was more than $1.5 million (AVMA 2004).

Raccoons have been associated with the spread of rabies throughout the eastern United States, including Tennessee (USDA 2005*b*). Rabies in raccoons was virtually unknown prior to the 1950s. It was first described in Florida and spread slowly during the next three decades into Georgia, Alabama, and South Carolina. It was unintentionally introduced into the Mid-Atlantic States, probably by translocation of infected animals (Krebs et al. 1998). The first cases appeared in West Virginia and Virginia in 1977 and 1978, respectively. Since then, raccoon variant rabies expanded to form the most intensive rabies outbreak in the United States. The variant is now enzootic in all of the eastern coastal states, as well as Alabama, Pennsylvania, Vermont, West Virginia, and most recently, parts of Ohio (Krebs et al. 2000). The raccoon rabies epizootic front reached Maine in 1994, reflecting a movement rate of about 30 to 35 miles per year. Raccoon variant rabies was first identified in 2003 in upper east Tennessee (USDA 2004)

and has since been found in nine Tennessee counties (USDA 2007). The westward movement of the raccoon rabies front has slowed, probably in response to both natural geographic and man-made barriers. The Appalachian Mountains and perhaps river systems flowing eastward have helped confine the raccoon variant to the eastern United States. In addition, the USDA has created an oral rabies vaccine (ORV) "*barrier*" of vaccinated wild animals on the western edge of the Appalachian Mountains (USDA 2005*b*). If this combined barrier is breached by raccoon variant rabies, research suggests that raccoon populations are sufficient for rabies to spread westward along a front at a rate similar to or greater than the rate at which this rabies strain has spread in the eastern United States (Sanderson and Huber, Jr. 1982, Glueck et al. 1988, Hasbrouck et al. 1992, Mosillo et al. 1999).

Raccoon variant rabies presents a human health threat through potential direct exposure to rabid raccoons, or indirectly through the exposure of pets that have an encounter with rabid raccoons. Additionally, the number of pets and livestock examined and vaccinated for rabies, the number of diagnostic tests requested, and the number of post exposure treatments are all higher when raccoon rabies is present in an area. Human and financial resources allocated to rabies-related human and animal health needs also increase, often at the expense of other important activities and services.

Skunks are also an important wildlife host for the rabies virus in North America and are second only to raccoons in being the most commonly reported rabid wildlife species in the United States (Majumdar et al. 2005). The skunk variant of rabies may be found in the Midwest and California; however, skunks found throughout North America may be infected with different variants of rabies such as the raccoon variant. The distribution of rabies in skunks extends from Georgia to Maine east of the Appalachians, Texas to the Canadian border, and throughout the northern two thirds of California (Majumdar et al. 2005). The fox is one of the four major maintenance hosts for rabies in North America. In the 1950s, rabies in red fox spread throughout Canada, parts of New England, and Alaska. The range has since decreased, but fox rabies persists in Alaska and parts of Texas. Clinical signs of rabies in fox are often manifested as the "*furious*" form of rabies (Majumdar et al. 2005).

In an effort to halt the westward spread of the raccoon variant of the rabies virus and to limit the spread of the canine variant from Texas, WS began participating in the distribution of ORV baits (fishmeal polymer containing Raboral V-RG® vaccine [Merial, Athens, Georgia, USA]). Currently, WS participates in the distribution of ORV baits and the surveillance of wildlife rabies vectors in 26 states, including Tennessee. ORV baits were first distributed by WS in Tennessee during the fall of 2002. A total of 130,494 baits were distributed (111,414 by air and 19,080 by hand) across a 4,113 km^2 area which included portions of Grainger, Greene, Hamblen, Hancock, Hawkins, Sullivan, and Washington Counties. Tennessee has increased the size of this barrier (now referred to as the Appalachian Ridge ORV zone) and moved it east, in five mile increments, to include large portions of Carter, Cocke, Greene, Hamblen, Hawkins, Sullivan, Unicoi and Washington Counties. In addition, a second barrier (called the Georgia-Alabama-Tennessee or GAT ORV zone) was created in 2003, in which, 67,202 baits were distributed (28,322 by air and 38,880 by hand) across a 1,014-km2 area that included portions of Hamilton, Marion, and Sequatchie Counties. Tennessee has increased the size of the GAT barrier significantly to extend further north into the State and east, in five-mile increments, to include large portions of Bradley, Hamilton, McMinn, Meigs, Monroe, and Polk Counties. Since the inception of the program in the fall of 2002 through the FY 2012 bait distribution in October 2011, approximately 6,000,311 ORV baits have been distributed in Tennessee. As part of surveillance of rabies vectors in Tennessee, from 2002 through 2011, WS collected 8,328 samples for rabies testing. Of those 8,328 samples, 45 samples tested positive for the Eastern United States raccoon rabies variant, 55 samples tested positive for north central plains skunk variant, and two tested positive for the big brown bat variant. WS' participation in the ORV program is further addressed in a separate EA (USDA 2005*b*) but will be addressed in this EA to evaluate potential cumulative effects of activities proposed in this EA and the capturing and releasing of target animals during surveillance activities associated with the ORV program (USDA 2005*b*).

Increasing populations of raccoons have been implicated in the outbreak of distemper in certain areas (Majumdar et al. 2005). Distemper has not been identified as transmissible to humans. However, cooperators who feel threatened by the possibility of disease transmission often request assistance after observing sick raccoons on their property. Symptoms of distemper often lead to abnormal behavior in raccoons that are similar to symptoms associated with rabies. Raccoons with distemper often lose their fear of humans and can act aggressively which increases the risk that people, livestock, or companion animals may be bitten. Distemper is also known to occur in coyotes, red fox, and gray fox with symptoms that are similar to those exhibited by animals infected with the rabies virus.

Diseases and parasites affecting feral cats and dogs can have particularly serious implications to human health given the close association of those animals with humans and companion animals. The topic of feral animals and their impacts on native wildlife and human health elicits a strong response in numerous professional and societal groups with an interest in the topic. Feral cats and dogs are considered by most professional wildlife groups to be a non-native species that has detrimental impacts to the native ecosystems especially in the presence of a human altered landscape. However, a segment of society views feral animals to be an extension of companion animals that should be cared for and for which affection bonds are often developed especially when societal groups feed and care for individual feral animals. Of special concern are those cats and dogs considered companion animals that are not confined indoors at all times but are allowed to range outside the home for extended periods. If interactions occur between companion animals and feral animals of the same species, companion animals could become exposed to a wide-range of zoonoses that could be brought back into the home where direct contact between the companion animal and people increases the likelihood of disease transmission. Feral animals that are considered companion animals are also likely to affect multiple people if disease transmission occurs since those animals are likely to come in direct contact with several members of families and friends before diagnosis of a disease occurs.

Several known diseases that are infectious to humans, including rabies, have been found in feral cats and dogs. A common zoonosis found in cats is ringworm. Ringworm (*Tinea* spp.) is a contagious fungal disease contracted through direct interactions with an infected person, animal, or soil. Other common zoonoses of cats are pasteurella, salmonella, cat scratch disease, and numerous parasitic diseases, including roundworms, tapeworms, and toxoplasmosis.

Most of the zoonoses known to infect cats and dogs that are infectious to humans are not life threatening if diagnosed and treated early. However, certain societal segments are at higher risks if exposed to zoonoses. Women who are pregnant, people receiving chemotherapy for immunologic diseases and organ transplants, and those with weakened immune systems are at increased risk of clinical disease if exposed to toxoplasmosis (AVMA 2004). In 1994, five Florida children were hospitalized with encephalitis that was associated with cat scratch fever (AVMA 2004). The daycare center at the University of Hawaii in Manoa was closed for two weeks in 2002 because of concerns about potential transmission of murine typhus (*Rickettsia typhi*) and flea (*Ctenocephalides felis*) infestations afflicting 84 children and faculty. The fleas at the facility originated from a feral cat colony that had grown from 100 cats to over 1,000, despite a trap, neuter, and release effort (AVMA 2004).

Feral swine can pose a threat to human safety from disease transmission, from aggressive behavior, and from being struck by vehicles and aircraft. Feral swine are potential reservoirs for at least 30 viral and bacterial diseases (Samuel et al. 2001, Williams and Barker 2001, Davidson 2006) and 37 parasites (Forrester 1991) that are transmissible to humans. Brucellosis, salmonellosis, toxoplasmosis, trichinosis, tuberculosis, and tularemia are some of the common diseases that can be carried by feral swine that are also known to infect humans (Stevens 1996, Hubalek et al. 2002, Seward et al. 2004). In addition, feral

swine can pose risks to domestic livestock through the potential transmission of diseases between feral swine populations and domestic livestock where interactions may occur.

Conflicts involving bats can include property damage, but primarily involve threats to human, pet, and livestock health. The buildup of bat droppings and urine in attics and between walls can result in odor problems and discoloration of walls and ceilings (Agency for Toxic Substances and Disease Registry 1998). In addition to the threat of rabies from direct contact or a bat entering the living area of a home, there are other threats associated with bat colonies including histoplasmosis, fungal spores, and mites.

Bat droppings, particularly when they accumulated for many years, are likely to be contaminated with the fungus, *Histoplasma capsulatum* or with fungi species such as molds, especially in warm, moist conditions. As long as people are not in contact with fungal spores, they are unlikely to be affected by them. When people inhale spores from *Histoplasma capsulatum*, they may become ill with a disease known as histoplasmosis. Symptoms of histoplasmosis include some combination of mild, flu-like respiratory illness, a general ill feeling, chest pain, fever, cough, headache, loss of appetite, shortness of breath, joint and muscle pains, chills, and hoarseness. Although there are other, more rare illnesses associated with exposure, the most likely is histoplasmosis. Similarly, mold spores released into the air may result in increases in asthma attacks (Agency for Toxic Substances and Disease Registry 1998).

Bat bugs (*Cimex adjunctus*) are free-living ectoparasites of bats that feed on blood. They will bite humans in the absence of their primary hosts. The main means of dispersal for bat bugs is by clinging to the fur of bats as bats move between locations. Typically, bat bug infestations originate from bat populations established in attics, wall voids, unused chimneys, or uninhabited portions of a house. Bat bugs typically do not wander far from occupied bat roosting sites where they have easy access to food. However, if their normal hosts were eliminated or the area was vacated, bat bugs are known to seek other sources of food and may crawl about and invade living areas within the house (Jones and Jordan 2004). Similarly, bat mites may enter the home and bite people. Although their bite is not particularly harmful, the person may experience an allergic reaction and develop a skin rash in response (Agency for Toxic Substances and Disease Registry 1998).

This discussion on zoonoses is intended to briefly address the more common known zoonoses found in the United States for those species specifically addressed in this EA but is not intended to be an exhaustive discussion of all potential zoonoses. The transmission of diseases from wildlife to humans is neither well documented nor well understood for most infectious zoonoses. Determining a vector for a human infected with a disease known to occur in wildlife populations is often complicated by the presence of the known agent across a broad range of naturally occurring sources. For example, a person with salmonella poisoning may have contracted salmonella bacterium from direct contact with an infected pet but may have also contracted the bacterium from eating undercooked meat or from other sources.

Disease transmission directly from wildlife to humans is uncommon. However, the infrequency of such transmission does not diminish the concerns of those individuals requesting assistance that are fearful of exposure to a diseased animal since disease transmissions have been documented to occur. WS actively attempts to educate the public about the risks associated with disease transmission from wildlife to humans through technical assistance and by providing technical leaflets on the risks of exposure.

In addition to disease transmission threats, requests are also received for assistance from perceived threats of physical harm from wildlife, especially from predatory wildlife. Human encroachment into wildlife habitat increases the likelihood of human-wildlife interactions. Those species that humans are likely to encounter are those most likely to adapt to and thrive in human altered habitat. Several predatory and omnivorous wildlife species thrive in urban habitat due to the availability of food, water, and shelter. Many people enjoy wildlife to the point of purchasing food specifically for feeding wildlife despite laws

prohibiting the act in many areas. The constant presence of human created refuse, readily available water supplies, and abundant rodent populations found in some areas often increases the survival rates and carrying capacity of wildlife species that are adaptable to those habitats. Often the only limiting factor of wildlife species in and around areas inhabited by people is the prevalence of diseases, which can be confounded by the overabundance of wildlife congregated into a small area that can be created by the unlimited amount of food, water, and shelter found within those habitats.

As people are increasingly living with wildlife, the lack of harassing and threatening behavior by people toward many species of wildlife has led to a decline in the fear wildlife have toward people. When wildlife species begin to habituate to the presence of humans and human activity, a loss of apprehension occurs that can lead to threatening behavior toward humans. This threatening behavior continues to increase as human populations expand and the populations of those species that adapt to human activity increase. Threatening behavior can be in the form of aggressive posturing, a general lack of apprehension toward people, or abnormal behavior. Although wildlife attacking people occurs rarely, the number of attacks appears to be on the increase. Timm et al. (2004) reported that coyotes attacking people have increased in California and the recent, highly publicized coyote attacks, including a fatal attack on a 19-year old woman in Nova Scotia (Canadian Broadcast Company 2009), have only heightened people's awareness of the threat of such encounters.

Black bears occasionally threaten human health and safety. Herrero (1985) documented 500 injuries to humans resulting from encounters with black bears from 1960 to 1980. Of those, 90% were minor injuries (minor bites, scratches, and bruises). Only 23 fatalities were recorded from 1900 to 1980 due to black bear attacks. The number of bear attacks could be considered low considering the geographic overlap of human and black bear populations. Of those fatalities, 90% were likely associated with habituated, food-conditioned bears. Black bear attacks on people in Tennessee have resulted in recent fatalities occurring at the Great Smoky Mountains National Park in 2000 and the Cherokee National Forest in 2006. Additional attacks on humans from gray fox, feral swine, feral cats, feral dogs, and raccoons have been documented in the last few years in Tennessee. Often, wildlife exhibiting threatening behavior or a loss of apprehensiveness to the presence of humans is a direct result and indication of an animal inflicted with a disease. So, requests for assistance are caused by both a desire to reduce the threat of disease transmission and from fear of aggressive behavior either from an animal that is less apprehensive of people or induced as a symptom of disease.

As part of the proposed program, WS could provide mammal damage management assistance, upon request, involving those mammal species addressed in this EA that pose a threat to human health and safety to any requester experiencing such a threat throughout Tennessee.

Disease Surveillance and Monitoring

Public awareness and health risks associated with zoonoses (*i.e.*, diseases of animals that can be transmitted to humans) have increased in recent years. Several zoonotic diseases associated with mammals are addressed in this EA. Those zoonotic diseases remain a concern and continue to pose threats to human safety where people encounter mammals. WS has received requests to assist with reducing damage and threats associated with several mammal species in Tennessee and could conduct or assist with disease monitoring or surveillance activities for any of the mammal species addressed in this EA. Most disease sampling would occur ancillary to other wildlife damage management activities (*i.e.*, disease sampling occurs after wildlife have been captured or lethally taken for other purposes). For example, WS may sample deer harvested during the annual hunting season or during other damage management programs for Chronic Wasting Disease or may collect ticks from raccoons that were lethally taken to alleviate damage occurring to property. WS could sample feral hogs taken by private landowners

or during damage management activities to test for classical swine fever, swine brucellosis, pseudorabies, or other diseases.

Emergency Response Efforts

Both large-scale natural disasters (*e.g*., hurricanes, tornadoes, and floods) and small-scale localized emergencies (*e.g*., release of exotic animals, traffic accidents involving animal transport vehicles) may occur in which WS' personnel could be requested to assist federal, state, and local governments in charge of responding to those situations. Those requests for assistance would be on extremely short notice and rare emergencies that would be coordinated by federal, state, and local emergency management agencies. For example, WS' personnel may be requested to participate in the lethal removal of cattle that were injured or were released from their transport vehicle at the scene of an accident to prevent those animals from endangering other drivers. WS could be asked to corral those animals that were uninjured and euthanize those animals that have been injured to reduce their suffering. In another example, WS' personnel may be requested to assist local and state law enforcement in immobilization or lethal control of exotic animals that have been accidentally released in the aftermath of a hurricane or tornado.

Need for Mammal Damage Management at Airports

Airports provide ideal conditions for many wildlife species due to the large open grassy areas adjacent to brushy, forested habitat used as noise barriers. Access to most airport properties is restricted so wildlife living within airport boundaries would be protected during hunting and trapping seasons and would be insulated from many other human disturbances.

The civil and military aviation communities have acknowledged that the threat to human health and safety from aircraft collisions with wildlife is increasing (Dolbeer 2000, MacKinnon et al. 2001, Dolbeer 2009). Collisions between aircraft and wildlife are a concern throughout the world because wildlife strikes threaten passenger safety (Thorpe 1996), result in lost revenue, and repairs to aircraft can be costly (Linnell et al. 1996, Robinson 1996, Thorpe 1997, Keirn et al. 2010). Aircraft collisions with wildlife can also erode public confidence in the air transport industry as a whole (Conover et al. 1995).

Between 1990 and 2010, there were 2,558 reported aircraft strikes involving terrestrial mammals in the United States (Dolbeer et al. 2012). The number of mammal strikes actually occurring is likely to be much greater, since Dolbeer (2009) estimated 39% of civil wildlife strikes are actually reported. Civil and military aircraft have collided with a reported 36 species of terrestrial mammals from 1990 through 2010, including raccoons, gray fox, red fox, cats, coyotes, opossums, and striped skunks. In addition, 13 species of bats have been identified as having been struck by aircraft in the United States (Dolbeer et al. 2012). Of the terrestrial mammals reported struck by aircraft, 33% were carnivores (primarily coyotes), causing nearly $3.2 million in damages (Dolbeer et al. 2012). Deer accounted for 39% of the reported strikes involving terrestrial mammals in the United States causing nearly $31 million in damages (Dolbeer et al. 2012). Data also indicates that a much higher percentage of mammal strikes resulted in aircraft damage compared to bird strikes (Dolbeer et al. 2012). Costs of those collisions vary, but the Federal Aviation Administration (FAA) data reveals that mammal strikes in the United States cost the civil aviation industry approximately 275,290 hours of down time and $41.1 million in direct monetary losses between 1990 and 2010 (Dolbeer et al. 2012).

About 59% of mammal strikes in the United States have resulted in damage compared to 13% for birds from 1990 through 2010 (Dolbeer et al. 2012). In addition to direct damage, an aircraft striking a mammal can pose serious threats to human safety if the damage from the strike causes a catastrophic failure of the aircraft leading to a crash. For example, damage to the landing gear during the landing roll and/or takeoff run can cause a loss of control of the aircraft, causing additional damage to the aircraft and

increasing the threat to human safety. Nearly 63% of the reported mammal strikes from 1990 through 2010 occurred at night, with 63% occurring during the landing roll or the takeoff run (Dolbeer et al. 2012).

Between 1990 and 2010, aircraft have struck five coyotes, one striped skunk, three red bats, two unidentified bats, and 16 white-tailed deer in Tennessee according to reports filed with the FAA (FAA 2010). Airports in Tennessee have requested assistance to manage the threat to human safety and damage to property caused by mammals present inside the area of operations of the airport. The infrequency of mammal strikes does not lessen the need to prevent threats to human safety and the prevention of damage to property. Preventing damage and reducing threats to human safety is the goal of those cooperators requesting assistance at airports in Tennessee given that a potential strike can lead to the loss of human life and considerable damage to property.

Wildlife populations near or found confined within perimeter fences at airports can be a threat to human safety and cause damage to property when struck by aircraft. Those wildlife confined inside the airport perimeter fence would not be considered distinct populations nor separate from those populations found outside the perimeter fence. Wildlife found within the boundaries of perimeter fences originate from populations outside the fence. Those individuals of a species inside the fence neither exhibit nor have unique characteristics from those individuals of the same species that occur outside the fence; therefore, those individuals of a species confined inside an airport perimeter fence do not warrant consideration as a unique population under this analysis.

Need for Mammal Damage Management to Protect Agricultural Resources

Black bears, red fox, gray fox, bobcats, coyotes, deer, and other mammals can cause losses or injury to crops, livestock (*e.g.*, sheep, goats, cattle, pigs, horses), and poultry (*e.g.*, chickens, turkeys, geese, ducks) through predation. During 2001, crop and livestock losses from wildlife in the United States totaled $944 million, with field crop losses totaling $619 million, livestock and poultry losses totaling $178 million, and losses of vegetables, fruits, and nuts totaling $146 million. Those losses include destruction of or damage to crops in the field and death or injury to livestock. In 2001, the National Agricultural Statistics Service (NASS) reported that raccoons were responsible for 6%, 3%, and 6% of the total damage to field crops; livestock and poultry; and vegetables, fruits, and nuts, respectively, in the United States (NASS 2002). In addition, white-tailed deer accounted for 58% of the total field crop damage and 33% of vegetable, fruit, and nut damage. Wild pigs accounted for 3% or $18.5 million in damages to field crops (NASS 2002).

In 2010, the NASS (2011) reported cattle and calf losses from animal predation totaled 219,900 head in the United States according to livestock producers. Animal predation represented 5.5% of the total cattle and calf losses reported by livestock producers in 2010 totaling $98.5 million in economic losses. Coyotes were indicated as the primary predator of livestock with 53.1% of cattle and calf losses attributed to coyotes. Livestock losses were also attributed to bobcats, bears, and dogs. Producers spent nearly $188.5 million dollars on non-lethal methods to reduce cattle and calf losses from predation by animals in 2010 (NASS 2011). The primary non-lethal method employed by livestock producers was the use of guard animals with a reported 36.9% of producers using guard animals. Producers also reported using exclusion fencing, frequent checking, and culling as additional employed methods for reducing predation (NASS 2011).

In Tennessee, the NASS (2011) reported 1,300 cattle and 7,800 calves were killed in 2010 by animal predators. The economic loss from animal predators in Tennessee was estimated at nearly $3.6 million in 2010 (NASS 2011). Coyotes were attributed to 62.1% of the cattle losses and 62.5% of the calves lost in Tennessee. Dogs accounted for 26.4% of the cattle reported lost while 16.0% of the calves lost were

attributed to dogs in the State (NASS 2011). Cattle producers in Tennessee reported using a number of non-lethal methods to reduce losses due to predators. The use of exclusion fencing was reported as being employed by 33.9% of Tennessee cattle producers along with 33.9% reporting the use of guard animals (NASS 2011).

NASS (2011) reported that 0.3% of the calves lost to animal predator were attributed to mountain lions and bobcat predation in Tennessee. Cattle producers in the United States indicated mountain lions and bobcats[6] caused 7.8% of the cattle and calf losses attributed to animal predators in 2010 (NASS 2011). Bobcats are also known to prey on other livestock.

Woodchucks (commonly referred to as groundhogs) can cause damage to field crops, such as row and forage crops, orchards, nursery plants, and commercial gardens. Cottontail rabbits and voles are reported to damage orchard trees by gnawing at the base of the tree. Trees are badly damaged or the bark is girdled and trees die when feeding by rabbits and voles is severe. Similar damage occurs in nurseries, which grow landscape ornamentals and shrubs.

River otters and to a lesser extent bears and raccoons may prey on fish and other cultured species at hatcheries and aquaculture facilities (Bevan et al. 2002). River otters may even prey on fish in marine aquaculture facilities (Goldburg et al. 2001).

The domestic cat has been found to transmit *Toxoplasma gondii* to both domestic and wild animal species. Cats have been found to be important reservoirs and the only species known to allow for the completion of the life cycle for the protozoan parasite *T. gondii* (Dubey 1973, Teutsch et al. 1979). Both feral and domiciled cats may be infected by this protozoan, but this infection is more common in feral cats. Fitzgerald et al. (1984) documented that feral cats transmitted *T. gondii* to sheep in New Zealand, resulting in ewes aborting fetuses. The authors also found *Sarcocystis* spp. contamination in the musculature of sheep. Dubey et al. (1995) found cats to be 68.3% positive for seroprevalence of *T. gondii* on swine farms in Illinois and the major reservoir for this disease. The main sources for infecting cats are thought to be birds and mice.

Agricultural damage and threats caused by feral swine in Tennessee occurs to crops, livestock, and other agricultural resources. Damage occurs from direct consumption of agricultural crops and from trampling, rooting, and/or wallowing that are common activities of feral swine (Beach 1993). Rooting is a common activity of feral swine during their search for food where they overturn sod and soil (Stevens 1996). Feral swine also wallow in water and mud to regulate body temperature and to ward off skin parasites.

Damage and threats to livestock associated with feral swine occurs from predation on livestock and the risks associated with disease transfer from feral swine to domestic livestock. Feral swine can also cause damage to other agricultural resources. For example, feral swine can cause damage to pastures and land used for hay by rooting and wallowing, can cause damage to ponds and water sources for livestock, and can cause damage from the consumption of livestock feed. Feral swine feeding activities in agricultural crops can also lead to increased erosion from the removal of vegetation that leaves the soil bare along with the overturning of soil caused by rooting.

In addition, feral swine also damage pastures, land used for hay, and sod farms from rooting and wallowing activities (Beach 1993). Rooting activities can also lead to increased erosion and soil loss. Wallowing and rooting activities in livestock watering areas can lead to a degradation in water quality

[6]The 2011 NASS cattle loss report groups mountain lion and bobcat predation into one category and does not separate losses attributed to the two species. Mountain lions, given their preference for larger prey, are likely the cause of most of the losses attributed to this category, especially to adult cattle. However, bobcats are known to prey upon calves though infrequently.

through an increase in turbidity, by causing algal blooms, by depleting dissolved oxygen, and increasing erosion (Beach 1993). Since feral hogs often travel in family groups, damages from rooting and wallowing can be extensive often encompassing several acres.

Additional risks associated with feral hogs are the potential for disease transmission from feral swine to domestic livestock, especially to domestic swine. Feral swine are potential reservoirs for several diseases that are known to be transmissible between feral swine and domestic livestock (Wood and Barrett 1979, Corn et al. 1986, Beach 1993, Davidson 2006). Corn et al. (1986) found feral swine tested in Texas were positive for pseudorabies, brucellosis, and leptospirosis. A study in Oklahoma found samples from feral swine tested positive for antibodies of porcine parvovirus, swine influenza, and porcine reproductive and respiratory syndrome virus (Saliki et al. 1998). Cholera, trichinosis, and African swine fever are additional diseases that can be transmitted between livestock and feral swine. Disease transmission is likely to occur where domestic livestock and feral swine have a common interface, such as at water sources and livestock feeding areas. The WS program in Tennessee could conduct disease surveillance in the feral swine population as part of the National Wildlife Disease Surveillance Program.

Although several diseases that are carried by swine are also transmissible to other livestock, the primary concern is the potential transmission of diseases from feral swine to domestic swine. Pseudorabies is a viral disease associated with an extremely contagious herpes virus that can have negative effects on reproduction in domestic swine. Brucellosis is a bacterial disease that can also have negative effects on reproduction in swine. Many of the other diseases associated with feral swine also negatively affect the health and marketability of domestic swine that can lead to economic losses to the livestock producer. The United States is one of the world's largest producers of pork and is the second largest exporter of pork. Pork production in the United States accounts for about 10% of the total world supply. The retail value of pork sold to consumers exceeds $30 billion annually. In addition, the pork industry supports more than 600,000 jobs. An economic analysis estimated that the annual cost of pseudorabies to pork producers in the United States at more than $30 million annually in lost production as well as testing and vaccination costs (USDA 2008).

Although the source of livestock disease outbreaks can be difficult to identify, a risk of transmission and the spreading of diseases to domestic swine and other livestock exists wherever feral swine and domestic livestock interact. A disease outbreak not only has negative economic implications to the individual livestock producer, but also can cause economic losses that can negatively affect the statewide swine industry.

In addition to the potential for disease transmission, feral swine are also known to predate on livestock. Feral swine are known to kill calves, kids, lambs, and poultry (Stevens 1996, West et al. 2009). Predation occurs primarily on young livestock but feral hogs can also kill weakened or injured livestock. If feral swine populations continue to increase, WS could be requested to address localized predation associated with feral swine.

Nutria depredation on crops has also been documented (LeBlanc 1994). Crops that have been damaged include corn, milo (grain sorghum), sugar and table beets, alfalfa, wheat, barley, oats, peanuts, various melons, and a variety of vegetables from home gardens and truck farms.

Examples of some of the requests for assistance to resolve or alleviate damage to agricultural resources that the WS' program in Tennessee has responded to include:

- Coyotes attacking and killing calves, lambs, chickens, and emus
- Raccoons digging up grass and sod while foraging for insects
- Gray squirrels feeding on strawberries, peaches, and pecans

- Gray fox killing chickens and domestic waterfowl
- Striped skunks killing chickens

Need for Mammal Damage Management to Resolve Damage Occurring to Natural Resources

Natural resources may be described as those assets belonging to the public and often managed and held in trust by government agencies as representatives of the people. Such resources may be plants or animals, including threatened and endangered (T&E) species; historic properties; or habitats in general. Examples of natural resources in Tennessee are historic structures and places; parks and recreation areas; natural areas, including unique habitats or topographic features; threatened and endangered plants or animals; and any plant or animal populations which have been identified by the public as a natural resource.

Mammals can also cause damage to natural resources. Mammals causing damage are often locally overabundant at the damage site and threaten the welfare of a species population identified as a natural resource. An example of this would be nest predation of a local ground-nesting bird population by mammalian carnivores, such as raccoons, opossum, feral swine, feral cats, coyotes, or fox.

Massey (1971) and Massey and Atwood (1981) found that predators can prevent endangered least terns (*Sterna antillarum*) from nesting or cause them to abandon previously occupied sites. In another study, mammalian predators were found to have adversely affected the nesting success of least terns on sandbars and sandpits (Kirsch 1996). In Tennessee, WS has conducted extensive predator removal efforts to protect endangered least tern nesting sites at the request of and in cooperation with the TVA. Personnel witnessed the predation of nests and nestlings by raccoons and coyotes and WS removed several of these animals in order to allow these terns to complete their nesting season.

Beaver activities can also destroy habitat (*e.g.*, free-flowing water, riparian areas, and bird roosting and nesting areas) which are important to many species. Patterson (1950) and Avery (1992) reported that the presence of beaver dams could negatively affect fisheries. Beaver dams may adversely affect stream ecosystems by increasing sedimentation in streams, and thereby affecting wildlife that depend on clear water such as certain species of fish and mussels. Stagnant water impounded by beaver dams can increase the water temperature of water impounded upstream of the dam which can negatively affect aquatic organism. Beaver dams can also act as barriers that inhibit movement of aquatic organisms and prevent the migration of fish to spawning areas.

The TWRA and TVA have identified several areas of west Tennessee that are considered Critically Imperiled Habitats for several species of wildlife. Those areas most likely contain plant, vertebrate, and invertebrate species that are either already endangered or threatened. At the request of the TVA, the major land manager in those areas, WS could be requested to provide assistance with managing beaver damage associated with water impounded by beaver dams to allow water levels to fluctuate seasonally. In addition, WS could be requested to manage beaver damage on State Natural Areas to protect the state endangered Lamance iris (*Iris brevicaulis*) and the state threatened lake-bank sedge (*Carex lacustris*). Beaver dams can impound water and flood areas where those sensitive species occur. The consistent water levels maintained by beaver dams can be detrimental to those species.

Scientists estimate that nationwide cats kill hundreds of millions of birds and more than a billion small mammals, such as rabbits, squirrels, and chipmunks, each year. The American Bird Conservancy (ABC) states that "*cats often kill common* [bird] *species such as cardinals, blue jays, and house wrens, as well as rare and endangered species such as piping plovers, Florida scrub-jays, and California least terns*" (ABC 2011). Some feral and free-ranging cats kill more than 100 animals each year. For example, at a wildlife experiment station, a roaming, well-fed cat killed more than 1,600 animals over 18 months, primarily small mammals (ABC 2011). Researchers at the University of Wisconsin coupled their four-

year cat predation study with the data from other studies, and estimated that rural feral and free-ranging cats kill at least 7.8 million and perhaps as many as 217 million birds a year in Wisconsin (Coleman et al. 1997). In some parts of Wisconsin, feral and free ranging cat densities reached 114 cats per square mile, outnumbering all similar-sized native predators (Coleman et al. 1997). Churcher and Lawton (1989) observed 77 well fed free-ranging cats in a British village for one year. Churcher and Lawton (1989) estimated that 30% to 50% of a cat's catch were birds and that the cats had adversely affected house sparrow populations within the village. Based on information acquired in the study, Churcher and Lawton (1989) estimated that more than 20 million birds are killed by cats in Britain each year with more than 70 million animals overall being taken by cats annually.

The diet of feral and free-ranging cats varies depending on availability, abundance, and geographic location. In a survey of New Zealand scientific literature, Fitzgerald (1990) concluded that prey selection of feral and free-ranging cats is dependent on availability. Fitzgerald (1990) found that cats on the mainland fed most heavily on mammals; whereas, cats on islands fed almost exclusively on birds (particularly seabirds). Feral and free-ranging cats are known to prey on birds as large as mallard ducks (Figley and VanDruff 1982) and young brown pelicans (Anderson et al. 1989) along with mammals as large as hares and rabbits. Many cat populations rely heavily on humans either for handouts and/or for garbage. Pearson (1971) found that cats were serious predators of California voles and that the greatest pressure on voles occurred when vole numbers were lowest. Liberg (1984) found that cats in southern Sweden fed predominantly on native mammals. Prey use was based more on availability than abundance. Langham (1990) found that mammals made up 74% of diets of New Zealand farmland feral cats, while 24% were birds. Cats fed most heavily on the most abundant species and groups. A study on a southern Illinois farmstead concluded that well-fed cats preferred microtine rodents; however, they also consumed birds (George 1974). Microtine rodents are particularly susceptible to over harvest by cats and other predators (Pearson 1964). Coman and Brunner (1972) found that small mammals were the primary food item for feral cats in Victoria, Australia. Prey selection was directly related to proximity of cats to human habitation. Pearson (1964) found rodents composed a large portion of a cat's diet. Some people view cat predation of rodents as beneficial, but native small mammals are important to maintaining biologically diverse ecosystems. Field mice and shrews are also important prey for birds, such as great horned owls and red-tailed hawks.

Childs (1986) and Childs (1991) found that urban cats use of rats is size limiting. Few rats of reproductive size or age were preyed on by domesticated cats. In rural areas, rats were more vulnerable to cat predation for longer periods. The duration of susceptibility of rats to predation was attributed to abundance of garbage and artificial food sources in the urban environment. Artificial feeding of cats also reduces predation to non-native rodents because of size differences in urban rats. In rural setting, cats can control rat populations for longer durations but ultimate suppression of population growth is achieved via chemicals (poisons). Jackson (1951) found feral and free-ranging cats in Baltimore, Maryland urban areas were insignificant predators of Norway rats. The largest percentage of ingested food was comprised of garbage. It was estimated that a cat in the study area would consume roughly 28 rats per year.

Reptiles are thought to provide an important food source to cats when birds and mammals are less abundant, and in some situations, cats have been observed to prey on threatened species of reptiles. Domesticated cats have been identified as significant nest and/or hatchling predators of sea turtles. A study on the Aldabra Atoll, Seychelles found feral cats had an adverse effect on green turtle hatchlings. Seabrook (1989) found a positive correlation in cat activity and green turtle nesting at Aldabra Atoll. Cats are known to have contributed to the near extirpation of the West Indian rock iguana (*Cyclura carinata*) on Pine Cay in the Caicos Islands (Iverson 1978).

Cats can adversely affect local wildlife populations, especially in habitat "*islands*", such as suburban and urban parks, wildlife refuges, and other areas surrounded by human development (Wilcove 1985). The

loss of bird species from habitat islands is well documented and nest predation is an important cause of the decline of neotropical migrants (Wilcove 1985). A two-year study was conducted in two parks with grassland habitat. One park had no cats but more than 25 cats were being fed daily in the other park. There were almost twice as many birds seen in the park with no cats as in the park with cats. California thrasher and California quail, both ground-nesting birds, were seen during surveys in the no-cat area; whereas, they were never seen in the cat area. In addition, more than 85% of the native seer mice and harvest mice trapped were in the no-cat area; whereas, 79% of the house mice, an exotic pest species, were trapped in the cat area. The researchers concluded, "*Cats at artificially high densities, sustained by supplemental feeding, reduce abundance of native rodent and bird populations, change the rodent species composition, and may facilitate the expansion of the house mouse into new areas*" (Hawkins et al. 1999).

WS could also be requested to provide assistance associated with mammal damage at historical sites within the State. WS has previously been requested to provide assistance associated with woodchucks burrowing into earthen embankments at Civil War national historic sites in Tennessee. Woodchucks can cause extensive damage by burrowing and denning in earthen levees and other mounds. Burrowing activities can threaten the integrity of the earthen embankments. In addition, burrows can be aesthetically displeasing to the public and can cause damage to mowing equipment. In addition, there are thousands of archaeological and historical sites on TVA retained properties, some of which are extremely sensitive that could be disturbed by the burrowing and activities of mammals. Many of those sites, especially earthen mounds, have been damaged by the burrowing of woodchucks and could be damage by similar activities associated with nine-banded armadillos.

Feral swine compete with over 100 species of native wildlife for important and limited natural food supplies. Some species including quail, turkey, endangered sea turtles, and shorebirds are at risk of predation by nest destruction and the consuming of eggs. Feral swine cause damage to natural areas such as parks and wildlife management areas in Tennessee. Those sites suffer erosion and local loss of critical ground plants and roots, as well as destruction of seedlings because of feral swine feeding and rooting (Barrett and Birmingham 1994). Many state and federal natural resource managers are now in the process of controlling hog numbers because of their known impact to endangered plants and animals (Thompson 1977, West et al. 2009).

Feral swine are not native to North America, and many native species have not evolved to deal with swine competition or predation. Feral hogs are known to feed on many smaller animals (some threatened or endangered), disrupt ecosystems via rooting, and feed on rare and endangered plants. Many experts in the fields of botany and herpetology have observed marked declines in some rare species of plants, reptiles, amphibians, and soil invertebrates in areas inhabited by feral swine (Singer et al. 1984). It has been well documented that feral swine disturb large areas of vegetation and soil through rooting, and it is documented that hogs inhabiting coastal, upland, and wetland ecosystems are uprooting, damaging, and feeding on rare native species of plants and animals (Means 1999). Feral swine can disrupt natural vegetative communities, eliminate rare plants and animals, alter species composition within a forest, including both canopy and low growing species (Lipscomb 1989, Frost 1993), increase water turbidity in streams and wetlands (reducing water quality and impacting native fish), and increase soil erosion and alter nutrient cycling (Singer et al. 1984, DeBenedetti 1986). Kaller and Kelso (2003) found that feral and free-ranging swine were linked to increased levels of fecal coliform and other potentially pathogenic bacteria in several watersheds in Louisiana. Additionally, some species of freshwater mussels and aquatic insects were negatively affected by feral swine (Kaller and Kelso 2006).

Deer overabundance can affect native vegetation and natural ecosystems in addition to ornamental landscape plantings. White-tailed deer selectively forage on vegetation (Strole and Anderson 1992), and thus can have substantial impacts on certain herbaceous and woody species and on overall plant community structure (Waller and Alverson 1997). These changes can lead to adverse impacts on other

wildlife species, which depend on these plants for food and/or shelter. Numerous studies have shown that over browsing by deer can decrease tree reproduction, understory vegetation cover, plant density, and diversity (Warren 1991). By one count, 98 species of threatened and endangered plants, many of them orchids and lilies, are disturbed by deer browsing (Ness 2003).

The alteration and degradation of habitat from over-browsing by deer can have a detrimental effect on deer herd health and may displace other wildlife communities (*e.g.*, neotropical migrant songbirds and small mammals) that depend upon the understory vegetative habitat destroyed by deer browsing (Virginia Department of Game and Inland Fisheries 1999). Similarly, deCalesta (1997) reported that deer browsing affected vegetation that songbirds need for foraging surfaces, escape cover, and nesting. Species richness and abundance of intermediate canopy nesting songbirds was reduced in areas with higher deer densities (deCalesta 1997). Intermediate canopy-nesting birds declined 37% in abundance and 27% in species diversity at higher deer densities. Five species of birds were found to disappear at densities of 38.1 deer per square mile and another two disappeared at 63.7 deer per square mile. Casey and Hein (1983) found that three species of birds could no longer be found in a research preserve stocked with high densities of ungulates and that the densities of several other species of birds were lower than in an adjacent area with lower deer density. Waller and Alverson (1997) hypothesize that by competing with squirrels and other fruit-eating animals for oak mast, deer may further affect many other species of animals and insects.

Nutria may also cause damage to natural resources in Tennessee. Nutria are a non-native species in the United States that were introduced from South America. Nutria primarily inhabit brackish or freshwater marshes, but are also found in swamps, rivers, ponds, and lakes. They live in dense vegetation, in abandoned burrows, or in burrows they dig along stream banks or shorelines (Wade and Ramsey 1986). Nutria are almost entirely herbivorous and eat animal material (mostly insects) incidentally. Freshwater mussels and crustaceans are occasionally eaten by nutria in some parts of their range.

The digging and feeding behavior of nutria can be destructive to marsh ecosystems. Nutria forage directly on the vegetative root mat, leaving marshes pitted with digging sites and fragmented with deeply cut swimming canals. The denuding of marsh vegetation can accelerate erosion associated with tidal currents and wave action. The loss of vegetation can also facilitate salt-water intrusion into marsh interiors. Nutria also cause damage by eating lawn grasses found adjacent to aquatic habitats. Nutria are opportunistic feeders and eat approximately 25% of their body weight daily (LeBlanc 1994).

Need for Mammal Damage Management to Protect Property

Mammals cause damage to a variety of property types in Tennessee each year. Raccoons, skunks, and armadillos can cause damage to property by digging under porches, buildings, homes, and many other places. Armadillos often cause damage to lawns and turf while digging for grubs and insects. Beaver flood land, roads, and railways. They also girdle large trees and consume landscaping. Feral hogs root up turf in neighborhoods and golf courses.

During the six-year period, FY 2006 through FY 2011, complainants reported to WS over $5.6 million in property damages caused by mammals. These damages were caused by numerous species including beavers, woodchucks, opossums, raccoons, bats, squirrels, muskrats, voles, moles, roof rats, feral swine, and white-tailed deer.

Feral swine can damage landscaping, golf courses, roads, drainage ditches and cause erosion by feeding in these areas. Feral swine dig or root in the ground with their nose in search of desired roots, grubs, earthworms, and other food sources. The rooting and digging activity of feral swine turns sod and grass over which often leaves the area bare of vegetation and susceptible to erosion. Feral swine also pose a threat to property from being struck by motor vehicles and aircraft.

Deer-vehicle collisions are a serious concern nationwide because of losses to property and the potential for human injury and death (Conover et al. 1995, Romin and Bissonette 1996, Conover 1997). The economic costs associated with deer-vehicle collisions include vehicle repairs, human injuries and fatalities, and picking up and disposing of deer (Drake et al. 2005). The Insurance Institute for Highway Safety (2005) estimated that 1.5 million deer-vehicle collisions occur annually in the United States causing approximately 150 fatalities and $1.1 billion in damage to property. In 1995, the damage to vehicles associated with vehicles striking deer was estimated at $1,500 per strike in damages (Conover et al. 1995). Damage costs associated with deer collisions in 2011 were estimated at $3,171 per incident, which was an increase of 2.2% over the 2010 estimate (State Farm Mutual Automobile Insurance 2011*a*). An estimated 20,039 deer-vehicle collisions occurred in Tennessee from July 1, 2010 through June 30, 2011 (State Farm Mutual Automobile Insurance 2011*b*). Based on the average repair costs associated with vehicle strikes estimated at $3,171 in 2010 and the number of strikes that have occurred in the State estimated at 20,039 from July 2010 through June 2011, deer-vehicle collisions resulted in over $63.5 million in damage to property in the State.

Incidences of deer-vehicle collisions on highways passing through TVA Dam Reservation properties have been reported to TVA personnel in recent years from public stakeholders. Some of these dam reservation properties have elevated deer populations and WS could be request to provide assistance to reduce local deer populations on TVA properties.

Often, deer-vehicle collisions in which a deer carcass was not recovered or little vehicle damage occurred go unreported. A Cornell University study estimated that the actual number of deer-vehicle collisions could be as high as six times the reported number (Decker et al. 1990).

Beaver are generally considered beneficial where their activities do not compete with human land use or human health and safety (Wade and Ramsey 1986). The opinions and attitudes of individuals, organizations, and communities vary greatly and are primarily influenced and formed by the benefits and/or damage directly experienced by each individual (Hill 1982). Woodward et al. (1976) found that 24% of landowners who reported beaver activity on their property indicated benefits to having beaver ponds on their land and desired assistance with beaver pond management (Hill 1976, Lewis 1979, Woodward et al. 1985).

In some situations, the damage and threats caused by beaver outweigh the benefits (Grasse and Putnam 1955, Woodward et al. 1985, Novak 1987). Damage to resources associated with beaver are most often a result of their feeding, burrowing, and dam building behaviors. It is estimated that beaver cause $75 to $100 million dollars in economic losses annually in the United States, with total losses in the southeastern United States over the past 40 years estimated to be $4 billion (Novak 1987).

Beaver often will gnaw through trees and other woody vegetation for use in dam building, food caches, and the buildings of lodges. The girdling and felling of trees and other woody vegetation can cause economic losses, can threaten human safety and property when trees fall, and the loss of trees can be aesthetically displeasing to property owners. Timber resources have the highest recorded damage caused by beaver (Hill 1976, Lewis 1979, Hill 1982, Woodward et al. 1985). In some southeastern states, losses from beaver damage have been estimated at $3 million to $5 million dollars annually (Miller and Yarrow 1994), with timber losses as the most common type of damage (Hill 1982). Tracts of bottomland hardwood timber up to several thousand acres in size may be lost to beaver activity (Miller and Yarrow 1994). Timber damage caused by beaver activity in the southeastern United States has been estimated at $2.2 million annually in Mississippi (Arner and Dubose 1982), $2.2 million in Alabama (Hill 1976), $45 million in Georgia (Godbee and Price 1975), and $14.5 million in Louisiana in 1993 (Fowler et al. 1994).

In addition to damage associated with beaver feeding and gnawing on trees, damages and threats can also occur from dam building activities. Beaver dams impound water, which can flood property resulting in economic damage. Flooding from beaver dams can cause damage to roads, impede traffic, inundate timber, weaken earthen embankments, and cause damage to residential and commercial utilities. In addition, beaver dams constructed on TVA property can result in flooding of adjacent private property, which often creates issues that demand a response from TVA.

Beaver often inhabit sites in or adjacent to urban/suburban areas and cut or girdle trees and shrubs in yards, undermine yards and walkways by burrowing, flood homes and other structures, destroy pond and reservoir dams by burrowing into levees, gnaw on boat houses and docks, and cause other damage to private and public property (Wade and Ramsey 1986). Additionally, roads and railroads may be damaged by saturation of the roadbed from beaver flooding or by beaver burrowing into the banks that comprise roadbeds and railroad beds.

Burrowing activities of nutria can severely damage levees, dikes, earthen dams, and other structures (Federal Emergency Management Agency 2005). Additionally, nutria burrows can weaken flood control levees that protect low-lying areas. In some cases, tunneling in levees is so extensive that water will flow unobstructed from one side to the other, necessitating their complete reconstruction. Nutria can also burrow into the Styrofoam flotation under boat docks and wharves, causing those structures to lean and sink. Nutria burrow under buildings, which may lead to uneven settling or failure of the foundations. Burrows can weaken road beds, stream banks, dams, and dikes, which may collapse when the soil is saturated by rain or high water or when subjected to heavy objects on the surface (such as vehicles, farm machinery, or grazing livestock). Rain and wave action can wash out and enlarge collapsed burrows, which compounds the damage.

Burrowing activities of woodchucks can also severely damage levees, dikes, earthen dams, landfills, and other structures (Federal Emergency Management Agency 2005). Woodchucks burrowing into roadbeds and embankments could potentially weaken or cause the collapse of these structures. Woodchucks also cause damage by chewing underground utility cables, sometimes resulting in power outages. Additionally, woodchuck burrows may cause damage to property when tractors and other equipment drop into a burrow or roll over due to a burrow.

1.3 SCOPE OF THIS ENVIRONMENTAL ASSESSMENT

Actions Analyzed

This EA evaluates the need for mammal damage management to reduce threats to human safety and to resolve damage to property, natural resources, and agricultural resources on federal, state, tribal, municipal, and private land within the State of Tennessee wherever such management is requested by a cooperator (USDA 2005*a*). This EA discusses the issues associated with conducting mammal damage management in the State to meet the need for action and evaluates different alternatives to meeting that need while addressing those issues.

The methods available for use or recommendation under each of the alternatives evaluated are provided in Appendix B[7]. The alternatives and Appendix B also discuss how methods would be employed to manage damage and threats associated with mammals in the State. Therefore, the actions evaluated in this EA are the use of those methods available under the alternatives and the employment of those methods by WS to

[7]A complete list of chemical and non-chemical methods available for use under the identified alternatives can be found in Appendix B. However, listing methods neither implies that all methods would be used by WS to resolve requests for assistance nor does the listing of methods imply that all methods would be used to resolve every request for assistance.

manage or prevent damage and threats associated with mammals from occurring when requested by the appropriate resource owner or manager. WS' activities that could involve the take of mammals under the alternatives would only occur when agreed upon by the requester and when permitted by the TWRA, when required, and only at levels permitted.

Federal, State, County, City, and Private Lands

Under two of the alternatives, WS could continue to provide damage management activities on federal, state, county, municipal, and private land in Tennessee when a request is received for such services by the appropriate resource owner or manager. In those cases where a federal agency requests WS' assistance with managing damage caused by mammals, the requesting agency would be responsible for analyzing those activities in accordance with the NEPA. However, this EA could cover such actions if the requesting federal agency determined the analyses and scope of this EA were appropriate for those actions and the requesting federal agency adopted this EA through their own Decision based on the analyses in this EA. Therefore, actions taken on federal lands have been analyzed in the scope of this EA.

Native American Lands and Tribes

The WS program in Tennessee would only conduct damage management activities on Native American lands when requested by a Native American Tribe. Activities would only be conducted after a Memorandum of Understanding (MOU) or cooperative service agreement had been signed between WS and the Tribe requesting assistance. Therefore, the Tribe would determine when WS' assistance was required and what activities would be allowed. Because Tribal officials would be responsible for requesting assistance from WS and determining what methods would be available to alleviate damage, no conflict with traditional cultural properties or beliefs would be anticipated. Those methods available to alleviate damage associated with mammals on federal, state, county, municipal, and private properties under the alternatives analyzed in this EA would be available for use to alleviate damage on Tribal properties when the use of those methods had been approved for use by the Tribe requesting WS' assistance. Therefore, the activities and methods addressed under the alternatives would include those activities that could be employed on Native American lands, when requested and when agreed upon by the Tribe and WS.

Period for which this EA is Valid

If the analyses in this EA indicate an Environmental Impact Statement (EIS) is not warranted, this EA would remain valid until WS and the TVA, in consultation with the TWRA, determined that new needs for action, changed conditions, new issues, or new alternatives having different environmental impacts must be analyzed. At that time, this analysis and document would be reviewed and supplemented pursuant to the NEPA. Review of the EA would be conducted to ensure that activities conducted under the selected alternative occur within the parameters evaluated in the EA. If the alternative analyzing no involvement in mammal damage activities by WS were selected, no additional analyses would occur based on the lack of involvement by WS. The monitoring of activities by WS would ensure the EA remained appropriate to the scope of damage management activities conducted by WS in Tennessee under the selected alternative, including activities conducted on TVA properties, when requested.

Site Specificity

As mentioned previously, WS would only conduct damage management activities when requested by the appropriate resource owner or manager. In addition, WS' activities that could involve the take of mammals under the alternatives would only occur when permitted by the TWRA, when required, and only at levels permitted.

This EA analyzes the potential impacts of mammal damage management based on previous activities conducted on private and public lands in Tennessee where WS and the appropriate entities entered into a MOU, cooperative service agreement, or other comparable document. The EA also addresses the potential impacts of mammal damage management in areas where additional agreements may be signed in the future. Because the need for action is to reduce damage and because the program's goals and directives would be to provide services when requested, within the constraints of available funding and workforce, it is conceivable that additional damage management efforts could occur. Thus, this EA anticipates receiving additional requests for assistance and analyzes the impacts of the efforts to resolve those additional requests as part of the alternatives.

Many of the mammal species addressed in this EA can be found statewide and throughout the year in the State; therefore, damage or threats of damage can occur wherever those mammals occur. Planning for the management of mammal damage must be viewed as being conceptually similar to the actions of other entities whose missions are to stop or prevent adverse consequences from anticipated future events for which the actual sites and locations where they would occur are unknown but could be anywhere in a defined geographic area. Examples of such agencies and programs include fire and police departments, emergency clean-up organizations, and insurance companies. Although some of the locations where mammal damage would occur can be predicted, not all specific locations or times where such damage would occur in any given year can be predicted. In addition, the threshold triggering an entity to request assistance from WS to manage damage associated with mammals is often unique to the individual; therefore, predicting where and when such a request for assistance would be received by WS is difficult. This EA emphasizes major issues as those issues relate to specific areas whenever possible; however, many issues apply wherever mammal damage and the resulting management actions occurs and are treated as such.

Chapter 2 of this EA identifies and discusses issues relating to mammal damage management in Tennessee. The standard WS Decision Model (Slate et al. 1992) would be the site-specific procedure for individual actions conducted by WS in the State (see Chapter 3 for a description of the Decision Model and its application). Decisions made using the model would be in accordance with WS' directives and Standard Operating Procedures (SOPs) described in this EA as well as relevant laws and regulations.

The analyses in this EA are intended to apply to any action that may occur in any locale and at any time within Tennessee. In this way, WS believes it meets the intent of the NEPA with regard to site-specific analysis and that this is the only practical way for WS to comply with the NEPA and still be able to accomplish its mission.

Summary of Public Involvement

Issues related to mammal damage management as conducted by WS in Tennessee were initially developed by WS in cooperation with the TVA and during consultation with the TWRA. Issues were defined and preliminary alternatives were identified through the scoping process. As part of this process, and as required by the Council on Environmental Quality (CEQ) and APHIS' NEPA implementing regulations, this document will be noticed to the public for review and comment. The public will be noticed through legal notices published in local print media, through direct mailings to parties that have requested to be notified, or have been identified to have an interest in the reduction of threats and damage associated with mammals in the State, and by posting the EA on the APHIS website at http://www.aphis.usda.gov/wildlife_damage/nepa.shtml.

WS will provide for a minimum of a 30-day comment period for the public and interested parties to provide new issues, concerns, and/or alternatives. Through the public involvement process, WS will

clearly communicate to the public and interested parties the analyses of potential environmental impacts on the quality of the human environment. New issues or alternatives raised after publication of public notices would be fully considered to determine whether the EA should be revisited and, if appropriate, revised prior to issuance of a final Decision or publication of a notice of intent to prepare an EIS.

1.4 RELATIONSHIP OF THIS EA TO OTHER ENVIRONMENTAL DOCUMENTS

WS' Programmatic Final Environmental Impact Statement: WS has developed a programmatic FEIS that addresses the need for wildlife damage management in the United States (USDA 1997). The FEIS contains detailed discussions of potential impacts to the human environment from methods that could be used by WS to alleviate wildlife damage. In addition, the FEIS contains risk assessments of many of the methods available to manage damage caused by mammals in the State (USDA 1997).

WS' Environmental Assessment - Mammal Damage Management in the Tennessee Wildlife Services Program: As was stated previously, WS, in cooperation with the TVA, previously developed an EA that addressed WS' activities to manage damage associated with mammals in the State (USDA 2005*a*). Based on the analyses in that EA, a Decision and Finding of No Significant Impact was signed by WS selecting the proposed action alternative. The proposed action alternative implemented a damage management program using a variety of methods in an integrated approach (USDA 2005*a*). Changes in the need for action and the affected environment have prompted WS and the TVA to initiate this new analysis to address mammal damage in the State. This EA will address more recently identified changes and will assess the potential environmental impacts of program alternatives based on a new need for action. Since activities conducted under the previous EA will be re-evaluated under this EA to address the new need for action and the associated affected environment, the previous EA that addresses mammal damage will be superseded by this analysis and the outcome of the Decision issued based on the analyses in this EA. However, information in the previous EA associated with the need for action relative to mammals continues to be appropriate to the need for action associated with this EA (USDA 2005*a*).

WS' Supplemental Environmental Assessment – Oral Vaccination to Control Specific Rabies Virus Variants in Raccoons, Gray Fox, and Coyotes in the United States: WS issued an EA that analyzed the environmental effects of WS' involvement in the funding of and participation in Oral Rabies Vaccination programs to eliminate or stop the spread of raccoon rabies in a number of eastern states (including Tennessee) and gray fox and coyote rabies in Texas (USDA 2005*b*). The EA has been supplemented to analyze changes in the scope and analysis area of the ORV program. The most recent Decision/FONSI was signed in 2010. WS determined the action would not have any significant impact on the quality of the human environment. Pertinent information has been incorporated by reference into this EA.

TVA's Natural Resource Plan (NRP): TVA has developed an extensive plan to strategically evaluate both renewable and nonrenewable resources and fulfill the responsibilities associated with good stewardship of TVA lands and resources. The NRP is designed to integrate the objectives of six resource areas (biological, cultural, recreation, water, public engagement and reservoir lands planning); provide optimum public use benefit; and balance competing and sometimes conflicting resource uses (TVA 2011*a*).

TVA's Environment Impact Statement Assessing the Natural Resource Plan: TVA has also prepared an EIS to assess the impacts of the NRP and its reasonable alternatives on the environment. It specifically describes the stewardship programs that are ongoing and are being evaluated for future implementation as part of the NRP; and assesses the potential environmental impacts associated with implementing the various alternatives. Pertinent information available in the FEIS has been incorporated by reference into this EA (TVA 2011*b*).

TWRA Comprehensive Wildlife Conservation Strategy (CWCS): TWRA has developed an extensive wildlife conservation plan that evaluates all species of plant and animal known to exist within the State. This plan identifies all of the species and habitats that are currently listed as endangered, threatened, or species of concern, both federally by the United States Fish and Wildlife Service (USFWS; USFWS 2011) and at the state level by the TWRA and the Tennessee Department of Environment and Conservation's (TDEC)-Natural Heritage Inventory Program (TDEC 2008, TDEC 2009). It also incorporates additional species of which little is known or with questionable population trends, and creates a comprehensive prioritized list of species in need of conservation. This CWCS was consulted as part of this analysis and no species found in the CWCS will be considered for management herein.

1.5 AUTHORITY OF FEDERAL AND STATE AGENCIES

The authorities of WS and other agencies as those authorities relate to conducting wildlife damage management activities are discussed by agency below:

WS' Legislative Authority

The primary statutory authority for the WS program is the Act of March 2, 1931 (46 Stat. 1468; 7 USC 426-426b) as amended, and the Act of December 22, 1987 (101 Stat. 1329-331, 7 USC 426c). The WS program is the lead federal authority in managing damage to agricultural resources, natural resources, property, and threats to human safety associated with wildlife. WS' directives define program objectives and guide WS' activities in managing wildlife damage.

Tennessee Valley Authority

The TVA is a federal corporation created by an Act of Congress in May 18, 1933 [48 Stat. 58-59, 16 USC Sec. 831, as amended]. TVA provides electricity to 9 million people, businesses and industries, and manages 293,000 acres of public land and 11,000 miles of reservoir shoreline in the seven-state Tennessee Valley Region (Tennessee, Alabama, Mississippi, Kentucky, Georgia, North Carolina, and Virginia – an area of 80,000 square miles). TVA also provides flood control, navigation, land management, and recreation for the Tennessee River system and works with local utilities and state and local governments to promote economic development across the region.

The TVA operates 21 hydroelectric dams, seven coal-fired power plants, two nuclear power plants, 10 solar facilities, one wind turbine facilities, and four combustion turbine sites in Tennessee. TVA also owns or maintains 262 substations and switching stations and nearly 9,400 circuit miles of transmission line and rights-of-way easements in Tennessee.

In addition, TVA manages 21 reservoirs in Tennessee with more than 7,500 miles of shoreline. Along and over most of these Tennessee River and tributary streams, TVA owns 175,000 acres of shore-land and manages various other land rights. The TVA conducts and requests assistance from WS to provide wildlife damage management on land and at facilities owned by the TVA. TVA also makes its public lands available for use for continuation and expansion of the WS Oral Rabies Vaccination program across the Tennessee River Basin and Valley states.

United States Environmental Protection Agency (EPA)

The EPA is responsible for implementing and enforcing the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA), which regulates the registration and use of pesticides.

Tennessee Wildlife Resources Agency

The TWRA authority in wildlife management is given within the Tennessee Code Annotated Section 70-1-1 et seq., the official regulations of the Tennessee Fish and Wildlife Commission and applicable federal laws. This legislation covers general provisions; licenses, permits and stamps; wildlife; fish; and wild animals.

Tennessee Department of Agriculture (TDA)

The Pesticide Program of the TDA enforces state laws pertaining to the use and application of pesticides. Under the Tennessee Application of Pesticide Act (Sections 62-21-101 through 62-21-131), this section monitors the use of pesticides in a variety of pest management situations. It also licenses private and commercial pesticide applicators and pesticide contractors. Under this Act (Section 62-21-115 through 62-21-127) the program licenses restricted use pesticide dealers and registers all pesticides for sale and distribution in the state of Tennessee.

Tennessee Department of Health (TDH)

The TDH authority in public health is given within the Tennessee Code Annotated Section 68-1-1 et seq., the official regulations of the Tennessee Department of Health. This legislation covers various aspects of disease management and prevention including rabies (Section 68-8-101 through 68-8-113) among other zoonotic diseases.

1.6 COMPLIANCE WITH LAWS AND STATUTES

Several laws or statutes would authorize, regulate, or otherwise affect WS' activities under the alternatives. WS would comply with all applicable federal, state, and local laws and regulations in accordance with WS Directive 2.210. Those laws and regulations related to activities conducted to reduce mammal damage in the State are addressed below:

National Environmental Policy Act

All federal actions are subject to the NEPA (Public Law 9-190, 42 USC 4321 et seq.). WS follows CEQ regulations implementing the NEPA (40 CFR 1500 et seq.) along with USDA (7 CFR 1b) and APHIS Implementing Guidelines (7 CFR 372) as part of the decision-making process. These laws, regulations, and guidelines generally outline five broad types of activities to be accomplished as part of any project: public involvement, analysis, documentation, implementation, and monitoring. The NEPA also sets forth the requirement that all major federal actions be evaluated in terms of their potential to significantly affect the quality of the human environment for the purpose of avoiding or, where possible, mitigating and minimizing adverse impacts. Federal activities affecting the physical and biological environment are regulated in part by CEQ through regulations in 40 CFR, Parts 1500-1508. In accordance with CEQ and USDA regulations, APHIS guidelines concerning Implementation of the NEPA Procedures, as published in the Federal Register (44 CFR 50381-50384) provide guidance to APHIS regarding the NEPA process.

Pursuant to the NEPA and CEQ regulations, this EA documents the analyses resulting from proposed federal actions, informs decision-makers and the public of reasonable alternatives capable of avoiding or minimizing adverse impacts, and serves as a decision-aiding mechanism to ensure that the policies and goals of the NEPA are infused into federal agency actions. This EA was prepared by integrating as many of the natural and social sciences as warranted, based on the potential effects of the proposed action. The direct, indirect, and cumulative impacts of the alternatives are analyzed.

Endangered Species Act

Under the ESA, all federal agencies will seek to conserve T&E species and will utilize their authorities in furtherance of the purposes of the Act (Sec.2(c)). WS conducts Section 7 consultations with the USFWS to use the expertise of the USFWS to ensure that "*any action authorized... funded or carried out by such an agency . . . is not likely to jeopardize the continued existence of any endangered or threatened species . . . Each agency will use the best scientific and commercial data available*" (Sec.7 (a)(2)).

Evaluation of the alternatives in regards to the ESA will occur in Chapter 4 of this EA.

National Historic Preservation Act (NHPA) of 1966, as amended

The NHPA and its implementing regulations (36 CFR 800) require federal agencies to initiate the section 106 process if an agency determines that the agency's actions are undertakings as defined in Sec. 800.16(y) and, if so, whether it is a type of activity that has the potential to cause effects on historic properties. If the undertaking is a type of activity that does not have the potential to cause effects on historic properties, assuming such historic properties were present, the agency official has no further obligations under section 106. None of the mammal damage management methods described in this EA that might be used operationally by WS would cause major ground disturbance, any physical destruction or damage to property, any alterations of property, wildlife habitat, or landscapes, nor involves the sale, lease, or transfer of ownership of any property. In general, such methods also do not have the potential to introduce visual, atmospheric, or audible elements to areas in which they are used that could result in effects on the character or use of historic properties. Therefore, the methods that would be used by WS under the alternatives would not generally be the types of activities that would have the potential to affect historic properties. If an individual activity with the potential to affect historic resources were planned under an alternative selected because of a decision on this EA, the site-specific consultation as required by Section 106 of the NHPA would be conducted as necessary.

Noise-making methods, such as firearms, that are used at or in close proximity to historic or cultural sites for the purposes of removing wildlife have the potential for audible effects on the use and enjoyment of historic property. However, such methods would only be used at a historic site at the request of the owner or manager of the site to resolve a damage problem, which means such use, would be to the benefit of the historic property. A built-in minimization factor for this issue is that virtually all the methods involved would only have temporary effects on the audible nature of a site and could be ended at any time to restore the audible qualities of such sites to their original condition with no further adverse effects. Site-specific consultation as required by the Section 106 of the NHPA would be conducted as necessary in those types of situations.

Environmental Justice in Minority and Low Income Populations - Executive Order 12898

Executive Order 12898 promotes the fair treatment of people of all races, income levels, and cultures with respect to the development, implementation, and enforcement of environmental laws, regulations, and policies. Environmental justice is the pursuit of equal justice and protection under the law for all environmental statutes and regulations without discrimination based on race, ethnicity, or socioeconomic status. Executive Order 12898 requires federal agencies to make environmental justice part of their mission, and to identify and address disproportionately high and adverse human health and environmental effects of federal programs, policies, and activities on minority and low-income persons or populations. All activities are evaluated for their impact on the human environment and compliance with Executive Order 12898.

WS would use only legal, effective, and environmentally safe damage management methods, tools, and approaches. All chemicals used by WS and the TVA would be regulated by the EPA through the FIFRA, by the TDA, by the Drug Enforcement Agency (DEA), by MOUs with land managing agencies, and by WS' Directives. WS would properly dispose of any excess solid or hazardous waste. It is not anticipated that the proposed action would result in any adverse or disproportionate environmental impacts to minority and low-income persons or populations. In contrast, the alternatives may benefit minority or low-income populations by reducing threats to public health and safety and property damage.

Protection of Children from Environmental Health and Safety Risks - Executive Order 13045

Children may suffer disproportionately for many reasons from environmental health and safety risks, including the development of their physical and mental status. WS and the TVA make it a high priority to identify and assess environmental health and safety risks that may disproportionately affect children. WS and the TVA have considered the impacts that this proposal might have on children. The proposed activities would occur by using only legally available and approved methods where it is highly unlikely that children would be adversely affected. For these reasons, WS concludes that it would not create an environmental health or safety risk to children from implementing this proposed action. Additionally, the need for action identified a need to reduce threats to human safety, including risks to children; therefore, it would be expected that health and safety risks to children posed by mammals would be reduced under the alternatives.

Invasive Species - Executive Order 13112

Executive Order 13112 establishes guidance to federal agencies to use their programs and authorities to prevent the spread or to control populations of invasive species that cause economic or environmental harm or harm to human health. The Order states that each federal agency whose actions may affect the status of invasive species shall, to the extent practicable and permitted by law: 1) reduce invasion of exotic species and the associated damages, 2) monitor invasive species populations and provide for restoration of native species and habitats, 3) conduct research on invasive species and develop technologies to prevent introduction, and 4) provide for environmentally sound control and promote public education of invasive species.

The Native American Graves and Repatriation Act of 1990

The Native American Graves Protection and Repatriation Act (Public Law 101-106, 25 USC 3001) requires federal agencies to notify the Secretary of the Department that manages the federal lands upon the discovery of Native American cultural items on federal or tribal lands. Federal projects would discontinue work until a reasonable effort has been made to protect the items and the proper authority has been notified.

Federal Insecticide, Fungicide, and Rodenticide Act

The FIFRA and its implementing regulations (Public Law 110-426, 7 USC 136 et. seq.) require the registration, classification, and regulation of all pesticides used in the United States. The EPA is responsible for implementing and enforcing the FIFRA. All chemical methods integrated into the WS' program in Tennessee, including the use of or recommendation of repellents are registered with and regulated by the EPA and the TDA, and used or recommended by WS in compliance with labeling procedures and requirements.

Occupational Safety and Health Act of 1970

The Occupational Safety and Health Act of 1970 and its implementing regulations (29 CFR 1910) on sanitation standards states that, "*Every enclosed workplace shall be so constructed, equipped, and maintained, so far as reasonably practical, as to prevent the entrance or harborage of rodents, insects, and other vermin. A continuing and effective extermination program shall be instituted where their presence is detected*." This standard includes mammals that may cause safety and health concerns at workplaces.

Federal Food, Drug, and Cosmetic Act (21 USC 360)

This law places administration of pharmaceutical drugs, including those used in wildlife capture and handling, under the Food and Drug Administration.

Controlled Substances Act of 1970 (21 USC 821 et seq.)

This law requires an individual or agency to have a special registration number from the federal DEA to possess controlled substances, including those that are used in wildlife capture and handling.

Animal Medicinal Drug Use Clarification Act of 1994

The Animal Medicinal Drug Use Clarification Act (AMDUCA) and its implementing regulations (21 CFR 530) establish several requirements for the use of animal drugs, including those used to capture and handle wildlife in damage management programs. Those requirements are: (1) a valid "*veterinarian-client-patient*" relationship, (2) well defined record keeping, (3) a withdrawal period for animals that have been administered drugs, and (4) identification of animals. A veterinarian, either on staff or on an advisory basis, would be involved in the oversight of the use of animal capture and handling drugs under any alternative where WS could use those immobilizing and euthanasia drugs. Veterinary authorities in each state have the discretion under this law to establish withdrawal times (*i.e.*, a period after a drug is administered that must lapse before an animal may be used for food) for specific drugs. Animals that might be consumed by a human within the withdrawal period must be identified. WS establishes procedures in each state for administering drugs used in wildlife capture and handling that must be approved by state veterinary authorities in order to comply with this law.

Clean Water Act (Section 404)

Section 404 (33 USC 1344) of the Clean Water Act prohibits the discharge of dredged or fill material into waters of the United States without a permit from the United States Army Corps of Engineers unless the specific activity is exempted in 33 CFR 323 or covered by a nationwide permit in 33 CFR 330. The breaching of most beaver dams is covered by these regulations (33 CFR 323 and 330).

Food Security Act

The Wetland Conservation provision (Swampbuster) of the 1985 (16 USC 3801-3862), 1990 (as amended by PL 101-624), and 1996 (as amended by Public Law 104-127) farm bills require all agricultural producers to protect wetlands on the farms they own. Wetlands converted to farmland prior to December 23, 1985 are not subject to wetland compliance provisions even if wetland conditions return because of lack of maintenance or management. If prior converted cropland is not planted to an agricultural commodity (crops, native and improved pastures, rangeland, tree farms, and livestock production) for more than 5 consecutive years and wetland characteristics return, the cropland is considered abandoned and then becomes a wetland subject to regulations under Swampbuster and Section 404 of the Clean

Water Act. The Natural Resource Conservation Service is responsible for certifying wetland determinations according to this Act.

Airborne Hunting Act

The Airborne Hunting Act, passed in 1971 (Public Law 92-159), and amended in 1972 (Public Law 92-502) added to the Fish and Wildlife Act of 1956 as a new section (16 USC 742j-l) that prohibits shooting or attempting to shoot, harassing, capturing or killing any bird, fish, or other animal from aircraft except for certain specified reasons. Under exception [16 USC 742j-l, (b)(1)], State and Federal agencies are allowed to protect or aid in the protection of land, water, wildlife, livestock, domesticated animals, human life, or crops using aircraft.

Title 70 of the Tennessee Code Annotated and Other State Regulations

Title 70 of the Tennessee Code Annotated (TCA) and the annual TWRA Hunting and Trapping Guide (TWRA 2012*a*) provide the state laws and regulations pertaining to take of mammals within the State. The individual laws and regulations that directly pertain to damage management that would regulate activities conducted by WS would be:

TCA § 70-1-101 (a)(10) defines the "*Cushion-hold trap*" as an approved trap that is spring-loaded with offset jaws and designed to capture an animal by closing upon one of its legs so that the edges designed to touch the animal are composed of a non-metallic substance that eliminates or mitigates injury to the trapped animal. Specific traps and sizes are to be identified by the TWRA commission in its annual Hunting and Trapping Guide. The most recent regulations printed in the 2012 Hunting and Trapping Guide state that all steel leg-hold traps must have an outside measurement of nine inches or less at the widest point. Steel square instant-kill traps must have an exterior jaw measurement of 16 inches or less, and steel circular instant-kill traps must have an exterior measurement of 12 inches or less. In addition, steel cable snares must have a cable diameter between 5/64 inch and 3/32 inch and have a tag affixed bearing the name of the owner (TWRA 2012*a*).

TCA § 70-4-120 outlines the laws pertaining specifically to trapping, snaring, and baiting. Namely, it is unlawful to place steel traps in the open, except in water sets. Cushion-hold traps can be used in the open during the open season for the target species and with written permission from the land owner [TCA § 70-4-120 (a)(1)(A)]. All traps must be inspected within each 36 hours and any animal caught must be removed [TCA § 70-4-120 (a)(1)(B)]. In addition, all traps must be stamped with the owner's name and fashioned so that the name is legible at all times [TCA § 70-4-120 (a)(1)(C)].

1.7 DECISIONS TO BE MADE

Based on agency relationships, MOUs, and legislative authorities, WS is the lead agency for this EA, and therefore, responsible for the scope, content, and decisions made. The TVA owns and operates numerous electrical power generation sites and transmission structures within Tennessee, including electrical substations and transmission lines. In addition, the TVA manages lands within the State for recreational, natural, and cultural resources. Many of these sites experience damage associated with mammals within the State. The TVA would be the primary decision-maker for mammal damage management activities occurring on sites owned or managed by the TVA. As the authority for the management of mammal populations in the State, the TWRA was involved in the development of the EA and provided input throughout the EA preparation process to ensure an interdisciplinary approach according to the NEPA and agency mandates, policies, and regulations. The TWRA is responsible for managing wildlife in the State of Tennessee, including those mammals addressed in this EA. The TWRA establishes and enforces regulated hunting and trapping seasons in the State. WS' activities to reduce and/or prevent mammal

damage in the State would be coordinated with the TWRA, which ensures WS' actions would be incorporated into population objectives established for mammal populations in the State.

Based on the scope of this EA, the decisions to be made are: 1) should WS, in cooperation with the TVA, conduct mammal damage management to alleviate damage to agriculture, property, natural resources, and threats to human safety, 2) should WS conduct disease surveillance and monitoring in mammal populations when requested by the TWRA and other agencies, 3) should WS, in cooperation with the TVA, implement an integrated wildlife damage management strategy, including technical assistance and direct operational assistance, to meet the need for mammal damage management in Tennessee, 4) if not, should WS attempt to implement one of the alternatives to an integrated damage management strategy as described in the EA, and 5) would the proposed action or the other alternatives result in adverse impacts to the environment requiring the preparation of an EIS.

CHAPTER 2: AFFECTED ENVIRONMENT AND ISSUES

Chapter 2 contains a discussion of the issues, including issues that will receive detailed environmental impact analysis in Chapter 4 (Environmental Consequences), issues that have driven the development of SOPs, and issues that will not be considered in detail, with rationale. Pertinent portions of the affected environment will be included in this chapter in the discussion of issues. Additional descriptions of the affected environment will be incorporated into the discussion of the environmental effects in Chapter 4.

2.1 AFFECTED ENVIRONMENT

Damage or threats of damage caused by those mammal species addressed in this EA can occur statewide in Tennessee wherever those mammals occur (USDA 2005*a*). However, mammal damage management would only be conducted by WS when requested by a landowner or manager and only on properties where a cooperative service agreement or other comparable document were signed between WS and a cooperating entity. Most species of mammals addressed in this EA can be found throughout the year across the State where suitable habitat exists for foraging and shelter. Those mammal species addressed in this EA are capable of utilizing a variety of habitats in the State. Since those mammal species addressed in this EA can be found throughout most of the State, requests for assistance to manage damage or threats of damage could occur in areas occupied by those mammal species. Additional information on the affected environment is provided in Chapter 4.

Upon receiving a request for assistance, activities to reduce mammal damage or threats could be conducted on federal, state, tribal, municipal, and private properties in Tennessee. Areas where damage or threats of damage could occur include, but would not be limited to agricultural fields, vineyards, orchards, farmyards, dairies, ranches, livestock operations, aquaculture facilities, fish hatcheries, grain mills, grain handling areas, railroad yards, waste handling facilities, industrial sites, natural resource areas, park lands, and historic sites; state and interstate highways and roads; railroads and their right-of-ways; property in or adjacent to subdivisions, businesses, and industrial parks; timberlands, croplands, and pastures; private and public property where burrowing mammals cause damage to structures, dikes, ditches, ponds, and levees; public and private properties in rural/urban/suburban areas where mammals cause damage to landscaping and natural resources, property, and are a threat to human safety through the spread of disease. The area would also include airports and military airbases where mammals are a threat to human safety and to property; areas where mammals negatively affect wildlife, including T&E species; and public property where mammals are negatively affecting historic structures, cultural landscapes, and natural resources.

In addition, mammal damage management could occur at facilities owned or managed by the TVA when those mammal species addressed in this assessment cause damage or pose threats of damage to property,

natural resources, pose a threat to human safety, or threaten the reliability of electric system transmission. Damage management activities could be conducted at any of the 48 TVA power generation facilities, 261 TVA electrical substations, or along any of the 10,200 circuit miles of transmission lines and right-of-way easements owned by the TVA in Tennessee. Mammal damage management activities could also be conducted on recreational, natural, or cultural lands owned or managed by the TVA, including 21 reservoirs with more than 7,500 miles of shoreline, and 175,000 acres of shore-land and reservoir property and various other land rights along and over most the Tennessee River and its tributaries.

Environmental Status Quo

As defined by the NEPA implementing regulations, the "*human environment shall be interpreted comprehensively to include the natural and physical environment and the relationship of people with that environment*" (40 CFR 1508.14). Therefore, when a federal action agency analyzes their potential impacts on the "*human environment*", it is reasonable for that agency to compare not only the effects of the federal action, but also the potential impacts that occur or could occur in the absence of the federal action by a non-federal entity. This concept is applicable to situations involving federal assistance to reduce damage associated with wildlife species.

Unprotected wildlife species, such as most non-native invasive species, are not protected under state or federal law. Most state-resident wildlife species are managed under state authority or law without any federal oversight or protection. In some situations, with the possible exception of restrictions on methods (*e.g.*, firearms restrictions, pesticide regulations), unprotected wildlife species and certain resident wildlife species are managed with little or no restrictions, which allows them to be killed or taken by anyone at any time when they are committing damage. For mammal damage management in Tennessee, the TWRA has the authority to manage and authorize the taking of mammals for damage management purposes.

When a non-federal entity (*e.g.*, agricultural producers, municipalities, counties, private companies, individuals, or any other non-federal entity) takes an action to alleviate mammal damage or threat, the action is not subject to compliance with the NEPA due to the lack of federal involvement in the action. Under such circumstances, the environmental baseline or status quo must be viewed as an environment that includes those resources as they are managed or impacted by non-federal entities in the absence of the federal action being proposed. Therefore, in those situations in which a non-federal entity has decided that a management action directed towards mammals should occur and even the particular methods that should be used, WS' involvement in the action would not affect the environmental status quo since the entity could take the action in the absence of WS' involvement. WS' involvement would not change the environmental status quo if the requestor had conducted the action in the absence of WS' involvement in the action.

A non-federal entity could lethally take mammals to alleviate damage without the need for a permit when those species are non-native or are unregulated by the TWRA. In addition, mammals could be removed to alleviate damage during the hunting and/or trapping season, and/or through the issuance of permits by the TWRA. In addition, most methods available for resolving damage associated with mammals would also be available for public use. Therefore, WS' decision-making ability would be restricted to one of three alternatives. WS could take the action using the specific methods as decided upon by the non-federal entity, provide technical assistance only, or take no action. If no action were taken by WS, the non-federal entity could take the action anyway using the same methods without the need for a permit, during the hunting or trapping season, or through the issuance of a permit by the TWRA. Under those circumstances, WS would have virtually no ability to affect the environmental status quo since the action would likely occur in the absence of WS' direct involvement.

Therefore, based on the discussion above, it is clear that in those situations where a non-federal entity has obtained the appropriate permit or authority, and has already made the decision to remove or otherwise manage mammals to stop damage with or without WS' assistance, WS' participation in carrying out the action would not affect the environmental status quo.

2.2 ISSUES ASSOCIATED WITH MAMMAL DAMAGE MANAGEMENT ACTIVITIES

Issues are concerns of the public and/or professional community raised regarding potential adverse effects that might occur from a proposed action. Such issues must be considered in the NEPA decision-making process. Issues related to managing damage associated with mammals in Tennessee were developed by WS in consultation with the TWRA and the TVA along with those issues addressed during the scoping process during the development of the previous mammal damage management EA (USDA 2005*a*). This EA will also be made available to the public for review and comment to identify additional issues.

The issues as those issues relate to the possible implementation of the alternatives, including the proposed action, are discussed in detail in Chapter 4. The issues analyzed in detail in the EA are the following:

Issue 1 - Effects of Mammal Damage Management Activities on Target Mammal Populations

A common issue when addressing damage caused by wildlife are the potential impacts of management actions on the populations of target species. Methods available to resolve damage or threats to human safety under the alternatives are categorized into lethal and non-lethal methods.

Non-lethal methods could disperse or otherwise make an area unattractive to target species causing damage, which reduces the presence of those species at the site and potentially the immediate area around the site where non-lethal methods were employed. Lethal methods would be employed to remove a mammal or those mammals responsible for causing damage or posing threats to human safety. The use of lethal methods would therefore result in local population reductions in the area where damage or threats were occurring. The number of target species removed from the population using lethal methods under the alternatives would be dependent on the number of requests for assistance received, the number of individuals involved with the associated damage or threat, and the efficacy of methods employed.

The analysis for magnitude of impact on populations from the use of lethal methods generally follows the process described in WS' programmatic FEIS (USDA 1997). Magnitude is described in WS' programmatic FEIS as "*...a measure of the number of animals killed in relation to their abundance*." Magnitude may be determined either quantitatively or qualitatively. Quantitative determinations are based on population estimates, allowable harvest levels, and actual harvest data. Qualitative determinations are based on population trends and harvest data when available. Generally, WS only conducts damage management on species whose population densities are high and usually only after they have caused damage. WS' take would be monitored by comparing numbers of animals killed with overall populations or trends in populations to assure the magnitude of take is maintained below the level that would cause significant adverse impacts to the viability of native species populations. Lethal take of mammals by WS under the alternatives would only occur at the request of a cooperator seeking assistance and only after the TWRA authorized the take, when required.

In addition, many of the mammal species addressed in this EA can be harvested in the State during annual hunting and/or trapping seasons and can be addressed using available methods by other entities in the State when those species cause damage or pose threats of damage when permitted by the TWRA. Therefore, any damage management activities conducted by WS under the alternatives addressed would be occurring along with other natural process and human-induced events such as natural mortality,

human-induced mortality from private damage management activities, mortality from regulated harvest, and human-induced alterations of wildlife habitat.

Methods available under each of the alternatives to resolve damage and reduce threats to human safety would be employed targeting an individual of a mammal species or a group of individuals after applying the WS' Decision Model (Slate et al. 1992) to identify possible techniques. The effects on the populations of target mammal populations in the State from implementation of the alternatives addressed in detail, including the proposed action, are analyzed in Chapter 4. Information on mammal populations and trends are often derived from several sources including the fur harvest reports, damage complaints, ground surveys, aerial surveys, and published literature.

Issue 2 - Effects of Mammal Damage Management Activities on Non-target Wildlife Species Populations, Including T&E Species

The issue of non-target species effects, including effects on T&E species, arises from the use of non-lethal and lethal methods identified in the alternatives. The use of non-lethal and lethal methods has the potential to inadvertently disperse, capture, or kill non-target wildlife. Methods available for use under the alternatives are described in Appendix B.

Concerns have also been raised about the potential for adverse effects to occur to non-target wildlife from the use of chemical methods. Chemical methods considered for use to manage damage or threats associated with those mammal species addressed in this EA include immobilizing drugs, euthanasia chemicals, reproductive inhibitors, fumigants, rodenticides, and taste repellents. Chemical methods being considered for use to manage damage and threats associated with mammals in Tennessee are further discussed in Chapter 4 and Appendix B.

The ESA states that all federal agencies *"...shall seek to conserve endangered and threatened species and shall utilize their authorities in furtherance of the purposes of the Act"* [Sec. 7(a)(1)]. WS conducts Section 7 consultations with the USFWS to ensure compliance with the ESA and to ensure that *"any action authorized, funded or carried out by such an agency...is not likely to jeopardize the continued existence of any endangered or threatened species…Each agency shall use the best scientific and commercial data available"* [Sec. 7(a)(2)].

As part of the scoping process for this EA, WS consulted with the USFWS pursuant to Section 7 of the ESA to facilitate interagency cooperation between WS and the USFWS. The potential effects of the alternatives on this issue are further discussed in Chapter 4.

Issue 3 - Effects of Mammal Damage Management Activities on Human Health and Safety

An additional issue often raised is the potential risks to human safety associated with employing methods to manage damage caused by target species. Both chemical and non-chemical methods have the potential to have adverse effects on human safety. WS' employees would use and recommend only those methods that were legally available under each of the alternatives. Still, some concerns exist regarding the safety of methods available despite their legality and selectivity. As a result, this EA will analyze the potential for proposed methods to pose a risk to members of the public. In addition to the potential risks to the public associated with the methods available under each of the alternatives, risks to WS' employees would also be an issue. WS' employees would potentially be exposed to damage management methods, as well as, subject to workplace accidents. Selection of methods, under the alternatives, would include consideration for public and employee safety.

The issue of using chemical methods as part of managing damage associated with wildlife relates to the potential for human exposure either through direct contact with the chemical or exposure to the chemical from wildlife that have been exposed. Under the alternatives identified, the use or recommendation of chemical methods would include immobilizing drugs, euthanasia chemicals, fumigants, reproductive inhibitors, rodenticides, and repellents.

Immobilizing drugs would include ketamine and telazol, which are anesthetics (*i.e.*, general loss of pain and sensation) used during the capture of wildlife to eliminate pain, calm fear, and reduce anxiety in wildlife when handling and transporting wildlife. Xylazine is a sedative that is often used in combination with ketamine to calm nervousness, irritability, and excitement in wildlife during the handling and transporting of wildlife. Euthanasia chemicals would include sodium pentobarbital, Beuthanasia®-D, Fatal-Plus™ and potassium chloride, which would be administered after an animal has been anesthetized.

Gonacon™ is the only product currently registered as a reproductive inhibitor and is only available to manage local deer populations. However, Gonacon™ is not currently registered for use in the State. If Gonacon™ becomes registered to manage a local deer population, the product would only be available for use by WS, the TWRA, or agents under their direct supervision. The application of Gonacon™ to manage local deer herds could only occur after a permit had been issued by the TWRA.

Rodenticides would include products containing the active ingredient zinc phosphide, warfarin, brodifacoum, or diphacinone, which could be available to address damage and threats associated with those small rodent species addressed in this EA. Rodenticides are pesticides that require a restricted-use pesticide applicators license from the TDA. Zinc phosphide when ingested reacts with the acids in the gut releasing phosphine gas, which interferes with cell respiration leading to the death of the animal (EPA 1998). Warfarin, brodifacoum, and diphacinone are anticoagulant rodenticides that prevent the clotting of blood. Products containing the active ingredients, warfarin, brodifacoum, or diphacinone are currently registered for use in Tennessee. Rodenticides containing those active ingredients are generally restricted-use pesticides, which, if available, can be purchased and applied by appropriately licensed people, and are not products that are restricted to use by WS only. Those active ingredients are discussed in this EA as possible methods that could be available under the alternatives, since products are available containing those active ingredients and are registered for use in the State.

Repellents for many mammal species contain different active ingredients with most ingredients occurring naturally in the environment. The most common ingredients of repellents are coyote urine, putrescent whole egg solids, and capsaicin. Repellents are generally restricted-use products that can only be purchased and applied by licensed applicators. Repellents are generally applied directly to affected resources and elicit an adverse taste response when ingested or cause temporarily sickness (*e.g.*, nausea). Products containing coyote urine or other odors associated with predatory wildlife are intended to elicit a fright response in target wildlife by imitating the presence of a predatory animal (*i.e.*, wildlife tend to avoid areas where predators are known to occur). WS would only employ or recommend for use those rodenticides and repellents that were registered for use pursuant to the FIFRA with the EPA and were registered for use in the State by the TDA.

Gas cartridges would be available to fumigate burrows and den sites in areas where damages were occurring. Gas cartridges act as a fumigant by producing carbon monoxide gas when ignited. The cartridges contain sodium nitrate, which when burnt, produces carbon monoxide gas. The cartridges would be placed inside active burrows and dens at the entrance, the cartridge would be ignited, and the entrance to the burrow or den would be sealed with dirt, which allows the burrow or den to fill with carbon monoxide.

Products containing the active ingredient aluminum phosphide are used either as a fumigant or as a rodenticide. Fumigants containing aluminum phosphide as the active ingredient are formulated as tablets, which are placed inside rodent burrows and the burrows are sealed up. The aluminum phosphide in the table reacts with the moisture in the soil releasing phosphine gas. Since burrows are sealed after placing the tablets, the burrow fills with toxic phosphine gas. When used as a rodenticide, products containing aluminum phosphide are formulated as pellets and are present as bait for ingestion. When the pellet is ingested, the aluminum phosphide reacts with the acid in the stomach releasing phosphine gas. Products containing the active ingredient aluminum phosphide are registered for use in the State, primarily as fumigants. Products containing aluminum phosphide are restricted use pesticides and would be available to any licensed applicator.

The issue of the potential for drugs used in animal capture, handling, and euthanasia to cause adverse health effects in humans that hunt and consume the species involved has been raised. Among the species to be captured and handled under the proposed action, this issue is expected to be of concern for wildlife that are hunted and sometimes consumed by people as food. Chemicals methods available for use under the relevant alternatives would be regulated by the EPA through FIFRA, by Tennessee laws, by the DEA, by the United States Food and Drug Administration (FDA), and by WS' Directives.

Most methods available to alleviate damage and threats associated with mammals are considered non-chemical methods. Non-chemical methods may include cultural methods, limited habitat modification, animal behavior modification, and other mechanical methods. Changes in cultural methods could include improved animal husbandry practices, altering feeding schedules, changes in crop rotations, or conducting structural repairs. Limited habitat modification would be practices that alter specific characteristics of a very localized area, such as removing bushes to eliminate shelter locations or planting vegetation that are less palatable to mammals. Animal behavior modification methods would include those methods designed to disperse mammals from an area through harassment or exclusion. Behavior modification methods could include pyrotechnics, propane cannons, barriers, electronic distress calls, effigies, Mylar tape, and lasers. Other mechanical methods could include live-traps, body-gripping traps, cable restraints, cannon nets, shooting, or the recommendation that a local population of mammals be reduced using hunting and/or trapping.

The primary safety risk of most non-chemical methods occurs directly to the applicator or those persons assisting the applicator. However, risks to others do exist when employing non-chemical methods, such as when using firearms, cannon nets, pyrotechnics, or body-gripping traps. Most of the non-chemical methods available to address mammal damage in Tennessee would be available for use under any of the alternatives and could be employed by any entity, when permitted. Risks to human safety from the use of non-chemical methods will be further evaluated as this issue relates to the alternatives in Chapter 4. A complete list of non-chemical methods available to alleviate damage associated with mammals is provided in Appendix B of this EA.

An issue identified is the concern for human safety from not employing methods or not employing the most effective methods to reduce the threats that mammals can pose. The risks to human safety from diseases associated with certain mammal populations were addressed previously in Chapter 1 under the need for action. The low risk of disease transmission from mammals does not lessen the concerns of cooperators requesting assistance to reduce threats from zoonotic diseases. Increased public awareness of zoonotic events has only heightened the concern of direct or indirect exposure to zoonoses. Not adequately addressing the threats associated with potential zoonoses could lead to an increase in incidences of injury, illness, or loss of human life.

Additional concern is raised with inadequately addressing threats to human safety associated with aircraft striking mammals at airports in the State. Mammals have the potential to cause severe damage to aircraft,

which can threaten the safety of passengers. Limiting or preventing the use of certain methods to address the potential for aircraft striking mammals could lead to higher risks to passenger safety. This issue will be fully evaluated in Chapter 4 in relationship to the alternatives.

Issue 4 - Effects of Mammal Damage Management Activities on the Socio-cultural Elements of the Human Environment

One issue is the concern that the proposed action or the other alternatives would result in the loss of aesthetic benefits of target mammals to the public, resource owners, or neighboring residents. Wildlife generally is regarded as providing economic, recreational, and aesthetic benefits (Decker and Goff 1987), and the mere knowledge that wildlife exists is a positive benefit to many people. Aesthetics is the philosophy dealing with the nature of beauty, or the appreciation of beauty. Therefore, aesthetics is truly subjective in nature, dependent on what an observer regards as beautiful.

The human attraction to animals has been well documented throughout history and started when humans began domesticating animals. The public share a similar bond with animals and/or wildlife in general and in modern societies, a large percentage of households have indoor or outdoor pets. However, some people may consider individual wild animals and mammals as "*pets*" or exhibit affection toward those animals, especially people who enjoy viewing wildlife. Therefore, the public reaction can be variable and mixed to wildlife damage management because there are numerous philosophical, aesthetic, and personal attitudes, values, and opinions about the best ways to manage conflicts/problems between humans and wildlife.

Wildlife populations provide a wide range of social and economic benefits (Decker and Goff 1987). Those include direct benefits related to consumptive and non-consumptive uses, indirect benefits derived from vicarious wildlife related experiences, and the personal enjoyment of knowing wildlife exists and contributes to the stability of natural ecosystems (Bishop 1987). Direct benefits are derived from a personal relationship with animals and may take the form of direct consumptive use (*i.e.*, using parts of or the entire animal) or non-consumptive use (*e.g.*, viewing the animal in nature or in a zoo, photographing) (Decker and Goff 1987).

Indirect benefits or indirect exercised values arise without the user being in direct contact with the animal and come from experiences such as looking at photographs and films of wildlife, reading about wildlife, or benefiting from activities or contributions of animals such as their use in research (Decker and Goff 1987). Indirect benefits come in two forms: bequest and pure existence (Decker and Goff 1987). Bequest is providing for future generations and pure existence is merely knowledge that the animals exist (Decker and Goff 1987).

Public attitudes toward wildlife vary considerably. Some people believe that all wildlife should be captured and translocated to another area to alleviate damage or threats to protected resources. Some people directly affected by the problems caused by wildlife strongly support removal. Individuals not directly affected by the harm or damage may be supportive, neutral, or totally opposed to any removal of wildlife from specific locations or sites. Some people totally opposed to wildlife damage management want WS to teach tolerance for damage and threats caused by wildlife, and that wildlife should never be killed. Some of the people who oppose removal of wildlife do so because of human-affectionate bonds with individual wildlife. Those human-affectionate bonds are similar to attitudes of a pet owner and result in aesthetic enjoyment.

Some individuals are offended by the presence of overabundant mammal species, such as raccoons, armadillos, gray squirrels, coyotes, or feral species, such as cats or dogs. To such people those species represent pests that are nuisances, which upset the natural order in ecosystems, and are carriers of diseases

transmissible to humans or other wildlife. Their overall enjoyment of other animals is diminished by what they view as a destructive presence of such species. They are offended because they feel that those mammal species proliferate in such numbers and appear to remain unbalanced.

Issue 5 - Humaneness and Animal Welfare Concerns of Methods

The issue of humaneness and animal welfare, as it relates to the killing or capturing of wildlife is an important but very complex concept that can be interpreted in a variety of ways. Schmidt (1989) indicated that vertebrate damage management for societal benefits could be compatible with animal welfare concerns, if "*…the reduction of pain, suffering, and unnecessary death is incorporated in the decision making process.*"

According to the AVMA (1987), suffering is described as a "*…highly unpleasant emotional response usually associated with pain and distress.*" However, suffering "*…can occur without pain…*" and "*…pain can occur without suffering…*". Because suffering carries with it the implication of a time frame, a case could be made for "*…little or no suffering where death comes immediately…*" (California Department of Fish and Game 1991). Pain and physical restraint can cause stress in animals and the inability of animals to effectively deal with those stressors can lead to distress. Suffering occurs when action is not taken to alleviate conditions that cause pain or distress in animals.

Defining pain as a component in humaneness appears to be a greater challenge than that of suffering. Pain obviously occurs in animals. Altered physiology and behavior can be indicators of pain, and identifying the causes that elicit pain responses in humans would "*…probably be causes for pain in other animals…*" (AVMA 1987). However, pain experienced by individual animals probably ranges from little or no pain to considerable pain (California Department of Fish and Game 1991).

Pain and suffering, as it relates to methods available for use to manage mammals has both a professional and lay point of arbitration. Wildlife managers and the public would be better served to recognize the complexity of defining suffering, since *"…neither medical nor veterinary curricula explicitly address suffering or its relief"* (California Department of Fish and Game 1991). Research suggests that some methods, such as restraint in foot-hold traps or changes in the blood chemistry of trapped animals, indicate "*stress*" (USDA 1997). However, such research has not yet progressed to the development of objective, quantitative measurements of pain or stress for use in evaluating humaneness.

The AVMA states "*...euthanasia is the act of inducing humane death in an animal*" and "*... the technique should minimize any stress and anxiety experienced by the animal prior to unconsciousness*" (Beaver et al. 2001). Some people would prefer AVMA accepted methods of euthanasia to be used when killing all animals, including wild and invasive animals. The AVMA states that "*For wild and feral animals, many of the recommended means of euthanasia for captive animals are not feasible. In field circumstances, wildlife biologists generally do not use the term euthanasia, but terms such as killing, collecting, or harvesting, recognizing that a distress- free death may not be possible*" (Beaver et al. 2001).

The decision-making process involves tradeoffs between the above aspects of pain and humaneness. Therefore, humaneness, in part, appears to be a person's perception of harm or pain inflicted on an animal, and people may perceive the humaneness of an action differently. The challenge in coping with this issue is how to achieve the least amount of animal suffering. The issue of humanness and animal welfare concerns will be further discussed in Chapter 4. SOPs to alleviate pain and suffering are discussed in Chapter 3.

Issue 6 - Effects of Mammal Damage Management Activities on the Regulated Harvest of Mammals

Another issue commonly identified is a concern that damage management activities conducted by WS would affect the ability of persons to harvest those species during the regulated hunting and trapping seasons either by reducing local populations through the lethal removal of mammals or by reducing the number of mammals present in an area through dispersal techniques. Those species that are addressed in this EA that also can be hunted and/or trapped during regulated seasons in the State include beaver, muskrat, woodchuck, cottontail rabbit, weasel, mink, Virginia opossum, raccoon, river otter, striped skunk, spotted skunk, coyote, gray fox, red fox, bobcat, gray squirrel, fox squirrel, black bear, elk, and white-tailed deer.

Potential impacts could arise from the use of non-lethal or lethal damage management methods. Non-lethal methods used to reduce or alleviate damage caused by those mammal species are used to reduce mammal densities through dispersal in areas where damage or the threat of damage is occurring. Similarly, lethal methods used to reduce damage associated with those mammals could lower densities in areas where damage is occurring resulting in a reduction in the availability of those species during the regulated harvest season. WS' mammal damage management activities would primarily be conducted on populations in areas where hunting access is restricted (*e.g.*, airports, TVA facilities, and, recreation areas) or has been ineffective. The use of non-lethal or lethal methods often disperses mammals from areas where damage is occurring to areas outside the damage area, which could serve to move those mammal species from those less accessible areas to places accessible to hunters.

Issue 7 – Effects of Beaver Removal and Dam Manipulation on the Status of Wetlands in the State

Wetlands are a valuable component of land-based ecosystems that provide numerous direct and indirect benefits to people and wildlife (*e.g.*, see Costanza et al. 1997, Millennium Ecosystem Assessment 2005). Between the 1780s and the 1980s, Dahl (1990) estimated 53% of the original wetland acres in the lower 48 states were lost, primarily from human development. Over that 200-year time span, Dahl (1990) estimated the wetland acres in Tennessee decreased from 1,937,000 acres to 787,000 acres, which represented a 59% decline. Beaver, through their building of dams and impounding water can have a unique role in establishing wetlands that not only provide benefit to the beaver, but to people and other wildlife. Beaver are often considered a "*keystone*" species for their ability to manipulate and create their own habitats, which can also provide benefits to other wildlife and people. Beaver may also be an inexpensive way of restoring wetlands or creating new wetlands (*e.g.*, see Hey 1995, Muller-Schwarze and Sun 2003, Buckley et al. 2011).

The issue of WS' potential impacts to wetlands could occur from activities conducted to alleviate damage or threats of damage associated with beaver, primarily from the breaching or removal of beaver dams. Beaver dam breaching or removal during activities to manage damage caused by beaver sometimes occurs in areas inundated by water from water impounded by beaver dams. Dam material usually consists of mud, sticks, and other vegetative material. Beaver dams obstruct the normal flow of water, which can change the preexisting hydrology from flowing or circulating waters to slower, deeper, more expansive waters that accumulate bottom sediment over time. The depth of the bottom sediment depends on the length of time an area is covered by water and the amount of suspended sediment in the water.

Beaver dams, over time, can establish new wetlands. The regulatory definition of a wetland stated by the United States Army Corps of Engineers and the EPA (40 CFR 232.2) is:

"Those areas that are inundated or saturated by surface or groundwater at a frequency and duration sufficient to support, and that under normal circumstances do support, a prevalence of vegetation

typically adapted for life in saturated soil conditions. Wetlands generally include swamps, marshes, bogs, and similar areas."

Therefore, the breaching or removal of a beaver dam could result in the degrading or removal of a wetland, if wetland characteristics exist at a location where a beaver dam occurs. The preexisting habitat (prior to the building of the dam) and the altered habitat (areas flooded by impounded water) have different ecological values to the fish and wildlife native to the area. Some species may benefit by the addition of a beaver dam that creates a wetland, while the presence of some species of wildlife may decline. For example, darters listed as federally endangered require fast moving waters over gravel or cobble beds, which beaver dams can eliminate; thus, reducing the availability of habitat. In areas where bottomland forests were flooded by beaver dams, a change in species composition could occur over time as trees die. Hardwood trees are often killed when flooding persists for extended periods, as soils become saturated. Conversely, beaver dams could be beneficial to some wildlife such as river otter, neotropical migratory birds, and waterfowl that require aquatic habitats.

If a beaver dam was not removed and water was allowed to stand, hydric soils and hydrophytic vegetation could eventually form. This process could take anywhere from several months to years depending on preexisting conditions. Hydric soils are those soils that are saturated, flooded, or ponded long enough during the growing season to develop anaerobic conditions. In general, hydric soils form much easier where wetlands have preexisted. Hydrophytic vegetation includes those plants that grow in water or on a substrate that is at least periodically deficient in oxygen because of excessive water content. If those conditions were met, then a wetland has developed that would have different wildlife habitat values than an area that has been more recently impounded by beaver dam activity.

In addition, concerns are often raised regarding the use of lethal methods to remove beaver to alleviate damage or threats. If beaver were lethally removed from an area and any associated beaver dam was removed or breached, the manipulation of water levels by removing/breaching the dam could prevent the establishment of wetlands in areas where water has been impounded for an extended period by beaver dams. If beaver were removed but the beaver dam was left undisturbed, the lack of maintenance to the dam by beaver would likely result in the eventual recession of the impounded water as weathering eroded the dam.

2.3 ISSUES CONSIDERED BUT NOT IN DETAIL WITH RATIONALE

Additional issues were also identified by WS, the TVA, and the TWRA during the scoping process of this EA. Those additional issues were considered but detailed analyses will not occur for the reasons provided. The following issues were considered but were not analyzed in detail:

Appropriateness of Preparing an EA (Instead of an EIS) for Such a Large Area

A concern was raised that an EA for an area as large as the State of Tennessee would not meet the NEPA requirements for site specificity. Wildlife damage management falls within the category of federal or other regulatory agency actions in which the exact timing or location of individual activities cannot usually be predicted well enough ahead of time to describe accurately such locations or times in an EA or EIS. Although WS can predict some of the possible locations or types of situations and sites where some kinds of wildlife damage could occur, the program cannot predict the specific locations or times at which affected resource owners would determine a damage problem has become intolerable to the point that they request assistance from WS. In addition, the WS program would not be able to prevent such damage in all areas where it might occur without resorting to destruction of wild animal populations over broad areas at a much more intensive level than would be desired by most people, including WS and other

agencies. Such broad scale population management would also be impractical or impossible to achieve within WS' policies and professional philosophies.

Lead agencies have the discretion to determine the geographic scope of their analyses under the NEPA (Kleppe v Sierra Club, 427 U.S. 390, 414 (1976), CEQ 1508.25). Ordinarily, according to APHIS procedures implementing the NEPA, WS' individual wildlife damage management actions could be categorically excluded (7 CFR 372.5(c)). The intent in developing this EA was to determine if the proposed action would potentially have significant individual and/or cumulative impacts on the quality of the human environment that would warrant the preparation of an EIS. This EA addresses impacts for managing damage and threats to human safety associated with mammals in the State to analyze individual and cumulative impacts and to provide a thorough analysis.

In terms of considering cumulative effects, one EA analyzing impacts for the entire State would provide a more comprehensive and less redundant analysis than multiple EAs covering smaller areas. If a determination were made through this EA that the proposed action or the other alternatives might have a significant impact on the quality of the human environment, then an EIS would be prepared. Based on previous requests for assistance, the WS program in Tennessee would continue to conduct mammal damage management in a very small area of the State where damage is occurring or likely to occur.

WS' Impact on Biodiversity

The WS program does not attempt to eradicate any species of native wildlife in the State. WS operates in accordance with federal and state laws and regulations enacted to ensure species viability. Methods available are employed to target individual mammals or groups of mammals identified as causing damage or posing a threat of damage. Any reduction of a local population or group is frequently temporary because immigration from adjacent areas or reproduction replaces the animals removed. WS operates on a small percentage of the land area of Tennessee and only targets those mammals identified as causing damage or posing a threat. Therefore, mammal damage management activities conducted pursuant to any of the alternatives would not adversely affect biodiversity in the State.

A Loss Threshold Should Be Established Before Allowing Lethal Methods

One issue identified through WS' implementation of the NEPA processes is a concern that a threshold of loss should be established before employing lethal methods to resolve damage and that wildlife damage should be a cost of doing business. Some damage and economic loss would likely be tolerated by cooperators until the damage reaches a threshold where the damage becomes an economic burden. The appropriate level of allowed tolerance or threshold before employing lethal methods would differ among cooperators and damage situations. In addition, establishing a threshold would be difficult or inappropriate to apply to human health and safety situations. For example, aircraft striking mammals can lead to property damage and can threaten passenger safety if a catastrophic failure of the aircraft occurs because of the strike. Therefore, addressing the threats of wildlife strikes prior to an actual strike occurring would be appropriate.

In a ruling for Southern Utah Wilderness Alliance, et al. vs. Hugh Thompson, Forest Supervisor for the Dixie National Forest, et al., the United States District Court of Utah denied the plaintiffs' motion for a preliminary injunction. In part, the court determined a need for wildlife damage management could be established if a forest supervisor could show that damage from wildlife was threatened (Civil No. 92-C-0052A January 20, 1993). Thus, there is judicial precedence indicating that it is not necessary to establish a criterion such as a percentage of loss of a particular resource to justify the need for damage management actions.

Mammal Damage Management Should Not Occur at Taxpayer Expense

An issue identified is the concern that wildlife damage management should not be provided at the expense of the taxpayer or that activities should be fee-based. Funding for damage management activities would be derived from federal appropriations and through cooperative funding. Activities conducted in the State for the management of damage and threats to human safety from mammals would be funded through cooperative service agreements with individual property owners or managers. A minimal federal appropriation is allotted for the maintenance of a WS program in Tennessee. The remainder of the WS program would mostly be fee-based. Technical assistance would be provided to requesters as part of the federally funded activities, but the majority of direct assistance in which WS' employees perform damage management activities would be funded through cooperative service agreements between the requester and WS.

Cost Effectiveness of Management Methods

The CEQ does not require a formal, monetized cost benefit analysis to comply with the NEPA. Consideration of this issue is not essential to making a reasoned choice among the alternatives being considered. However, the methods determined to be most effective to reduce damage and threats to human safety caused by mammals and that prove to be the most cost effective would likely receive the greatest application. As part of an integrated approach and as part of the WS Decision Model, evaluation of methods would continually occur to allow for those methods that were most effective at resolving damage or threats to be employed under similar circumstance where mammals were causing damage or posing a threat. Additionally, management operations may be constrained by cooperator funding and/or objectives and needs. The cost effectiveness of methods and the effectiveness of methods would be linked. The issue of cost effectiveness as it relates to the effectiveness of methods is discussed in the following issue.

Effectiveness of Mammal Damage Management Methods

The effectiveness of any damage management program could be defined in terms of losses or risks potentially reduced or prevented. The effectiveness can also be dependent upon how accurately practitioners diagnose the problem, the species responsible for the damage, and how actions were implemented to correct or mitigate risks or damages. To determine that effectiveness, WS must be able to complete management actions expeditiously to minimize harm to non-target animals and the environment, while at the same time, using methods as humanely as possible. The most effective approach to resolving any wildlife damage problem would be to use an adaptive integrated approach, which may call for the use of several management methods simultaneously or sequentially (Courchamp et al. 2003).

The purpose behind integrated management is to implement methods in the most effective manner while minimizing the potentially harmful effects on humans, target and non-target species, and the environment[8]. Efficacy is based on the types of methods employed, the application of the method, restrictions on the use of the method(s), the skill of the personnel using the method and, for WS' personnel, the guidance provided by WS' directives and policies.

The goal would be to reduce damage, risks, and conflicts with wildlife as requested and not to reduce/eliminate populations. Localized population reduction could be short-term with new individuals immigrating into the area or born to animals remaining at the site (Courchamp et al. 2003). The ability of

[8]The cost of management may sometimes be secondary because of overriding environmental, legal, human health and safety, animal welfare, or other concerns.

an animal population to sustain a certain level of removal and to eventually return to pre-management levels does not mean individual management actions were unsuccessful, but that periodic management may be necessary. The return of wildlife to pre-management levels also demonstrates that limited, localized damage management methods have minimal impacts on species' populations.

Comments are often received that lethal methods would be ineffective because additional mammals would likely to return to the area. In addition, comments also claim that because mammals return to an area after initial removal efforts were complete, the use of lethal methods gives the impression of creating a financial incentive to continue the use of only lethal methods. Those statements assume mammals only return to an area where damage was occurring if lethal methods were used. However, the use of non-lethal methods would also often be temporary, which could result in mammals returning to an area where damage was occurring once those methods were no longer used. The common factor when employing any method would be that mammals would return if suitable conditions continue to exist at the location where damage was occurring and mammal densities were sufficient to occupy all available habitats to the extent that damage occurs. Therefore, any reduction or prevention of damage from the use of methods addressed in Appendix B would be temporary if habitat conditions continue to exist that attract mammals to an area where damage was occurring.

Therefore, any method that disperses or removes mammals from areas would only be temporary if habitat containing preferred habitat characteristics continues to exist. Dispersing mammals using non-lethal methods addressed in Appendix B often requires repeated application to discourage mammals from returning to locations, which increases costs, moves mammals to other areas where they could cause damage, and would be temporary if habitat conditions that attracted those mammals to damage areas remained unchanged. Dispersing and translocating mammals could be viewed as moving a problem from one area to another, which would require addressing damage caused by those mammals at another location, which increases costs and could be perceived as creating a financial incentive to continue the use of those methods since mammals would have to be addressed annually and at multiple locations. WS' recommendation of or use of techniques to modify existing habitat or making areas unattractive to mammals is discussed in Appendix B. WS' objective would be to respond to requests for assistance with the most effective methods and to provide for the long-term solution to the problem using WS' Decision Model.

Managing damage caused by mammals can be divided into short-term redistribution approaches and long-term population and habitat management approaches. Short-term approaches focus on redistribution and dispersal of mammals to limit use of an area where damage or threats were occurring. Short-term redistribution approaches may include prohibiting feeding, the use of pyrotechnics, propane cannons, effigies, and other adverse noise, erecting access barriers such as fencing, and taste aversion chemicals. Population reduction by limiting survival or reproduction, removing mammals, and habitat modification would be considered long-term solutions to managing damage caused by wildlife.

Redistribution methods would often be employed to provide immediate resolution to damage occurring until long-term approaches can be implemented or have had time to reach the desired result. Dispersing mammals can often be a short-term solution that moves those mammals to other areas where damages or threats could occur. Some short-term methods may become less effective in resolving damage as a mammal population increases, as mammals become more acclimated to human activity, and as mammals become habituated to harassment techniques. Non-lethal methods often require a constant presence at locations when mammals are present and must be repeated every day or night until the desired results are achieved, which can increase the costs associated with those activities. Non-lethal methods may also require constant monitoring and maintenance to insure proper results. For example, fencing could be used to prevent access to a resource; however, constant monitoring of the fencing would be required and necessary repairs completed to ensure the use of fencing would be successful in preventing access to

resources. Long-term solutions to resolving mammal damage often require management of the population and identifying the habitat characteristics that attract mammals to a particular location.

Based on the evaluation of the damage situation, the most effective methods would be employed individually or in combination based on prior evaluations of methods or combinations of methods in other damage management situations. Once employed, methods would be further evaluated for effectiveness based on a continuous evaluation of activities by WS. Therefore, the effectiveness of methods would be considered as part of the decision making-process under WS' use of the Decision Model described in Chapter 3 for each damage management request based on the continual evaluation of methods and results.

Mammal Damage Should be Managed by Private Nuisance Wildlife Control Agents

Private wildlife control agents could be contacted to reduce mammal damage for property owners or property managers when deemed appropriate by the resource owner. Some resource owners would prefer to use a private nuisance wildlife control agent because the nuisance wildlife agent is located in closer proximity and thus could provide the service at less expense, or because they prefer to use a private business rather than a government agency. However, some resource owners would prefer to enter into an agreement with a government agency. In particular, those persons seeking assistance may prefer to use WS because of security and safety issues. WS further clarifies interfacing with private business and establishing cooperative projects in WS Directive 3.101.

Effects from the Use of Lead Ammunition in Firearms

Questions have arisen about the deposition of lead into the environment from ammunition used in firearms to lethally take mammals. As described in Appendix B, the lethal removal of mammals with firearms by WS to alleviate damage or threats could occur using a handgun, rifle, or shotgun. In an ecological risk assessment of lead shot exposure in non-waterfowl birds, ingestion of lead shot was identified as the concern rather than just contact with lead shot or lead leaching from shot in the environment (Kendall et al. 1996).

The take of mammals by WS using firearms in the State would occur primarily from the use of rifles. However, the use of shotguns or handguns could be employed to lethally take some species. To reduce risks to human safety and property damage from bullets passing through mammals, the use of rifles would be applied in such a way (*e.g.*, caliber, bullet weight, distance) to ensure the bullet does not pass through mammals. Mammals that are removed using rifles would occur within areas where retrieval of all mammal carcasses for proper disposal is highly likely (*e.g.*, at an airport). With risks of lead exposure occurring primarily from ingestion of bullet fragments, the retrieval and proper disposal of mammal carcasses would greatly reduce the risk of scavengers ingesting or being exposed to lead that may be contained within the carcass.

However, deposition of lead into soil could occur if, during the use of a rifle, the projectile passes through a mammal, if misses occur, or if the mammal carcass was not retrieved. Laidlaw et al. (2005) reported that, because of the low mobility of lead in soil, all of the lead that accumulates on the surface layer of the soil is generally retained within the top 20 cm (about 8 inches). In addition, concerns occur that lead from bullets deposited in soil from shooting activities could contaminate ground water or surface water from runoff. Stansley et al. (1992) studied lead levels in water that was subjected directly to high concentrations of lead shot accumulation because of intensive target shooting at several shooting ranges. Lead did not appear to "*transport*" readily in surface water when soils were neutral or slightly alkaline in pH (*i.e.*, not acidic), but lead did transport more readily under slightly acidic conditions. Although Stansley et al. (1992) detected elevated lead levels in water in a stream and a marsh that were in the shot "*fall zones*" at a shooting range, the study did not find higher lead levels in a lake into which the stream

drained, except for one sample collected near a parking lot. Stansley et al. (1992) believed the lead contamination near the parking lot was due to runoff from the lot, and not from the shooting range areas. The study also indicated that even when lead shot was highly accumulated in areas with permanent water bodies present, the lead did not necessarily cause elevated lead levels in water further downstream. Muscle samples from two species of fish collected in water bodies with high lead shot accumulations had lead levels that were well below the accepted threshold standard of safety for human consumption (Stansley et al. 1992).

Craig et al. (1999) reported that lead levels in water draining away from a shooting range with high accumulations of lead bullets in the soil around the impact areas were far below the "*action level*" of 15 parts per billion as defined by the EPA (*i.e.*, requiring action to treat the water to remove lead). The study found that the dissolution (*i.e.*, capability of dissolving in water) of lead declines when lead oxides form on the surface areas of the spent bullets and fragments (Craig et al. 1999). Therefore, the transport of lead from bullets or shot distributed across the landscape was reduced once the bullets and shot formed crusty lead oxide deposits on their surfaces, which served to reduce naturally the potential for ground or surface water contamination (Craig et al. 1999). Those studies suggest that, given the very low amount of lead being deposited and the concentrations that would occur from WS' activities to reduce mammal damage using firearms, as well as most other forms of dry land small game hunting in general, lead contamination of water from such sources would be minimal to nonexistent.

A secondary concern surrounding lead ammunition surrounds the issue of lead deposition in meat, particularly meat that is donated to various charities. Stewart and Veverka (2011) documented that white-tailed deer that were shot with lead ammunition in the head or extreme upper neck in sharpshooting situations showed no deposition of lead fragments in the meat of the animals that would have been processed for human consumption. Lower neck shots do frequently experience lead fragmentation in the loin muscle and the authors recommend removing the loins prior to processing to ensure that these fragments are not ingested. WS' personnel would be trained to shoot and target the head and upper neck of white-tailed deer. Any deer that were shot in the lower neck would not be donated but would be disposed of properly to avoid potential human ingestion of lead fragments.

Since those mammals removed by WS using firearms could be lethally removed by the entities experiencing damage using the same method in the absence of WS' involvement, WS' assistance with removing those mammals would not be additive to the environmental status quo. The amount of lead deposited into the environment could be lowered by WS' involvement in damage management activities due to efforts by WS to ensure projectiles do no pass through but are contained within the mammal carcass, which limits the amount of lead potentially deposited into soil from projectiles passing through the carcass. The proficiency training received by WS' employees in firearm use and accuracy would increase the likelihood that mammals were lethally removed humanely in situations that ensure accuracy and that misses occur infrequently which further reduces the potential for lead to be deposited in the soil from misses or from projectiles passing through carcasses. In addition, WS' involvement ensures mammal carcasses lethally removed using firearms would be retrieved and disposed of properly to limit the availability of lead in the environment and ensures mammal carcasses were removed from the environment to prevent the ingestion of lead in carcasses by scavengers. Based on current information, the risks associated with lead bullets that could be deposited into the environment from WS' activities due to misses, the bullet passing through the carcass, or from mammal carcasses that may be irretrievable would be below any level that would pose any risk from exposure or significant contamination of water.

Effects on Human Health from Consumption of Deer Meat Donated by WS

Of concern under this issue would be the consumption of deer meat donated to a charitable organization after being lethally taken by WS. Of recent concern is the potential for lead and other contaminants to be

present in meat that has been processed for human consumption. The potential for the spreading of zoonotic diseases in deer processed and donated for human consumption is also a concern. Under the proposed action alternative, meat from deer lethally taken during damage management activities could be donated to charitable organizations for human consumption. Only meat from deer would be donated under the proposed action alternative. WS could recommend the donation or consumption of meat under the technical assistance only alternative but would not be directly involved with damage management activities under that alternative.

If WS donated deer for human consumption, WS' policies pertaining to the testing or labeling of meat would be followed in order to address potential health concerns. Deer donated for human consumption may be tested for exposure to substances such as organophosphate and carbamate insecticides, lead, mercury, arsenic, organochlorines, and organic chemicals prior to distribution. Deer immobilized using immobilizing drugs or euthanized using euthanasia chemicals would not be donated for human consumption with disposal of carcasses occurring pursuant to WS Directive 2.515. Deer taken by any method for disease sampling or in an area where zoonotic diseases of concern are known to be prevalent and of concern to human health after consuming processed deer meat would not be donated for consumption and would be disposed of by deep burial or incineration. WS' adherence to policy would not result in adverse effects to human health from the donation of deer meat.

A Site Specific Analysis Should be made for Every Location Where Mammal Damage Management Would Occur

The underlying intent for preparing an EA is to determine if a proposed action might have a significant impact on the human environment. WS' EA development process is issue driven, meaning issues that were raised during the interdisciplinary process and through public involvement that were substantive, were used to drive the analysis and determine the significance of the environmental impacts of the proposed action and the alternatives. Therefore, the level of site specificity must be appropriate to the issues listed.

The analysis in this EA was driven by the issues raised during the scoping process during the development of the EA. In addition to the analysis contained in this EA, WS' personnel use the WS Decision Model (Slate et al. 1992) described in Chapter 3 as a site-specific tool to develop the most appropriate strategy at each location. The WS Decision Model is an analytical thought process used by WS' personnel for evaluating and responding to requests for assistance.

As discussed previously, one EA analyzing impacts for the entire State would provide a more comprehensive and less redundant analysis that allows for a better cumulative impact analysis. If a determination were made through this EA that the alternatives developed to meet the need for action could result in a significant impact on the quality of the human environment, then an EIS would be prepared.

CHAPTER 3: ALTERNATIVES

Chapter 3 contains a discussion of the alternatives that were developed to meet the need for action discussed in Chapter 1 and to address the identified issues discussed in Chapter 2. Alternatives were developed for consideration based on the need for action and issues using the WS Decision model (Slate et al. 1992). The alternatives will receive detailed environmental impacts analysis in Chapter 4 (Environmental Consequences). Chapter 3 also discusses alternatives considered but not analyzed in detail, with rationale. SOPs for mammal damage management in Tennessee are also discussed in Chapter 3.

3.1 DESCRIPTION OF THE ALTERNATIVES

The following alternatives were developed to meet the need for action and address the identified issues associated with managing damage caused by mammals in the State:

Alternative 1 - Continue the Current Adaptive Integrated Mammal Damage Management Program (No Action/Proposed Action)

The proposed action/no action alternative would continue the current implementation of an adaptive integrated approach utilizing non-lethal and lethal techniques, when requested, as deemed appropriate using the WS Decision Model, to reduce damage and threats caused by mammals in Tennessee. A major goal of the program would be to resolve and prevent damage caused by mammals and to reduce threats to human safety. To meet this goal, WS would continue to respond to requests for assistance with, at a minimum, technical assistance, or when funding was available, operational damage management. Funding could occur through federal appropriations or from cooperative funding. The adaptive approach to managing damage associated with mammals would integrate the use of the most practical and effective methods to resolve a request for damage management as determined by a site-specific evaluation to reduce damage or threats to human safety for each request. City/town managers, agricultural producers, property owners, and others requesting assistance would be provided information regarding the use of appropriate non-lethal and lethal techniques.

Under this alternative, WS could respond to requests for assistance by: 1) taking no action, if warranted, 2) providing only technical assistance to property owners or managers on actions they could take to reduce damages caused by mammals, or 3) providing technical assistance and direct operational assistance to a property owner or manager experiencing damage. The take of many of the mammal species addressed in this EA can only legally occur through the issuance of a permit by the TWRA and only at levels specified in the permit, unless those mammal species are afforded no protection in which case no permit for take is required.

Property owners or managers requesting assistance from WS would be provided with information regarding the use of effective and practical non-lethal and lethal techniques. Preference would be given to non-lethal methods when practical and effective under this alternative (see WS Directive 2.101). Property owners or managers may choose to implement WS' recommendations on their own (*i.e.*, technical assistance), use contractual services of private businesses, use volunteer services of private organizations, use the services of WS (*i.e.*, direct operational assistance), take the management action themselves, or take no further action.

WS would work with those persons experiencing mammal damage in addressing those mammals responsible for causing damage as expeditiously as possible. To be most effective, damage management activities should begin as soon as mammals begin to cause damage. Mammal damage that has been ongoing can be difficult to resolve using available methods since mammals would be conditioned to an area and would be familiar with a particular location. Subsequently, making that area unattractive using available methods could be difficult to achieve once damage was ongoing. WS would work closely with those entities requesting assistance to identify situations where damage could occur and begin to implement damage management activities under this alternative as early as possible to increase the likelihood of those methods achieving the level of damage reduction requested by the cooperating entity.

WS' Decision Model would be the implementing mechanism for a damage management program under the proposed action alternative that would be adapted to an individual damage situation that allows for the broadest range of methods to be used to address damage or the threat of damage in the most effective, most efficient, and most environmentally conscious way available. When WS receives a request for

direct operational assistance, WS would conduct site visits to assess damage or threats, would identify the cause of the damage, and would apply the Decision Model described by Slate et al. (1992) and WS Directive 2.201 to determine the appropriate methods to resolve or prevent damage. The use of the Decision model by WS' employees under the proposed action is further discussed below. In addition, preference would be given to non-lethal methods when practical and effective (WS Directive 2.101).

Non-lethal methods that would be available for use by WS under this alternative include, but are not limited to minor habitat modification, behavior modification, visual deterrents, live traps, translocation, exclusionary devices, frightening devices, immobilizing drugs, reproductive inhibitors, and chemical repellents (see Appendix B for a complete list and description of potential methods). Lethal methods that would be available to WS under this alternative include body-gripping traps, cable restraints, the recommendation of take during hunting and/or trapping seasons, fumigants, rodenticides, and shooting, including the use of firearms from aircraft. In addition, target mammal species live-captured using non-lethal methods (*e.g.*, live-traps, immobilizing drugs) could be euthanized. The lethal control of target mammals would comply with WS Directive 2.505.

Listing methods implies neither that all methods would be used or recommended by WS to resolve requests for assistance nor does listing of methods imply that all methods would be used to resolve every request for assistance. The most appropriate response would often be a combination of non-lethal and lethal methods, or there could be instances where application of lethal methods alone would be the most appropriate strategy. For example, if an entity requesting assistance had already attempted to alleviate damage using non-lethal methods, WS would not necessarily employ those same non-lethal methods, since those methods were proven ineffective at reducing damage or threats to an acceptable level to the requester.

Many lethal and non-lethal methods are intended to be short-term attempts at reducing damage occurring at the time those methods are employed. Long-term solutions to managing mammal damage would include limited habitat manipulations and changes in cultural practices, which are addressed further below and in Appendix B.

Non-lethal methods can disperse or otherwise make an area unattractive to mammals causing damage; thereby, reducing the presence of mammals at the site and potentially the immediate area around the site where non-lethal methods were employed. Non-lethal methods would be given priority when addressing requests for assistance (WS Directive 2.101). However, non-lethal methods would not necessarily be employed to resolve every request for assistance if deemed inappropriate by WS' personnel using the WS Decision Model, especially when the requesting entity had used non-lethal methods previously and found those methods to be inadequate to resolving the damage or threats of damage. Non-lethal methods would be used to exclude, harass, and disperse target wildlife from areas where damage or threats were occurring. When effective, non-lethal methods would disperse mammals from an area resulting in a reduction in the presence of those mammals at the site where those methods were employed. For any management methods employed, the proper timing would be essential in effectively dispersing those mammals causing damage. Employing methods soon after damage begins or soon after threats were identified, increases the likelihood that those damage management activities would achieve success in addressing damage. Therefore, coordination and timing of methods would be necessary to be effective in achieving expedient resolution of mammal damage.

Under the proposed action alternative, WS could employ only non-lethal methods when determined to be appropriate for each request for assistance to alleviate damage or reduce threats of damage using the WS Decision Model. In many situations, the cooperating entity has tried to employ non-lethal methods to resolve damage prior to contacting WS for assistance. In those cases, the methods employed by the requester were either unsuccessful or the reduction in damage or threats had not reached a level that was

tolerable by the requesting entity. In those situations, WS could employ other non-lethal methods, attempt to apply the same non-lethal methods, or employ lethal methods. In many situations, the implementation of non-lethal methods such as exclusion-type barriers would be the responsibility of the requestor, which means that, in those situations, WS only function would be to implement lethal methods, if determined to be appropriate using the WS Decision Model.

Lethal methods could be employed to resolve damage associated with those mammals identified by WS as responsible for causing damage or threats to human safety under this alternative; however, WS would only employ lethal methods after receiving a request for the use of those methods. The use of lethal methods would result in local population reductions in the area where damage or threats were occurring since mammals would be removed from the population. Lethal methods would often be employed to reinforce non-lethal methods and to remove mammals that were identified as causing damage or posing a threat to human safety. The use of lethal methods would result in local reductions of mammals in the area where damage or threats were occurring. The number of mammals removed from the population using lethal methods under the proposed action would be dependent on the number of requests for assistance received, the number of mammals involved with the associated damage or threat, and the efficacy of methods employed.

Often of concern with the use of lethal methods is that mammals that were lethally taken would only be replaced by other mammals either during the application of those methods (*e.g.*, mammals that relocate into the area) or by mammals the following year (*e.g.*, increase in reproduction and survivability that could result from less competition). As stated previously, the use of lethal methods would not be used as population management tools over broad areas. The use of lethal methods would be intended to reduce the number of mammals present at a specific location where damage was occurring by targeting those mammals causing damage or posing threats. Since the intent of lethal methods would be to manage only those mammals causing damage and not to manage entire mammal populations, those methods would not be ineffective because mammals return.

Most lethal and non-lethal methods currently available provide only short-term benefits when addressing mammal damage. The use of those methods would be intended to reduce damage occurring at the time those methods were employed but do not necessarily ensure mammals would not return once those methods were discontinued. Long-term solutions to resolving mammal damage would often be difficult to implement and can be costly. In some cases, long-term solutions involve exclusionary devices, such as fencing, or other practices that would not be costly or difficult to implement such as closing garbage cans. When addressing mammal damage, long-term solutions generally involve modifying existing habitat or making conditions to be less attractive to mammals. To ensure complete success, alternative sites in areas where damage was not likely to occur would often be required to achieve complete success in reducing damage and to avoid moving the problem from one area to another. Modifying a site to be less attractive to mammals would likely result in the dispersal of those mammals to other areas where damage could occur or could result in multiple occurrences of damage situations.

WS may recommend mammals be harvested during the regulated hunting and/or trapping season for those species in an attempt to reduce the number of mammals causing damage. Managing mammal populations over broad areas could lead to a decrease in the number of mammals causing damage. Establishing hunting or trapping seasons and the allowed take during those seasons is the responsibility of the TWRA. WS does not have the authority to establish hunting or trapping seasons or to set allowed harvest numbers during those seasons.

A complete list of chemical and non-chemical methods available for use under the identified alternatives, except the alternative with no damage management (Alternative 3), can be found in Appendix B. However, listing methods neither implies that all methods would be used by WS to resolve requests for

assistance nor does the listing of methods imply that all methods would be used to resolve every request for assistance. As part of an integrated approach, WS may provide technical assistance and direct operational assistance to those persons experiencing damage associated with mammals.

Technical Assistance Recommendations

Under the proposed action, WS would provide technical assistance to those persons requesting assistance with managing damage as part of an integrated approach. Technical assistance would occur as described in Alternative 2 of this EA. From FY 2005 through FY 2011, WS conducted 3,062 technical assistance projects that involved mammal damage to agricultural resources, property, natural resources, and threats to human safety (see Table 1.1).

Operational Damage Management Assistance

Operational damage management assistance would include damage management activities that were directly conducted by or supervised by personnel of WS. Operational damage management assistance may be initiated when the problem cannot effectively be resolved through technical assistance alone and there was a written MOU, cooperative service agreement, or other comparable document signed between WS and the entity requesting assistance. The initial investigation by WS' personnel would define the nature, history, and extent of the problem; species responsible for the damage; and methods available to resolve the problem. The professional skills of WS' personnel could be required to effectively resolve problems, especially if chemical methods were necessary or if the problems were complex.

The following examples serve as illustrations of WS' operational damage management assistance projects. The examples are intended to present realistic examples of on-going projects only and are not an inclusive or all-encompassing list of all projects conducted by WS in Tennessee.

Management of Wildlife Hazards to Aircraft and Air Passengers in Tennessee

Upon receiving a request for assistance, WS evaluates wildlife hazards at an airport, prepares a Wildlife Hazard Assessment that identifies wildlife hazards, and assists the airport in developing a Wildlife Hazard Management Plan to address those hazards and threats.

Direct operational activities consist of various harassment techniques, and live capture and lethal removal techniques aimed at removing potentially injurious wildlife. WS' personnel also provide ongoing technical advice to airport managers regarding methodologies to reduce the presence of wildlife in airport environments, including providing technical advice on various habitat management projects implemented by airport personnel. In addition, WS promotes improved mammal strike record keeping, maintains a program of mammal identification, and monitors mammal numbers at participating airports to assist in developing an effective damage management program.

Management of Mammals That Threaten Power Generation and Transmission in Tennessee

Upon receiving a request for assistance, WS conducts site visits and evaluates damage caused by various mammals on properties and facilities owned or managed by the TVA. Once WS' evaluation is complete and mammal damage is addressed through technical assistance or direct operational activities, each damage site then becomes part of a damage monitoring schedule. Periodic monitoring of these known damage sites allows WS to better manage wildlife damage issues for the TVA by quickly identifying the repopulation of mammals that have caused specific damage in the past, identifying signs that damage is about to occur again (*e.g.*, beaver may rebuild dams, woodchucks may open up burrows), and removing the offending species prior to damage occurring. Direct operational activities may consist of utilizing

snares, body-grip traps, and padded foothold traps to remove aquatic rodents and woodchucks burrowing in earthen levees or flooding sensitive areas; live traps to capture mammals such as raccoons, fox, and skunks that pose threats to power transmission by chewing, denning, or otherwise shorting out electrical circuits; identifying species and installing exclusion mechanisms to keep bats from roosting in high human traffic areas of power production facilities; and shooting or otherwise lethally removing aggressive mammals that threaten TVA personnel safety or are depredating on protected T&E species that are nesting on property owned or managed by the TVA.

Management of Aquatic Rodents in Tennessee

WS conducts site visits and evaluates damage caused by aquatic rodents when requested. WS' personnel provide technical assistance and demonstration of techniques available for use by the requestor. Direct operational activities may consist of utilizing snares, body grip traps and padded foothold traps to remove the rodents causing damage. WS' personnel then determine if beaver dams can be removed in accordance with the Clean Water Act (see Appendix E). Once the determination that dam removal is appropriate and legal, it is conducted manually or when safe and appropriate, with use of explosives. In some instances, WS' personnel install hardware cloth to protect specific trees most susceptible to loss by beavers or install water flow devices to maintain water levels at appropriate levels instead of lethally removing beaver colonies.

Management of Feral Swine in Tennessee

WS evaluates agricultural damage or disease transmission caused by feral swine. Direct operational activities consist of various lethal removal techniques, including corral trapping, snaring, and shooting. In some cases, WS works with adjoining landowners to establish large cooperative land masses suitable for aerial operations to pursue feral swine with aircraft. WS' personnel demonstrate techniques for excluding feral swine from specific areas and utilize harassment techniques to provide time for agricultural crops to mature and become less attractive.

Management of Domestic and Exotic Mammals in Tennessee

Upon request for assistance, WS participates in emergency response situations where there is a need to capture or lethally control domestic or exotic mammals due to natural disasters, accidental releases, or disease outbreaks. Direct operational activities include various lethal and non-lethal removal techniques, including corral trapping, snaring and shooting. While these cases are rare, WS' personnel are specially trained to respond to emergency response situations and have the skills and tools necessary to complement and support efforts of various state agencies that would take the lead in responding to these situations.

Educational Efforts

Education is an important element of activities because wildlife damage management is about finding balance and coexistence between the needs of people and needs of wildlife. This is extremely challenging as nature has no balance, but rather is in continual flux. In addition to the routine dissemination of recommendations and information to individuals or organizations, WS provides lectures, courses, and demonstrations to producers, homeowners, state and county agents, colleges and universities, and other interested groups. WS frequently cooperates with other entities in education and public information efforts. Additionally, technical papers have been and would continue to be presented at professional meetings and conferences so that other wildlife professionals and the public were periodically updated on recent developments in damage management technology, programs, laws and regulations, and agency policies.

Research and Development

The National Wildlife Research Center (NWRC) functions as the research unit of WS by providing scientific information and the development of methods for wildlife damage management, which are effective and environmentally responsible. Research biologists with the NWRC work closely with wildlife managers, researchers, and others to develop and evaluate methods and techniques for managing wildlife damage. For example, research biologists from the NWRC were involved with developing and evaluating the reproductive inhibitor known under the trade name of Gonacon™. Research biologists with the NWRC have authored hundreds of scientific publications and reports based on research conducted involving wildlife and methods.

WS' Decision Making Procedures

WS' personnel would use a thought process for evaluating and responding to damage complaints that is depicted by the WS Decision Model (WS Directive 2.201) and described by Slate et al. (1992). WS' personnel would assess the problem and then evaluate the appropriateness and availability (legal and administrative) of strategies and methods based on biological, economic, and social considerations. Following this evaluation, methods deemed to be practical for the situation would be incorporated into a damage management strategy. After this strategy was implemented, monitoring would be conducted and evaluation would continue to assess the effectiveness of the strategy. If the strategy were effective, the need for further management would be ended. In terms of the WS Decision Model, most efforts to resolve wildlife damage consist of continuous feedback between receiving the request and monitoring the results of the damage management strategy. The Decision Model is not a written documented process, but a mental problem-solving process common to most, if not all, professions, including WS.

Community-based Decision Making

WS could receive requests for assistance from community leaders and/or representatives. In those situations, the WS program in Tennessee, under this alternative, would follow the "*co-managerial approach*" to solve wildlife damage or conflicts as described by Decker and Chase (1997) when receiving a request for assistance from a community leader or representative. Under a community based decision-making process, WS would provide information, demonstration, and discussion on available methods to the appropriate representatives of the community for which services were requested to ensure a community-based decision was made. By involving decision-makers in the process, damage management actions could be presented to allow decisions on damage management to involve those individuals that the decision-maker(s) represents. As addressed in this EA, WS would provide technical assistance to the appropriate decision-maker(s) to allow for information on damage management activities to be presented to those persons represented by the decision-maker(s), including demonstrations and presentation by WS at public meetings to allow for involvement of the community. Requests for assistance to manage damage caused by mammals often originate from the decision-maker(s) based on community feedback or from concerns about damage or threats to human safety. As representatives of the community, the decision-maker(s) would be able to provide the information to local interests either through technical assistance provided by WS or through demonstrations and presentation by WS on damage management activities. This process would allow decisions on damage management activities to be made based on local input. The community leaders could implement management recommendations provided by WS or others, or may request management assistance from WS, other wildlife management agencies, local animal control agencies, or private businesses or organizations.

Community Decision-Makers

The decision-maker for the local community would be elected officials or representatives of the communities. The elected officials or representatives are popularly elected residents of the local community or appointees who oversee the interests and business of the local community. This person or persons would represent the local community's interest and make decisions for the local community or bring information back to a higher authority or the community for discussion and decision-making. Identifying the decision-maker for local business communities can be more complex because building owners may not indicate whether the business must manage wildlife damage themselves, or seek approval to manage wildlife from the property owner or manager, or from a governing Board. WS could provide technical assistance and make recommendations for damage reduction to the local community or local business community decision-maker(s). Direct assistance could be provided by WS only if requested by the local community decision-maker, funding is provided, and if the requested direct control was compatible with WS' recommendations.

Private Property Decision-Makers

WS could also receive requests for assistance from private property owners. In the case of private property owners, the decision-maker is the individual that owns or manages the affected property. The decision-maker has the discretion to involve others as to what occurs or does not occur on property they own or manage. Due to privacy issues, WS cannot disclose cooperator information to others. Therefore, in the case of an individual property owner or manager, the involvement of others and to what degree others are involved in the decision-making process would be a decision made by that individual. Direct control could be provided by WS if requested, funding was provided, and the requested management was in according with WS' recommendations.

Public Property Decision-Makers

Request for WS' assistance could originate from public property owners or managers. The decision-maker for local, state, or federal property would be the official responsible for or authorized to manage the public land to meet interests, goals, and legal mandates for the property. WS could provide technical assistance to this person and recommendations to reduce damage. Direct control could be provided by WS if requested, funding provided, and the requested actions were within the recommendations made by WS.

Alternative 2 – Mammal Damage Management by WS through Technical Assistance Only

Under this alternative, WS would provide those cooperators requesting assistance with technical assistance only. Similar to Alternative 1, WS could receive requests for assistance from community representatives, private individuals/businesses, or from public entities. Technical assistance would provide those cooperators experiencing damage or threats associated with mammals with information, demonstrations, and recommendations on available and appropriate methods available. The implementation of methods and techniques to resolve or prevent damage would be the responsibility of the requester with no direct involvement by WS. In some cases, WS may provide supplies or materials that are of limited availability for use by private entities (*e.g.*, loaning of propane cannons). Technical assistance may be provided through a personal or telephone consultation, or during an on-site visit with the requester. Generally, several management strategies would be described to the requester for short and long-term solutions to managing damage; these strategies would be based on the level of risk, need, and the practicality of their application. WS would use the Decision Model to recommend those methods and techniques available to the requester to manage damage and threats of damage. Those persons receiving

technical assistance from WS could implement those methods recommended by WS, could employ other methods not recommended by WS, could seek assistance from other entities, or take no further action.

Under a technical assistance only alternative, WS would recommend an integrated approach similar to the proposed action alternative (Alternative 1) when receiving a request for assistance; however, WS would not provide direct operational assistance under this alternative. Preference would be given to non-lethal methods when practical and effective under this alternative (see WS Directive 2.101). Recommendation of methods and techniques by WS to resolve damage would be based on information provided by the individual seeking assistance using the WS Decision Model. In some instances, wildlife-related information provided to the requestor by WS would result in tolerance/acceptance of the situation. In other instances, damage management options would be discussed and recommended. Only those methods legally available for use by the appropriate individual would be recommend or loaned by WS. Similar to Alternative 1, those methods described in Appendix B would be available to those persons experiencing damage or threats associated with mammals in the State except for immobilizing drugs, euthanasia drugs, reproductive inhibitors, and shooting from aircraft. Immobilizing drugs and euthanasia drugs would only be available to WS or appropriately licensed veterinarians. Under this alternative, the reproductive inhibitor available under the trade name of Gonacon™ would only be available for use by the TWRA. At the time this EA was developed, Gonacon™ was not registered for use in the State.

The WS program in the State regularly provides technical assistance to individuals, organizations, and other federal, state, and local government agencies for managing mammal damage. Technical assistance includes collecting information about the species involved, the nature and extent of the damage, and previous methods that the cooperator has attempted to resolve the problem. WS then provides information on appropriate methods that the cooperator may consider to resolve the damage themselves. Types of technical assistance projects may include a visit to the affected property, written communication, telephone conversations, or presentations to groups such as homeowner associations or civic leagues. Between FY 2006 and FY 2011, WS has conducted 3,062 technical assistance projects that involved mammal damage to agricultural resources, property, natural resources, and threats to human safety.

This alternative would place the immediate burden of operational damage management work on the resource owner, other governmental agencies, and/or private businesses. Those persons experiencing damage or are concerned with threats posed by mammals could seek assistance from other governmental agencies, private entities, or conduct damage management on their own. Those persons experiencing damage or threats could take action using those methods legally available to resolve or prevent mammal damage as permitted by federal, state, and local laws and regulations or those persons could take no action.

Alternative 3 – No Mammal Damage Management Conducted by WS

This alternative would preclude all activities by WS to reduce threats to human health and safety, and to alleviate damage to agricultural resources, property, and natural resources. WS would not be involved with any aspect of mammal damage management in the State. All requests for assistance received by WS to resolve damage caused by mammals would be referred to the TWRA, other governmental agencies, and/or other private entities.

Despite no involvement by WS in resolving damage and threats associated with mammals in the State, those persons experiencing damage caused by mammals could continue to resolve damage by employing those methods legally available since the take of mammals to alleviate damage or threats can occur despite the lack of involvement by WS. The take of mammals by other entities could occur through the issuance of permits by the TWRA, when required, and during the hunting or trapping seasons. Landowners, agricultural leaseholders and their designated agents can lethally take coyotes, fox, skunks,

and feral swine at any time when those species are causing damage in accordance with appropriate regulations (TWRA 2012*a*). All methods described in Appendix B would be available for use by those persons experiencing damage or threats except for the use of immobilizing drugs and euthanasia chemicals. Gonacon™ would not be used by WS under this alternative but would be available to the TWRA, if registered for use in the State. Immobilizing drugs and euthanasia chemicals can only be used by WS or appropriately licensed veterinarians.

Therefore, under this alternative, those persons experiencing damage or threats of damage could contact WS but WS would immediately refer the requester to the TWRA and/or other entities, the requester could contact other entities for information and assistance with managing damage, could take actions to alleviate damage without contacting any entity, or could take no action.

3.2 ALTERNATIVES CONSIDERED BUT NOT ANALYZED IN DETAIL WITH RATIONALE

In addition to those alternatives analyzed in detail, several additional alternatives were identified by WS and the TVA. However, those alternatives will not receive detailed analyses for the reasons provided. Those alternatives considered but not analyzed in detail include:

Non-lethal Methods Implemented Before Lethal Methods

This alternative would require that non-lethal methods or techniques described in Appendix B be applied to all requests for assistance to reduce damage and threats to safety from mammals in the State. If the use of non-lethal methods fails to resolve the damage situation or reduce threats to human safety at each damage situation, lethal methods would be employed to resolve the request. Non-lethal methods would be applied to every request for assistance regardless of severity or intensity of the damage or threat until deemed inadequate to resolve the request. This alternative would not prevent the use of lethal methods by other entities or by those persons experiencing mammal damage but would only prevent the use of those methods by WS until non-lethal methods had been employed.

Those persons experiencing damage often employ non-lethal methods to reduce damage or threats prior to contacting WS. Verification of the methods used would be the responsibility of WS. No standard exists to determine requester diligence in applying those methods, nor are there any standards to determine how many non-lethal applications are necessary before the initiation of lethal methods. Thus, only the presence or absence of non-lethal methods can be evaluated. The proposed action (Alternative 1) and the technical assistance only alternative (Alternative 2) are similar to a non-lethal before lethal alternative because WS would use or recommend non-lethal methods before lethal methods (WS Directive 2.101). Adding a non-lethal before lethal alternative and the associated analysis would not add additional information to the analyses in the EA.

Use of Non-lethal Methods Only by WS

Under this alternative, WS would be required to implement non-lethal methods only to resolve damage caused by mammals in the State. Only those methods discussed in Appendix B that are considered non-lethal would be employed by WS. No lethal take of mammals would occur by WS. The use of lethal methods could continue to be used under this alternative by other entities or by those persons experiencing damage by mammals. The non-lethal methods used or recommended by WS under this alternative would be identical to those non-lethal methods identified in any of the alternatives.

In situations where non-lethal methods were impractical or ineffective to alleviate damages, WS could refer requests for information regarding lethal methods to the TWRA, local animal control agencies, or

private businesses or organizations. Under this alternative, however, property owners/managers might be limited to using non-lethal methods only as they may have difficulty obtaining permits for lethal methods.

Property owners or managers could conduct management using any method that was legal. Property owners or managers might choose to implement WS' non-lethal recommendations, implement lethal methods, or request assistance from a private or public entity other than WS. Property owners/managers frustrated by the lack of WS' assistance with the full range of mammal damage management techniques may try methods not recommended by WS or use illegal methods (*e.g.*, poisons). In some cases, property owners or managers may misuse some methods or use some methods in excess of what is necessary, which could then become hazardous and pose threats to the safety of humans and non-target species.

The proposed action, using an integrated damage management approach, incorporates the use of non-lethal methods when addressing requests for assistance. In those instances where non-lethal methods would effectively resolve damage from mammals those methods would be used or recommended under the proposed action. Since non-lethal methods would be available for use under the alternatives analyzed in detail, this alternative would not add to the analyses. Those mammals that could be lethally removed by WS under any of the alternatives could be removed by those persons experiencing damage or threats even if WS was not involved.

Use of Lethal Methods Only by WS

This alternative would require the use of lethal methods only to reduce threats and damage associated with mammals. However, non-lethal methods can be effective in preventing damage in certain instances. Under WS Directive 2.101, WS must consider the use of non-lethal methods before lethal methods. Non-lethal methods have been effective in alleviating mammal damage. For example, the use of one-way exclusion devices can be effective at allowing bats to exit a structure but prevent re-entry. Once bats have exited the structure, structural repairs could be completed to permanently prevent re-entry of bats. In those situations where damage could be alleviated using non-lethal methods deemed effective, those methods would be employed or recommended as determined by the WS Decision Model. Therefore, this alternative was not considered in detail.

Trap and Translocate Mammals Only

Under this alternative, all requests for assistance would be addressed using live-capture methods or the recommendation of live-capture methods. Mammals would be live-captured using immobilizing drugs, live-traps, cannon nets, or rocket nets. All mammals live-captured through direct operational assistance by WS would be translocated. Translocation sites would be identified and have to be approved by the TWRA and/or the property owner where the translocated mammals would be placed prior to live-capture and translocation. The translocation of mammals could only occur under the authority of the TWRA. The translocation of mammals by WS would only occur as directed by the TWRA. When requested by the TWRA, WS could translocate mammals or recommend translocation under any of the alternatives analyzed in detail, except under the no involvement by WS alternative (Alternative 3). However, other entities could translocate mammals under Alternative 3.

The translocation of mammals that have caused damage or pose threats of damage to other areas following live-capture generally would not be effective or cost-effective. Translocation is generally ineffective because problem mammal species are highly mobile and can easily return to damage sites from long distances, habitats in other areas are generally already occupied, and translocation would most likely result in mammal damage problems at the new location. Also, hundreds of mammals would need to be captured and translocated to solve some damage problems (*e.g.*, deer confined within a perimeter fence); therefore, translocation would be unrealistic. Translocation of wildlife is also discouraged by WS

policy (WS Directive 2.501) because of the stress to the translocated animal, the potential for spreading diseases, poor survival rates, and the difficulties that translocated wildlife have with adapting to new locations or habitats (Nielsen 1988). Since WS does not have the authority to translocate mammals in the State unless permitted by the TWRA, this alternative was not considered in detail.

Reducing Damage by Managing Mammal Populations through the Use of Reproductive Inhibitors

Under this alternative, the only method that would be available to resolve requests for assistance by WS would be the recommendation and the use of reproductive inhibitors to reduce or prevent reproduction in mammals responsible for causing damage. Reproductive inhibitors are often considered for use where wildlife populations are overabundant and where traditional hunting or lethal control programs are not publicly acceptable (Muller et al. 1997). Use and effectiveness of reproductive control as a wildlife population management tool is limited by population dynamic characteristics (*e.g.*, longevity, age at onset of reproduction, population size and biological/cultural carrying capacity), habitat and environmental factors (*e.g.*, isolation of target population, cover types, and access to target individuals), socioeconomic, and other factors.

Reproductive control for wildlife could be accomplished through either sterilization (permanent) or contraception (reversible). Sterilization could be accomplished through: 1) surgical sterilization (vasectomy, castration, and tubal ligation), 2) chemosterilization, and 3) through gene therapy. Contraception could be accomplished through: 1) hormone implantation (synthetic steroids such as progestins), 2) immunocontraception (contraceptive vaccines), and 3) oral contraception (progestin administered daily).

Population modeling indicates that reproductive control is more efficient than lethal control only for some rodent and small bird species with high reproductive rates and low survival rates (Dolbeer 1998). Additionally, the need to treat a sufficiently large number of target animals, multiple treatments, and population dynamics of free-ranging populations place considerable logistic and economic constraints on the adoption of reproduction control technologies as a wildlife management tool for some species.

Currently, reproductive inhibitors are not available for use to manage most mammal populations. Given the costs associated with live-capturing and performing sterilization procedures on mammals and the lack of availability of chemical reproductive inhibitors for the management of most mammal populations, this alternative was not evaluated in detail. If a reproductive inhibitor becomes available to manage a large number of mammal populations that proves to be effective in reducing localized mammal populations, the use of the inhibitor could be evaluated under the proposed action as a method available that could be used in an integrated approach to managing damage. This EA would be reviewed and supplemented to the degree necessary to evaluate the use of the reproductive inhibitor as part of an integrated approach described under the proposed action. Currently, the only reproductive inhibitor that is registered with the EPA is Gonacon™, which is registered for use on white-tailed deer only. However, Gonacon™ is not currently registered for use in the State. Reproductive inhibitors for the other mammal species addressed in this EA do not currently exist.

Compensation for Mammal Damage

The compensation alternative would require WS to establish a system to reimburse persons impacted by mammal damage and to seek funding for the program. Under such an alternative, WS would continue to provide technical assistance to those persons seeking assistance with managing damage. In addition, WS would conduct site visits to verify damage. Evaluation of this alternative indicates that a compensation only alternative has many drawbacks. Compensation would: 1) require large expenditures of money and labor to investigate and validate all damage claims, and to determine and administer appropriate

compensation, 2) compensation most likely would be below full market value, 3) give little incentive to resource owners to limit damage through improved cultural or other practices and management strategies, and 4) not be practical for reducing threats to human health and safety.

Short Term Eradication and Long Term Population Suppression

An eradication alternative would direct all WS' program efforts toward total long-term elimination of mammal populations wherever a cooperative program was initiated in Tennessee. Eradication as a general strategy for managing mammal damage was not considered in detail because State and federal agencies with interest in, or jurisdiction over, wildlife oppose eradication of any native wildlife species and eradication is not acceptable to most people.

Suppression would direct WS' program efforts toward managed reduction of certain problem populations or groups. In areas where damage can be attributed to localized populations of mammals, WS could decide to implement local population suppression using the WS' Decision Model. However, it is not realistic or practical to consider large-scale population suppression as the basis of the WS' program. Problems with the concept of suppression are similar to those described above for eradication. Typically, WS' activities in Tennessee would be conducted on a very small portion of the sites or areas inhabited or frequented by problem species.

Bounties

Payment of funds (bounties) for killing some mammals suspected of causing economic losses have not been supported by state fish and game agencies, such as the TWRA, as well as most wildlife professionals for many years (Latham 1960, Hoagland 1993). WS concurs with those agencies and wildlife professionals because of several inherent drawbacks and inadequacies in the payment of bounties. Bounties are often ineffective at controlling damage over a wide area, such as across the entire State. The circumstances surrounding the take of animals are typically arbitrary and completely unregulated because it is difficult or impossible to assure animals claimed for bounty were not taken from outside the area where damage was occurring. In addition, WS does not have the authority to establish a bounty program.

Trap-Neuter-Release Program for Feral and Free Ranging Cats and/or Dogs

This topic has undergone considerable debate in animal welfare and scientific communities for a number of years. The debate focuses on whether controlling feral, free-ranging, or invasive animal populations through Trap-Neuter-Release (TNR) programs are effective and alleviate problems (*i.e.*, diseases, predation, agricultural damage, and human safety).

Theoretically, TNR would work if all animals of one sex or both were sterilized. However, the probability of controlling invasive species in the wild with this technique is not currently reasonable; especially, with the animals being self-sufficient and not relying on humans to survive. Additionally, some individuals within a population can be trap-shy. Capturing or removing trap-shy individuals often requires implementing other methods.

In addition, the National Association of State Public Health Veterinarians and the AVMA oppose TNR programs based on health concerns and threats (AVMA 2003). Of major concern would be the potential for disease and parasite transmission to humans from direct contact during sterilization or the risk of exposure after the animal was released. Once live-captured, performing sterilization procedures during field operations on anesthetized animals would be difficult. Sanitary conditions would be difficult to maintain when performing surgical procedures in field conditions. To perform operations under appropriate conditions, live-captured animals would need to be transported from the capture site to an

appropriate facility, which increases the threat from handling and transporting. A mobile facility could be used; however, a mobile facility would still require additional handling and transporting of the live-captured animals to the facility. Once the surgical procedure was completed, the animal would have to be held to ensure recovery and transported back to the area where capture occurred.

TNR programs are often not as successful as desired and needed to reduce immediate threats posed by wildlife, especially when human safety is a concern (AVMA 2003, Barrows 2004, Levy and Crawford 2004, Jessup 2004, Winter 2004, AVMA 2009). Feral animals subjected to a TNR program would continue to cause the same problems[9] they caused before the TNR program was initiated because of slow attrition. TNR programs can take a decade or longer to reduce target species populations (Barrows 2004, Winter 2004); especially, when acute issues need rapid solutions (Levy and Crawford 2004, Stoskopf and Nutter 2004). Several studies report that target species' populations often remain stable or increase following TNR programs due to immigration and reproduction from other members of the groups (Castillo and Clarke 2003, Levy and Crawford 2004, Winter 2004) with little to no resolution of threats to human safety or damages (Barrows 2004, Slater 2004, Winter 2004).

Other concerns arise when considering the legality of TNR programs given the documented damage caused by target species, especially to native wildlife (Barrows 2004, Levy and Crawford 2004, Jessup 2004). Some people have questioned whether TNR programs are violating the Migratory Bird Treaty Act and the ESA because released animals may continue to kill migratory birds and/or endangered species (Barrows 2004, Levy and Crawford 2004, Jessup 2004). Because of the continued threat to human safety created by TNR programs and the continued threat to T&E wildlife and native wildlife in general, this alternative will not be considered further.

3.3 STANDARD OPERATING PROCEDURES FOR MAMMAL DAMAGE MANAGEMENT

SOPs improve the safety, selectivity, and efficacy of activities intended to resolve wildlife damage. The WS program in Tennessee uses many such SOPs. Those SOPs would be incorporated into activities conducted by WS under the appropriate alternatives when addressing mammal damage and threats in the State.

Some key SOPs pertinent to resolving mammal damage in the State include the following:

- The WS Decision Model, which is designed to identify effective strategies to managing wildlife damage and their potential impacts, would be consistently used and applied when addressing mammal damage.

- EPA-approved label directions would be followed for all pesticide use. The registration process for chemical pesticides is intended to assure minimal adverse effects occur to the environment when chemicals are used in accordance with label directions.

- Immobilizing and euthanasia drugs would be used according to the DEA, FDA, and WS' directives and procedures.

- All controlled substances would be registered with the DEA or the FDA.

[9] Brickner (2003), Levy et al. (2003), Barrows (2004), and Jessup (2004) reported that sterilized cats that do not spend any time on courting and mating are left with more time to hunt than non-sterilized cats and therefore, continue to remain as potential reservoirs of animal and human disease, a social nuisance, and continue to hunt and kill protected species.

- WS' employees would follow approved procedures outlined in the WS' Field Manual for the Operational Use of Immobilizing and Euthanizing Drugs (Johnson et al. 2001).

- WS' employees that use controlled substances would be trained to use each material and would be certified to use controlled substances.

- WS' employees who use pesticides and controlled substances would participate in State-approved continuing education to keep current of developments and maintain their certifications.

- Pesticide and controlled substance use, storage, and disposal would conform to label instructions and other applicable laws and regulations, and Executive Order 12898.

- Material Safety Data Sheets for pesticides and controlled substances would be provided to all WS' personnel involved with specific damage management activities.

- All personnel who use firearms would be trained according to WS' Directives.

- WS' employees participating in any aspect of aerial wildlife operations would be trained and/or certified in their role and responsibilities during the operations. All WS' personnel would follow the policies and directives set forth in WS' Directive 2.620; WS' Aviation Operations Manual; WS' Aviation Safety Manual and its amendments; Title 14 CFR; and Federal Aviation Regulations (FAR), Part 43, 61, 91, 119, 133, 135, and 137.

- The use of non-lethal methods would be considered prior to the use of lethal methods when managing mammal damage.

- The take of mammals by WS under the proposed action alternative would only occur when authorized by the TWRA, when applicable, and only at levels authorized.

- Management actions would be directed toward localized populations, individuals, or groups of target species. Generalized population suppression across Tennessee, or even across major portions of Tennessee, would not be conducted.

- Non-target animals live-captured in traps would be released unless it is determined that the animal would not survive and/or that the animal could not be released safely.

3.4 ADDITIONAL STANDARD OPERATING PROCEDURES SPECIFIC TO THE ISSUES

Several additional SOPs are applicable to the alternatives and the issues identified in Chapter 2 including the following:

Issue 1 - Effects of Mammal Damage Management Activities on Target Mammal Populations

- Lethal take of mammals by WS would be reported and monitored by WS and the TWRA to evaluate population trends and the magnitude of WS' take of mammals in the State.

- WS would only target those individuals or groups of target species identified as causing damage or posing a threat to human safety.

- The WS' Decision Model, designed to identify the most appropriate damage management strategies and their impacts, would be used to determine strategies for resolving mammal damage.

- WS would monitor activities to ensure those activities do not adversely affect mammal populations in the State.

- Preference would be given to non-lethal methods, when practical and effective.

Issue 2 - Effects of Mammal Damage Management Activities on Non-target Wildlife Species Populations, Including T&E Species

- When conducting removal operations via shooting, identification of the target would occur prior to application.

- As appropriate, suppressed firearms would be used to minimize noise impacts.

- Personnel would use lures, trap placements, and capture devices that would be strategically placed at locations likely to capture a target animal and minimize the potential of non-target animal captures.

- Any non-target animals captured in cage traps, nets, or any other restraining device would be released whenever it is possible and safe to do so.

- Live-traps would be checked frequently to ensure non-target species would be released in a timely manner to ensure survival.

- Carcasses of mammals retrieved after damage management activities have been conducted would be disposed of in accordance with WS Directive 2.515.

- WS has consulted with the USFWS and the TWRA to evaluate activities to resolve mammal damage and threats to ensure the protection of T&E species.

- WS would monitor activities conducted under the selected alternative, if activities are determined to have no significant impact on the environment and an EIS is not required, to ensure those activities do not negatively impact non-target species.

Issue 3 - Effects of Mammal Damage Management Activities on Human Health and Safety

- Damage management activities would be conducted professionally and in the safest manner possible. Whenever possible, damage management activities would be conducted away from areas of high human activity. If this were not possible, then activities would be conducted during periods when human activity is low (*e.g.*, early morning).

- Shooting would be conducted during times when public activity and access to the control areas was restricted. Personnel involved in shooting operations would be fully trained in the proper and safe application of this method.

- All personnel employing chemical methods would be properly trained and certified in the use of those chemicals. All chemicals used by WS would be securely stored and properly monitored to ensure the safety of the public. WS' use of chemicals and training requirements to use those chemicals are outlined in WS Directive 2.401 and WS Directive 2.430.

- All chemical methods used by WS or recommended by WS would be registered with the EPA, DEA, FDA, and/or the TDA, as appropriate.

- WS would adhere to all established withdrawal times for mammals when using immobilizing drugs for the capture of mammals that are agreed upon by WS, the TWRA, and veterinarian authorities. Although unlikely, in the event that WS was requested to immobilize mammals either during a time when harvest of those mammal species is occurring or during a time where the withdrawal period could overlap with the start of a harvest season, WS would euthanize the animal or mark the animal with ear tags labeled with a "*do not eat*" warning and appropriate contact information.

- Carcasses of mammals retrieved after damage management activities would be disposed of in accordance with WS Directive 2.515.

Issue 4 - Effects of Mammal Damage Management Activities on the Socio-cultural Elements of the Human Environment

- Management actions to reduce or prevent damage caused by mammals would be directed toward specific individuals identified as responsible for the damage, identified as posing a threat to human safety, or identified as posing a threat of damage.

- All methods or techniques applied to resolve damage or threats to human safety would be agreed upon by entering into a cooperative service agreement, MOU, or comparable document prior to the implementation of those methods.

- Preference would be given to non-lethal methods, when practical and effective.

Issue 5 - Humaneness and Animal Welfare Concerns of Methods

- Personnel would be well trained in the latest and most humane devices/methods for removing mammals causing damage.

- WS' personnel would check methods frequently to ensure mammals captured would be addressed in a timely manner to minimize the stress of being restrained.

- When deemed appropriate using the WS' Decision Model, WS' use of lethal methods would comply with WS' directives (WS Directive 2.505, WS Directive 2.430).

- The NWRC is continually conducting research to improve the selectivity and humaneness of wildlife damage management devices used by personnel in the field.

- The use of non-lethal methods would be considered prior to the use of lethal methods when managing mammal damage.

Issue 6 - Effects of Mammal Damage Management Activities on the Regulated Harvest of Mammals

- Management actions to reduce or prevent damage caused by mammals in the State would be directed toward specific individuals identified as responsible for the damage, identified as posing a threat to human safety, or identified as posing a threat of damage.

- WS' activities to manage damage and threats caused by mammals would be coordinated with the TWRA.

- WS' lethal take (killing) of mammals would be reported to and monitored by the TWRA to ensure WS' take has been considered as part of management objectives for those mammal species in the State.

- WS would monitor activities to ensure those activities do not adversely affect mammal populations in the State.

Issue 7 – Effects of Beaver Removal and Dam Manipulation on the Status of Wetlands in the State

- WS' personnel would remove beaver dams in accordance with federal and state laws and regulations for environmental protection. Beaver dam removal would be conducted to restore drainage or the stream channel for an area, or if an area has an established silvicultural or other agricultural, commercial/industrial activity, and where such an area has not become an established wetland.

- Upon receiving a request to remove beaver dams, WS would visually inspect the dam and the associated water impoundment to determine if characteristics exist at the site that would meet the definition of a wetland under section 404 of the Clean Water Act (40 CFR 232.2; see Issue 5 in Section 2.2 of this EA). If wetland conditions were present at the site, the entities requesting assistance from WS would be notified that a permit might be required to remove the dam and to seek guidance from the TDEC and the United States Army Corps of Engineers pursuant to Tennessee State Law and the Clean Water Act.

CHAPTER 4: ENVIRONMENTAL CONSEQUENCES

Chapter 4 provides information needed for making informed decisions in selecting the appropriate alternative to address the need for action described in Chapter 1 and the issues described in Chapter 2. This chapter analyzes the environmental consequences of each alternative as that alternative relates to the issues identified. The following resource values in the State are not expected to be significantly impacted by any of the alternatives analyzed: soils, geology, minerals, water quality/quantity, flood plains, critical habitats (areas listed in T&E species recovery plans), visual resources, air quality, prime and unique farmlands, aquatic resources, timber, and range. Those resources will not be analyzed further.

The activities proposed in the alternatives would have a negligible effect on atmospheric conditions, including the global climate. Meaningful direct or indirect emissions of greenhouse gases would not occur because of any of the proposed alternatives. Those alternatives would meet the requirements of applicable laws, regulations, and Executive Orders including the Clean Air Act and Executive Order 13514.

4.1 ENVIRONMENTAL CONSEQUENCES FOR ISSUES ANALYZED IN DETAIL

This section analyzes the environmental consequences of each alternative in comparison to determine the extent of actual or potential impacts on the issues. Therefore, the proposed action/no action alternative (Alternative 1) serves as the baseline for the analysis and the comparison of expected impacts among the alternatives. The analysis also takes into consideration mandates, directives, and the procedures of WS, the TWRA, and the TVA.

Issue 1 - Effects of Mammal Damage Management Activities on Target Mammal Populations

A common issue is whether damage management actions would adversely affect the populations of target mammal species, especially when lethal methods were employed. WS would maintain ongoing contact with the TWRA to ensure activities occurred within management objectives for those species. WS would submit annual activity reports to the TWRA. The TWRA would monitor the total take of mammals from all sources and would factor in survival rates from predation, disease, and other mortality data. Ongoing contact with the TWRA would assure local, state, and regional knowledge of wildlife population trends would be considered. As discussed previously, the analysis for magnitude of impact from lethal take can be determined either quantitatively or qualitatively. Quantitative determinations are based on population estimates, allowable harvest levels, and actual harvest data. Qualitative determinations are based on population trends and harvest trend data. Information on mammal populations and trends are often derived from several sources, including published literature and harvest data.

Methods available to address mammal damage or threats of damage in the State that would be available for use or recommendation under Alternative 1 (proposed action/no action alternative) and Alternative 2 (technical assistance only alternative) would either be lethal methods or non-lethal methods. Many of the methods would also be available to other entities under Alternative 3 (no involvement by WS alternative). The only methods that would not be available for use by other entities under Alternative 2 and Alternative 3 would be immobilizing drugs, euthanasia chemicals, and Gonacon™. Under Alternative 2, WS could recommend lethal and non-lethal methods as part of an integrated approach to resolving requests for assistance. Alternative 1 would address requests for assistance received by WS through technical and/or operational assistance where an integrated approach to methods would be employed and/or recommended. Non-lethal methods that would be available to WS under Alternative 1 would include, but would not be limited to habitat/behavior modification, pyrotechnics, visual deterrents, live traps, translocation, cable restraints, exclusionary devices, frightening devices, nets, immobilizing drugs, reproductive inhibitors, and chemical repellents (see Appendix B for a complete list and description of potential methods).

Non-lethal methods that would be available under all of the alternatives can disperse or otherwise make an area unattractive to mammals causing damage; thereby, reducing the presence of mammals at the site and potentially the immediate area around the site where non-lethal methods were employed. Non-lethal methods would be given priority by WS when addressing requests for assistance under Alternative 1 and Alternative 2 (see WS Directive 2.101). However, non-lethal methods would not necessarily be employed or recommended to resolve every request for assistance if deemed inappropriate by WS' personnel using the WS Decision Model. For example, if a cooperator requesting assistance had already used non-lethal methods, WS would not likely recommend or continue to employ those particular methods since their use had already been proven ineffective in adequately resolving the damage or threat.

Many non-lethal methods would be used to exclude, harass, and disperse target wildlife from areas where damage or threats were occurring. When effective, non-lethal methods would disperse mammals from the area resulting in a reduction in the presence of those mammals at the site where those methods were employed. However, mammals responsible for causing damage or threats would be dispersed to other areas with minimal impact on those species' populations. Non-lethal methods would not be employed over large geographical areas or applied at such intensity that essential resources (*e.g.*, food sources, habitat) would be unavailable for extended durations or over a wide geographical scope that long-term adverse effects would occur to a species' population. Non-lethal methods would generally be regarded as having minimal impacts on overall populations of wildlife since individuals of those species were unharmed. The use of non-lethal methods would not have adverse impacts on mammal populations in the State under any of the alternatives.

The continued use of non-lethal methods often leads to the habituation of mammals to those methods, which can decrease the effectiveness of those methods. For any management methods employed, the proper timing would be essential in effectively dispersing those mammals causing damage. Employing methods soon after damage begins or soon after threats were identified would increase the likelihood that those damage management activities would achieve success in addressing damage. Therefore, the coordination and timing of methods would be necessary to be effective in achieving expedient resolution of mammal damage.

In addition to non-lethal methods that would be used to disperse, exclude, or harass wildlife, another non-lethal method available under the alternatives would be the reproductive inhibitor commonly known as Gonacon™. The reproductive inhibitor Gonacon™ is currently not registered for use in Tennessee. However, it is discussed in this assessment to evaluate the potential use of the chemical if it becomes registered for use in the future. Gonacon™ has been classified as a restricted-use pesticide by the EPA. Restricted-use pesticides can only be purchased and/or applied by those persons who have successfully completed an applicators course to use restricted-use pesticides. The TDA administers training and testing required for applicators to purchase and apply restricted-use pesticides in the State. Gonacon™ could be employed by WS and the TWRA, if registered for use in the State, under Alternative 1. Only the TWRA or their designated agents could use Gonacon™ if Alternative 2 or Alternative 3 were selected.

In addition to non-lethal methods, lethal methods would also be available for use under all the alternatives by WS and/or by other entities. Lethal methods that would be available to address mammal damage include live-capture followed by euthanasia, shooting, body-gripping traps, fumigants, cable restraints, rodenticides, and the recommendation of hunting and/or trapping, where appropriate. All of those methods would be available for use by WS or for recommendation by WS under Alternative 1. Lethal methods would be employed by WS under Alternative 1 to resolve damage only after receiving a request for the use of those methods. Those same methods would also be available for WS to recommend and for other entities to use under Alternative 2. Under Alternative 3, those same lethal methods would continue to be available for use by other entities despite the lack of involvement by WS in damage management activities.

When live-captured target animals are to be lethally taken under Alternative 1, take would occur pursuant to WS Directive 2.505 and WS Directive 2.430. Under alternative 2, WS would recommend the use of methods to lethally take live-captured or restrained target animals in accordance with WS Directive 2.505. No assistance would be provided by WS under Alternative 3; however, many of those methods available to lethally take live-captured or restrained animals would continue to be available for use by other entities under Alternative 3.

The use of lethal methods by any entity could result in local population reductions in the area where damage or threats were occurring since mammals would be removed from the population. Lethal methods could be employed or recommended to remove mammals that have been identified as causing damage or posing a threat to human safety. Therefore, the use of lethal methods could result in local reductions of mammals in the area where damage or threats were occurring. The number of mammals removed from the population by WS using lethal methods under Alternative 1 would be dependent on the number of requests for assistance received, the number of mammals involved with the associated damage or threat, and the efficacy of methods employed. The number of mammals removed by other entities under Alternative 2 and Alternative 3 would be unknown but would likely be similar to the take that could occur under Alternative 1.

Chemical methods that could be available under the alternatives to manage damage associated with certain mammal species would include zinc phosphide, aluminum phosphide, gas cartridges, warfarin, brodifacoum, diphacinone. Those chemical methods are not restricted to use by WS only; therefore,

when registered for use in the State as a restricted-use pesticide, those products would be available for use by licensed pesticide applicators under any of the alternatives. Rodenticides containing warfarin, brodifacoum, and diphacinone are not currently registered for use in the State. However, products containing those active ingredients have been registered with the EPA and could be registered for use in the State in the future.

Most lethal methods are intended to reduce the number of mammals present at a location since a reduction in the number of mammals at a location could lead to a reduction in damage, which would be applicable whether using lethal or non-lethal methods. The intent of non-lethal methods would be to harass, exclude, or otherwise make an area unattractive to mammals, which disperses those mammals to other areas leading to a reduction in damage at the location where those mammals were dispersed. Similarly, the use of a reproductive inhibitor would be to reduce a local population of target mammals, which could reduce the damage occurring since fewer individuals in a localized population can lead to more tolerable damage levels. The intent of using lethal methods would be similar to the objective trying to be achieved when using non-lethal methods, which would be to reduce the number of mammals in the area where damage was occurring; thereby, reducing the damage occurring at that location.

Often of concern with the use of lethal methods is that mammals that were lethally taken would only be replaced by other mammals either during the application of those methods (*e.g.*, mammals that relocate into the area) or by mammals the following year (*e.g.*, increase in reproduction and survivability that could result from less competition). As stated previously, WS would not use lethal methods during direct operational assistance as population management tools over broad areas. Lethal methods would be employed under Alternative 1 to reduce the number of mammals present at a location where damage was occurring by targeting those mammals causing damage or posing threats. Since the intent of using lethal methods would be to manage those mammals causing damage and not to manage entire mammal populations, those methods would not be ineffective because mammals could be replaced by other mammals later.

Most lethal and non-lethal methods currently available provide only short-term benefits when addressing mammal damage. Those methods would be employed to reduce damage occurring at the time those methods were employed but do not necessarily ensure mammals would not return once those methods were discontinued or after the reproductive season (when young disperse and occupy vacant areas). Long-term solutions to resolving mammal damage can often be difficult to implement and can be costly. In some cases, long-term solutions involve exclusionary devices, such as fencing, or other practices such as structural repairs. When addressing mammal damage, long-term solutions generally involve modifying existing habitat or making conditions to be less attractive to mammals. To ensure complete success, alternative sites in areas where damage was not likely to occur would often times be required to achieve complete success in reducing damage and to avoid moving the problem from one area to another. Modifying a site to be less attractive to mammals would likely result in the dispersal of those mammals to other areas where damage could occur or could result in multiple occurrences of damage situations.

WS may recommend under Alternative 1 and Alternative 2 that property owners or managers, that request assistance, allow mammals to be harvested during the regulated hunting and/or trapping season for those species in an attempt to reduce the number of mammals causing damage on their properties. Managing localized mammal populations by allowing hunting and/or trapping could lead to a decrease in the number of mammals causing damage. Establishing hunting and trapping seasons and the allowed take during those seasons is the responsibility of the TWRA. WS does not have the authority to establish hunting or trapping seasons or to set allowed harvest numbers during those seasons. However, the harvest of those mammals during hunting and/or trapping seasons in the State would be occurring in addition to any take that could occur by WS under the alternatives or recommended by WS. In addition, mammals could also be lethally removed by other entities to alleviate damage or threats of damage under

all the alternatives. The total number of individuals from each species that were lethally removed by other entities to alleviate damage or threats of damage is currently not available.

As discussed previously, the analysis to determine the magnitude of impact from lethal take can be determined either quantitatively or qualitatively. Quantitative determinations are based on population estimates, allowable harvest levels, and actual harvest data. Qualitative determinations are based on population trends and harvest trend data. The issue of the potential impacts of conducting the alternatives on the populations of those mammal species addressed in this assessment is analyzed for each alternative below.

Alternative 1 - Continue the Current Adaptive Integrated Approach to Managing Mammal Damage (Proposed Action/No Action)

Under the proposed action, WS would continue to provide both technical assistance and direct operational assistance to those persons requesting assistance with managing damage and threats associated with mammals in the State. WS could employ those methods described in Appendix B in an adaptive approach that would integrate methods to effectively reduce damage and threats associated with mammals in the State.

The 2006 National Land Cover Dataset (NLCD)[10] was created from satellite imagery that was digitally rendered to produce a land-cover database comprised of three elements: land cover, impervious surface, and canopy density. Those elements were then combined to create a digital map of the United States in which each pixel, covering 30 square meters each, is classified into one of 29 different land classes. By adding all of the individually quantified pixels together, the resulting number is the most accurate number currently available for quantifying land within the state in any given land class. The State of Tennessee includes 15 of the 29 possible classes and this information is presented in Table 4.1. These numbers serve as the basis for current population estimates as presented in each species' population information and effects analysis.

Table 4.1: Tennessee Land Cover Area by Class as calculated from 2006 NLCD with Definitions.

Land Cover Class	Area in Tennessee (km^2)	Area in Tennessee (mi^2)	Definition
Open Water	2,435.9	940.5	All areas of open water, generally with less than 25% cover of vegetation or soil.
Developed, Open Space	6,298.1	2,431.8	Includes areas with a mixture of some constructed materials, but mostly vegetation in the form of lawn grasses. Impervious surfaces account for less than 20% of total cover.
Developed, Low Intensity	2,368.6	914.5	Includes areas with a mixture of constructed materials and vegetation. Impervious surfaces account for 20% to 49% of total cover.
Developed, Medium Intensity	803.4	310.2	Includes areas with a mixture of constructed materials and vegetation. Impervious surfaces account for 50% to 79% of the total cover.

[10]Habitat availability was calculated using the 2006 NLCD created through a cooperative project conducted by the Multi-Resolution Land Characteristics (MRLC) Consortium. The MRLC Consortium is a partnership of federal agencies, consisting of the United States Geological Survey (USGS), the National Oceanic and Atmospheric Administration, the EPA, USDA-Natural Resources Conservation Service (NRCS), the United States Forest Service (USFS), the National Park Service (NPS), the USFWS, the Bureau of Land Management, the National Aeronautics and Space Administration, and the Office of Surface Mining.

Land Cover Class	Area in Tennessee (km^2)	Area in Tennessee (mi^2)	Definition
Developed, High Intensity	303.6	117.2	Includes highly developed areas where people reside or work in high numbers. Impervious surfaces account for 80% to 100% of the total cover.
Barren Land	211.8	81.8	Barren areas of bedrock, desert pavement, scarps, talus, slides, volcanic material, glacial debris, sand dunes, strip mines, gravel pits and other accumulations of earthen material. Generally, vegetation accounts for less than 15% of total cover.
Deciduous Forest	46,148.0	17,818.2	Areas dominated by trees generally greater than 5 meters tall, and greater than 20% of total vegetation cover. More than 75% of the tree species shed foliage simultaneously in response to seasonal change.
Evergreen Forest	4,792.5	1,850.4	Areas dominated by trees generally greater than 5 meters tall, and greater than 20% of total vegetation cover. More than 75% of the tree species maintain their leaves all year. Canopy is never without green foliage.
Mixed Forest	3,970.5	1,533.1	Areas dominated by trees generally greater than 5 meters tall, and greater than 20% of total vegetation cover. Neither deciduous nor evergreen species are greater than 75% of total tree cover.
Shrub/Scrub	2,785.4	1,075.5	Areas dominated by shrubs; less than 5 meters tall with shrub canopy typically greater than 20% of total vegetation. This class includes true shrubs, young trees in an early successional stage or trees stunted from environmental conditions.
Grassland/ Herbaceous	3,745.3	1,446.1	Areas dominated by grammanoid or herbaceous vegetation, generally greater than 80% of total vegetation. These areas are not subject to intensive management such as tilling, but can be utilized for grazing.
Pasture/Hay	19,816.9	7,651.5	Areas of grasses, legumes, or grass-legume mixtures planted for livestock grazing or the production of seed or hay crops, typically on a perennial cycle. Pasture/hay vegetation accounts for greater than 20% of total vegetation.
Cultivated Crops	12,261.8	4,734.4	Areas used for the production of annual crops, such as corn, soybeans, vegetables, tobacco, and cotton, and also perennial woody crops such as orchards and vineyards. Crop vegetation accounts for greater than 20% of total vegetation. This class also includes all land being actively tilled.
Woody Wetlands	3,082.6	1,190.2	Areas where forest or scrubland vegetation accounts for greater than 20% of vegetative cover and the soil or substrate is periodically saturated with or covered with water.
Emergent Herbaceous Wetlands	126.9	49.0	Areas where perennial herbaceous vegetation accounts for greater than 80% of vegetative cover and the soil or substrate is periodically saturated with or covered with water.
Total Area of All Land Cover Classes	**109,151.5**	**42,144.4**	

The analysis of potential impacts on each of the species populations includes an estimate of annual take by WS as compared to statewide population estimates of the species. The statewide population has been estimated using the most current reliable information possible. Frequently, there is no current reliable information available for a species and conservative estimates are calculated based upon habitat availability and species use of those habitats.

WS' take that could occur to alleviate damage or threats of damage under the proposed action would be monitored by comparing numbers of animals killed with overall populations or trends in populations to assure the magnitude of take was maintained below the level that would cause adverse effects to the viability of native species' populations. The potential impacts on the populations of target mammal species from the implementation of the proposed action are analyzed for each species below.

Beaver Population Information and Effects Analysis

The North American beaver is a semi-aquatic mammal occurring in rivers, streams, lakes, reservoirs, and wetlands across North America. Beaver are large, bulky rodents whose most prominent features include a large scaly, paddle-shaped tail and nearly orange colored incisors (Hill 1982). Most adults weigh from 15.8 to 38.3 kg (35 to 50 lbs) with some occasionally reaching more than 45 kg (100 lbs), and are the largest North American rodents (Miller and Yarrow 1994). They range throughout most of Canada and the United States, with the exception of portions of Florida and the desert southwest. Beaver are active throughout most of the year and are primarily nocturnal, but it is not uncommon to see them during daylight hours. Beaver living along a river or large stream generally make bank burrows with multiple underwater entrances. Those in quiet streams, lakes, and ponds usually build dams and a lodge (National Audubon Society 2000). Sign of beaver in an area include gnawing around the bases of trees and trees, which have fallen because of this gnawing. Tree parts are stripped of bark, which is a primary beaver food.

Beaver are unique in their ability to create and modify their habitat by building dams (Boyle and Owens 2007). Beaver have a wide range and are extremely abundant, being found widely distributed over much of North America, including most of the United States. Beaver were trapped extensively during the 19th and part of the 20th centuries, and as a result, disappeared from much of their range (Novak 1987). Now reestablished over most of the continent, and protected from overexploitation, the beaver population has exceeded the societal carrying capacity in some areas. Dams built and maintained by beaver may flood stands of commercial timber, highways, and croplands. However, the dams also help reduce erosion, and the ponds formed by dams may create a favorable habitat for many forms of life (Hill 1982).

Beaver occur mostly in family groups that are comprised of two adult parents with two to six offspring from the current or previous breeding season. The average family group has been documented as ranging from 3.0 to 9.2 individuals (Novak 1987). Beaver abundance has been reported in terms of families per kilometer of stream or per square kilometer of habitat. Novak (1987) summarized reported beaver family abundance as ranging from 0.31 to 1.5 families per kilometer of stream, which converts to 0.5 to 2.4 families per mile of stream. Densities in terms of families per square kilometer have been reported to range from 0.15 to 3.9 (Novak 1987), which is the same as 0.24 to 6.3 per square mile. Novak (1987) indicates that rates of beaver populations are density dependent, which means that rates of increase generally occur as a population is reduced and become less as a population increases toward its carrying capacity[11]. This natural function of most wildlife populations helps to mitigate population reductions. Logan et al. (1996) indicated that wildlife populations being held at a level below carrying capacity could

[11]Carrying capacity is the maximum number of animals that the environment can sustain and is determined by the availability of food, water, cover, and the tolerance of crowding by the species in question.

sustain a higher level of harvest because of the compensatory mechanisms that cause higher rates of increase in such populations.

Beaver have a relatively low biotic potential due to their small litter size and a long juvenile development period. Population matrix models showed that survival of kits (1st year juveniles) and yearlings (2nd year juveniles) is the most critical factor in population viability. Survival of those age classes is partly dependent on the ability of beaver to successfully disperse and re-colonize habitats. Beaver are strong dispersers, and populations can recover quickly from local reductions when dispersal corridors are maintained (Boyle and Owens 2007).

Coyotes, black bears, bobcats, fishers (*Mustela pennanti*), red fox, river otters, mink, and large raptors such as hawks and owls have been documented preying on beaver (Tesky 1993, Baker and Hill 2003, Jackson and Decker 2004). With the exception of coyote, bear, and bobcat predation, most predation likely occurs to kits, yearlings, and young adults. With little exception, those predator species do not appear to exert significant predation pressure on beaver populations (Baker and Hill 2003).

The current population of beaver in the State is unknown; however, beaver are present in all 95 Tennessee counties, and their population is considered stable to increasing across the State (G. Anderson, TWRA, pers. comm. 2011). Beaver population estimates are often derived from density estimates for beaver based on the number of beaver colonies per a linear unit of measure (*e.g.*, stream miles) or per unit of area (*e.g.*, habitat type) (Baker and Hill 2003). Beaver densities specific to Tennessee are currently unavailable. Beaver densities by habitat calculated from other studies in the United States and Canada have ranged from 0.4 beaver colonies per square mile to a high of 12 beaver colonies per square mile (Novak 1987). Density estimates in the United States and Canada based on stream miles have ranged from 0.5 beaver colonies per stream mile to two beaver colonies per stream mile (Novak 1987). To derive a population estimate the number of beaver per colony must also be known; however, the average number of beaver per colony in Tennessee is currently unknown. From other studies, the average size of beaver colonies has ranged from 3.2 beaver to 9.2 beaver per colony (Novak 1987). In the southeastern United States, the average number of beaver per colony in Alabama was estimated at 4.6 beaver (Wilkinson 1962) and the average beaver per colony in Georgia was estimated at 5.3 beaver (Parrish 1960). Wetland estimates in Tennessee range from 632,000 acres (Hefner et al. 1994) to 761,000 acres (see Table 4.1) including an estimated 60,394 miles of streams (TDEC 2010).

Using the lowest beaver colony density per linear measure derived from other studies of 0.5 beaver colonies per stream mile and using the assumption that all stream miles in Tennessee are suitable beaver habitat and occupied by beaver colonies, a statewide population of beaver using the lowest calculated number of beaver per colony of 4.6 beaver in the southeastern United States, a statewide population of beaver inhabiting rivers and streams could be estimated at nearly 139,000 beaver. Of the total miles of streams and rivers in the State, 53,872 miles are considered perennial streams where water is present throughout the year (American Rivers 2011). Using only those river miles with water present throughout the year, a beaver population in the State could be estimated at approximately 124,000 beaver using the lowest densities of colonies and the lowest number of beaver per colony. The actual statewide population of beaver is likely much larger than 124,000 beaver since the population estimate was only based on perennial stream miles using the lowest density information and did not include beaver that could inhabit other aquatic habitats or create their own habitats by impounding water in areas associated with water runoff or storage (*e.g.*, drainage ditches, irrigation canals, storm water storage facilities).

The authority for management of resident mammal species in Tennessee, including beaver, is the responsibility of the TWRA. Beaver are considered a furbearer in the State that can be harvested annually during hunting and trapping seasons (TWRA 2012*a*). The TWRA collects and compiles information on beaver population trends and take, and uses this information to manage beaver populations in the State.

The primary tool for the management of beaver populations in Tennessee is through adjusting the allowed lethal take during the hunting and trapping season in the State, which is determined and regulated by the TWRA. When beaver are causing damage or about to cause damage, beaver can be taken without a permit during anytime of the year. In addition, the hunting and trapping season for beaver in the State is open throughout the year with no limit on the number of beaver that can be harvested.

During the six-year period, FY 2006 through FY 2011, WS continued to respond to requests for assistance associated with beaver in which those persons requesting assistance reported or WS verified $8,200,795 in damages to resources. Beaver damaged natural resources such as timber and reclamation sites, agricultural crops, commercial timber, roadways, drainage and irrigation structures, and other property through flooding, damming, and feeding. As part of those requests for assistance, WS lethally removed 3,448 beaver to alleviate damage and threats of damage, which is an average of 575 beaver lethally removed per year. The highest level of lethal take by WS occurred during FY 2011 when 917 beaver were lethally removed to alleviate damage or threats of damage. Based on the number of requests received previously by WS and in anticipation of receiving additional requests for assistance, WS could lethally remove up to 1,800 beaver annually under all damage management activities.

As stated previously, beaver can also be harvested annually during continuously open hunting and trapping seasons with no limit on the number of beaver that can be harvested. In addition, beaver can be lethally removed at any time to alleviate damage or threats of damage. The number of beaver harvested in the State and lethally removed to alleviate damage is currently unknown. The TWRA allowing beaver to be lethally removed at any time throughout the year with no limit on the number of beaver that can be harvested or removed to alleviate damage provides an indication that population levels in the State are sufficient to sustain the level of harvest occurring and that overharvest is not likely to occur. An allowable harvest level for beaver has been estimated at 30% of the population (Novak 1987).

Based on a statewide population estimated at 124,000 beaver, the annual take by WS of up to 1,800 beaver would represent 1.5% of the population. As indicated previously, the actual statewide population of beaver is likely much larger than 124,000 beaver since the population estimate was only based on perennial stream miles using the lowest density information. Therefore, the proposed take of up to 1,800 beaver annually by WS is likely a much lower percentage of the actual statewide population.

Under the proposed action alternative, WS could also be requested to breach or remove beaver dams to alleviate or prevent flooding damage. In addition, WS could be requested to install devices to control the water flow through dams to alleviate flooding or install exclusion devices to prevent damming. WS would only utilize manual methods (*e.g.*, hands and hand tools) to breach dams. To remove dams, WS could also use manual methods but could employ explosives in some cases. When dams are breached or removed, the building material used to create the dam (*e.g.*, sticks, logs, and other vegetative matter) is discarded on the bank or is released to flow downstream. Mud and small materials, such as bark and other plant debris, also escape downstream and tend to settle out within a short distance of the dam. Small to medium limbs, along with sediments, may drift further distances downstream. Dam breaching and removal would generally be conducted in conjunction with the removal of beaver responsible for constructing the dam since beaver would likely repair and/or rebuild dams quickly if dams were breached or removed prior to the beaver being removed. Therefore, the removal or breaching of beaver dams would not adversely affect beaver populations in the State since those activities would be conducted in association with removing beaver from the site; therefore, the take would be included in the estimated annual take levels of beaver addressed previously.

Muskrat Population Information and Effects Analysis

Muskrat are fairly large rodents with dense, glossy fur, dark brown above, lighter on the sides, paler below, to nearly white on the throat. They have long scaly tails that are nearly naked and laterally flattened, tapering to a point but not paddle shaped as the beaver. The muskrat spends its life in aquatic habitats and is well adapted for swimming. Its large hind feet are partially webbed, stiff hairs align the toes, and its laterally flattened tail is almost as long as its body. The muskrat has a stocky appearance, with small eyes and very short, rounded ears. Its front feet, which are much smaller than its hind feet, are adapted primarily for digging and feeding (Miller 1994).

They build houses, or lodges of aquatic plants, especially cattails, up to 2.4 m (8 feet) in diameter and 1.5 m (5 feet) high. Those structures are usually built atop piles of roots, mud, or similar support in marshy areas, streams, lakes, or along water banks. They also burrow in stream or pond banks with entrances often above the water line. Other sign of the presence of muskrat includes feeding platforms built of cut vegetation in water or on ice, marked by discarded or uneaten grasses or reed cuttings, and floating blades of cattails, sedges, and similar vegetation near banks. This species is most active at dusk, dawn, and at night, but may be seen at any time of the day in all seasons, especially spring. Muskrat are excellent swimmers and spend much of their time in the water. They inhabit fresh, salt, and brackish waters throughout most of Canada and the United States; except for the Arctic regions (National Audubon Society 2000). They can be found in marshes, ponds, sloughs, lakes, ditches, streams, and rivers (Boutin and Birkenholz 1987).

Muskrat are highly prolific and produce three to four litters per year that average five to eight young per litter (Wade and Ramsey 1986), which makes them relatively immune to overharvest (Boutin and Birkenholz 1987). Gestation period varies between 25 and 30 days. Young muskrats can reproduce the spring after their birth. Harvest rates of three to eight animals per acre have been reported to be sustainable in muskrat populations (Boutin and Birkenholz 1987). Muskrat home ranges have been shown to vary from 529 square feet to 11,970 square feet (0.1 to 0.25 acres), with the size of home ranges occupied by muskrats dependent upon habitat quality and population density (Boutin and Birkenholz 1987).

Young muskrats are especially vulnerable to predation. Adult muskrats may also be subject to predation, but rarely in numbers that would lower populations. Predation cannot be relied upon to solve damage problems caused by muskrats (Miller 1994). Predators of muskrat include great horned and barred owls, red-tailed hawks, bald eagles, raccoons, mink, river otter, red fox, gray fox, coyotes, bobcat, Northern pike (*Esox lucius*), largemouth bass (*Micropterus salmoides*), snapping turtles (*Chelydra serpentine*), and bullfrogs (*Rana catesbeiana*). The young are also occasionally killed by adult muskrats (Miller 1994).

No population estimates are available in Tennessee for muskrats; however, muskrats can be found statewide in suitable habitat. As stated previously, wetland estimates in Tennessee range from 632,000 acres (Hefner et al. 1994) to 761,000 acres (see Table 4.1), including an estimated 60,394 miles of streams (TDEC 2010).

Since population estimates are not currently available, a population estimate will be derived based on the best available information for muskrats to provide an indication of the magnitude of take proposed by WS to alleviate damage and threats of damage. Using the lowest acreage of wetlands in Tennessee of 632,000 acres and using a single muskrat home range of 0.25 acres and assuming only one muskrat occupies a home range with no overlap of ranges, a statewide population could be estimated at 2.5 million muskrats. However, not all wetlands likely provide suitable habitat for muskrats. If only 25% of the wetland acreage in the State were suitable habitat for muskrats, the population would be estimated at 632,000 muskrats.

Muskrats are classified as regulated furbearers in Tennessee, and seasons and limits for take are set by the TWRA. Muskrats can be harvested during annual hunting and trappings seasons in the State with no limit on the number of muskrats that can be harvested. The number of muskrats harvested annually during the hunting and trapping season is currently unknown.

To alleviate damage at the request of a cooperator, WS lethally removed 340 muskrats in the State during the six-year period, FY 2006 through FY 2011, which is an average of 57 muskrats lethally removed annually. Those persons requesting assistance reported or WS verified damages associated with muskrats totaling $115,450. Damages occurred to earthen dams, golf courses, gardens, and turf. Based on the number of muskrats lethally taken during this period, and a reasonable anticipation of an increase in the number of requests for assistance, WS could lethally take up to 600 muskrats per year as part of an integrated damage management program. WS anticipates an increase in the need to address damage and threats associated with muskrats on federal, State, municipal and private property, landfills, along road and railways, and to protect T&E species from predation and habitat manipulation.

Using a population estimated at 632,000 muskrats, the lethal take of up to 600 muskrats annually would represent 0.1% of the statewide population. Although the number of muskrats harvested annually in the State during the hunting and trapping season is unknown, the cumulative take is not likely to reach a magnitude where adverse effects would occur to the muskrat population. The unlimited take allowed by the TWRA provides an indication that the statewide densities of muskrats are sufficient that overharvest is not likely to occur. In addition, most muskrats would probably be taken in habitats where little or no trapping by fur harvesters is done. Damage management activities associated with muskrats would target single animals or localized populations at sites where their presence was causing unacceptable damage to agriculture, human health and safety, natural resources, or property.

Nutria Population Information and Effects Analysis

The nutria is a large, dark colored, semi-aquatic rodent that is native to South America. It was introduced to the United States in the late 1930s (Whitaker, Jr. and Hamilton, Jr., 1998). The nutria is somewhat similar to the native muskrat in appearance. Nutria have small eyes and ears with a tail that is long, scaly, sparsely haired, and round (National Audubon Society 2000). Nutria weigh on average about 12 pounds (Whitaker, Jr. and Hamilton, Jr., 1998).

Nutria primarily inhabit brackish or freshwater marshes, but are also found in swamps, rivers, ponds, and lakes. They live in dense vegetation, in abandoned burrows, or in burrows they dig along stream banks or shorelines (Wade and Ramsey 1986). The burrowing activity of nutria can severely damage levees, dikes, earthen dams, and other structures. Nutria feed on terrestrial or aquatic green plants, but also feed on crops adjacent to their habitat. Nutria will consume approximately 25% of their own weight in food each day (Whitaker, Jr. and Hamilton, Jr. 1998).

Nutria females begin breeding in their first year. Breeding can occur at any time during the year. In the right conditions, nutria can produce up to 15 young per year (Whitaker, Jr. and Hamilton, Jr. 1998). In the wild, the life expectancy of nutria is approximately two years. Home ranges for nutria are estimated to be from 12 to 445 acres, and densities range up to 10 nutria per acre (Whitaker, Jr. and Hamilton, Jr. 1998).

Nutria are not considered a native wildlife species in Tennessee and are considered an unprotected species by the TWRA. The first reported sighting of nutria in Tennessee occurred in 1996, as populations began expanding from introduced populations in nearby States (Kennedy and Kennedy 1998, Carter and Leonard 2002). A small population has become established in the State and appears to be expanding

(Bounds 2000, Carter and Leonard 2002). The current population of nutria in the State is unknown, but appears to be sporadic based on reported sightings. The nutria management objective of the TWRA is to eradicate the species in Tennessee, if possible, or keep the population at the lowest level possible (G. Anderson, TWRA, pers. comm. 2011).

WS has received two requests for assistance associated with nutria during the six-year period, FY 2006 through FY 2011, both requests occurred during FY 2011. WS lethally removed four nutria in response to those requests. Nutria are known to have established extensive populations within the United States. In addition, nutria are considered an introduced species within the State that can cause extensive damage to native vegetation and often competes with native wildlife species for resources. Based on the presence of established populations within the United States and the expanding population occurring within the State, WS could receive additional requests for assistance to manage damage associated with nutria.

The number of nutria addressed by WS would be dependent on the number of requests received, the number of nutria associated with causing damage or the threat of damage, and the efficacy of methods employed to resolve the damage. WS anticipates that up to 50 nutria could be lethally taken annually to resolve requests for assistance. Activities would only be conducted when requested by a property owner or property manager. In addition, nutria could be lethally taken as non-targets during damage management activities conducted targeting other mammal species, primarily beaver damage management activities. During the six-year period, FY 2006 through FY 2011, three nutria were lethally taken as unintentional non-targets by WS.

Nutria can be lethally taken throughout the year in the State with no limit on the number that can be lethally removed. The number of nutria lethally removed by other entities in the State is unknown. As stated previously, the TWRA has expressed a desire to keep the nutria population at the lowest possible level, including the complete elimination of the species from the State, if possible. Executive Order 13112 states that each federal agency whose actions may affect the status of invasive species shall, to the extent practicable and permitted by law; 1) reduce invasion of exotic species and the associated damages, 2) monitor invasive species populations, provide for restoration of native species and habitats, 3) conduct research on invasive species and develop technologies to prevent introduction, and 4) provide for environmentally sound control and promote public education on invasive species. WS' activities would be conducted pursuant to Executive Order 13112.

Woodchuck Population Information and Effects Analysis

The woodchuck, also known as the "*groundhog*," is a large rodent, often seen in pastures, meadows, and fields in Tennessee. They dig large burrows, generally eight to 12 inches at the opening, sometimes five feet deep and 30 feet long with more than one entrance to a spacious grass-filled chamber. Green vegetation such as grass, clover, and alfalfa forms its diet; at times, the woodchuck will feed heavily on corn and can cause extensive damage in a garden to other crops (National Audubon Society 2000). Woodchucks may also jeopardize the integrity of earthen dams, present hazards to livestock and farm equipment because of burrowing, gnaw electrical cables, and damage hoses and other accessories on automobiles by gnawing (Bollengier, Jr. 1994, USDA 2007).

The breeding season for woodchucks is usually from March through April (Bollengier, Jr. 1994). Female woodchucks usually produce from four to six young (Chapman and Feldhamer 2003). The offspring breed at one year of age and live four to five years. Mammal species with high mortality rates, such as rodents (*e.g.*, woodchucks) and lagomorphs (*e.g.*, rabbits), typically possess high reproductive rates, and produce large and frequent litters of young (Smith 1996). For example, if a pair of woodchucks and their offspring all survived to breed as soon as possible, with an average litter size of four with a 1:1 sex ratio; they could produce over 645 woodchucks through their lifetime. Woodchucks range in the United States

extends throughout the East, northern Idaho, northeastern North Dakota, southeastern Nebraska, eastern Kansas, northeastern Oklahoma, and south to Virginia and Alabama.

Both sexes are similar in appearance, but the male is slightly larger, weighing an average of five to 10 pounds (2.2 to 4.5 kg). The total length of the head and body averages 16 to 20 inches (40 to 51 cm). The tail is usually four to seven inches (10 to 18 cm) long. Like other rodents, woodchucks have white or yellowish-white, chisel-like incisor teeth. Their eyes, ears, and nose are located toward the top of the head, which allows them to remain concealed in their burrows while they check for danger over the rim or edge. Although they are slow runners, woodchucks are alert and scurry quickly to their dens when they sense danger (Bollengier, Jr. 1994).

Woodchucks seldom stray far from their home dens. Chapman and Feldhamer (2003) estimated that distances of daily travel ranged from 100 m in colonies occupying good habitat to 400 m in somewhat lacking habitat which makes a home range of seven to 124 acres in size. Groundhog colonies have not been extensively studied to determine the social structure of a typical colony. However, in order for the species to survive, a colony would have to be comprised of at minimum two adults and the young of that year, totaling at least six to eight individuals.

Woodchucks are classified as a small game, furbearing species in Tennessee, with a year-round statewide open season and no limits on take. However, woodchuck populations in Tennessee are not monitored by the TWRA and population estimates are not available. As stated previously, woodchucks are typically associated with pastures, meadows and fields, open woodlands, and clearings (Chapman and Feldhamer 2003). The land cover categories most likely to encompass those habitats include pasture; shrub/scrub, and mixed forest, and cumulatively total approximately 26,573 km^2 in Tennessee (see Table 4.1). If only 50% of those land classes supported woodchucks, under a worst case scenario, with an estimate of a single woodchuck colony home range at 124 acres and assuming that only one woodchuck occupies a home range and no home ranges overlap, the statewide woodchuck population could be estimated to be approximately 26,000 woodchucks in Tennessee. This would be a worst-case scenario since the woodchuck population is likely to inhabit a much larger portion of these lands, woodchuck colonies likely consist of six to eight individuals, and some portion of most other land cover categories support woodchuck populations.

WS continues to receive requests for assistance to manage damage associated with woodchucks in the State. Requests for assistance received by WS from FY 2006 through FY 2011 were primarily associated with woodchuck damage occurring to gardens, vegetables, turf and flowers, electrical utilities, and drainage and irrigation structures. As part of those requests for assistance, WS employed lethal methods to lethally remove 141 woodchucks, which is an average of 24 woodchucks lethally removed by WS annually. Those persons requesting assistance reported to WS or WS verified $35,700 in damage associated with woodchucks in the State during this period.

In addition, WS treated 104 woodchuck burrows with gas cartridges during the same time, averaging approximately 17 burrow entrances treated per year. It is possible that WS could be requested to provide assistance to address woodchuck damage at any location in the State. Based on previous requests for assistance and in anticipation of receiving additional requests in the future, it is estimated that up to 500 woodchucks could be lethally removed to alleviate damage by WS. In addition, up to 500 burrow entrances could be treated using gas cartridges annually by WS.

Gas cartridges could be employed to fumigate woodchuck burrows in areas where damages were occurring. Gas cartridges act as a fumigant by producing carbon monoxide when ignited. The cartridges contain sodium nitrate which when burnt, produces carbon monoxide gas. The cartridges would be

placed inside active burrows at the entrance, the cartridge would be ignited, and the entrance to the burrow would be sealed with dirt, which allows the burrow to fill with carbon monoxide.

The number of entrances to burrow systems used by woodchucks varies. Twichell (1939) found the number of entrances to burrow systems used by woodchucks ranged from two to six entrances in Missouri with the average number being 2.8 entrances. Other studies note the number of entrances per burrow system ranged from one to five entrances (Grizzell, Jr. 1955) to a high of 11 entrances per system (Merriam 1971). Merriam (1971) found the mean number of entrances per burrow system was 2.98 entrances. The use of burrow systems is usually restricted to a male and a reproductive female (Swihart 1992, Armitage 2003). The number of woodchucks lethally removed when using gas cartridges to fumigate burrows would be based on the mean number of entrances per burrow system of approximately three entrances (Twichel 1939, Merriam 1971) and each burrow system occupied by a male and a female (Swihart 1992, Armitage 2003). The take of woodchucks would also occur using other methods, such as shooting, live traps, and body-gripping traps. However, the number of woodchucks lethally taken using gas cartridges and by other methods would not be expected to exceed 500 woodchucks.

Damage management activities associated with woodchucks would target single animals or local populations of the species at sites where their presence was causing unacceptable damage to agriculture, human health or safety, natural resources, or property. Some local populations may be temporarily reduced because of damage management activities conducted under the proposed action alternative aimed at reducing damage at a local site. If WS' annual take reached 500 woodchucks, the take would represent 1.9% of a statewide population estimated at 26,000 woodchucks, if the population remains at least stable. However, WS' annual take of woodchucks would likely represent a smaller percentage of the actual population given the population estimate derived represents a worst-case scenario. The unlimited take and continuous open season for woodchucks provides an indication that densities are sufficient that overharvest is unlikely to occur.

Fox and Gray Squirrel Population Information and Effects Analysis

Fox squirrels and gray squirrels are similar in behavior and appearance with a few exceptions. The fox squirrel is considerably larger than the gray squirrel and shows more color variation within its populations than do gray squirrels; however, both species exhibit melanistic and albino phases. Fox squirrels measure 48 to 73 cm (19 to 29 inches) long and weigh from 544 to 1,362 g (1 1/5 to 3 lbs). Gray squirrels measure 41 to 51 cm (16 to 20 inches) long and weigh from 567 to 794 g (1 ¼ to 1 ¾ lbs) (National Audubon Society 2000).

Both squirrel species are found throughout most of the eastern United States, including Tennessee. They inhabit mixed hardwood forests, especially those containing nut trees such as oak and hickory. While they are commonly referred to as tree squirrels, they spend quite a bit of time on the ground foraging. Squirrels feed on a wide variety of foods and adapt quickly to unusual food sources. Typically, they feed on wild tree fruits and nuts in fall and early winter. Acorns, hickory nuts, walnuts, and Osage orange fruits are favorite fall foods. Nuts are often cached for later use. In late winter and early spring, they prefer tree buds. In summer, they eat fruits, berries, and succulent plant materials. Fungi, corn, and cultivated fruits are taken when available. They may also chew bark during high population peaks, when food is scarce and may eat insects and other animal matter (Jackson 1994*a*).

Gray squirrels produce young during early spring but may actually produce at any time until early September (National Audubon Society 2000). Older adults may produce two litters per year (Burt and Grossenheider 1976, Jackson 1994*a*). The gestation period is 42 to 45 days, and about three young comprise a litter. Young begin to explore outside the nest at about 10 to 12 weeks of age (Jackson 1994*a*). Home ranges of squirrels range from 1.2 to over 40 acres in size (Flyger and Gates 1982) with

gray squirrels generally occupying home ranges up to seven acres and fox squirrels occupying areas from seven to 40 acres in size.

Both fox and gray squirrel populations periodically increase and decline. Gray squirrels have been documented to have mass emigrations of thousands or millions of individuals moving simultaneously and during which time many die. Although fox squirrels have been described as participating in these migrations, they are not as frequent or extensive in number. Squirrels are vulnerable to numerous parasites and diseases such as ticks, mange mites, fleas, and internal parasites. Squirrel hunters often notice bot fly larvae, called "*wolves*" or "*warbles*," protruding from the skin of animals killed. Larvae do not impair the quality of the meat for eating. In addition to being a food source for some people, squirrels are also prey for hawks, owls, snakes, and several mammalian predators. Predation seems to have little effect on squirrel populations. Typically, about half the squirrels in a population die each year and wild squirrels over four years old are rare, while captive individuals may live 10 years or more (Jackson 1994*a*).

Gray squirrel densities fluctuate based on available food sources but long-term densities tend to be stable (Gurnell 1987). Manski et al. (1981) found gray squirrel densities were typically less than 1.2 squirrels per acre in continuous areas of woodlands in North Carolina. Doebel and McGinnes (1974) found gray squirrel densities in small woodlots of less than 10 ha in area can be as high as 16 squirrels per ha. In urban parks, Manski et al. (1981) found gray squirrel densities can be more than 8.4 squirrels per acre. A three acre park in Washington, D.C. had a density of 50 squirrels per ha (20 per acre) (Hadidian et al. 1987).

Population estimates for the gray squirrel and the fox squirrel are currently not available. Therefore, the best available information will be used to estimate statewide populations. The rural land cover classifications most likely to encompass suitable squirrel habitats are deciduous, evergreen, and mixed forests, which cumulatively total approximately 55,000 km^2 in Tennessee (see Table 4.1). If only 50% of those land classes supported squirrels, under a worst case scenario, with a conservative estimate of 1 fox squirrel per every 40 acres and 1 gray squirrel per every 7 acres, and assuming that only one squirrel of each species occupies a home range and no home ranges overlap, the conservative statewide populations could be estimated to be approximately 170,000 fox squirrels and 970,770 gray squirrels in Tennessee. This would be a worst case scenario since both squirrel populations are likely to inhabit a much larger portion of the land classes in the State, squirrels typically occur at much higher densities, and no urban or suburban lands were included in these calculations where squirrel densities are likely to be high.

Fox and gray squirrels are considered small game species by the TWRA in Tennessee with a regulated hunting season. The daily bag limit on the number of squirrels that can be harvested during the season is 10, with no limit on the number of squirrels that can be possessed during the length of the season (TWRA 2012*a*). During the 2009 hunting season, an estimated 817,092 squirrels were harvested in the State (Applegate 2010). Harvest data provided by Applegate (2010) combines the three squirrel species (gray, fox, and red) that can be harvested during the squirrel season. There is currently no harvest total that only represents gray squirrels or fox squirrels.

WS has previously received requests for direct operational assistance associated with gray squirrels and fox squirrels. Requests for assistance were primarily associated with squirrels causing damage to gardens, vegetables, turf and flowers, electrical utilities, and residential buildings. Requestors reported $10,475 in damage caused by squirrels from FY 2006 through FY 2011. Based on the requests for assistance, WS employed lethal methods to remove 13 gray squirrels and fox squirrels to alleviate damage or threats of damage during this period. Squirrels could also be unintentionally lethally removed as non-targets during other damage management activities targeting other mammal species. Based on previous

requests for assistance and in anticipation of receiving an increasing number of requests for assistance, up to 100 gray squirrels and up to 100 fox squirrels could be lethally removed by WS annually.

If up to 100 fox squirrels were lethally remove by WS, the take would represent 0.06% of the fox squirrel population estimated at 170,000 squirrels under a worst-case scenario. If up to 100 gray squirrels were lethally removed by WS, the lethal take would represent 0.01% of the estimate gray squirrel population in the State under a worst-case scenario. If the proposed take were combined, the total take of 200 squirrels would represent 0.03% of the estimated number of squirrels harvested in the State during the 2009 hunting season.

Eastern Chipmunk Population Information and Effects Analysis

The eastern chipmunk is a ground-dwelling squirrel, typically 13 to 15 cm (5 to 6 inches) long and weighing 90 g (3 oz). Their tail is eight to 10 cm (3 to 4 inches) long and hairy, but it is not bushy (Williams and Corrigan 1994). Generally found in forested habitats, chipmunks have large, fur-lined internal cheek pouches for carrying nuts and seeds. They have black and white facial stripes, and five dark stripes separated by four pale ones, on the back and sides of their bodies. Chipmunks cache a great deal of food in the form of seeds, nuts, fruits, and sometimes, green vegetation and insects. They hibernate in the winter, but awaken about every two weeks to feed, since they do not store body fat before hibernation.

They are daytime animals, and are usually most active in early morning and late afternoon. They live mostly on the ground, but their nests may be either an underground burrow, or a hollow tree limb (National Audubon Society 2000). Burrows are often well hidden near objects, such as stumps, woodpiles, brush piles, basements, and garages or other buildings. The burrow entrance is usually about five cm (2 inches) in diameter with no obvious mounds of soil around them (Williams and Corrigan 1994, National Audubon Society 2000). Chipmunks can be found across most of the eastern United States, except the extreme south and along the southeastern seaboard (Williams and Corrigan 1994, National Audubon Society 2000).

Eastern chipmunks have two mating periods; those being during early spring and again during the summer or early fall. There is a 31-day gestation period, producing two to five young (Williams and Corrigan 1994). First year females not breeding in early spring may produce litters in late July or August (National Audubon Society 2000). The young are sexually mature within one year. Adults may live up to three years (Williams and Corrigan 1994).

Population densities of eastern chipmunks typically are five to 10 animals per hectare (2 to 4 per acre) (Burt and Grossenheider 1976), and may be as high as 24 chipmunks per hectare (10 per acre) if sufficient food and cover are available. Home ranges often overlap among individuals and are usually less than 92 m (100 yards) across (Williams and Corrigan 1994).

Throughout their North American range, chipmunks are considered minor agricultural pests. Most conflicts with chipmunks are nuisance problems. However, when chipmunks are present in large numbers they can cause structural damage by burrowing under patios, stairs, retention walls, or foundations. They may also consume flower bulbs, seeds, or seedlings as well as birdseed, grass seed, and pet food that is not stored in rodent-proof storage containers.

Chipmunks are not classified as a protected non-game species in Tennessee and can be lethally removed when causing damage. No statewide population estimates are currently available for chipmunks. Therefore, the best available information will be used to estimate statewide populations. Chipmunks are most likely to utilize deciduous forest as preferred habitat. There are approximately 46,000 km^2 of

deciduous forest in Tennessee (see Table 4.1). If only 50% of those lands supported chipmunks, under a worst case scenario, with a conservative estimate of two chipmunks per acre, and assuming that only one chipmunk occupies a home range and no home ranges overlap, the conservative statewide populations could be estimated to be approximately 5.7 million chipmunks. This would be a worst case scenario since chipmunk populations are likely to inhabit a much larger portion of those lands, they are typically at much higher densities, and no urban or suburban lands were included in the calculations where chipmunks are frequently found.

Based upon anticipated requests for WS' assistance, it is possible that WS could kill up to 100 chipmunks each year in the State under the proposed action alternative. Removing 100 chipmunks would represent 0.002% of the estimated statewide population annually and would be of low magnitude when compared to the actual statewide population of chipmunks. WS anticipates the annual take of up to 100 chipmunks to represent a much smaller percentage of the statewide population, since the population estimate was derived from a small portion of the land area occupied by chipmunks in the State.

Meadow, Pine, and Prairie Vole Population Information and Effects Analysis

Voles, also called meadow mice or field mice are compact rodents with stocky bodies, short legs, and short tails. Their eyes are small and their ears are partially hidden. Their underfur is generally dense and covered with thicker, longer guard hairs. They are usually brown or gray, though many color variations exist (O'Brien 1994, National Audubon Society 2000).

There are 23 species of voles in the United States (O'Brien 1994), and the genus *Microtus* to which meadow, pine, and prairie voles belong, contains 15 (National Audubon Society 2000) or 17 (Jones et al. 1992) species. Of those species, the meadow, pine, and prairie voles are relatively abundant, with meadow voles being the most widely distributed *Microtus* species in the United States. Meadow voles measure 14 to 19 cm (5 ½ to 7 ½ inches) and their fur is gray to yellow-brown, obscured by black-tipped hairs. Northern subspecies may also have some red in their fur. The underparts are gray, at times washed with silver or buff (O'Brien 1994). Pine voles measure 10.5 to 14.5 cm (4 1/8 to 5 ¾ inches) and have reddish brown short soft fur above with grayish, washed with buff coloration below (National Audubon Society 2000). The pine vole utilizes tunnel systems that are usually one to several inches below ground. Prairie voles measure 13 to 18 cm (5 to 7 inches), and have gray to dark brown fur mixed with gray, yellow or hazel-tipped hairs, giving a peppery appearance. Underparts are gray to yellow-gray. It is the most common vole in prairie habitats.

Meadow voles range from Alaska and Canada south and east to northern Washington, Idaho, Utah, New Mexico, Wyoming, Nebraska, northern Missouri, northern Illinois, Kentucky, Tennessee, northeastern Georgia, and South Carolina. Prairie voles range from southeastern Alberta, southern Saskatchewan, and southern Manitoba south to northwestern New Mexico, northern Oklahoma, northern Arkansas, Tennessee, and western West Virginia. There have been isolated populations documented in southeastern Texas and southwestern Louisiana. Both species are found together in some habitats, since meadow voles live chiefly in lush, grassy fields, marshes, swamps, woodland glades, and mountaintops, while prairie voles live principally in dry grass prairie or mixed grassy-weedy situations. As a rule when these two species occur together, meadow voles inhabit the moist areas, while prairie voles inhabit the drier (National Audubon Society 2000). Meadow and prairie voles are active day and night, constructing many tunnels and surface runways with numerous burrow entrances. A single burrow system may contain several adults and young (O'Brien 1994).

The pine vole is found in the eastern United States ranging from New England to central Iowa, north to central Wisconsin, and in the southern states except for most coastal areas. Habitat for the pine vole is

deciduous woodlands with thick leaf mold or thick herbaceous ground cover and sometimes park-like grassy areas (National Audubon Society 2000).

Voles feed on a variety of green vegetation, including grasses, forbs, and tubers, and in late summer they store seeds, tubers, bulbs, and rhizomes (O'Brien 1994, National Audubon Society 2000). They eat bark at times, primarily in fall and winter, and will eat crops, especially when their populations are high. Occasional food items include snails, insects, and animal remains. Voles may cause extensive damage to orchards, ornamentals, and tree plantings due to their girdling of seedlings and mature trees while feeding on bark. This usually occurs in fall and winter. Field crops such as alfalfa, clover, grain, potatoes, and sugar beets may be damaged by voles. Their activities in such crops may interfere with crop irrigation by displacing water and causing levees and checks to wash out. They can also ruin lawns, golf courses, and ground covers. Voles can also act as attractants for raptors and other predators at airports, where those species can pose aircraft strike hazards.

Voles may breed throughout the year, but most commonly in spring and summer (O'Brien 1994), and may do so all year long in the south (National Audubon Society 2000). In the field, they have one to five litters per year, and meadow voles have produced up to 17 litters per year in a laboratory (O'Brien 1994). Litter sizes range from one to 11, but usually average three to six. The gestation period is about 21 days and young are weaned by the time they are 21 days old. Females mature in 35 to 40 days and life spans of voles are short; probably two to 16 months. In one population studied, there was 88% mortality during the first month of life (O'Brien 1994).

Large population fluctuations are characteristic of voles and generally peak every two to five years, but cycles are not predictable. Population densities are variable. Cole and Batzli (1979) found that prairie vole populations averaged 38 per hectare (15 per acre) in prairie, 132 per hectare (52 per acre) in bluegrass, and 244 per hectare (99 per acre) in alfalfa. Among meadow voles, an Ontario, Canada population ranged from 80 to 400 per hectare (32 to 162 per acre) over one year, while an Illinois population ranged from five to 15 per hectare (two to six per acre), also over one year. Much higher densities may be reached during population irruptions. A wide variety of predators feed on voles. Voles are relatively easy for most predators to catch and are active, and therefore available, day and night year-round. Despite their vulnerability and availability, voles are not usually "*controlled*" by predators, because they have a high reproductive potential. Postpartum breeding is common and females may breed as early as one week of age. Synchronous breeding also occurs. These factors enable voles to increase at a faster rate than predators are able to reduce them (Pearson 1985).

WS responded to 14 requests (2.3 per year) for assistance with vole damage from FY 2006 through FY 2011. Most complaints were handled by providing technical assistance advice on methods for addressing damage. Complainants reported $1,050 in damages resulting from voles during this period. In many cases, damage costs were not recorded because complainants were attempting to prevent damages from occurring. Resources of concern included landscaping, turf and flowers, levees and irrigation ditches, as well as, damage to machinery from digging, gnawing, nesting, and girdling.

Determination of numbers of voles killed during damage management activities can be difficult when rodenticides, such as zinc phosphide treatments, are employed. This is because most animals killed by those methods die underground. Most activities to alleviate or prevent damage conducted by WS previously associated with voles have been addressed using zinc phosphide placed into vole burrow entrances. Those activities have been associated with damage to earthen dams as the integrity of those dams was threatened by the burrowing of a high density of voles. In addition, WS has also been requested to reduce vole densities at airports that could act as attractants for avian and mammalian predators that pose aircraft strike risks and to reduce damage to landscape plants, seedlings, trees, shrubs, and other ground cover.

Based upon the above information, WS' lethal take of voles would have no adverse impacts on overall populations of the species in Tennessee. Due to relatively high reproductive rates of vole species and because management activities would be restricted to specific local sites, WS' activities under the proposed action would have minimal impacts on overall populations of voles. The TWRA has determined that there is no evidence to suggest that human mediated mortality resulting from damage management, including removal by WS, would be detrimental to the survival of the vole populations in the State of Tennessee (G. Anderson TWRA, pers. comm., 2011).

Old World Rodent Population Information and Effects Analysis

Old World rodents are those rodents introduced to North America during colonization by Europeans. They are frequently referred to as "*commensal*" rodents due to their habits of "*living with or in close association with humans*" and are all part of the Muridae family. In Tennessee, this group of rodents includes the Norway rat, the roof rat, and the house mouse. All three of those species are non-native, invasive species.

The Norway rat is a stocky burrowing rodent that weighs an average of 454 g (1 lb). Their fur is coarse and usually brownish or reddish gray above and whitish gray on the belly. They are also called the brown rat, house rat, barn rat, sewer rat, gray rat, or wharf rat and are slightly larger than the roof rat. Blackish individuals occur in some locations. Norway rats make a network of interconnecting tunnels for nesting and are colonial. They may burrow to make nests under buildings and other structures, beneath concrete slabs, along stream banks, around ponds, in garbage dumps, and at other locations where suitable food, water, and shelter are present (Timm 1994). Norway rats live in close association with people (Burt and Grossenheider 1976, Timm 1994, National Audubon Society 2000), and in urban areas they live in and around residences, in cellars, warehouses, stores, slaughterhouses, docks and sewers. On farms, rats may inhabit granaries, barns, livestock buildings, silos, and kennels (Timm 1994). In summer, rats may inhabit cultivated fields (National Audubon Society 2000), developing extensive colonies of several hundred individuals at such sites (B. Dunlap, USDA, pers. comm. 2011).

Norway rats are found throughout the United States and southern Canada, and the Pacific Coast north to Alaska. They may be found in this range wherever humans live (Timm 1994, National Audubon Society 2000). They are primarily nocturnal, and usually become active about dusk, when they begin to forage for food and water. Some individuals may be active during daylight hours when rat populations are high. They have poor eyesight, relying more on hearing and the sense of smell, taste, and touch. They are considered color blind (Timm 1994).

Norway rats will eat nearly any type of food. When given a choice, they will select a nutritionally balanced diet, choosing fresh, wholesome items over stale or contaminated food. They prefer cereal grains, meats and fish, nuts, and some types of fruit. Food items in household garbage offer a fairly balanced diet and also satisfy their water requirements (Timm 1994). Rats often contaminate food they do not eat with droppings. They can also kill chickens and eat their eggs. They also eat wild plants, insects, and seeds (National Audubon Society 2000).

Norway rats breed year round (National Audubon Society 2000) but often have peaks in spring and fall with reproductive activity declining during the heat of summer and often stopping completely in winter, depending upon habitat. The average female rat has four to six litters per year with two to 22 young per litter (Timm 1994, National Audubon Society 2000). Twelve litters per year are possible (Burt and Gossenheider 1976, National Audubon Society 2000). Gestation is 21 to 26 days. Female rats may breed again within a day or two after a litter is born (Timm 1994).

The roof rat is similar in appearance to the Norway rat, but has a longer tail and a shorter nose (Whitaker, Jr. and Hamilton, Jr. 1998). While the roof rat is most abundant in coastal areas, in the eastern United States it can be found inland to east Arkansas, west Kentucky, north Alabama and Georgia, and in North Carolina and Virginia (National Audubon Society 2000). Within its range, the roof rat is commonly found inhabiting buildings. When found with Norway rats in the same building, roof rats will generally be found higher in the building, due to their ability to climb better than Norway rats. Roof rats generally nest and live in the walls of buildings (Whitaker, Jr. and Hamilton, Jr. 1998). Roof rats are capable of breeding when two to three months old. A female roof rat will typically have from four to six litters per year, and wean approximately 20 young (Whitaker, Jr. and Hamilton, Jr. 1998). The roof rat is omnivorous, but prefers to feed on grain and seeds if they are available. Considerable damage to stored grains in the form of consumption and contamination is done by roof rats (National Audubon Society 2000). Because of their ability to climb, they often can do damage to nuts and fruits while still on the tree (Whitaker, Jr. and Hamilton, Jr. 1998).

The house mouse is a small grayish-brown mouse with a gray or buff belly and a scaly tail that is uniformly colored. It ranges in length from 81 to 86 mm (3 ¼ to 3 ½ inches) and weighs 11 to 22 g (1/3 to ½ oz). Although it is occasionally found in fields, it is primarily associated with buildings. The house mouse will eat anything edible and they are found throughout all of North America, wherever there are concentrations of humans (National Audubon Society 2000). House mice are primarily nocturnal but can be found active during the day. House mice often build nests under floors and in the walls of buildings using fibrous material that may include paper, burlap, or fabric. They breed year-round with three to 11 young in every litter. They have a gestation period of 18 to 21 days and females can breed at six weeks of age (National Audubon Society 2000). House mice are capable of producing up to 13 litters each year and their populations can expand rapidly wherever food is abundant and shelter is adequate.

WS received three requests for assistance associated with Old World rodent damage in Tennessee from FY 2006 through FY 2011. Those persons requesting assistance from WS reported $9,750 in damages. Requests for assistance were associated with gnawing of structures and vehicle hoses and belts, as well as damage threats to aviation. WS addressed requests by providing technical assistance. No direct operational assistance was provided by WS from FY 2006 through FY 2011 and no lethal take of Norway rats, roof rats, or house mice occurred by WS.

Norway rats, roof rats, and house mice may be taken by WS during wildlife hazard management, assessment, and monitoring at airports and airbases. Although those species do not cause direct hazards to aviation safety, they serve as prey attractants to raptors and mammalian predators that may pose serious threats to aircraft safety. Removal of those species by WS would occur primarily at airports by methods that may include trapping and the use of registered rodenticides (see Appendix B for a description of the rodenticides). Typically, any lethal take would be associated with small mammal trapping surveys at airports/airbases or with operational prey base management activities to reduce hazards created by avian or mammalian predators in the aircraft operations area. Removal could also occur to alleviate agricultural damage at feedlots or other agricultural facilities.

The level of WS' involvement in those activities may vary considerably from year to year depending on the number of airports/airbases and agricultural facilities requesting assistance from WS. Determination of the number of rats and mice killed during damage management activities can be difficult when rodenticides are employed. This is because most rats and mice killed by those methods die underground or in structures. Although population estimates are not available, Norway rats, roof rats, and house mice are generally prolific breeders and are generally abundant throughout their range. Additionally, populations of those species fluctuate greatly over time. Due to the species' relatively high reproductive rates and because management activities would be restricted to specific local sites, WS' activities under

the proposed action would have minimal impacts on overall populations of Norway rats, roof rats, and house mice in the State. WS' activities would be conducted pursuant to Executive Order 13112.

New World Rodent Population Information and Effects Analysis

The New World rodents are those rodents that do not live in close association with humans and inhabited North America prior to colonial settlement by Europeans. New World rodents found in Tennessee that WS could receive requests for assistance with include the eastern harvest mouse, deer mouse, white-footed mouse, cotton mouse, rice rat, and the hispid cotton rat. Those native species are all from families other than the Muridae family from which all Old World rodents originate. Some of the New World rodents will occasionally nest in buildings, do cause some damage to reseeding programs, consume and contaminate agricultural crops on a limited basis and play a role in the transmission of zoonotic diseases. However, they typically live in areas away from human activity, do not live in large colonies, and do not cause the same level of damage that Old World rodents can. WS has previously addressed requests for assistance associated with those species through technical assistance. Technical assistance was provided through the identification of species causing damage and collection for research purposes in small mammal studies, primarily conducted in association with airports and airport safety.

The eastern harvest mouse is a small mouse that weighs nine to 14 g (1/3 to ½ oz), with a rich brown color above and a paler belly and underside of the tail (National Audubon Society 2000). It is found primarily in old fields, roadside ditches, honeysuckle thickets, and wet meadows with spotty dense vegetation (Gottschang 1965). Nests of harvest mice are made of finely shredded grasses and are globular in shape. They are usually built on the ground in thick clumps of grass (Kaye 1961). The eastern harvest mouse can breed at three months of age and will have multiple litters of two to five young each year (National Audubon Society 2000). Population densities vary greatly and range from 8.75 to 44.4 per hectare (3 to 17 per acre) (Stalling 1997).

The white-footed mouse is a medium-sized mouse that weighs 15 to 30 g (1/2 to 1 oz), with rich reddish brown on upperparts and white belly and feet. They are found primarily in warm, dry forests and brush lands, and may be found in brushy areas adjacent to agricultural crops (National Audubon Society 2000). They will nest nearly anywhere including, abandoned bird or squirrel nests, old logs, underground, in stumps, and in buildings. White-footed mice have two to four litters per year with two to six young in each. Young females can breed at 10 weeks of age. Population densities vary from ten to 30 per hectare (4 to 12 per acre) (National Audubon Society 2000).

The deer mouse is sometimes difficult to distinguish from the white-footed mouse. It is similar in size at 18 to 35 g (2/3 to 1 ¼ oz) and is extremely variable in color ranging from pale grayish buff to deep reddish brown, but will always have a distinct difference between dark upper and light undersides. They occupy nearly every dry land habitat within their range (National Audubon Society 2000). Nesting occurs in burrows that occur in trees, stumps, buildings, and underground. Deer mice have two to four litters per year with three to six young. Females can breed at five to six weeks of age and population densities vary from 25 to 37 per hectare (10 to 15 per acre) (National Audubon Society 2000).

The cotton mouse is a large mouse that weighs 28 to 51 g (1 to 1 ¾ oz), with a dark brown upper body and lighter underside. They are found in swamplands and forested areas along streams and agricultural borders (National Audubon Society 2000). Nesting occurs in trees, buildings, and under logs. They regularly climb trees. Cotton mice have more than four litters per year with three to four young per litter. Population densities range from two to four per hectare (0.8 to 1.6 per acre) (Wolfe and Linzey 1977).

The rice rat is a medium sized rat that measures 226 to 305 mm and weighs 40 to 80 g (1.4 to 2.8 oz). The body is grayish-brown, the belly is gray or tawny, and the tail is long, scaly, and paler above than

below. Rice rats are primarily nocturnal and semi-aquatic (National Audubon Society 2000). A fair portion of their diets includes freshwater mussels, baby turtles, fiddler crabs, snails, and insects, as well as seeds and succulent plants (Wolfe 1982). They inhabit marshy areas primarily of grasses and sedges. Nesting occurs on high ground at the base of debris. Rice rats have from one to seven young several times per year and females can breed at seven weeks of age (National Audubon Society 2000).

The hispid cotton rat is a medium sized rat that weighs 113 to 198 g (4 to 7 oz) and measures 80 to 320 mm. The body is covered in long coarse hair mixed with buff and black above, whitish below (National Audubon Society 2000). This rat prefers dense, grassy areas, which may include salt marshes, grasslands, brushy pastures, canal banks, roadsides, agricultural crop edges with areas of dense broomsedge, and honeysuckle. Hispid cotton rats usually have five to six young per litter and as many as nine litters per year. The young become sexually mature at 40 days old. Overall populations of cotton rats fluctuate from year to year but are found regularly at 25 to 30 per hectare (10 to 12 per acre) (National Audubon Society 2000).

Of all of the New World rodents, the hispid cotton rat is the most likely species to cause damage to crops including cotton, rice, alfalfa, grains, vegetables, fruits, squash, sugarcane, corn, sweet potatoes, and melons. Large populations can cause damage to canal banks as well (Espinoza and Rowe 1979). The hispid cotton rat has been extending its range northward over the last 50 to 100 years. They were captured in 2008 in central Virginia – the first record of this species in the Ridge and Valley province (Francl and Meikle 2009).

From FY 2006 through FY 2011, WS provided technical assistance to address New World rodent damage, in particular deer mice, at three sites. In each case, deer mice populations were of concern on airports with the potential for those rodents to attract mammalian and avian predators to active runways, threatening both the aircraft and its passengers. WS' personnel in Tennessee killed 26 deer mice during FY 2011 as part of a small mammal study and an airport Wildlife Hazard Assessment to determine rodent densities.

Although those species do not cause direct hazards to aviation safety, they serve as prey attractants to raptors and mammalian predators that may pose serious threats to aircraft safety. Typically, any lethal take by WS would be associated with small mammal trapping surveys at airports/airbases or with operational prey base management activities to reduce hazards created by avian or mammalian predators in the aircraft operations area. Removal could also occur to alleviate agricultural damage at agricultural facilities. The level of WS' involvement in those activities may vary considerably from year to year depending on the number of requests for assistance received. Based upon anticipated requests for WS' assistance, it is possible that WS could kill up to 100 individuals each year of any of the New World rodents under the proposed action alternative.

Although population estimates are not available, New World rodents are generally prolific breeders and are generally abundant throughout their range. Due to the species' relatively high reproductive rates, the limited take that could occur, and because management activities would be restricted to specific local sites, WS' activities under the proposed action would have minimal impacts on overall populations of those species in the State.

Cottontail Rabbit Population Information and Effects Analysis

There are nine species of cottontail rabbits in North America, north of Mexico. The eastern cottontail is the most abundant and widespread of all these. Although, in Tennessee, there are swamp rabbits (*Sylvilagus aquaticus*), Allegheny cottontails (*Sylvilagus obscurus*), and eastern cottontails, damage associated with rabbits is almost, without exception, caused by the eastern cottontail (Craven 1994).

The eastern cottontail is approximately 37 to 48 cm (15 to 19 inches) in length and weighs 0.9 to 1.8 kg (2 to 4 lbs). Males and females are nearly the same size and color. These animals do not distribute themselves evenly across the landscape, but tend to concentrate in favorable habitats such as brushy fence rows or field edges, gullies filled with debris, brush piles, areas of dense briars invaded with Japanese honeysuckle, or landscaped backyards where food and cover are suitable. They are rarely found in dense forest or open grasslands, but fallow crop fields, such as those in the Conservation Reserve Program may provide suitable habitat. Within these habitats, they spend their entire lives in an area of 10 acres or less. Occasionally they may move a mile or so from summer range to winter cover or to a new food supply. In suburban areas, rabbits are numerous and mobile enough to fill any "*empty*" habitat created when other rabbits are removed. Population densities vary with habitat quality, but one rabbit per 0.4 hectares (1 acre) is a reasonable average (Craven 1994). Rabbits live only 12 to 15 months, yet make the most of time available reproductively. They can raise as many as six litters per year of one to nine young (usually four to six), having a gestation period of 28 to 32 days. If no young were lost, a single pair together with their offspring could produce 350,000 rabbits in five years (National Audubon Society 2000).

From FY 2006 through 2011, WS provided technical assistance in Tennessee to address eastern cottontail rabbit damage at only a few sites. A total of $300 damage to turf and flowers was reported during this time. To address aircraft strike risks associated with avian and mammalian predators caused by high densities of rabbits at airports, WS lethally removed 60 cottontail rabbits from FY 2006 through FY 2011. Those rabbits were removed as part of an integrated management plan at airports to protect human health and safety and aircraft.

Cottontails are considered a small game species by the TWRA in Tennessee, with a regulated hunting season with a daily bag limit of five animals and no limit on the number that can be possessed during the length of the season. Population estimates and annual cottontail harvest information for Tennessee is currently unavailable. Therefore, the best available information will be used to estimate a statewide population. The rural land cover classifications most likely to encompass suitable cottontail rabbit habitats are developed, open space, mixed forest, shrub/scrub, hay/pasture, and cultivated crops, which cumulatively total approximately 45,000 km^2 in Tennessee (see Table 4.1). If only 50% of those lands supported cottontails, under a worst case scenario, with a conservative estimate of one rabbit per acre, and assuming that only one rabbit occupies a home range and no home ranges overlap, the conservative statewide populations could be estimated at over 11 million rabbits. This would be a worst case scenario since rabbit populations are likely to inhabit a much larger portion of those lands, rabbits are typically at much higher densities, and no urban or suburban lands were included in the calculations where rabbit populations are likely to be fairly high.

Based on previous requests for assistance and in anticipation of receiving additional requests for assistance in the State, WS could lethally take up to 500 rabbits annually to alleviate damage or threats of damage. If the population of cottontail rabbits in the State remains at least stable, WS' take of up 500 cottontails annually would represent 0.01% of the minimum statewide population of 11 million rabbits. However, WS anticipates that take of up to 500 rabbits annually would represent a much smaller percentage of the actual statewide population since the population is likely much higher than 11 million rabbits. Studies show that even if hunters take as many as 40% of the rabbits available in autumn, the rabbit population the following year would not be adversely affected because of the tremendous reproductive potential of rabbits (Fergus 2006). Therefore, WS' proposed take would not adversely affect the ability to harvest rabbits during the annual regulated hunting season in the State.

Black Bear Population Information and Effects Analysis

The American black bear is the smallest and most widely distributed of the three North American bear species (Pelton 1982). This species is a compact, heavily structured mammal with relatively massive legs and feet. Adult male black bear weigh from 120 to 280 kg (265 to 617 lbs) and measure from 130 to 190 cm (51 to 61.7 inches) in length from the tip of the nose to the tip of the tail. Adult females weigh from 45 to 192 kg (100 to 400 lbs) and measure from 110 to 170 cm (45 to 67 inches) in total length. The normal color is black with a brownish muzzle and frequently, a white V-shape across the throat or chest (Kolenosky and Strathearn 1987). Black bears mate in June and July and sow (female) bear produce litters of one to five cubs, with two cubs usually the normal litter size. Although black bears are primarily nocturnal, they may be seen at any time. They occupy ranges of 20 to 25 km^2 (8 to 10 mi^2), and sometimes up to 40 km^2 (15 mi^2). The home range of the male black bear is about double that of the female.

Black bear are powerful swimmers and climb trees for protection or food. Although they are in the order Carnivora, their diet includes all types of vegetation including twigs, buds, leaves, nuts, roots, fruit, corn, berries, and newly sprouted plants. Black bear will rip open bee trees to feast on honey, honeycomb, bees, and larvae. They will also tear apart rotting logs for grubs, beetles, crickets, and ants. The black bear wades in streams or lakes, snagging fish with its jaws or pinning them with a paw. The black bear's diet also includes small to medium-size mammals or other vertebrates, and even livestock such as cattle, sheep, and goats. Bear are often a problem around open dumps, becoming dangerous as they lose their fear of humans. Occasionally, people have been killed by black bear (National Audubon Society 2000, Herrero et al. 2011). It has been suggested that habituated, food-conditioned bears pose the greatest threat to humans and such bears are usually found in association with campgrounds and sites where people regularly feed them (Herrero 1985, Kolenosky and Strathearn 1987). However, Herrero et al. (2011) found that most encounters with bears drawn into close contact with humans by food or garbage only result in threat displays and rarely lead to physical attacks.

In North America, black bear densities range from 0.1 to 1.3 bears per km^2 depending on region and habitat. Densities are greatest in highly diverse forests at a relatively early stage of development. In unhunted and lightly hunted populations, the annual survival rate of adult female black bear is about 80% to 90% with adult male survival slightly less. As hunting pressure increases, the number of males decreases more rapidly than that of females because of their greater vulnerability to hunting (Fraser et al. 1982). The statewide population of black bear for Tennessee is estimated to be between 750 to 1,500 bears and is believed to be increasing (Vaughan and Pelton 2003).

Black bear are classified as a big game species in Tennessee and seasons and harvest limits are set annually by the TWRA. From the 2006 through 2010 bear hunting seasons, between 300 and 570 bear were legally harvested in Tennessee each year (TWRA 2010). These numbers have been rising steadily since 1999.

No black bear have been killed by WS in Tennessee. WS' personnel have occasionally received requests for assistance, but WS has previously provided only technical assistance and referred people to the TWRA or the Great Smoky Mountains National Park for further action. Any direct damage management actions conducted by WS to address black bear damage in Tennessee under the proposed action alternative would be conducted as part of a coordinated effort with the TWRA to meet state wildlife resource management objectives. Such projects would usually involve live-capture and translocation of bear causing damage. Translocation would occur to locations determined by the TWRA with suitable habitat; therefore, the translocation of bears would not be expected to adversely affect bear populations in the State. Some bears could be killed in actions to protect human health and safety, or livestock. Any lethal control of bear would be coordinated by the TWRA.

In anticipation of receiving requests for assistance, up to five bears could be lethally taken annually by WS to alleviate damage, when requested by the TWRA. Based upon the above information, WS limited lethal take of black bear would have no adverse impacts on overall black bear populations in the State. The TWRA has determined that there is no evidence to suggest that human mediated mortality resulting from regulated hunting and damage management, including any removal by WS, would be detrimental to the survival of the black bear population in the State of Tennessee (G. Anderson TWRA, pers. comm., 2011).

Raccoon Population Information and Effects Analysis

The raccoon is a stocky mammal about 61 to 91 cm (2 to 3 feet) long, weighing 4.5 to 13.5 kg (10 to 30 lbs). It is distinctly marked, with a prominent black mask over the eyes and a heavily furred, ringed tail. The animal is a grizzled salt-and-pepper gray and black above, although some individuals are strongly washed with yellow (Boggess 1994*a*).

The raccoon is one of the most omnivorous of animals. It will eat carrion, garbage, birds, mammals, insects, crayfish, mussels, other invertebrates, and a wide variety of grains, various fruits, other plant materials, and most foods prepared for human or animal consumption (Sanderson 1987). They occasionally kill poultry (Boggess 1994*a*).

The raccoon is found throughout most of the United States, with the exception of the higher elevations of mountainous regions and some areas of the arid southwest (Boggess 1994*a*, National Audubon Society 2000). Raccoons are more common in the wooded eastern portions of the United States than in the more arid western plains (Boggess 1994*a*), and are frequently found in cities or suburbs as well as rural areas (National Audubon Society 2000). Movements and home ranges of raccoons vary according to sex, age, habitat, food sources, season, and other factors. In general, males have larger home ranges then females. Home range diameters of raccoons have been reported as being one to three km (0.6 to 2.9 mi) maximum, with some home range diameters of dense suburban populations to be 0.3 to 0.7 km (0.2 to 0.4 mi).

In Tennessee, raccoons cause damage to gardens, residential and non-residential buildings, fish, domestic fowl, and pets, as well as general property damage. Results of their feeding may be the total loss of ripened sweet corn in a garden. Damage to buildings generally occurs when they seek to gain entry or begin denning in those structures. Raccoons may den in uncapped chimneys, or may tear off shingles or fascia boards to gain access to attics or wall spaces. They may also damage or destroy sod by rolling it up in search of earthworms and other invertebrates (Boggess 1994*a*).

The public are also concerned about health and safety issues associated with raccoons. Those diseases include, but are not limited to, canine distemper and rabies, and the roundworm *Baylisascaris procyonis*, the eggs of which survive for extremely long periods in raccoon feces and soil contaminated by them. Ingestion of those eggs can result in serious or fatal infections in other animals as well as humans (Davidson and Nettles 2006) (see Table 1.3).

Absolute raccoon population densities are difficult or impossible to determine because of the difficulty in knowing what percentage of the population had been counted or estimated with the additional difficulty of knowing how large an area the raccoons were using (Sanderson 1987). Due to their adaptability, raccoon densities reach higher levels in urban areas than that of rural areas. Relative raccoon population densities have been variously inferred by take of animals per unit area. For instance, Twichell and Dill (1949) reported removing 100 raccoons from tree dens in a 41 hectares (101 acres) waterfowl refuge area, while Yeager and Rennels (1943) studied raccoons on 881 hectares (2,177 acres) in Illinois and reported trapping 35 to 40 raccoons in 1938-1939, 170 in 1939-1940, and 60 in 1940-1941. Slate (1980) estimated

one raccoon per 7.8 ha (19.3 acres) in New Jersey in predominantly agricultural land on the inner coastal plain. Raccoon densities of 100 per square mile (1 raccoon per 6.4 acres) have been attained around abundant food sources (Kern 2002). Riley et al. (1998) summarized rural raccoon densities based on published literature that ranged from two to 650 per square mile in rural habitats, with an average of 10 to 80 raccoons per square mile. Relative density studies conducted in eastern Tennessee report raccoon densities ranging from three to 26 per square kilometer (seven to 67 raccoons per square mile) (USDA unpublished data 2010).

WS continues to provide assistance in efforts to contain the spread of raccoon rabies in Tennessee. Those activities are part of the national rabies barrier program addressed under separate environmental analyses (USDA 2005*b*). Other rabies monitoring or control activities may occur as part of this program. Raccoons killed under the ORV program are addressed in a separate EA (USDA 2005*b*) but are included in this EA for cumulative impact analysis.

Raccoons are classified as small game furbearers in Tennessee with a regulated hunting and trapping season with unlimited take allowed during the length of those seasons (TWRA 2012*a*). Neither population estimates nor annual raccoon harvest information are available. Therefore, the best available information will be used to estimate statewide population. The rural land cover classifications most likely to encompass suitable raccoon habitats are deciduous forest, mixed forest, shrub/scrub, and woody wetlands which cumulatively total approximately 56,000 km^2 in Tennessee (see Table 4.1). If only 50% of those lands supported raccoons, under a worst-case scenario, with a conservative estimate of three raccoons per km^2, the conservative statewide population could be estimated at 84,000 raccoons. This would be a worst case scenario since raccoon populations are likely to inhabit a much larger portion of these lands, raccoons are frequently found at much higher densities, and no urban or suburban lands were included in these calculations where raccoon populations are likely to be at their highest.

During all damage management activities conducted by WS from FY 2006 through FY 2011, WS lethally remove 744 raccoons, which is an annual take of 124 raccoons. Potential impacts to the raccoon population and to non-targets from the ORV program were discussed in a separate EA (USDA 2005*b*). WS' activities conducted under the ORV program are primarily non-lethal and do not involve the lethal take of raccoons for monitoring purposes.

Based on previous requests for assistance received by WS and in anticipation of receiving additional requests for assistance with managing raccoon damage, up to 1,200 raccoons could be lethally removed by WS annually when requested under the proposed action. Using the lowest population estimate of 84,000 raccoons, the take of 1,200 raccoons under the proposed action would represent 1.4% of the estimated population. Activities conducted to prevent the further spread of raccoon rabies in the State generally do not result in the lethal take of raccoons. Raccoons are live-captured, sampled, and released on-site as part of the post-baiting protocols (USDA 2005*b*). However, if raccoons were visibly injured or exhibit signs of disease, those raccoons are often euthanized and processed for rabies testing. The number of raccoons lethally taken in the State during the post-baiting trapping varies, but is not likely to exceed 50 individuals annually. Therefore, the statewide cumulative take of raccoons by WS in Tennessee under all damage and disease management activities would not exceed 1,250 raccoons annually, which would represent 1.5% of the lowest population estimate of raccoons in the State.

Raccoon populations can remain relatively abundant if annual harvest levels are below 49% (Sanderson 1987). In addition, the statewide population is likely much higher than 84,000 raccoons. As with many of the other mammals species harvested for fur in the State, the unlimited harvest levels allowed by the TWRA provides an indication that overharvest of raccoons is not likely to occur during annual harvest seasons and from damage management activities. Although the statewide population of raccoons and the annual harvest levels are unknown, the cumulative take of raccoons would be of low magnitude when

compared to the actual statewide population. In addition, the live-capture and subsequent release of raccoons would not likely result in adverse effects to the statewide population since those animals would be released unharmed (USDA 2005*b*).

River Otter Population Information and Effects Analysis

Historically, river otter inhabited aquatic ecosystems throughout much of North America, excluding the frozen Arctic and arid Southwest (Hall and Kelson 1959). Information on historic numbers and distribution is limited. As its broad geographic distribution suggests, the river otter is able to adapt to diverse aquatic habitats. Otter are found in both marine and freshwater environments, ranging from coastal to high elevation mountainous habitat. Riparian vegetation adjacent to lakes, streams, and other wetland areas is a key component of otter habitat.

Human encroachment, habitat destruction, and overharvest have eliminated river otter from marginal portions of their range. However, present distribution spans the North American continent from east to west and extends from southern Florida to northern Alaska (Melquist and Dronkert 1987). In the 1950s, the heaviest river otter populations in Tennessee were reported along the Hatchie River. The river otter was virtually non-existent in the remainder of the state (TWRA 2001). In 1975, the river otter was placed on the Tennessee threatened list through the enactment of Proclamation 75-15. The river otter population increased in west Tennessee to the point where in 1989 the Wildlife Commission removed the river otter from the threatened list in all drainages west of Kentucky and Pickwick Lakes (TWRA 2001).

In the 1990s, the TWRA initiated a large-scale project to reintroduce river otters in middle and east Tennessee. Those efforts have been so successful that trapping seasons exist statewide now and river otter are currently a regulated furbearer with no take limit (TWRA 2012*a*). Information regarding the total number of river otter killed in Tennessee annually during the harvest season is not available.

The current statewide otter population is currently unknown. Densities of river otter in linear waterways have been reported ranging from one otter per 0.7 miles in southeast Alaska (Woolington 1984) to one otter per 10.6 miles (Reid 1984) in northeastern Alberta. Melquist and Dronkert (1987) summarized studies estimating river otter densities, which showed that densities were about 1 per 175 to 262 acres in Texas coastal marshes, and ranged from 1 per 1.8 miles to 1 per 3.6 miles of waterway (stream or river). The results of a Missouri study found 1 otter per 2.5 to 5.0 miles of linear waterways (Erickson et al. 1984).

Wetland estimates in Tennessee range from 632,000 acres (Hefner et al. 1994) to 761,000 acres (see Table 4.1), including an estimated 60,394 miles of streams (TDEC 2010). As was discussed previously, otter are closely associated with aquatic habitats where they forage and den along shorelines. Using 60,394 miles of streams in Tennessee and the range of 1 otter per 2.5 to 5.0 miles of waterway would result in a statewide population estimate ranging from 12,000 otter to 24,150 otter. If only 50% of those streams supported river otter, the minimum statewide river otter population could be estimated to range from 6,000 to 12,000 river otter in Tennessee. This would be considered a worst-case scenario since the otter population is likely to inhabit a much larger portion of the streams and wetlands of Tennessee.

WS responded to 12 requests for assistance associated with river otter damage from FY 2006 through FY 2011. Most complaints were handled by providing technical assistance advice on methods for addressing damage. Complainants reported $3,150 in damages resulting from river otter during the period. Loss values were not obtained for all reports. Resources affected include boat docks, watercraft, and fish.

From FY 2006 through FY 2011, WS killed 141 river otters (average = 23.5/year) in Tennessee. Of those otters taken by WS from FY 2006 through FY 2011, 140 otter were taken as unintentional non-targets

during aquatic rodent damage management activities. The highest unintentional take occurred during FY 2011 when 49 otters were unintentionally taken during other damage management activities. Non-target take of otters during aquatic rodent damage management activities are discussed here to evaluate cumulative take. Based on previous requests for assistance and anticipating future requests, WS reasonably expects the intentional take of otter would not exceed 25 otters annually in Tennessee to resolve requests to manage damage to resources. WS anticipates receiving requests primarily from aquaculture producers that are experiencing unacceptable predation of fish stock by river otters.

As was discussed previously, river otters are also likely to be lethally removed by WS as unintentional non-targets during other activities to alleviate wildlife damage. To evaluate the potential cumulative impacts on the river otter population from the activities proposed under this alternative, WS will evaluate cumulative take using the highest annual non-target take of otter that occurred during previous activities. As stated previously, the highest annual non-target take by WS occurred during FY 2011 when 49 otter were killed unintentionally. Based upon the aforementioned population estimate, WS' limited lethal take of 25 river otters annually under the proposed action would represent 0.4% of the otter population in Tennessee estimated at 6,000 otters and 0.2% of a statewide population estimated at 12,000 otters. When WS' average annual unintentional take is combined with the proposed take evaluated in this EA, the cumulative take by WS would represent 0.8% of a statewide population estimated at 6,000 otter and would represent 0.4% of a statewide population estimated at 12,000 otter. If the highest unintentional take by WS of 49 otters was combined with the proposed take under this proposed action of up to 25 otters, WS' cumulative take would represent 1.2% of a statewide population estimated at 6,000 otters and 0.6% of a statewide population estimated at 12,000 otters.

The proposed take and the cumulative take of otters in the State by WS would be of low magnitude when compared to the actual statewide population estimates. The unlimited take allowed by the TWRA also provides an indication that harvest and damage management activities are not sufficient to cause the overharvest of otters.

Long-tailed Weasel Population Information and Effects Analysis

The long-tailed weasel is the largest true weasel in North America and is about 30 to 40 cm (11 to 16 inches) in length, including the slender black-tipped tail. They weigh from 28 to 170 g (1 to 6 oz). Males are distinctly larger than females but both sexes are dark brown with white underparts and feet in summer and turn white in the winter in northern climates (National Audubon Society 2000).

They exhibit the typical mustelid form, which is a long, slender body with short legs. The tail is 44% to 70% of the length of the head and body (Sheffield and Thomas 1997). Northern long-tailed weasels have white winter coats and a brown summer coat with light-colored underparts from the chin to the inguinal region, with lateral margins tinged with buff or yellow. Long-tailed weasels have single annual litters averaging four to five with a maximum of nine after a 205- to 337-day gestation period due to delayed implantation of embryos. The long-tailed weasel has the widest distribution of any mustelid in the Western Hemisphere from Canada south to Venezuela, Ecuador, Peru, and Bolivia (Eisenberg 1989, King 1989, Emmons and Feer 1990). The long-tailed weasel inhabits all life zones, with the exception of desert throughout its range (Hall 1981). Long-tailed weasels are active in both winter and summer. In addition, peak activity occurs during the day when sunlight is at its height (Chapman and Feldhamer 2003).

Between 50% and 80% of the yearly food intake of weasels consists of small mammals, especially rodents. In particular, long-tailed weasels prefer voles, cottontail rabbits, mice, rats, shrews, squirrels and chipmunks (Hamilton, Jr. 1933). They will vary their diet based on season, availability, and individual preferences. Despite altered diets, long-tailed weasel population densities fluctuate considerably with

year-to-year changes in small mammal abundances (MacLean et al. 1974, Fitzgerald 1977). Long-tailed weasel population densities range from as low as 0.004 to as high as 0.38 per ha (Glover 1943, Quick 1951) and populations occasionally crash, requiring several years to recover (Osgood 1935). In favorable habitat, maximum densities of long-tailed weasel may reach six to seven animals per km^2 (Glover 1942, Quick 1944, Jackson 1961, King 1975). In general, the long-tailed weasel has a home range of 12 to 16 ha (29 to 40 acres) and males have larger home ranges than females during the summer (Svendsen 1990).

Population estimates for long-tailed weasels in Tennessee are currently not available. Given that these weasels inhabit all cover types throughout Tennessee, except open water, barren land, and developed spaces the total cumulative area of the remaining classifications in Tennessee is about 96,730 km^2 (9,673,00 hectares) (see Table 4.1). If only 50% of the land area of the State has sufficient habitat to support long-tailed weasels and weasel densities are between 0.004 and 0.38 long-tailed weasels per ha (Glover 1943, Quick 1951), a statewide long-tailed weasel population could be estimated at between 19,300 and 1.8 million long-tailed weasels. The population of long-tailed weasels within the State is likely higher than 19,300 long-tailed weasels since weasels can be found statewide and across multiple habitat types.

No requests for assistance associated with long-tailed weasels were received by WS in Tennessee from FY 2006 to FY 2011. Based on an anticipation of receiving requests for assistance, WS may be requested to address damage being caused by long-tailed weasel in Tennessee to protect resource being damaged or threatened. Based upon receiving for requests for WS' assistance, as many as 100 long-tailed weasels could be killed each year by WS to address such damage.

Long-tailed weasel densities are sufficient to allow for an annual harvest season within the State. Long-tailed weasels are classified as small game furbearers in Tennessee with a regulated hunting and trapping season with no limit on take (TWRA 2012*a*). The number of long-tailed weasels harvested annually in the State is currently unknown.

If the statewide population of long-tailed weasel were estimated at 19,300 individuals, the cumulative take of 100 long-tailed weasels would represent 0.5% of the estimated population. If the statewide population of long-tailed weasel were estimated at 1.8 million individuals, the cumulative take of 100 long-tailed weasels would represent 0.01% of the estimated population.

Damage management activities associated with long-tailed weasels would target single animals or local populations of the species at sites where their presence was causing unacceptable damage to agriculture, human health or safety, natural resources, or property. Some local populations may be temporarily reduced because of WS' activities under this alternative aimed at reducing damage at a local site. Like other furbearing species in the State, the unlimited take allowed by the TWRA during the harvest season provides an indication that population densities of weasels in the State are sufficient that over harvest is not likely to occur.

Mink Population Information and Effects Analysis

The mink is a member of the weasel family and is about 46 to 61 cm (18 to 24 inches) in length, including the somewhat bushy tail. These animals weigh about 0.7 to 1.4 kg (1.5 to 3 lbs). Females are about three-fourths the size of males. Both sexes are a rich chocolate-brown color, usually with a white patch on the chest or chin, and scattered white patches on the belly. The fur is relatively short with the coat consisting of a soft, dense under-fur concealed by glossy, lustrous guard hairs. Mink also have anal musk glands common to the weasel family, and can discharge a disagreeable musk if frightened or disturbed (Boggess 1994*b*). They also mark their hunting territory with this fetid musk, which is as malodorous as a skunk's musk, although it does not carry as far (National Audubon Society 2000).

Mink are found throughout North America, with the exception of the desert southwest and tundra areas (Eagle and Whitman 1987). They are shoreline dwellers and their one basic habitat requirement is a suitable permanent water area. This may be a stream, river, pond, marsh, swamp, or lake. They make their dens in muskrat houses, bank burrows, holes, crevices, logjams, or abandoned beaver lodges. They are active mainly at night and are active year-round except for brief intervals during periods of low temperature or heavy snow (Boggess 1994*b*). However, they may adjust hunting times to prey availability (National Audubon Society 2000).

Eagle and Whitman (1987) indicated mink population densities varied spatially based on habitat, and could be influenced temporally by weather, trapping, and intraspecific aggression. Mink are most abundant in those areas with stable aquatic habitat. Mink densities in Louisiana were found to be highest in swamps, followed by marshes, and drained bottomlands (Linscombe et al. 1982). In Montana, 280 mink were found inhabiting a 33 km^2 (12.8 mi^2) area during the initial year of a two-year study, which resulted in a population density of one mink per 11.8 ha (29.2 acres) (Mitchell 1961). However, Mitchell (1961) found only 109 mink in the same area the following year, resulting in a density of one mink per 30.3 ha (74.7 acres). Using mink tracks in snow, Marshall (1936), found 0.6 females in one km^2 (1.5/mi^2) of riverbank with a 1:1 sex ratio following heavy trapping in Michigan. During a study conducted in Iowa from 1933 to 1938, one to five mink families were found inhabiting a 180-ha (450 acres) marsh (Errington 1943). In 1939, Errington (1943) found no families present in the same marsh. Over-trapping was suggested as the reason for the decline in the number of mink families found in the marsh (Errington 1943). Intraspecific aggression between mink may have been the limiting factor for the upper limit of mink present at the marsh (Errington 1943).

At a refuge in Wisconsin, McCabe (1949) estimated 24 mink inhabited 446 ha (1,100 acres) in 1944, which resulted in a density of one mink per 18.8 ha (46.3 acres). Over the next four years (1945 to 1948), McCabe (1949) found the number of mink ranged from seven to 10 individuals at the refuge. McCabe (1949) also suggested that the lower population estimates found after the initial year of the study in 1944 were due to higher levels of mink trapping and excessive poaching along the refuge borders. The number of mink observed during the study conducted by McCabe (1949) at the refuge was inversely related to the duration and depth of snow cover; however, the number observed was poorly related to food availability (rabbits [*Sylvilagus* spp.] and mice [*Peromyscus* spp.]). During a two-year study in Sweden, Gerell (1971) estimated the number of mink present in a 10,000-ha (25,000 acres) area at 11 and 16, respectively, which resulted in a density of one mink per 909 ha (2,245 acres) during the first year of the study and one mink per 625 ha (1,545 acres) in the second year. Along 1.9 km (1.2 miles) of stream in British Columbia, Ritcey and Edwards (1956) caught 11, six, and five mink over three years, respectively, which were similar densities of 1.5 to 3 mink per km (2.5 to 5 mink per mile) found along the coastal shoreline on Vancouver Island reported by Hatler (1976).

No population estimates or density estimates were available for mink in Tennessee. Therefore, the best available information was used to estimate a statewide population. There are approximately 60,394 miles of streams in Tennessee (TDEC 2010), with 632,000 acres (Hefner et al. 1994) to 761,000 acres of wetlands (see Table 4.1). If only 50% of the 632,000 acres of wetlands (lowest estimate) present in the State supported mink and if the population density of mink in the State was one mink per 74.7 acres, the number of mink inhabiting wetlands in the State would 4,200 mink. If only 50% of the 60,394 miles of streams in the State supported mink and if the population density of mink were five mink per 1.2 miles of stream, the population inhabiting shoreline could be estimated at 126,000 mink. Combining the number of mink inhabiting wetlands and streams, the total statewide mink population could be estimated at 130,000 mink.

Mink are classified as small game furbearers in Tennessee with a regulated hunting and trapping season (TWRA 2012*a*). During the open hunting and trapping season, there is no limit on the number of mink that can be harvested (TWRA 2012*a*). The number of mink harvested annually in the State during the hunting and trapping seasons is currently not available.

WS did not receive requests for assistance associated with mink from FY 2006 to FY 2011; however, WS anticipates receiving requests for assistance to alleviate damage. Based upon an anticipated increase for requests for WS' assistance, up to 100 mink each year could be killed by WS to address damage or threats of damage. Take of up to 100 mink would represent 0.1% of the estimated statewide population of 130,000 mink in Tennessee.

Activities conducted under the proposed action alternative would target individual mink or local populations of mink at sites where they were causing damage to agriculture, human health or safety, natural resources, or property. Some local populations may be temporarily reduced because of WS' activities aimed at reducing damage at a local site. The unlimited harvest levels allowed by the TWRA during the hunting and trapping seasons provides an indication that mink densities within the State are sufficient that overharvest from the hunting/trapping season and activities to alleviate damage would not likely occur.

Striped Skunk Population Information and Effects Analysis

Although easily recognized by their black and white fur, striped skunks may be most readily recognized by the odiferous smell of their musk. They are common throughout the United States and Canada (Rosatte 1987). Striped skunk are primarily nocturnal and do not have a true hibernation period; although, during extremely cold weather they may become temporarily dormant. The striped skunk is an omnivore, feeding heavily on insects such as grasshoppers, crickets, beetles, bees, and wasps (Chapman and Feldhamer 2003). The striped skunk's diet also includes small mammals, the eggs of ground-nesting birds, and amphibians. Striped skunks are typically not aggressive and attempt to flee when approached by humans (Rosatte 1987). However, when provoked, skunks will give a warning and assume a defensive posture prior to discharging their foul-smelling musk. This musk is comprised of sulfur-alcohol compounds known as butylmercaptan (Chapman and Feldhamer 2003).

Adult skunks begin breeding in late February. Yearling females (born in the preceding year) mate in late March. Gestation usually lasts about seven to 10 weeks. Litters commonly consist of five to nine young with two litters per year possible (Hall and Kelson 1959). The home range of striped skunks is usually not consistent. It appears to be in relation to life history requirements such as winter denning, feeding activities, dispersal, and parturition (Rosatte 1987). According to Chamberlain and Leopold (2001), very little information regarding striped skunk densities in the southeast exists other than those densities based on harvest numbers and trapper/hunter observations. During the breeding season, males may travel larger areas in search of females. Skunk densities vary widely according to season, food sources, and geographic area. Densities have been reported to range from one skunk per 77 acres to one skunk per 10 acres (Rosatte 1987).

Population estimates for striped skunks in Tennessee are currently not available. Striped skunks can be found in a variety of habitats across the State. Given that striped skunks inhabit all rural, upland cover types throughout Tennessee, the cumulative area of these classifications in Tennessee is about 96,730 km^2 (23.9 million acres) (see Table 4.1). If skunks only inhabit 50% of those land classifications in the State and densities occur at one skunk per 77 acres, the statewide population could be estimated at nearly 155,200 skunks. Similar to other furbearing species, skunks can be found throughout the State and the estimate is intended to evaluate the magnitude of take proposed under the proposed action. The statewide population of skunks is likely higher than 155,200 skunks.

Striped skunks can be lethally taken using hunting methods during a continuous open season on private property in the State with no limit on the number that can be taken. In addition, skunks can be trapped during an annual season that places no limit on the number of skunks that can be harvested daily and no limit on the number of skunks that can be possessed throughout the trapping season. Like other mammal species addressed, the number of skunks harvested during the annual hunting and trapping season is currently not available. In addition, the number of skunks lethally removed annually in the State to alleviate damage is currently unknown.

From FY 2006 through FY 2011, WS responded to 143 requests for assistance (average 23.8/year) associated with striped skunks. Most requests were addressed by providing technical assistance on methods the requester could employ to alleviate damage or threats without any direct involvement by WS. Complainants reported $14,075 in damages caused by skunks from FY 2006 through FY 2011. Requests were received regarding damage or threats of damage to property, residential buildings, non-residential buildings, pets, turf, flowers, and human safety. Damage and threats occurred primarily from the burrowing/digging behavior of skunks, the odor associated with skunks spraying, and rabies threats. Most requests for assistance received were associated with threats to human safety associated with skunks, primarily risks of disease transmission.

From FY 2006 through FY 2011, WS lethally removed 348 striped skunks during all damage management activities in the State. Of those 348 skunks lethally removed, one skunk was lethally removed unintentionally during other damage management activities. In addition, six skunks were captured unintentionally but released unharmed.

Based on previous requests for assistance received by WS and in anticipation of receiving additional requests for assistance with managing striped skunk damage in Tennessee, up to 500 skunks could be lethally removed by WS annually under the proposed action, when requested. Using the lowest population estimate of 155,200 skunks, the take of 500 skunks would represent 0.3% of the estimated statewide population.

The unlimited take allowed by the TWRA with no closed season provides an indication that skunk densities in the State are sufficient to maintain a sustained harvest level and adverse effects from harvest and damage management purposes are not likely to cause overharvest of the species leading to population declines.

Spotted Skunk Population Information and Effects Analysis

Spotted skunk are one of the smallest skunks, (about ½ the size of a house cat). Their legs are short and their tail is long and bushy with a white tip. It has a black pelage with broken white stripes, and a white patch on the nose and front of the ears. Their average total length is 403 to 610 mm and tail length is 193 to 280 mm. Adult males weigh from 444 to 999 grams (1 to 2.5 lbs), and females weigh from 363 to 367 grams (0.8 to 1.25 lbs) (Whitaker, Jr. and Hamilton, Jr. 1998).

The eastern spotted skunk ranges from Costa Rica and northeastern Mexico through the Great Plains of the central United States to the Canadian border. It is also found throughout the southeastern United States. The eastern spotted skunk has been found in open lowlands, mountainous country, and at altitudes of 2,400 m (7,875 feet) (Baker and Baker 1975). Few studies have been published on the home range, population density, and mortality of spotted skunks. In Iowa, Crabb (1948) found that the western spotted skunk maintained densities of 2.2 skunks per km^2 (5.7 per mi^2). Crab (1948) also found skunks had a home range of 64.8 ha (160 acres) but noted movements of 4.8 km (3 mi) per night. Spotted skunks

appear to be somewhat nomadic without occupying a specific territory, and do not defend a home range (Crabb 1948).

Spotted skunk mate by April with a gestation period that has been reported to range from 50 to 70 days with an average litter size of two to six young. Young are blind when born and average 1/3 ounce each. Some males become sexually mature and breed at five months. The young develop teeth after approximately 40 days with weaning occurring after 55 days. Once weaned, the young forage with their mother until late fall when they disperse (National Audubon Society 2000).

The male provides no care to the young. The eyes of the young open at around one month and they can emit musk at about 46 days. This species is nocturnal and they climb trees more than other skunks. They are quicker and more alert, also. There is no true hibernation, just short inactive periods in the winter to conserve body fat. Several may den together in the winter. Populations up to 13 or more can be found per square mile. The males may wander farther, and dens distributed over the area seem to belong to the whole population, except during the breeding season. They have a characteristic handstand defense mechanism that makes the skunk appear larger.

Spotted skunk feed on a variety of items including rabbits, rodents, beetles, worms, crickets, grasshoppers, grubs, carrion, bird eggs, frogs, crayfish, lizards, and fruit (Rosatte 1987). They are more predaceous than other species of skunk, and mammals appear to be a more important food source than arthropods (Howard and Marsh 1982). For example, Crabb (1941) found mammals in 90% of scats collected during winter in Iowa. Eastern cottontails appeared to be a major food item, along with meadow and prairie voles.

Adult spotted skunks can stay in burrows for several weeks during cold spells losing up to 30% of their body weight with no ill effects. Underground dens used by spotted skunks either are excavated by the skunk or are dens abandoned by other animals. Dens have two to five entrances with one to three nest chambers. Dens can have up to 60 feet of tunnels. Sections below the frost line are used in winter when all but one entrance may be sealed. Deserted woodchuck and other small animal's burrows are frequently used as dens. Occasionally owls prey upon spotted skunk.

Spotted skunks are classified as small game furbearers in Tennessee with a regulated hunting and trapping season with no limits on take (TWRA 2012*a*). The number of spotted skunks harvested annually during the annual hunting and trapping seasons is currently not available. Similar to the other mammals classified as furbearers in the State, the statewide population is also not available. Given that spotted skunks inhabit most wooded cover types throughout Tennessee, the cumulative area of these classifications in Tennessee is about 55,000 km^2 (13.6 million acres) (see Table 4.1). If only 50% of those land classifications supported spotted skunks and if the statewide population density of spotted skunks was 2.2 skunks per km^2, assuming that one skunk occupies a home range, and no home ranges overlap, the minimum statewide striped skunk population could be estimated at 60,500 spotted skunks. Similar to striped skunks, spotted skunks can be found throughout the State and the estimate is intended to evaluate the magnitude of take proposed under the proposed action.

WS has not previously been requested to provide assistance with damage or threats of damage caused by spotted skunks in the State. No lethal take of spotted skunks by WS has occurred from FY 2006 through FY 2011. However, in anticipation of receiving requests for assistance associated with spotted skunks, WS could lethally remove up to 50 spotted skunks annually in the State. No unintentional take of spotted skunks has occurred by WS from FY 2006 from FY 2011. However, spotted skunks could be lethally removed unintentionally during other damage management activities. If take of up to 50 spotted skunks occurred by WS, the take would represent 0.1% of the estimated statewide population. Given the unlimited harvest allowed during the annual hunting and trapping seasons within the State, the cumulative

take of spotted skunks, including any take by WS, would not likely reach a level where adverse effects to the species' population would occur.

Coyote Population Information and Effects Analysis

Coyotes are a familiar mammal to most people. Their coloration is blended, primarily gray mixed with a reddish tint. The belly and throat are a paler color than the rest of the body (Beckoff 1982). Coyotes have long, rusty or yellowish legs with dark vertical lines on the lower foreleg. They are similar in appearance to gray and red wolves (National Audubon Society 2000). Color varies greatly; however, from nearly black to red or nearly white in some individuals and local populations. Most have dark or black guard hairs over their back and tail (Green et al. 1994). They sometimes breed with domestic dogs producing hybrids called "*coydogs*" (National Audubon Society 2000). The size of coyotes varies from about 20 to 40 lbs (9 to 18 kg) (Voigt and Berg 1987).

Coyotes range throughout the United States with the highest densities occurring on the Plains and in the south-central United States, including Texas. The distribution of coyotes in eastern North America began to expand beginning around 1900. Now, all eastern states and Canadian provinces have at least a small population of coyotes (Voigt and Berg 1987).

Coyotes often include many items in their diet. Rabbits are one of their most common prey. Other items in the coyote's diet include carrion, rodents, deer (usually fawns), insects (such as grasshoppers), as well as livestock and poultry. Coyotes readily eat fruits such as watermelons, berries, persimmons and other vegetative matter when it is available. In some areas, coyotes feed on human refuse at dumpsites and take small domestic pets such as cats and dogs (Voigt and Berg 1987).

Coyotes breed between January and March and are able to breed prior to reaching one year of age (Kennely and Johns 1976), but the percentage of yearlings having litters varies from zero to 80% in different populations (Gier 1968). This variation is influenced by a number of factors causing large annual variations in total number of coyotes breeding. In a study in Texas, the percentage of females having litters varied from 48% to 81% (Knowlton 1972). Pups are born after a gestation period of 60 to 63 days, with litter sizes varying primarily with prey availability. Gier (1968) reported average litter sizes of 4.8 to 5.1 in years with low rodent numbers, but litters of 5.8 to 6.2 during years with high rodent numbers. Litter sizes of one to 19 pups have been reported (National Audubon Society 2000).

Many references indicate that coyotes were originally found in relatively open habitats, particularly grasslands and sparsely wooded areas of the western United States. Today, coyotes have adapted to, and now exist in virtually every type of habitat, arctic to tropic, in North America. Coyotes live in deserts, swamps, tundra, grasslands, brush, dense forests, from below sea level to high mountain ranges, and at all intermediate altitudes. High densities of coyotes also appear in the suburbs of major cities (Green and Gipson 1994).

The coyote is probably the most extensively studied carnivore (Bekoff 1982), and considerable research has been conducted on population dynamics. Coyote densities as high as two per km^2 (5 per mi^2) have been reported in the southwestern and west-central United States, but are lower in other portions of the country, including eastern North America; although, few studies have accurately determined densities (Voigt and Berg 1987). Although coyote densities vary based on local habitat quality, Knowlton (1972) published that density estimates of 0.5 to 1.0 coyotes per mi^2 would likely be applicable to coyote densities across much of their range. However, methods for estimating carnivore populations are crude and often produce estimates with broad confidence intervals (Crawford et al. 1993).

Actual population estimates for coyotes in Tennessee are not available. Coyotes are common throughout the State and inhabit a variety of habitats. Because determinations of absolute coyote densities are frequently unknown (Knowlton 1972), many researchers have estimated coyote populations using various methods (Clark 1972, Knowlton 1972, Camenzind 1978, USDI 1979, Pyrah 1984). The cost to accurately determine absolute coyote densities over large areas is prohibitive (Connolly 1992) and would not appear to be warranted given the coyote's overall relative abundance. The presence of unusual food concentrations and the assistance provided to a breeding pair by non-breeding coyotes at the den can influence coyote densities and complicate efforts to estimate abundance (Danner and Smith 1980). Coyote densities are lowest in late winter prior to whelping, highest immediately after whelping, followed by a continued decline to the next whelping season (Parker 1995).

Predator abundance indices suggest that densities of coyotes in North America increase from north to south (Knowlton and Stoddart 1985, Parker 1995). Coyote densities range from 0.2 per square mile when populations are low (pre-whelping) to 3.6 coyotes per square mile when populations are high (post-whelping) (Knowlton 1972, USDI 1979). Although coyote densities vary considerably between habitat types and vary based on numerous environmental variables, Knowlton (1972) concluded that coyote densities might approach a high of five to six coyotes per square mile under extremely favorable conditions with densities of 0.5 to 1.0 per square mile possible over the entire range of the coyote in the United States. Such an estimate is speculative but represents some the best available information for estimating coyote populations.

Population modeling information suggests that a viable coyote population can withstand an annual removal of 70% of their population without causing a decline in the population (Connolly and Longhurst 1975, Connolly 1995). The unique resilience of the coyote, its ability to adapt, and its perseverance under adverse conditions is commonly recognized among biologists and land managers. Despite intensive historical damage management efforts in livestock production areas and despite sport hunting and trapping for fur, coyotes continue to thrive and expand their range, occurring widely across North and Central America (Miller 1995). Connolly and Longhurst (1975) determined that, "*...if 75% of the coyotes are killed each year, the population would be exterminated in slightly over 50 years*." However, Connolly and Longhurst (1975) go on to explain that their "*...model suggests that coyotes, through compensatory reproduction, can withstand an annual population mortality of 70%*" and that coyote populations would regain pre-control densities (through recruitment, reproduction, and migration) by the end of the fifth year after control was terminated even though 75% mortality had occurred for 20 years. In addition, other researchers (Windberg and Knowlton 1988) recognized that immigration, (not considered in the Connolly and Longhurst (1975) model) could result in rapid occupancy of vacant territories, which helps to explain why coyotes have thrived in spite of intensive damage management activities (Connolly 1978).

The statewide population of coyotes in Tennessee is currently not available. Given that coyotes inhabit every type of rural habitat within the State, the cumulative area of these classifications in Tennessee is 103,340 km^2 (39,899 mi^2) (see Table 4.1). If coyotes only occupy 50% of the rural habitat in Tennessee and the density of coyotes in the State ranges from 0.5 coyotes per square mile to five coyotes per square mile, the statewide population could be estimated to range from 10,000 coyotes to a high of 99,800 coyotes.

Coyotes are classified as a small game furbearer species in Tennessee with a continuous open harvest season and no limit on the number of coyotes that can be lethally taken (TWRA 2012*a*). The number of coyotes harvested annually in the State is currently not available. The number of coyotes taken to alleviate damage in the State is also currently unavailable. Between FY 2006 and FY 2011, WS conducted 120 technical assistance projects (average 20/year) associated with damage and threats of damages caused by coyotes, which includes only those projects where WS provided information on

managing damage or threats caused by coyotes. In most cases, WS refers requests for assistance associated with coyotes to the TWRA. Requests for assistance were primarily associated with threats to human safety and predation of animals. While providing technical assistance, people reported $23,695 in damages associated with coyotes from FY 2006 through FY 2011. WS also provide direct operational assistance associated with coyotes from FY 2006 through FY 2011, primarily at airports where coyotes posed strike risks to aircraft. During direct operational assistance projects, WS lethally removed 97 coyotes from FY 2006 through FY 2011, which is an average annual take of 16.2 coyotes.

Based on the number of requests for assistance received previously and the number of coyotes killed by WS to resolve damage, WS could take up to 200 coyotes annually under the proposed action to alleviate damage. Using a statewide coyote population ranging from 10,000 to 99,800 coyotes, take of up to 200 coyotes annually would represent from 0.2% to 2.0% of the estimated population. Although exact population estimates for coyotes in Tennessee and annual harvest rates are not available, the unlimited take allowed by the TWRA for the species during hunting and trapping seasons and as a nuisance species indicates the species is not at risk of overharvesting. Since the statewide population could reasonably be expected to be higher than 10,000 coyotes, the proposed take of 200 coyotes annually could actually be a smaller percentage of the actual statewide population.

Gray Fox Population Information and Effects Analysis

The gray fox is common in many parts of the United States where deciduous woodlands provide habitat. Yet, this secretive carnivore is seldom seen. This species is somewhat smaller in stature than the red fox, having shorter legs and extremities. Gray fox exhibit striking pelage that has grizzled upper parts resulting from individual guard hairs being banded with white, gray, and black. A predominance of black-tipped hairs in the middle of the back forms a dark longitudinal stripe that extends into a conspicuous black mane of coarse hair at the top of the black -tipped tail. Portions of the neck, sides, and limbs are cinnamon-colored. The ventral areas of a gray fox are buff colored. White shows on the ears, throat, chest, belly, and back legs, and the black, white, and reddish facial markings provide distinctive accents (Fritzell 1987).

Gray fox adults weigh about three to seven kilograms (6.5 to 15 lbs), with males being slightly larger than females. Generally, adult gray fox measure 80 to 113 cm (31.5 to 44 inches) from the tip of the nose to the tip of the tail. They inhabit wooded, brushy, and rocky habitats from extreme southern Canada to northern Venezuela and Colombia, excluding portions of the mountainous northwestern United States, the Great Plains, and eastern Central America. Gray fox occur over most of North America, north and east from southern California, Arizona, and central Texas (Fritzell 1987).

Gray fox prefer habitat with dense cover such as thickets, riparian areas, swampland, or rocky pinyon-cedar ridges. In eastern North America, this species is closely associated with edges of deciduous forest. They can also be found in urban areas where suitable habitat exists (Phillips and Schmidt 1994).

Gray fox mate from January through March and produce litters of one to seven kits after a gestation period of 53 days (National Audubon Society 2000). They rear young in a maternity den, commonly located in woodpiles, rocky outcrops, hollow trees, or brush piles (Phillips and Schmidt 1994). The male parent helps tend to the young but does not den with them. The young are weaned at three months and hunt for themselves at four months, when they weigh about 3.2 kg (7 lbs). Rabies and distemper are associated with this species (National Audubon Society 2000).

Accurate estimates of carnivore populations are rare and those for gray fox populations are no exception. Estimates based on knowledge of the species, experience, and intuition may be as accurate as those estimates based on recognized methods such as mark-recapture studies. Published estimates of gray fox

density vary from 1.2 to 2.1 per km^2 (3.1 to 5.4 per mi^2) depending on location, season, and method of estimation (Errington 1933, Gier 1948, Lord 1961, Trapp 1978). Over areas larger than 5,000 km^2 (1,930 mi^2) in which habitat quality varies, densities are likely lower. However, exceptionally high fox densities have been recorded in some situations (Grinnell et al. 1937, Hallberg and Trapp 1984).

Home ranges for gray fox vary throughout the year. Both males and females travel over larger areas during fall and winter, probably in response to increased energy demands and a declining food base (Follmann 1973, Nicholson 1982). During April, when young fox require regular feeding, a female's home range is less extensive than it is without the demands of those young (Follmann 1973). Although exceptions exist, eastern gray fox generally have larger home ranges than western animals (Fritzell 1987). For instance, 16 adult fox were tracked for more than one month in Alabama (Nicholson 1982) and Missouri (Haroldson and Fritzell 1984) and it was determined that they all had home ranges larger than 200 hectares (500 acres), and many exceeded 500 hectares (1,235 acres).

Gray fox feed on a wide variety of plant and animal matter, but feed on a wider variety of plant and animal matter than other North American canids (Fritzell 1987). Although active primarily at twilight and at night, the gray fox is sometimes seen foraging by day in brush, thick foliage, or timber. The only American canid with true climbing ability, gray fox occasionally forage in trees and often take refuge in them, especially leaning or thickly branched trees. The gray fox feeds heavily on cottontail rabbits, mice, voles, other small mammals, birds, insects, and plant material including corn, apples, persimmons, nuts, cherries, grapes, pokeweed fruit, grass, and blackberries. Grasshoppers and crickets are often a very important part of the diet in late summer and autumn (National Audubon Society 2000).

Gray fox can be found statewide in Tennessee but current population estimates are not available. Given the habitat preferences of gray fox, the most likely land cover types that would support gray fox are developed open space, deciduous forest, mixed forest, shrub/scrub, and woody wetlands. The cumulative area of those classifications in Tennessee is 62,000 km^2 (24,000 mi^2) (see Table 4.1). If gray fox only occupied 50% of those land classifications in the State and the density of gray fox in the State were 3.1 gray fox per square mile, the statewide population could be estimated at 37,200 gray fox. Gray fox can be found in a variety of habitats, including urban areas, so gray fox occupying only 50% of the land area of the State is unlikely since fox can be found almost statewide. However, similar to the other furbearing species, gray fox occupying only 50% of the land area was used to provide a minimum population estimate to evaluate the magnitude of the proposed take by WS.

Gray fox are classified as furbearers in Tennessee with annual hunting and trapping season in most areas of the State; however, in some areas, the take of gray fox during the hunting and trapping season is prohibited (TWRA 2012*a*). Throughout most of the State, gray fox can be harvested during annual open hunting and trapping seasons without a limit on the number that can be harvested during the open season. In Dyer, Lauderdale, Smith, and Wilson Counties, gray fox can be harvested using hunting and trapping methods throughout the year (no closed season) without a limit on the number that can be harvested. In Washington County, there is no open harvest season for gray fox (TWRA 2012*a*). The number of gray fox harvested annually in the State is currently not available. Gray fox are also likely taken to alleviate damage and threats of damage; however, the number of fox lethally taken annually to alleviate damage or threats of damage is currently unknown.

From FY 2006 through FY 2011, WS conducted 26 technical assistance projects (4.3 per year) associated with gray fox. Projects conducted by WS were primarily associated with disease threats and threats to aviation. WS also provided direct operational assistance associated with gray fox when requested from FY 2006 through FY 2011. During direct operational assistance projects to alleviate aircraft strike risks at airports, WS lethally remove eight gray fox from FY 2006 through FY 2011.

Based on previous requests received by WS to reduce damage and in anticipation of future requests, WS could intentionally remove up to 100 gray fox annually under the proposed action to address requests to alleviate damage and threats of damage. Using the lowest population estimate of 37,200 fox, the take of 100 gray fox by WS would represent 0.3% of the population.

Since the statewide population of gray fox is likely higher than 37,200 fox, WS' take of gray fox would represent a lower percentage of the actual statewide population. Like other mammal species addressed in this EA, the unlimited take allowed by the TWRA during the hunting and trapping seasons and allowing the take to alleviate damage by the TWRA provides an indication that gray fox populations maintain sufficient densities within the State to sustain unlimited harvest and that overharvest is unlikely.

Red Fox Population Information and Effects Analysis

The red fox is a typically proportioned member of the dog family. The bushy and unusually long tail, pointed ears, slender muzzle, and slanted eyes coupled with its small dog size and typical reddish coloration, make the red fox instantly recognizable to most people. This species is also the most common and well-known species in the genus *Vulpes*, which includes about 10 other species worldwide (Honacki et al. 1982). Typically, black-tipped ears, black cheek patches, white throat parts, a lighter underside, and black "*leg stockings*" are found on most red fox. The white tip of the tail (which is much more prominent in North American fox than elsewhere) can be used to distinguish brownish fox pups from similarly colored coyote pups, which lack a white tail tip (Voigt 1987).

In North America, the red fox weighs about 3.5 to 7 kg (7.7 to 15.4 lbs), with males averaging about one kg (2.2 lbs), which is heavier than females. Generally, adult fox measure 100 to 110 cm (39 to 43 inches) from the tip of the nose to the tip of the tail. Juveniles in their first autumn are similar in size to adults (Voigt 1987). Red fox occur throughout most of North America. They are found throughout most of the United States with the exception of a few isolated areas. Prehistoric fossil records suggest that the red fox may not have inhabited much of the United States; however, they were plentiful in many parts of Canada. Voigt (1987) has suggested climatic factors, interbreeding with the introduced European red fox, extirpation of the gray and red wolf, and clearing of land for agriculture has possibly contributed to the present-day expansion and range of this species in North America.

Red fox are adaptable to most habitats within their range, but usually prefer open country with moderate cover. Some of the highest fox densities reported are in the north-central United States and occur where woodlands are interspersed with farmlands. The range of the species has expanded in recent years to fill habitats formerly occupied by coyotes. The reduction of coyotes in many sagebrush/grassland areas of Montana and Wyoming has resulted in increased fox numbers. Red fox have also demonstrated their adaptability by establishing breeding populations in many urban areas of the United States, Canada, and Europe (Phillips and Schmidt 1994). In many areas, competition with other canids and the availability of suitable year-round food resources limit fox survival. Habitat determines the availability of year-round food resources and the presence or absence of other canids. Because those two factors strongly influence red fox survival, habitat limits fox numbers but seldom limits distribution (Voigt 1987).

Red fox mate from January through March and produce litters of one to 10 kits after a gestation period of 51 to 53 days. They rear young in a maternity den, commonly an enlarged woodchuck or badger den, usually in sparse ground cover on a slight rise, with a good view of all approaches (National Audubon Society 2000). Juvenile fox are able to breed before reaching a year old, but in areas of high red fox densities, most yearlings do not produce pups (Harris 1979, Voigt and MacDonald 1984, Voigt 1987). Gier (1968) reported average litter sizes of 4.8 to 5.1 in years with low rodent numbers, but litters of 5.8 to 6.2 during years with high rodent numbers. Litter sizes of one to 19 pups have been reported (National Audubon Society 2000). Offspring disperse from the denning area during the fall and establish breeding

areas in vacant territories, sometimes dispersing considerable distances. Red fox are generally solitary animals as adults, except when mating (Phillips and Schmidt 1994). Rabies and distemper are associated with this species.

The red fox is a skilled nonspecific predator, foraging on a variety of prey. It is also an efficient scavenger, and in parts of the world, garbage and carrion are extremely important to its diet (Voigt 1987). They are opportunists, feeding mostly on rabbits, mice, bird eggs, insects, and native fruit. They usually kill animals smaller than a rabbit, although fawns, pigs, kids, lambs, and poultry are sometimes taken (Phillips and Schmidt 1994). They also feed on squirrels, woodchucks, crayfish, and even grasses (National Audubon Society 2000).

Population densities are difficult to determine because of the secretive and elusive nature of fox. Estimates are prone to error even in open areas with good visibility. Methods used to estimate numbers have included aerial surveys, questionnaires to rural residents and mail carriers, scent post surveys, intensive ground searches, and indices derived from hunting and trapping harvest (Voigt 1987). Home ranges for red fox in the eastern United States are usually from 500 to 2,000 ha (1,235 to 4,940 acres) in rural settings, such as farmland (Voigt and Tinline 1980), but such sizes may not apply among fox populations in urban settings. In Great Britain, where food is abundant in many urban areas, densities as high as 30 fox per km^2 (78 per mi^2) have been reported (Harris 1977, MacDonald and Newdick 1982, Harris and Rayner 1986), while in southern Ontario, densities of about 1 fox per km^2 (2.6 per mi^2) occur during spring. This includes both pups and adults. In small areas of the best habitat, three times as many fox have been observed (Voigt 1987). However, those densities rarely occur extensively because of the dispersion of unsuitable habitat, high mortality, or from competition with coyotes (Voigt and Earle 1983). Cyclical changes in fox numbers occur routinely and complicate density estimates as well as management. Those cycles can occur because of changes in prey availability, or disease outbreaks, especially rabies, among red fox. For fox populations to remain relatively stable, mortality and reproduction must balance approximately.

Red fox can be found statewide in Tennessee; however, the statewide population in currently unknown. Given that red foxes are distributed through all rural habitats, the cumulative area of these classifications in Tennessee is about 97,000 km^2 (37,000 mi^2) (see Table 4.1). If red fox only occupy 50% of those land classifications in the State and the density of red fox in the State was 2.6 red fox per square mile, the statewide population could be estimated at 48,100 red fox. Similar to gray fox, red fox can be harvested during annual hunting and trapping seasons in the State; however, some exceptions occur. Like gray fox, red fox can be harvested during open hunting and trapping seasons throughout most of the State; however, fox can be harvested using hunting and trapping methods throughout the year in Dyer, Lauderdale, Smith, and Wilson Counties. There is no limit to the number of fox that can be harvested daily and no possession limit during the length of the season for red fox. However, in Washington County, there is no open harvest season for red fox. The number of red fox harvested annually in the State is currently unknown. Red fox could also be taken to alleviate damage and threats of damage. However, the number of fox lethally taken annually in the State to alleviate damage or threats of damage is also currently unknown.

WS conducted 56 technical assistance projects associated with red fox from FY 2006 through FY 2011, which is an average of 9.3 projects per year. Most requests for technical assistance were associated with disease threats, primarily rabies. One person requesting assistance reported $14,000 in damages to turf at a golf course from fox excavating a den on the course and from fox digging up turf and dirt to catch small mammals. During direct operational assistance projects, WS employed lethally removed 49 red fox from FY 2006 through FY 2011, which is an average take of 8.2 red fox per year.

Based on the number of requests for assistance received previously and based on the number of red fox addressed as part of those requests for assistance, WS could take up to 80 red fox annually under the proposed action. Using a statewide population estimate of 48,100 red fox, take of up to 80 red fox annually would represent 0.2% of the estimated population. Although exact population estimates for red fox in Tennessee are not available, the unlimited take allowed by the TWRA for the species during hunting and trapping seasons indicates the species is not at risk of overharvesting. The proposed take of red fox to alleviate damage would be a small component of the overall harvest of red fox in the State. The overall take would be of low magnitude when compared to the statewide population and the number of fox harvested during the annual hunting and trapping seasons.

Bobcat Population Information and Effects Analysis

The bobcat, also called "*wildcat*," is a medium-sized member of the North American cat family, and may be mistaken for a large bob-tailed domestic cat by some people. This species is actually two to three times larger than most domestic cats and appears more muscular and fuller in body. Their fur is dense, soft, short and generally yellowish to reddish-brown in color with numerous black spots and black-tipped guard hairs on the back and white with black spots on the belly. Forelegs are tawny with black bars. The species gets its common name from its characteristic stubby or "*bobbed*" tail. The tail is generally only 9 to 20 cm (3.5 to 8 inches) in length with two or three black bars and a black tip above, while the underside is pale or white (Larivière and Walton 1997). Their upper legs have dark horizontal bands. The face has thin, black lines stretching onto broad cheek ruff and their ears are tufted. Males are generally larger than females. The length of bobcats ranges from 47.5 to 125 cm (19 to 49 inches), while their weight ranges between 4.1 and 18.3 kg (9 to 40 lbs) (Larivière and Walton 1997).

Bobcats are capable of hunting and killing prey that range from the size of a mouse to that of a deer. Rabbits, tree squirrels, ground squirrels, woodrats, porcupines, pocket gophers, and ground hogs comprise most of their diet. Opossums, raccoon, grouse, wild turkey, and other ground nesting birds are also eaten. Occasionally, insects and reptiles can be part of a bobcat's diet. They also resort to scavenging. They are opportunistic predators, and may feed on livestock and domestic animals such as poultry, sheep, goats, house cats, small dogs, exotic birds and game animals, and rarely, calves (Virchow and Hogeland 1994). McCord and Cardoza (1982) reported the cottontail rabbit to be the principal prey of bobcats throughout their range.

Ruell et al. (2009) reported bobcat densities ranged from 0.65 to 1.09 bobcats per square mile (0.25 to 0.42 bobcats per km^2) in coastal southern California in both large open habitat and in habitat surrounded by human developments. Lawhead (1984) reported bobcat densities of 0.66 per square mile (0.26 bobcats per km^2) in Arizona with a preference for riparian habitat. Bobcats in southern Illinois were reported to have a population density of 0.70 bobcats per square mile (0.27 bobcats per km^2) (Nielsen and Woolf 2001), while Anderson (1987) provided population density estimates of 0.13 to 0.26 bobcats per square mile (0.05 to 0.10 bobcats per km^2). Bobcats reach densities of about one per 0.7 km^2 (1 per ¼ mi^2) on some islands in the Gulf Coast of the southeastern United States. Densities vary from about one per 1.3 km^2 (1 per ½ mi^2) in coastal plains to about one cat per 10.7 km^2 (1 per 4 mi^2) in portions of the Appalachian foothills. Mid-Atlantic and mid-western states usually have scarce populations of bobcats (Virchow and Hogeland 1994). Populations are stable in many northern states and reviving in other states where intensive trapping formerly decimated the species (National Audubon Society 2000). Rates of natural mortality reported for adult bobcats in protected populations appear to be quite low. Crowe (1975) estimated a 3% mortality rate in a protected population, based on Bailey's (1972) study of bobcats in southeastern Idaho. Causes of natural mortality for adult bobcats include starvation (Hamilton 1982), disease and predation (Lembeck 1978), and injuries inflicted by prey (Fuller et al. 1985).

Bobcats can be found statewide in Tennessee in suitable habitat. The statewide bobcat population is currently unknown. Given that bobcats are distributed through all rural habitats, the cumulative area of these classifications in Tennessee is about 97,000 km^2 (37,000 mi^2) (see Table 4.1). If bobcats only occupied 50% of those land classifications in the State and the density of bobcats in the State was estimated at a low of 0.13 bobcats per square mile to a high of 1.09 bobcats per square mile, the statewide population could be estimated at between approximately 2,400 and 20,200 individuals. Bobcats can be found in a variety of habitats, including developed areas, so bobcats occupying only 50% of the land area of the State would be unlikely since bobcats can be found almost statewide. However, similar to the other furbearing species, bobcat occupying only 50% of certain land classifications was used to provide a minimum population estimate to determine the magnitude of the proposed take by WS.

Bobcats are classified as a furbearer in Tennessee and may be taken during hunting and trapping seasons. During the hunting season, only one bobcat can be harvested per day with a possession limit of two bobcats (TWRA 2012*a*). During the trapping season, the TWRA allows an unlimited number of bobcats to be harvested during the length of the season (TWRA 2012*a*). Bobcats could also be lethally removed to alleviate damage in the State; however, the number of bobcats removed to alleviate damage is unknown.

WS conducted four technical assistance projects from FY 2006 through FY 2011 associated with damage or threats caused by bobcats. Those requests for assistance were primarily associated with threats to aviation safety and threats to family pets. From FY 2006 through FY 2011, WS intentionally removed four bobcats to reduce strike risks at airports. In anticipation of receiving requests for assistance, WS reasonably expects the total take of bobcats would not exceed 50 bobcats annually in Tennessee to resolve requests to manage damage to resources and threats to human health and safety.

Based upon the aforementioned population estimate, WS' limited lethal take of up to 50 bobcats annually under the proposed action would represent 0.3% to 2.1% of a statewide population estimated to be between 2,400 and 20,200 individuals. The proposed take of bobcats by WS in the State would be of low magnitude when compared to the actual statewide population.

Feral and Free-ranging Cat Population Information and Effects Analysis

Feral cats are domesticated cats living in the wild. They are small in stature, weighing from three to eight pounds (1.4 to 3.6 kg), standing eight to 12 inches (20 to 30.5 cm) high at the shoulder, and 14 to 24 inches (35.5 to 61 cm) long. The tail adds another 20 to 30.5 cm (8 to 12 inches) to their length. Colors range from black to white to orange, and a variety of combinations of those colors. Other hair characteristics also vary greatly (Fitzwater 1994).

Feral cats are found in commensal relationships wherever people are found. In some urban and suburban areas, cat populations equal human populations. In many suburban and eastern rural areas, feral cats are the most abundant predators. They are opportunistic predators and scavengers that feed on rodents, rabbits, shrews, moles, birds, insects, reptiles, amphibians, fish, carrion, garbage, vegetation, and leftover pet food (Fitzwater 1994).

Feral cats produce two to 10 kittens during any month of the year. An adult female may produce three litters per year where food and habitat are sufficient. Cats may be active during the day but typically are more active during twilight or night. House cats have been reported to live up to 27 years, but feral cats probably average only three to five years. They are territorial and move within a home range of roughly 4 km^2 (1.5 mi^2). After several generations, feral cats can be considered wild in habits and temperament (Fitzwater 1994).

Where it has been documented, the impact of feral cats on wildlife populations in suburban and rural areas, directly by predation, and indirectly by competition for food, has been enormous (Coleman and Temple 1989). In the United Kingdom, one study determined that house cats might take an annual toll of some 70 million animals and birds (Churcher and Lawton 1987). American birds face an estimated 117 to 157 million exotic predators in the form of free-ranging domestic cats, which are estimated to kill at least one billion birds every year in the United States. Cats have contributed to declines and extinctions of birds worldwide, with feral cats considered one of the most important drivers of global bird extinctions (Dauphine and Cooper 2009).

Feral and free-ranging cats also pose a health and safety threat to household pets. Feral and stray cats are at increased risk of feline immunodeficiency virus, feline leukemia, feline panleukopenia virus, also known as feline distemper, and rabies. All of these diseases can be transmitted to unvaccinated pet cats allowed to free-range. Feline panleukopenia virus is highly contagious, may survive in the environment for up to a year, and may be transmitted to indoor cats through indirect routes, such as on shoes (Berthier et al. 2000, Truyen et al. 2009). In addition, feral and free-ranging cats serve as a reservoir for wildlife and human diseases, including cat scratch fever, histoplasmosis, leptospirosis, mumps, plague, rabies, ringworm, salmonellosis, toxoplasmosis, tularemia, and various parasites (Fitzwater 1994).

The number of feral cats in Tennessee is unknown. Feral and free-ranging cats are considered by many wildlife biologists and ornithologists to be a detriment to native wildlife species. Feral cats prey upon native wildlife species and compete with native predators for prey. Thus, removing feral cats could be considered as providing some benefit to the natural environment by eliminating predation and competition from an introduced species.

Between FY 2006 and FY 2011, WS addressed feral or free-ranging cat damage or threats of damage during 19 projects. Individuals requesting assistance reported $20,050 in damages associated with feral or free-ranging cats. Requests for assistance were primarily associated with human safety threats associated with cats and damage to property. During direct operational assistance projects conducted by WS from FY 2006 through FY 2011, WS intentionally removed using lethal methods 23 feral cats across the State. In addition, one feral cat was lethally taken unintentionally during other damage management activities conducted between FY 2006 and FY 2011. Feral and free-ranging cats were also live-captured and released by WS between FY 2006 through FY 2011. In addition, 94 feral or free-ranging cats were intentionally live-captured and released unharmed or were relinquished to a local animal control facility for care and determine their adoptability. Between FY 2006 and FY 2011, 154 cats were unintentionally live-captured by WS across the State. Those cats unintentionally live-captured were released unharmed or relinquished to a local animal control facility.

In most cases, WS would employ live-capture methods to alleviate damage or threats of damage associated with feral or free-ranging cats. Once live-captured, WS would transfer custody of the cats to a local animal control facility. After relinquishing the feral cats to a local animal control facility, the care and the final disposition of the cat would be the responsibility of the animal control facility. However, in some cases, WS may be requested to lethally remove feral cats to alleviate damage or threats. In anticipation of receiving requests to lethally remove feral cats, up to 200 feral cats could be lethally removed by WS annually. Based upon the above information, WS' limited removal of feral cats would have minimal effects on local or statewide populations Tennessee. Some local populations may be temporarily reduced at a local site if cats were removed using non-lethal or lethal methods. In those cases where feral cats were causing damage or were creating a nuisance and complete removal of the local population could be achieved, this could be considered as providing some benefit to the natural environment since feral cats are not considered part of the native ecosystem.

Feral Dog Population Information and Effects Analysis

Like domestic dogs, feral dogs (sometimes referred to as wild or free-ranging dogs) manifest themselves in a variety of shapes, sizes, colors, and even breeds. McKnight (1964) noted German shepherds, Doberman pinschers, and collies as breeds that often become feral. Most feral dogs today are descendants of domestic dogs that appear similar to dog breeds that are locally common (Green and Gipson 1994). The primary feature that distinguishes feral from domestic dogs is the degree of reliance or dependence on humans, and in some respect, their behavior toward people. Feral dogs survive and reproduce independently of human intervention or assistance. While it is true that some feral dogs use human garbage for food, others acquire their primary subsistence by hunting and scavenging like other wild canids.

Feral and domestic dogs often differ markedly in their behavior toward people. Scott and Causey (1973) based their classification of those two types by observing the behavior of dogs while confined in cage traps. Domestic dogs usually wagged their tails or exhibited a calm disposition when a human approached; whereas, most feral dogs showed highly aggressive behavior, growling, barking, and attempting to bite. Some dogs were intermediate in their behavior and could not be classified as either feral or domestic based solely on their reaction to humans. Since many feral dogs have been pursued, shot at, or trapped by people, their aggressive behavior toward humans is not surprising. Gipson (1983) described the numerous lead pellets imbedded under the skin of a feral dog caught in Arkansas as a testament to its relationship with people (Green and Gipson 1994).

Feral dogs are usually secretive and wary of people. Thus, they are active during dawn, dusk, and at night, much like other wild canids. They often travel in packs or groups and may have rendezvous sites, similar to wolves. Travel routes to and from gathering sites or den sites may be well defined. Food scraps and other evidence of concentrated activity may be observed at gathering sites.

The appearance of tracks left by feral dogs varies with the size and weight of the animal. Generally, dog tracks are more round and show more prominent nail marks than those of coyotes, and they are usually larger than those of fox. Since a pack of feral dogs likely consists of animals in a variety of sizes and shapes, the tracks from a pack of dogs will be correspondingly varied, unlike the tracks of a group of coyotes (Green and Gipson 1994).

Feral dogs may occur where people permit their dogs to roam free or where people abandon unwanted dogs. Feral dogs probably occur in all of the 50 states, Canada, and Central and South America. They are also common in Europe, Australia, Africa, and on several remote ocean islands, such as the Galapagos. Home ranges of feral dogs vary considerably in size, with size likely influenced by the availability of food. Dog packs that are primarily dependent on garbage may remain in the immediate vicinity of a landfill, while other packs that depend on livestock or wild game may forage over an area of 130 km^2 (50 mi^2) or more (Green and Gipson 1994).

Feral dogs are often found in forested areas or scrublands near human habitation. Some people will not tolerate feral dogs in close proximity to human activity; thus, they take considerable effort to eliminate them in such areas. Feral dogs may be found on lands where human access is limited, such as military reservations and large airports. They may also live in remote sites, where they feed on wildlife and native fruits. The only areas that do not appear to be suitable for feral dogs are places where food and escape cover are not available, or where large native carnivores, particularly wolves, are common and prey on dogs (Green and Gipson 1994).

Like coyotes, feral dogs are best described as opportunistic feeders. They can be efficient predators, preying on small and large animals, including domestic livestock. Many rely on carrion, particularly

road-killed animals, crippled waterfowl, green vegetation, berries, and other fruits, and refuse at garbage dumps (Green and Gipson 1994).

Feral dogs are highly adaptable, social carnivores. Gipson (1983) suggested that family groups of feral dogs are more highly organized than previously believed. Pup rearing may be shared by several members of a pack. Survival of pups born during autumn and winter has been documented, even in areas with harsh winter weather. Gipson (1983) found that only one female in a pack of feral dogs studied in Alaska gave birth during two years of study, even though other adult females were present in the pack. The breeding female gave birth during late September or early October during both years. Gipson (1983) indicated that all pups from both litters had similar color markings, suggesting that the pups had the same father. Adult males of different colors were present in the pack.

Nesbitt (1975) commented on the rigid social organization of a pack of feral dogs where nonresident dogs were excluded, including females in estrus. In one instance, Nesbitt (1975) used three separate female dogs in estrus as bait (dogs were chained in the back of a corral-type trap) over a 59-day period and captured no feral dogs. Nesbitt (1975) then baited the same trap with carrion, and a pack of feral dogs, including four adult males, entered the trap within one week (Green and Gipson 1994).

Hybridization between feral dogs and other wild canids can occur, but non-synchronous estrus periods and pack behavior (that is, excluding non-resident canids from membership in the pack) may preclude much interbreeding. Dens may be burrows dug in the ground or sheltered spots under abandoned buildings or farm machinery. Feral dogs commonly use former fox or coyote dens (Green and Gipson 1994).

Feral dogs can cause damage by preying on livestock, poultry, house cats, or domestic dogs. They may also feed on fruit crops including melons, berries, grapes, and native fruit. They may also attack people, especially children. This is especially true where they feed at and live around landfills near human dwellings (Green and Gipson 1994). In some locales, they may present a serious threat to deer (Lowry 1978) and other valuable wildlife (Green and Gipson 1994).

WS provided technical assistance to requesters associated with dogs during 14 projects from FY 2006 through FY 2011. However, most requests for assistance were referred to a local animal control facility since requesters were usually unable to determine if a dog is feral or a pet. From FY 2006 through FY 2011, WS lethally removed 16 feral dogs during damage management activities in Tennessee. WS has also live-captured and released 11 feral dogs during damage management activities conducted from FY 2006 through FY 2011. In addition, WS live-captured eight feral dogs unintentionally between FY 2006 and FY 2011, which were released unharmed. WS employed non-lethal harassment methods to disperse 34 dogs between FY 2006 and FY 2011. Based on previous requests for assistance and in anticipation of receiving additional requests for assistance, WS could lethally remove up to 200 feral dogs per year under the proposed action alternative. In most cases, WS would employ live-capture methods to alleviate damage or threats of damage associated with dogs. Once live-captured, WS would transfer custody of the dogs to a local animal control facility. After relinquishing the dogs to a local animal control facility, the care and the final disposition of the dog would be the responsibility of the animal control facility.

Based upon the above information, WS' limited lethal removal of feral dogs should have no adverse effects on overall populations in Tennessee. Any activities involving lethal control actions by WS would be restricted to isolated individual sites. Some local populations may be temporarily reduced as a result of removals aimed at reducing damage at a local site. In those cases where feral dogs were causing damage or posing as a nuisance and complete removal of the local population could be achieved, this would be considered as providing some benefit to the natural environment since feral dogs are not considered part of the native ecosystem.

Nine-banded Armadillo Population Information and Effects Analysis

The nine-banded armadillo is easily recognized due to its unique appearance. An opossum-sized animal, the armadillo has a "*shell*", which is composed of ossified dermal plates covered by a leathery epidermis (Whitaker, Jr. and Hamilton, Jr. 1998). The armadillo is the only North American mammal that has heavy bony plates (National Audubon Society 2000). Female armadillos produce one litter of young per year, which are identical quadruplets (National Audubon Society 2000).

Originally thought to occur in Central and South America, including Mexico, the nine-banded armadillo has undergone a northward and eastward expansion into the United States since the late-1800s, likely through natural dispersal from Mexico and release of captive armadillos (Layne 2003). Today, the armadillo can be found across the southern portion of the United States with additional dispersal northward and eastward in the United States likely in the future (Layne 2003). Range expansion is likely only limited by the reduced food availability and the colder temperatures experienced during the winter months.

Armadillos do not tolerate extended periods of cold weather, which may limit their expansion northward. Armadillos do not hibernate and must feed every couple of days during winter months since they do not store food nor accumulate efficient amounts of body fat to survive through the winter. The presence of snow or frozen soils limits the availability of food sources, primarily the availability of insects, during winter months. The lack of food available often causes armadillos to starve during winter months. However, in Tennessee, winter temperatures are relatively sufficient to maintain armadillo populations, though periods of extreme cold or prolonged periods of cold temperatures may temporarily affect populations.

Armadillos occupy and exploit a variety of natural and human-modified terrestrial habitats in the United States and across their range, including those armadillos found Tennessee. Layne (2003) summarizes the natural habitat types occupied by armadillos throughout their range as "*...pine-oak woodlands, oak-elm woodlands, pine forests, mixed pine-hardwood forests, bottomland forests, riparian woodlands, mesic hardwood forests, scrub, chaparral-mixed grass, inland and coastal prairies, salt marsh, coastal dunes, and coastal strand*." Human-modified habitats where armadillos can be found which has been summarized by Layne (2003) included "*...pastures, parkland, cemeteries, golf courses, citrus groves, pine plantations, plant nurseries, cut-over pineland, and various croplands*." The ability of armadillos to exploit a wide variety of habitat types is likely one of the main components facilitating the range expansion of the armadillo into and across the United States (Layne 2003). Habitat suitability is likely more of a function of soil substrate rather than vegetative type due to the foraging and digging behavior of armadillos (Layne 2003).

Armadillos are opportunistic feeders and will often forage by digging and probing the soil, leaf litter, and decaying wood for invertebrates, primarily insects. One study found at least 488 different food items in the stomachs of 281 armadillo with insects and other invertebrates comprising 92% of the stomach contents (Kalmbach 1943). Armadillos are also known to forage on plant material and small vertebrates with food preferences often driven by the availability of food sources (Layne 2003).

The other limiting factor in armadillo expansion and for maintaining populations is the presence of sandy or clay soils. Armadillos are prolific diggers and damages attributed to armadillos are often associated with their digging behavior. Armadillos will dig out shelters and dig while rooting out invertebrates in the soil and leaf litter. This digging and rooting behavior are the most common complaints from resource owners in Tennessee. Damage to landscaping is the most common resource being damaged by armadillos

in Tennessee. Sandy soils are conducive to digging and armadillos can be found in those areas in Tennessee where sandy soils are present.

Armadillos were first casually observed in Tennessee in the 1990s. Today, armadillos can be found in portions of west and middle Tennessee with populations slowly expanding eastward. While armadillos are not rare in Tennessee, they are not considered common. Armadillos in Tennessee are showing a general increasing trend (B. Robbins, USDA pers. comm. 2010). Population estimates for armadillos in the United States range from 30 to 50 million armadillos (Gilbert 1995). However, population estimates in Tennessee are not currently available. Armadillos can be harvested during a continuous open hunting season in the State, which places no limits on the number of armadillos that can be harvested (TWRA 2012a).

Therefore, a population estimate will be derived based on the best available information for armadillos to provide an indication of the magnitude of take proposed by WS to alleviate damage and threats of damage. Population densities for armadillos are reported to range from 0.004 to 1.4 armadillos per acre with an average of 0.25 armadillos per acre (Mengak 2005). Based on the natural habitat types occupied by armadillos throughout their range summarized by Layne (2003), the cumulative area of those classifications in Tennessee is about 93,500 km^2 (36,100 mi^2) (see Table 4.1). Using a population density estimated at 0.004 to 1.4 armadillos per acre and if armadillos only inhabited 25% of those land classifications in the State, the statewide population could range from approximately 23,100 armadillos to approximately 8 million armadillos. With an average of 0.25 armadillos per acre, the statewide population could be estimated at 1.4 million armadillos. As stated previously, the actual number of armadillos in the State is currently unknown. The range of armadillos only encompassing 25% of the land area in the State was used to provide a minimum population estimate to determine the magnitude of the proposed take by WS to alleviate or prevent nest predation.

Since FY 2006, the WS program in Tennessee has conducted nine technical assistance projects associated with armadillos. During those projects, people reported $1,900 in damages associated with armadillos in the State. No direct operational assistance projects were conducted by WS associated with armadillos from FY 2006 through FY 2011. Based on previous requests for assistance received by WS and in anticipation of additional requests for assistance, WS could lethally remove up to 200 armadillos annually in the State as part of efforts to alleviate and prevent damage. Given the range of population estimates in the State, the take of 200 armadillos by WS annually would represent 0.9% of the statewide population based on a population estimated at 23,100 armadillos if the overall population remains at least stable. Although the number of armadillos lethally taken by other entities in the State to alleviate damage is unknown, the cumulative take of armadillo, including the proposed take of up to 200 armadillos annually by WS, is likely of low magnitude when compared to the statewide population of armadillos.

Virginia Opossum Population Information and Effects Analysis

Opossums are the only marsupials (possess a pouch in which young are reared) found north of Mexico (Seidensticker et al. 1987). They frequent most of the eastern and central United States, except Minnesota, northern Michigan, and New England, extending west to Wyoming, Colorado, and central New Mexico (National Audubon Society 2000). They are also found in parts of the southwestern United States, California, Oregon, and Washington (Jackson 1994*b*). Adults range in size from less than 1 kg (2.2 lbs) to about 6 kg (13 lbs), depending on sex and time of year. They grow throughout life (Seidensticker et al. 1987). They have a fairly broad range of pelage colors, but are usually considered as "*gray*" or "*black*" phase. Their fur is grizzled white above; long white hairs cover black tipped fur below. They climb well and feed on a variety of foods, including carrion, which forms much of its diet. In addition, opossum eat insects, frogs, birds, snakes, small mammals, earthworms, and berries and other fruits; persimmons, apples, and corn are favorite foods (National Audubon Society 2000). They use a

home range of four to 20 hectares (10 to 50 acres), foraging throughout this area frequently (Jackson 1994*b*), but concentrating on a few sites where fruits abound, when they are in season (Seidensticker et al. 1987).

The reproductive season of the Virginia opossum typically occurs from December to February, depending on latitude (Gardner 1982). Gestation is short (average of 12.8 days) with one to 17 young born in an embryonic state that climb up the mothers belly to the marsupium (pouch), attach to teats, and begin to suckle (Gardner 1982, National Audubon Society 2000). Those young remain in the pouch for about two months. After two months, the young begin to explore and may be found traveling on their mother's back with their tails grasping hers (Whitaker, Jr., and Hamilton, Jr. 1998). Opossums live for only one to two years, with as few as 8% of a population of those animals surviving into the second year in a study in Virginia conducted by Seidensticker et al. (1987). In the five-year study conducted by Seidensticker et al. (1987), there was a wide variation in opossum numbers in what was considered excellent habitat for the species. Those variations were observed seasonally and in different years. However, the mean density during the study was 10.1 opossum per square mile with a range of 1.3 opossum per square mile to 20.2 opossum per square mile (Seidensticker et al. 1987). This was comparable to other opossum population densities in similar habitats in Virginia. Verts (1963) found a density estimate of 10.1 opossum per square mile in farmland areas in Illinois while Wiseman and Hendrickson (1950) found a density of 6.0 opossum per square mile in mixed pasture and woodlands in Iowa. However, VanDruff (1971) found opossum densities in waterfowl nesting habitat as high as 259 opossum per square mile.

Opossum are common throughout Tennessee in appropriate habitat. Population estimates for opossum in the State are not available. Therefore, a population estimate will be derived based on the best available information for opossum to provide an indication of the magnitude of take proposed by WS to alleviate damage and threats of damage. Given that opossums typically prefer rural areas but likely avoid hay/pasture, the cumulative area of suitable classifications in Tennessee is about 77,000 km^2 (29,700 mi^2) (see Table 4.1). If opossum were only found on 50% of those land classifications in the State and using a mean density of 10.1 opossum per square mile found by Seidensticker et al. (1987) in Virginia, the population would be estimated at nearly 150,000 opossum. Using the range of opossum densities found by Seidensticker et al. (1987) of 1.3 opossum per square mile to 20.2 opossum per square mile and only 50% of those land classification in the State being occupied by opossum, the statewide population would range from a low of 19,300 opossum to a high of nearly 300,000 opossum.

Opossum can be found in a variety of habitats, including urban areas, so opossum occupying only 50% of those land classifications in the State is unlikely since opossum can be found almost statewide. However, opossum occupying only 50% of the land area was used to provide a minimum population estimate to determine the magnitude of the proposed take by WS to alleviate or prevent damage.

Opossum are considered a furbearing species in the State and can be harvested during annual hunting and trapping seasons (TWRA 2012*a*). During the development of the EA, opossum could be harvested during hunting and trapping seasons with no limit on the number that could be taken during those seasons. The number of opossum harvested during the annual hunting and trapping season in the State is currently not available. In addition, opossum could be lethally removed to alleviate damage; however, the number of opossum lethally taken in the State to alleviate damage is also unknown.

As part of damage management activities conducted by WS in the State, 214 opossum have been lethally taken by WS from FY 2006 through FY 2011 (209 target take; 5 non-target take). On average, WS has lethally removed 35 opossum annually from FY 2006 through FY 2011 to alleviate damage and threats in the State. In addition, WS has purposefully live-captured and released 64 opossum from FY 2006 through FY 2011 with an additional 950 opossum live-captured unintentionally during other damage management activities and released unharmed. Opossum are primarily live-captured as non-targets

during surveillance activities relating to the ORV rabies program (USDA 2005*b*). Based on previous requests for assistance received by WS and in anticipation of additional requests for assistance, WS could lethally remove up to 250 opossum annually in the State as part of efforts to reduce or eliminate damage under the proposed action alternative. Based on a statewide population ranging from 19,300 opossum to 300,000 opossum, the lethal take of up to 250 opossum annually by WS under the proposed action alternative, would represent 0.1% to 1.3% of the estimated population.

Although the total number of opossum lethally taken in the State during the annual hunting and trapping seasons and for damage management is unknown, the cumulative take of opossum, including the proposed take of up to 500 opossum annually by WS, would be of a low magnitude when compared to the statewide population. The unlimited harvest allowed by the TWRA during the harvest seasons provides an indication that population densities of opossum in the State are sufficient that overharvest is not likely to occur, including lethal take to alleviate or prevent damage. In addition, the live-capture and subsequent release of opossum would not likely result in adverse effects to the statewide population since those animals would be released unharmed.

Bat Population Information and Effects Analysis

Bats are the only mammals that can truly fly. They are exceeded only by rodents as the most numerous mammals, both in number of species and number of individuals (Greenhall and Frantz 1994, National Audubon Society 2000). Bat bodies are generally well furred and forelimbs are enlarged and developed as wings with membranes attached to four greatly elongated fingers, which spread when in flight and draw together when at rest. The "*thumb*" projects from the end of the "*forearm*" as a small but sharp claw that is used as the animal crawls about. Wing membranes are often naked and translucent (National Audubon Society 2000). The motion of bats in the air appears to be more of a swimming motion, where they rotate their wings to catch air with the membrane, as opposed to birds that flap their wings (National Audubon Society 2000). Although most North American bats have small eyes, their visual acuity is good (Humphrey 1982). However, insectivorous bats locate food and avoid objects by means of echolocation, which is similar to radar or sonar (Humphrey 1982). While flying, the bat emits through its nose or mouth a continuous series of supersonic sounds. These sounds bounce off objects and are picked up by the bats complex ears (National Audubon Society 2000).

Bats are nocturnal, leaving the roosts around dusk and usually flying to a stream, pond, or lake, where they obtain a drink by skimming the surface and dipping their lower jaw into the water. Bats in North America are virtually all insectivorous, feeding on a variety of flying insects, many of which are harmful to humans (Greenhall and Frantz 1994). Insectivorous bats obtain food by various means of capturing their prey mostly while in flight. During these feeding flights, they often fly close to animals, including humans and sometimes cause alarm (National Audubon Society 2000).

Among the 40 species of bats found north of Mexico, only a few cause problems for humans. Bats congregating in groups are called colonial bats; those that live alone are known as solitary bats. The colonial species most often encountered in and around human buildings in the United States are the little brown bat, the Brazilian free-tailed bat (*Tadarida brasiliensis*), the pallid bat (*Antrozous pallidus*), the Yuma myotis (*Myotis yumanensis*), and the evening bat. Solitary bats typically roost in tree foliage or under bark. However, occasionally solitary bats are found associated with buildings, some only as transients during migration (Greenhall and Frantz 1994).

Conflicts involving bats can include property damage, but primarily involve threats to human, pet, and livestock health. The buildup of bat droppings and urine in attics and between walls can result in odor problems and discoloration of walls and ceilings (Agency for Toxic Substances and Disease Registry 1998). Bat rabies has always occurred at low levels within bat populations and researchers estimate that

less than 1% of all bats are rabid. About 6% of all bats submitted for rabies testing were submitted because they could be easily captured, were obviously weak or sick, or had been captured by a cat or dog (CDC 2011). In addition to the threat of rabies from direct contact or a bat entering the living area of a home, there are other threats associated with bat colonies including histoplasmosis, fungal spores, and mites.

Bat droppings, particularly when they are thick, are likely to be contaminated with the fungus, *Histoplasma capsulatum* or with fungi species such as molds, especially in warm, moist conditions. As long as people are not in contact with fungal spores, they are unlikely to be affected by them. When people inhale spores from *Histoplasma capsulatum*, they may become ill with a disease known as histoplasmosis. Symptoms of histoplasmosis include some combination of mild, flu-like respiratory illness, a general ill feeling, chest pain, fever, cough, headache, loss of appetite, shortness of breath, joint and muscle pains, chills, and hoarseness. Although there are other, more rare illnesses associated with exposure, the most likely is histoplasmosis. Similarly, mold spores released into the air may result in increases in asthma attacks (Agency for Toxic Substances and Disease Registry 1998).

Bat bugs (*Cimex adjunctus*) are free-living ectoparasites of bats that feed on blood. They will bite humans in the absence of their primary hosts. The main means of dispersal for bat bugs is phoresy, hitching a ride on a bat to a new location by clinging to the fur of their host animal. Typically, bat bug infestations originate from bat populations established in attics, wall voids, unused chimneys, or uninhabited portions of a house. Bat bugs typically do not wander far from occupied bat roosting sites where they have easy access to food. However, if their normal hosts are eliminated or have vacated an area, bat bugs will seek other sources of food and may crawl about and invade living areas within a house (Jones and Jordan 2004). Similarly, bat mites may enter the home and bite people. Although their bite is not particularly harmful, the person may experience an allergic reaction and develop a skin rash in response (Agency for Toxic Substances and Disease Registry 1998).

At least sixteen species of bats are known to occur in Tennessee (Tennessee Bat Working Group 2012). Some of those are federally listed as threatened, endangered, or as species of management concern. Additionally, those bats and others are State listed as endangered, threatened, or as species of special concern by the TWRA. Table 4.2 identifies the bat species found in Tennessee. Further, information related to occurrence and roosting/nesting/hibernating information is given in Table 4.2. Several bat species in Tennessee are known to roost, raise young, or hibernate in various human structures. Such behavior sometimes causes human/bat conflicts, especially perceived or actual threats of rabies, by people who encounter bats in such locations.

Several bat species in Tennessee are known to roost, raise young, or hibernate in various human structures. Such behavior sometimes causes human/bat conflicts, especially perceived or actual threats of rabies, by people who encounter bats in such locations, especially when bats enter the living space of a home. From FY 2006 to FY 2011, WS responded to 59 requests for assistance associated with bats. Those persons requesting assistance reported $13,000 in damages caused by bats, primarily from damage to property. Most requests for assistance were addressed through technical assistance. More than 90% of requests for assistance were associated with bats that had wandered into the living or working spaces of buildings, or were roosting in various structures. Most requests for assistance were resolved by providing an escape route for the intruding bat, by capturing the bat and releasing it, or by excluding bats from buildings. From FY 2006 through FY 2011, one bat was hand captured inside a building and euthanized by WS in Tennessee for rabies testing.

In future program activities, bat damage would be handled by WS primarily through various technical assistance projects or referral to other entities. Program activities would continue to recommend the use of non-lethal methods, such as exclusion and live capture/release. To reduce the possibility of adversely

affecting a bat maternity colony, WS would implement and recommended to persons receiving technical assistance that all exclusion be conducted from September to early November, when practicable. Many bat species, except big brown bats, would have migrated at that time, and the rearing of young would have been completed. Therefore, activities conducted after this date would be highly unlikely to disturb maternity colonies of any species during critical young-rearing periods. Conducting exclusionary and other projects during those months would also give big brown bats, or other species that overwinter in Tennessee, an opportunity to find alternate roost sites before the onset of extremely cold weather.

Table 4.2: Bats Found in Tennessee (From TWRA's Tennessee Bat Working Group)

Common Name	Scientific Name	Occurrence	Roosting/Rearing/Hibernating Habitat	Status in TN*
Big Brown Bat	*Eptesicus fuscus*	year-round, statewide	buildings during the spring, summer, and winter, caves in the winter	PN
Eastern Pipistrelle	*Pipistrellus subflavus*	common during summer and migration	hollow trees, buildings, caves	PN
Evening Bat	*Nycticeius humeralis*	common in western and central portion of the State	caves, hollow trees, buildings, barns, under bridges	PN
Gray Bat	*Myotis grisescens*	year round, probably most of state recorded in 49 counties	caves, cave like habitats; migrate between summer and winter caves	PN, FE, SE
Hoary Bat	*Lasiurus cinereus*	summer, uncommon, most of State	trees, under clusters of leaves except in winter when may roost in hollow trees, abandoned buildings	PN
Indiana Bat	*Myotis sodalis*	year round; central and eastern portions of State	Limestone caves; floodplain and riparian forests, other habitats are used	PN, FE, SE, CHD
Little Brown Bat	*Myotis lucifugus*	year round, eastern two-thirds of State; restricted to locales with caves in winter	caves, mines, underground quarries in winter; barns, buildings (especially attics) in summer	PN
Northern Long-eared Bat	*Myotis septentrionalis*	year round, eastern two-thirds of State	caves, under loose tree bark, rock shelters in cliff lines and abandoned mines, bridges and abandoned buildings	PN
Rafinesque's Big-eared Bat	*Corynorhinus rafinesquii*	year round, statewide, except northeastern portion of State	caves, old mines, protected rock shelters along cliff lines, abandoned buildings, bridges, cisterns, large hollow trees	PN
Townsend's Big-eared Bat	*Corynorhinus townsendii*	year round, rare, isolated colonies, unconfirmed but suspected presence within the State	caves, old mines where temperatures are extremely stable and above the required 54 degrees Fahrenheit	PN, FE

Common Name	Scientific Name	Occurrence	Roosting/Rearing/Hibernating Habitat	Status in TN*
Eastern Red Bat	*Lasiurus borealis*	year round (primarily winter), statewide	forests, beneath clusters of leaves, except winter, then hollow trees, fallen logs, and even leaf litter of forest floor	PN
Seminole Bat	*Lasiurus seminolus*	year round, southern one half of State	trees (beneath clusters of leaves), in clumps of Spanish moss	PN
Silver-haired Bat	*Lasionycteris noctivagans*	winter migrant, locally across State	hollow trees, tree cavities, crevices beneath peeling bark during the summer, rock fissures of cliff lines and cave entrances in winter	PN
Eastern Small-footed Bat	*Myotis leibii*	year round (mostly in winter), majority of the State, except upper third of middle portion of State	caves, rock shelters, cliff fissures, old mines, quarries, abandoned buildings, bridges	PN
Southeastern Bat	*Myotis austroriparius*	year round, distributed western portion of the State	caves, large hollow trees, abandoned buildings	PN
Brazilian Free-tailed Bat	*Tadarida brailiensis*	year round, distributed in central portion of the State	caves in the southwestern U.S., in the eastern U.S. exclusively found in attics	PN

*Codes: F = Federal listing, S = State listing, E = Endangered, PN = Protected Non-game, CHD = Critical Habitat Determined

Most requests for WS' operational assistance would likely occur in relation to bats inhabiting human-occupied buildings. Bat species that may be removed include the big brown bat, Eastern pipistrelle, evening bat, Brazilian free-tailed bat, and little brown bat. Those species of bats can be found in buildings and other man-made structures. Bat species that are listed by the USFWS pursuant to the ESA and by the TWRA are not generally associated with man-made structures and so it is unlikely that any federally or state listed bat species would be encountered by WS during activities to address bats. In most cases, a single bat found in a building would be provided an escape route (*e.g.*, opening a door or window) or would be live captured and released outside on site if there was no possibility of an exposure to people or pets. If the bat appeared sick, acted unusually, or if there was a known bite or possible exposure to people or pets, the bat would be euthanized and submitted for rabies testing. Those bats euthanized by WS for disease testing would likely be euthanized and submitted for testing by other entities in the absence of WS' involvement given the risk to human safety associated with exposure. Therefore, take by WS would not be additive to take that would likely occur in the absence of involvement by WS.

Based on previous requests for assistance and in anticipation of receiving requests for assistance in the future, it is possible that WS could kill up to five bats each year statewide, in any species combination, consisting of big brown bats, Eastern pipistrelles, evening bats, Brazilian free-tailed bats, and little brown bats. If the need arises, WS would consult with a qualified biologist to identify positively bats prior to removing them in order to eliminate any chance of addressing a T&E species. If a threatened or endangered bat were encountered, WS would contact the USFWS and/or the TWRA to determine the appropriate action. WS would continue to provide escape routes or live-capture and release bats in those

instances where no human or pet exposure could be assured. Based upon this information, WS is not likely to adversely affect any bat populations in the State.

Regionally, some bats species are being adversely impacted by the fungal disease white-nose syndrome, an emerging disease causing unprecedented morbidity and mortality among bats in eastern North America. The disease is characterized by cutaneous infection of hibernating bats by the psychrophilic fungus *Geomyces destructans*. However, WS' limited lethal take of bats would not adversely affect overall populations of bat species in the State. Impacts to bats would be minimal because any bat removal would be localized and limited in scope. In addition, euthanizing and submitting bats for testing would likely occur in the absence of WS' participation due to the risks to human safety.

Mole Population Information and Effects Analysis

Moles may be distinguished from voles or shrews with which they are often confused by noting certain characteristics. They have a hairless, pointed snout extending nearly 13 mm (0.5 inches) in front of the mouth opening. The small eyes and the openings of the ear canals are concealed in the fur and there are no external ears. The forefeet are very large and broad, with palms wider than they are long and the toes are webbed to the base of the claws, which are broad and depressed. The hind feet are small and narrow, with slender, sharp claws. In North America, there are seven species of moles. Three of those species occur east of the Rocky Mountains. All three of the species found east of the Rocky Mountains occur in Tennessee. Those species of moles found in Tennessee are the star-nosed mole, hairy-tailed mole, and Eastern mole.

The star-nosed mole measures between 132 to 230 mm (5.2 to 9.1 inches) and weighs between 40 to 85 g (1.4 to 3.0 ounces) (Smithsonian National Museum of Natural History 2012). The star-nosed mole has a rose-colored ring of fleshy, retractable tentacles surrounding its nose. This nasal disc is bilaterally symmetrical with 11 projections on each side. Equally distinct is the scaly, fleshy tail that is covered with concentric rings and short, coarse hairs. The tail, nearly as long as the combined length of the head and body, is constricted at the base, tapered at the tip, and during the winter swollen in size, when it serves as a fat storage organ. The eyes of this species are larger than those of the hairy-tailed mole, and its blackish-brown to black fur is longer, the metallic sheen absent. The limbs are short, and the front feet are paddle-like with long, stout claws. The surfaces of the pinkish-colored feet possess dark scales (Saunders 1988). Litter size ranges from three to seven, averaging five. The star-nosed mole ranges from southeastern Manitoba to Labrador and Nova Scotia, south and east to southeastern Georgia. The star-nosed mole prefers damp to saturated soils, and often lives in the organic muck adjacent to water, in grassy meadows, marshes, swamps, and deciduous, coniferous, and mixed forests. Because of their habitat preference, they are seldom a problem to people (Whitaker, Jr. and Hamilton, Jr. 1998). Population densities may range from 25 to 30 or more star-nosed moles per ha (10 to 12 per acre) (Saunders 1988).

The hairy-tailed mole measures between 151 to 173 mm (5.9 to 6.8 inches) and weighs between 41.0 to 62.8 g (1.45 to 2.22 ounces) (Smithsonian National Museum of Natural History 2012). Hairy-tailed mole fur is black or blackish-brown and has a purplish-brown sheen. The tail is short, less than 25% of total length, fleshy, slightly constricted at the base and as the name implies, covered with long coarse hairs (Hallett 1978). Litter size averages four to five (Eadie 1948, Conner 1960), but may be as high as eight (Richmond and Roslund 1949). They range throughout extreme southeastern Canada and New England, southwest through the mountains of North Carolina and Tennessee, and west through eastern Ohio. The hairy-tailed mole is found in mountainous and foothill habitat in extreme eastern Tennessee where it inhabits well-drained, light soil, brushy areas, and occasionally lawns or golf courses adjacent to woods. Soils, rather than plant communities, determine its distribution. Its occurrence coincides with the presence of dry to moist, but never wet, sandy loams with good surface cover and sufficient moisture

(Saunders 1988, Hallett 1978). Eadie (1939) found average population densities of three per ha with a maximum of 27 per ha in New Hampshire and in years of unusually high densities, 25 to 30 moles per ha (10 to 12 per acre) have been reported in New York (Hamilton, Jr. 1939).

The Eastern mole exhibits distinct sexual dimorphism, males are generally larger than females. Male Eastern moles measure 103 to 208 mm (4.06 to 8.19 inches); females measure 129 to 168 mm (5.08 to 6.61 inches). Male Eastern moles weigh 40 to 140 g (1.41 to 4.94 ounces); females weigh from 32 to 90 g (1.13 to 3.17 ounces) (Smithsonian National Museum of Natural History 2012). The Eastern mole is grayish brown, darker above and paler or browner beneath and when viewed from different angles, the hair often has a silvery sheen. There is an orange strip on the belly due to skin gland secretions that is usually brighter in males (Schwartz and Schwartz 2001). The fur is short, fine, and directionless, it can lay flat facing forward or backward, depending on whether the animal is moving forward or backward through a tunnel (Smithsonian National Museum of Natural History 2012). Gestation is from four to six weeks and litter size ranges from two to five, young leave the nest at four weeks of age (Yates and Schmidly 1978, Schwartz and Schwartz 2001). The Eastern mole has the largest range of any North American mole, occurring from Northeastern Mexico north to the upper Midwest and Southern Ontario, Canada and east from Florida to Southern New England where favorable soils are found (Yates and Schmidly 1978, Smithsonian National Museum of Natural History 2012). Eastern mole habitat includes meadows, pastures, lawns, open woodlands, gardens, and stream banks where the soil is loose, contains humus and is well drained but moist (Schwartz and Schwartz 2001). The Eastern mole is found throughout all of Tennessee where it inhabits open fields, waste areas, lawns, gardens, and sometimes forests in well-drained loose soils. Population densities ranging from 4.9 to 7.4 Eastern moles per ha (2 to 3 moles per acre) were reported by Schwartz and Schwartz (2001) as being high, while Hartman and Krentz (1993) reported 1.3 to 3.0 Eastern moles per ha (0.5 to 1.2 per acre) in South Carolina.

Moles eat several kinds of invertebrates including earthworms, grubs, beetles, beetle larvae, centipedes, ants, wasps, spiders, and flies, among others. They also eat seeds and some other plant materials. The most common mole damage are tunnels in gardens, lawns, golf courses and other grassy areas, resulting from their incessant search for food. They can eat between 70% and 100% of their body weight each day (Godfrey and Crowcroft 1960, Holbrook and Timm 1986, Henderson 1994). Moles can also store food as well (Henderson 1994).

Eastern moles live in the seclusion of underground burrows, coming to the surface only rarely, and then often by accident. Hairy-tailed moles stay in burrows by day but may emerge at night to feed (National Audubon Society 2000). Researchers believe moles are solitary. On several occasions, two or even three moles have been trapped at the same spot, but that does not necessarily mean they had been living together in a particular burrow. Networks of runways made independently occasionally join otherwise separate burrows (Godfrey and Crowcroft 1960, Henderson 1994).

During the mating season, male moles will seek out a female in her burrow to mate (National Audubon Society 2000). Moles have few natural enemies, which allows them to maintain populations by producing only one annual litter of two to six (National Audubon Society 200) or three to five offspring (Henderson 1994) each year. The gestation period of moles is approximately 42 days, with young being born mainly in March and early April (Henderson 1994). Home range estimates for moles range from 1,385 to 114,486 square feet (Yates and Pedersen 1982).

No population estimates are currently available for moles in Tennessee. Since moles target non-forested areas and if moles were only found in rural areas, the cumulative area of the most likely habitat classifications would be approximately 45,000 km^2 (11,000,000 acres) (see Table 4.1). If only 50% of those land classifications supported moles, with a density estimate of one mole per 2.6 acres, assuming

that only one mole occupies each home range, and home ranges do not overlap, a statewide mole population could be estimated at over 2 million moles.

From FY 2006 through FY 2011, WS responded to 40 requests for assistance associated with moles. Most requests were addressed through technical assistance by providing information on methods and mole damage. Those persons reported $9,900 in damages associated with moles from FY 2006 through FY 2011. Damages were reported to landscaping, turf and flowers, and property caused by the burrowing and digging behavior of moles. No moles were lethally taken by WS in Tennessee from FY 2006 through FY 2011.

WS anticipates continuing to address most requests for assistance associated with moles by providing people with technical assistance. However, WS could be requested to address moles directly through operational assistance. If WS was requested to provide operational assistance with managing damage associated with moles, WS anticipates that up to 100 moles could be lethally removed annually. Although three species of moles are present in Tennessee, most damage and the associated requests for assistance would involve the Eastern mole. However, damage can also occur from the activities of star-nosed moles and the hairy-tailed mole. Those species could also be lethally removed in limited numbers.

If mole populations totaled 2 million individuals statewide, the potential lethal take of up to 100 moles by WS annually would represent a small percentage of the population and would be of low magnitude. The number of moles lethally removed by other entities to alleviate damage in the State is not currently available. However, the cumulative take of moles is not expected to reach a level where adverse effects would occur to the populations of those mole species found in Tennessee.

Feral Swine Population Information and Effects Analysis

Feral swine, also known as "*wild pigs*", "*wild boars*," and "*feral hogs*", are medium-sized hoofed mammals that look like domestic pigs. They usually have coarser and denser coats than their domestic counterparts and exhibit modified canine teeth called "*tusks*" which are usually 7.5 to 12.5 cm (3 to 5 inches) long but may be up to 23 cm (9 inches) long. These tusks curl out and up along the sides of the mouth. Lower canines are also prominent but smaller. Young feral hogs have pale longitudinal stripes on the body until they are six weeks of age. Adults of the species average 90 cm (3 feet) in height and 1.32 to 1.82 m (4 feet 6 inches to 6 feet). Males may attain a weight of 75 to 200 kg (165 to 440 lbs), while females may weigh 35 to 150 kg (77 to 330 lbs). Feral swine mate any time of year but peak breeding times usually occur from January through February and again in early summer. Litter sizes are usually three to 12 piglets (National Audubon Society 2000). Given adequate nutrition, a wild pig population can double in just four months. Feral hogs may begin to breed before six months of age and sows can produce two litters per year (Barrett and Birmingham 1994). Feral swine are found in variable habitat in much of the southern United States, as well as most of the United States. Populations are usually clustered around areas with ample food and water supplies. Evidence of the presence of feral swine may be rooted up earth, tree rubs at ground level to 900 cm (36 inches) high, with clinging hair or mud, and muddy wallows.

Damage in areas supporting feral swine populations is sometimes a serious natural resource management concern for land managers. Substantial damage has occurred to natural resources, including destruction of fragile plant communities, killing tree seedlings, and erosion of soils (Barrett and Birmingham 1994). Food sources for feral hogs includes acorns, hickory nuts, pecans, beech nuts, and a wide variety of vegetation including roots, tubers, grasses, fruit, and berries, but feral hogs also eat crayfish, frogs, snakes, salamanders, mice, eggs and young of ground-nesting birds, young rabbits, and any other easy prey or carrion encountered. Feral swine have been known to kill and eat deer fawns (National Audubon Society 2000). They have also been reported to kill considerable numbers of domestic livestock,

especially young animals, in some areas (Barrett and Birmingham 1994). Several diseases are associated with feral swine populations (see Table 1.3).

In Tennessee, feral swine are classified as a non-protected nuisance species and may be taken at any time on private lands, with some restrictions on some public lands (TWRA 2012*a*). The management goal of the TWRA is to eradicate feral swine from Tennessee or keep the population at the lowest possible level (TWRA 2011*a*). The total feral swine population in Tennessee is unknown; however, reports of feral swine have been increasing. It is anticipated that feral swine populations in Tennessee will continue to increase due to their prolific breeding behavior, adaptability, and additional swine being illegally released into the wild. Given the unregulated status of feral swine in the State, the number of feral swine lethally removed annually is currently unknown.

Feral swine damage may be addressed by WS in response to requests by federal agencies, state agencies, or the public at any location in Tennessee. Agricultural producers may request assistance with managing damage to standing crops or disease threats to domestic livestock. Natural resource managers may request assistance to protect natural areas, parks or recreation areas, or T&E species. Public health agencies may request assistance in reducing feral swine densities where disease threats to humans may be present (see Table 1.3). WS may use any legal methods among those outlined by Barrett and Birmingham (1994) and West et al. (2009) as suitable for feral swine damage management, including the use of aircraft to shoot feral swine.

Between FY 2006 and FY 2011, WS responded to 167 requests for assistance associated with feral swine in Tennessee. Those persons requesting assistance reported $790,690 in damage to agricultural crops, landscaping, turf, and golf courses. Damages occur primarily from the rooting and wallowing behaviors of feral swine. From FY 2006 through FY 2011, WS removed a total of 260 feral swine to reduce damage and for disease surveillance in Tennessee. Removal of a small number of feral swine or a single individual will sometimes reduce damage considerably where natural resources, agriculture, or property is affected (Barrett and Birmingham 1994). However, damage may increase dramatically in areas where feral swine have ample resources and opportunity to expand.

Based on previous requests for assistance and the likely continued spread of feral swine in Tennessee, WS anticipates that up to 2,000 feral swine could be killed annually in the State to alleviate damage associated with requests for assistance and for disease surveillance. However, such population reduction is not expected to affect the overall statewide population feral swine because of the high reproductive rates exhibited by these animals (Barrett and Birmingham 1994). Damage management activities associated with feral swine would target single animals or local populations of feral swine at sites where their presence was causing unacceptable damage or threats to agriculture, human health and safety, natural resources, or property. Feral swine are not native to North America, including Tennessee. The National Invasive Species Council specifically lists feral swine as an invasive species pursuant to Executive Order 13112. Executive Order 13112 directs federal agencies to address invasive species to the extent practicable and permitted by law. In addition, activities requested of WS to reduce feral swine damage under this alternative would further the population objective set by TWRA of keeping the feral swine population at the lowest possible level in the State.

Based upon the above information, WS' limited lethal take of feral swine would not adversely affect overall feral swine populations in the State. Any damage management activities involving lethal methods by WS would be restricted to isolated individual sites. Some local populations may be temporarily reduced because of damage management activities aimed at reducing damage at a local site. Since feral swine are classified as a nuisance species in Tennessee, in those cases where feral swine are causing damage or are a nuisance and complete removal of the local population could be achieved, this could be

considered as providing some benefit to the natural environment since feral swine are not considered part of the native ecosystem.

White-tailed Deer Population Information and Effects Analysis

White-tailed deer are small to medium-sized mammals with tan or reddish brown pelts above in summer and grayish brown in winter. The belly, throat, noseband, eye-ring, and inside of the ears are white and their tail is brown with white above, often with a dark stripe down the center and white below. Deer are known for raising their tail while alarmed and in flight, called "*flagging*," in which the tail appears as a large, bright flash of white. This communicates danger to other deer and helps young follow their mothers in flight (National Audubon Society 2000). The range in size of white-tailed deer is extreme. White-tailed deer in the northern extremes of its range, where there is good habitat, will achieve weights of greater than 136 kg (300 lbs). By comparison, the tiny Florida Keys subspecies (*O.v. clavium*) commonly weighs less than 23 kg (50 lbs) (National Audubon Society 2000).

Male white-tailed deer are called bucks. They exhibit antlers, which are a pair of bony outgrowths of the frontal bone that normally are shed annually. The antlers begin growing in the early summer at which time they are covered with a skin that grows as the antlers do. The skin has short fine hairs called "*velvet*", containing a network of blood vessels, which nourish the growing bone beneath. By late summer, the antlers are fully developed, and the "*velvet*" is rubbed off against small saplings by the animal as the bone hardens. The antlers then serve as sexual ornaments and rival males may use them as weapons in courtship battles during the breeding season, which is called the "*rut*". After the mating season, the antlers decalcify and detach from the frontal bone within two to three days of each other, fall to the ground, and are often quickly found and gnawed on by various rodents for the calcium (National Audubon Society 2000). Antler size depends upon nutrition, age, and genetics (Craven and Hygnstrom 1994).

The white-tailed deer reproductive season varies according to geographic range. It may occur by the first two weeks in November in the north, but occurs as late as January or February in the south. Females, called "*does*", may have one to three young, or "*fawns*", after a gestation period of approximately 202 days (Craven and Hygnstrom 1994). A young doe bred for the first time will usually have only one fawn, older does two or three. The female remains near the fawns, returning to feed them only once or twice a day. Twin fawns are separated, which serves to protect them. Weaning occurs between 1 and 2 ½ months. Fawns stay with the mother until fall or winter, sometimes up to two years, but the doe generally drives off her young the previous year shortly before giving birth (National Audubon Society 2000).

When compared to other land mammals in North America, the white-tailed deer currently occupies the largest geographic range of any other mammal (Pagel et al. 1991). White-tailed deer range throughout most of the United States, except the far southwest, and inhabit the southern half of the southern tier of Canadian provinces. This species inhabits farmlands, brushy areas, forests, suburbs, and gardens. Rural areas containing a matrix of forest and agricultural crops can contain the highest deer densities (Roseberry and Woolf 1998). Biologists and resource managers in Tennessee have been challenged with managing escalating populations of deer in many urban/suburban areas and in some rural areas. As deer populations increase, there is an increasing occurrence of damage from white-tailed deer to agricultural crops (DeVault et al. 2007), increasing incidences of Lyme disease (Fernandez 2008), a rise in deer-vehicle collisions (Conover et al. 1995), and a disruption in forest health, regeneration, and forest dependent species (Tilghman 1989). Additionally, white-tailed deer are ranked as the second most hazardous species to aviation according to the percentage of strikes that caused damage from 1990 through 2010 (Dolbeer et al. 2012).

White-tailed deer are present in all Tennessee counties, and occupy almost all land types that contain suitable habitat. Due to restoration efforts, the white-tailed deer has increased from a low of 2,000 deer in the 1940s to approximately 900,000 deer 2005 (TWRA 2012*b*). The statewide deer population is expected to increase 1% to 2% annually, with the greatest increases observed in the Mississippi River valley and in eastern Tennessee (TWRA 2012*b*). The authority for management of resident wildlife species, including deer, is the responsibility of the TWRA. The TWRA collects and compiles information on white-tailed deer population trends and harvest and uses this information to manage deer populations. The primary tool for the management of deer populations in Tennessee is through adjusting the allowed lethal take during the deer harvest season in the State. White-tailed deer are classified as a big-game species in Tennessee with annual hunting seasons. During the 2009-2010 hunting season, the TWRA reported that 156,590 deer were harvested (TWRA 2010). The number of deer allowed to be harvested by individual hunters during the length of the deer hunting season varies across the State. However, during the development of this assessment, up to three female deer could be harvested daily with a take limit of one antlered deer per day (three total during the length of the season) allowed over much of the central and western portions of the State as part of efforts to decrease or stabilize deer populations (TWRA 2012*a*, TWRA 2012*b*).

Mortality can also occur from vehicle collisions, dogs, illegal take, tangling in fences, disease, and other causes (Crum 2003). Annual deer mortality in Tennessee from other sources (*e.g.*, illegal take, disease, and predation) is currently unknown. From July 1, 2010 through June 30, 2011, State Farm Mutual Automobile Insurance Company (2011*b*) estimated 20,039 deer-vehicle collisions occurred in Tennessee.

From FY 2006 through FY 2011, WS responded to 26 requests for assistance associated with white-tailed deer in the State. Most requests for assistance were addressed by providing technical assistance. Those persons requesting assistance reported $63,500 in damages, primarily from damage to aircraft, gardens, golf courses, cotton crops, trees, and shrubs. Between FY 2006 and FY 2011, WS has lethally removed one deer to alleviate damage and used non-lethal harassment methods to disperse three deer to alleviate damage in the State.

After review of previous activities conducted by WS and in anticipation of addressing requests for lethal take, WS' future lethal take could reach 1,000 deer annually. In addition, WS may be requested by the TWRA and/or the TDA to assist with sampling and managing the spread of diseases found in free-ranging and/or captive deer populations. If a disease outbreak occurred, WS could be requested to lethally take white-tailed deer for sampling and/or to prevent further spread of diseases. However, WS' total annual take would not exceed 1,000 deer annually under the proposed action.

If requested, WS could also assist with sampling and removing deer from captive facilities where deer were confined inside a perimeter fence. The detection of a disease at a captive facility often raises concerns for the potential spread of diseases to free-ranging herds. The spread of diseases among deer inside those facilities is often increased due to their close contact with one another. Often, once a disease is detected in a confined deer herd, the entire herd is destroyed to ensure the containment of the disease. Any involvement with the depopulation of deer confined inside a perimeter fence by WS would be at the request of the TWRA and/or the TDA. As proposed in this alternative, in those cases where WS was requested to assist with the removal of a captive deer herd in Tennessee, the take would not exceed 1,000 deer for purposes of disease monitoring or surveillance. Deer confined inside perimeter fences for the purposes of non-traditional farming, including confined for hunting, are not included in statewide deer population estimates. However, since take of deer by WS for disease surveillance or monitoring could occur in free-ranging or captive herds, the potential take of up to 1,000 deer for disease surveillance and monitoring by WS would be considered as part of the impact analysis on the statewide free-ranging deer population.

From 2006 through 2009, 648,542 deer were harvested in Tennessee during the annual hunting season, with the highest harvest level occurring in 2006 when 174,937 deer were harvested. The lowest harvest level of deer between 2006 and 2009 occurred in 2009 when 156,590 deer were harvested (TWRA 2010). If WS' take reached 1,000 deer during the highest harvest of deer in the State that occurred in 2006, WS' take of 1,000 deer would have represented 0.6% of the harvest. If WS' take reached 1,000 deer during the lowest harvest total of deer in the State that occurred in 2009, WS' take of 1,000 deer would have represented 0.6% of the total harvest.

As stated previously, the deer population in Tennessee was estimated at 900,000 deer during 2005 (TWRA 2012*b*). The total deer mortality in the State in 2009 could be estimated at 176,629 deer, based on harvest and vehicle collision data. If the deer population estimate provided by the TWRA included recruitment of deer born that year, then the take of deer from all known sources in 2009 would represent 19.6% of the deer population, if the deer population remained at least stable. If WS had taken 1,000 deer in 2009, the total mortality of deer would have been estimated at 177,629 deer. When combined with the total known mortality in the State during 2009, WS' take of up to 1,000 deer would have represented 19.7% of the population. If WS had lethally removed 1,000 deer in 2009, WS' take would have represented an increase of 0.1% when compared to the total mortality in 2009 if no take by WS had occurred (*i.e.*, 19.6% without take by WS compared to 19.7% if WS' take had been 1,000 deer in 2009).

With oversight of the TWRA, the magnitude of take of deer by WS annually to resolve damage and threats would be low. TWRA has determined that there is no evidence to suggest that human mediated mortality resulting from regulated harvest and damage management, including removal by WS, would be detrimental to the survival of the white-tailed deer population in the State of Tennessee (G. Anderson, TWRA, pers. comm. 2011).

GonaCon™ was officially registered by the EPA in 2009 for use in reducing fertility in female white-tailed deer. According to the label, only WS or state wildlife management agency personnel or individuals working under their authority can use the reproductive inhibitor. Additionally, in order for GonaCon™ to be used in any given state, the product must also be registered with the state and approved for use by the appropriate state agency responsible for managing wildlife. The reproductive inhibitor Gonacon™ is currently not registered for use in Tennessee. However, if Gonacon™ becomes available to manage deer in the State, the use of the inhibitor could be evaluated under the proposed action as a method available that could be used in an integrated approach to managing damage.

Population management from the use of reproductive inhibitors to induce a decline in a localized deer population occurs through a reduction in the recruitment of fawns into the population by limiting reproductive output of adults. A reduction in the population occurs when the number of deer being recruited into the population cannot replace those individuals that die from other causes each year, which equates to a net loss in the number of individuals in the population and a reduction in the overall population. Although not generally considered a lethal method since no direct take occurs, reproductive inhibitors can result in the reduction of a target species' population. WS' use of GonaCon™ would target a local deer population identified as causing damage or threatening human safety. Although a reduction in a local deer population would likely occur from constant use of GonaCon™, the actual reduction in the local population annually would be difficult to derive prior to the initiation of the use of the vaccine.

One of the difficulties in calculating and analyzing any actual reduction that could occur from the use of the vaccine in a targeted population prior to application of the vaccine is the variability in the response of deer to the vaccine. Previous studies on GonaCon™ as a reproductive inhibitor have shown variability in the immune response of deer to the vaccine (Miller et al. 2000). Not all deer injected with GonaCon™ develop sufficient antibodies to neutralize the GnRH produced in the body. Those deer continue to enter into a reproductive state and produce fawns even after vaccination. The number of deer that do not

develop sufficient antibodies after the initial vaccination cannot be predicted beforehand. In one study, 88% of the deer vaccinated with GonaCon™ did not produce fawns the following reproductive season while 12% of the deer injected with GonaCon™ produced fawns (Gionfriddo et al. 2009). The year following the initial vaccination, the number of deer that were vaccinated the first year that did not produce fawns declined to 47% while the number of deer producing fawns increased to 53% (Gionfriddo et al. 2009) demonstrating the diminishing results that are likely over time if deer are not provided a booster shot periodically.

Since the effects of GonaCon™ are reversible if deer are not provided with a booster shot periodically, the reduction in a local population of deer from the use of GonaCon™ can be maintained at appropriate levels where damages or threats are resolved by increasing or decreasing the number of deer receiving booster injections. Although localized deer populations would likely be reduced from the use of GonaCon™, the extent of the reduction would be variable. For example, not all vaccinated deer are prevented from entering into a reproductive state and those deer that were initially prevented from entering into a reproductive state often become reproductively active in subsequent years as the antibody levels neutralizing the GnRH hormone diminish over time. Therefore, the actual decline in the number of deer in a localized population achieved from the use of GonaCon™ would be difficult to predict prior to the use of the reproductive inhibitor. However, since the decline would occur through attrition over time and since the ability of the inhibitor to prevent reproduction diminishes with time, the actual decline in a localized population would be gradual and could be monitored. In addition, the reduction in a local deer population could be fully reversed if deer are no longer vaccinated or provided booster shots and other conditions (*e.g.*, food, disease) are favorable for population growth.

Turner et al. (1993) noted that although contraception in white-tailed deer may be used to limit population growth, it would not reduce the number of deer in excess of the desired level in many circumstances. Turner et al. (1993) further contend that initial population reductions by various other means may be necessary to achieve management goals, and that reproduction control would be one facet of an integrated program. Although immunocontraceptive technology has been effective in laboratories, pens, and in island field applications, it has not been effective in reducing populations of free-ranging white-tailed deer over large geographical areas.

The magnitude of WS' activities to alleviate damage and threats associated with deer in the State would be low with the oversight and permitting of WS' activities occurring by the TWRA. If take by WS had reached 1,000 deer during 2009 when the lowest known deer harvest occurred in the State, WS' take would have represented 0.6% of the statewide harvest. In 2009, if WS' take had reached 1,000 deer, the total known mortality would have increased only 0.1% when compared to total known mortality if 1,000 deer had not be taken by WS. Based on the 2005 deer population estimate, take of up to 1,000 deer by WS would have represented 0.1% of the estimated population. WS would annually report to the TWRA and monitor take to ensure WS' activities do not adversely affect deer. The permitting of all WS' take by the TWRA would ensure WS' take would meet the objectives of the statewide wildlife management plan.

Elk Population Information and Effects Analysis

Elk are very large cervids with thick necks and slender legs. They are brown or tan above with darker underparts and a rump patch and tail of yellowish brown. Males, or "*bulls*", have dark brown manes on the throat and large many-tined antlers. They have a huge ponderous muzzle and a dewlap. These animals breed in late August-November and a single calf is born after a gestation period of approximately 250 days (deCalesta and Witmer 1994). They live in variable habitats and in mountainous areas, use high open mountain pastures in summer and lower wooded slopes in winter. Elk are very gregarious with the species main social unit being a herd of cows and calves. The size of these herds varies greatly and is sometimes composed of up to 400 individuals, with larger herds occurring in open areas. Bulls herd

separately, remaining on the outskirts of cow-dominated herds. During rutting season, adult bulls join cow/calf herds, and may assemble a cow harem of females during breeding activities (National Audubon Society 2000).

Elk range from eastern British Columbia, central Alberta, central Saskatchewan, and southern Manitoba in Canada south to central New Mexico and Arizona, with great numbers in Washington, Montana, Wyoming, and Colorado. They are also found along the west coast from Vancouver Island to northern California. Isolated populations are found elsewhere in California, Nevada, Utah, Arizona, New Mexico, Oklahoma, South Dakota, Minnesota, and Michigan. There are smaller numbers in several eastern states (National Audubon Society 2000).

Before colonists settled in Tennessee, eastern elk were fairly common in the State. Mixed forests and grasslands provided adequate habitat for this large cervid. However, by the middle of the 19th century, elk had disappeared because of unregulated hunting, encroaching civilization, and loss of habitat.

Elk were recently reintroduced into two distinct areas of Tennessee after being gone for some 150 years. In 2000, the TWRA began releasing elk in the Royal Blue Wildlife Management Area and the Sundquist Wildlife Management Areas. These areas cover portions of Scott, Morgan, Campbell, Anderson, and Claiborne counties. Over the next eight years, the TWRA released 201 elk in these areas (TWRA 2011*b*). Almost simultaneously, the Great Smoky Mountains National Park (GSMNP) released an experimental herd of 52 elk within their boundaries. The park straddles the state line between Tennessee and North Carolina and includes portions of Blount, Sevier, and Cocke counties. In both cases, elk that wander outside of the elk restoration zones are captured and moved back into the restoration zone, if possible or are destroyed if capture is not possible. The current population estimate in the Royal Blue/Sundquist Wildlife Management areas is about 300 animals and the GSMNP estimates 125 elk in its herd (TWRA 2011*b*, GSMNP 2011). Tennessee now has a limited hunting season for elk in the State (TWRA 2012*a*).

WS has not previously been requested to provide assistance with elk or elk damage in the State. No lethal take of elk has occurred by WS. Most individuals seeking assistance with elk that have wandered off the elk restoration areas likely contact the TWRA and/or the National Park Service. However, WS could be requested to provide assistance with the live-capture and translocation of elk and/or to lethally remove elk in the State. Most activities conducted by WS would involve the live-capture and translocation of elk back to restoration areas. All activities associated with elk would be coordinated with the TWRA and/or the National Park Service. Requests for assistance would likely be received from the TWRA or the National Park Service. Most requests for assistance received by WS from private entities would be referred to the TWRA and/or the National Park Service. However, in the event WS was requested to lethally remove elk, WS anticipates that up to five elk could be lethally removed annually in the State. The limited lethal take of elk proposed and the permitting of the take by the TWRA and/or the National Park Service would ensure elk populations remain viable and productive.

Additional Target Species

Target species, in addition to the mammal species analyzed previously, have been lethally taken in small numbers by WS and are typically feral animals. For example, a feral rabbit was lethally taken by WS in Tennessee during the implementation of a wildlife hazard program at an airport in the State. However, additional species including feral burros, feral cattle, feral goats, feral horses, fallow deer, and other non-native mammals may, on occasion, cause damage that WS could be requested to assist with. While WS does not currently expect to lethally take any of those species, WS could be requested by the TWRA and/or the TDA to assist with unique situations where a small number of those mammals have escaped or were released. Those occasions could include the accidental release of feral animals onto airport properties or animals that have escape from fenced enclosures. In addition, WS could be requested to

assist the TWRA or TDA as part of an incident response, such as the accidental release of domestic or exotic mammals from vehicle wrecks. There may also be additional need for taking other mammal species in the event of an animal disease outbreak to limit the spread of the disease. As part of the proposed program, WS could provide assistance, upon request, involving exotic and domestic mammals not specifically listed in this EA in emergency situations to alleviate threats to human health and safety. Any take requested would target specific individual mammals and take would not reach a magnitude where adverse effects would occur to a species' population based on the limited scope of the removal. In most cases, the take would be limited to a few individuals and removal would likely occur by other entities in the absence of WS' involvement.

Wildlife Disease Surveillance and Monitoring

The ability to efficiently conduct surveillance for and detect diseases is dependent upon rapid detection of the pathogen if it is introduced. Effective implementation of a surveillance system would facilitate planning and execution at regional and state levels, and coordination of surveillance data for risk assessment. It would also facilitate partnerships between public and private interests, including efforts by federal, state, and local governments as well as non-governmental organizations, universities, and other interest groups.

Sampling strategies that could be employed include sampling live-captured mammals that could be released on site after sampling occurs. This sampling (*e.g.*, drawing blood, swabbing nasal cavities, collecting fecal samples) and the subsequent release of live-captured mammals would not result in adverse effects since those mammals are released unharmed on site. In addition, sampling of sick, dying, or hunter harvested mammals would not result in the additive lethal take of mammals that would not have already occurred in the absence of a disease sampling program. Therefore, the sampling of mammals for diseases would not adversely affect the populations of any of the mammals addressed in this EA nor would result in any take of mammals that would not have already occurred in the absence of disease sampling (*e.g.*, hunter harvest).

Alternative 2 – Mammal Damage Management by WS through Technical Assistance Only

Mammal populations in the State would not be directly impacted by WS from a program implementing technical assistance only. However, persons experiencing damage or threats from mammals may implement methods based on WS' recommendations. Under a technical assistance only alternative, WS would recommend and demonstrate for use both non-lethal and lethal methods legally available for use to resolve mammal damage. Methods and techniques recommended would be based on WS' Decision Model using information provided from the requestor or from a site visit. Requestors may implement WS' recommendations, implement other actions, seek assistance from other entities, or take no action. However, those persons requesting assistance are likely people that would implement damage abatement methods in the absence of WS' recommendations.

Under a technical assistance only alternative, those persons experiencing threats or damage associated with mammals in the State could lethally take mammals despite WS' lack of direct involvement in the management action. Therefore, under this alternative the number of mammals lethally taken would likely be similar to the other alternatives since take could occur through the issuance of a permit by the TWRA, take of non-native mammal species can occur without the need for a permit from the TWRA, and take would continue to occur during the harvest season for those species. WS' participation in a management action would not be additive to an action that would occur in the absence of WS' participation.

With the oversight of the TWRA, it is unlikely that mammal populations would be adversely impacted by implementation of this alternative. Under this alternative, WS would not be directly involved with

damage management actions and therefore, direct operational assistance could be provided by other entities, such as the TWRA, private entities, and/or municipal authorities. If direct operational assistance was not available from WS or other entities, it is hypothetically possible that frustration caused by the inability to reduce damage and associated losses could lead to illegal take, which could lead to real but unknown effects on other wildlife populations. People have resorted to the illegal use of chemicals and methods to resolve wildlife damage issues (White et al. 1989, USFWS 2001, FDA 2003).

Alternative 3 – No Mammal Damage Management Conducted by WS

Under this alternative, WS would not conduct damage management activities in the State. WS would have no direct involvement with any aspect of addressing damage caused by mammals and would provide no technical assistance. No take of mammals by WS would occur under this alternative. Mammals could continue to be lethally taken to resolve damage and/or threats occurring either through permits issued by the TWRA, during the regulated hunting or trapping seasons, or in the case of non-native species, take can occur anytime using legally available methods. Management actions taken by non-federal entities would be considered the *environmental status quo*.

Local mammal populations could decline, stay the same, or increase depending on actions taken by those persons experiencing mammal damage. Some resource/property owners may take illegal, unsafe, or environmentally harmful action against local populations of mammals out of frustration or ignorance. While WS would provide no assistance under this alternative, other individuals or entities could conduct lethal damage management resulting in lethal take levels similar to the proposed action.

Since mammals could still be taken under this alternative, the potential effects on the populations of those mammal species in the State would be similar to the other alternatives for this issue. WS' involvement would not be additive to take that could occur since the cooperator requesting WS' assistance could conduct mammal damage management activities without WS' direct involvement. Therefore, any actions to resolve damage or reduce threats associated with mammals could occur by other entities despite WS' lack of involvement under this alternative.

Issue 2 - Effects of Mammal Damage Management Activities on Non-target Wildlife Species Populations, Including T&E Species

As discussed previously, a concern is often raised about the potential impacts to non-target species, including T&E species, from the use of methods to resolve damage caused by mammals. The potential effects on the populations of non-target wildlife species, including T&E species, are analyzed below.

Alternative 1 - Continue the Current Adaptive Integrated Mammal Damage Management Program (No Action/Proposed Action)

The potential for adverse effects to non-targets occurs from the employment of methods to address mammal damage. Under the proposed action, WS could provide both technical assistance and direct operational assistance to those persons requesting assistance. The risks to non-targets from the use of non-lethal methods as part of an integrated direct operational assistance program would be similar to those risks to non-targets discussed in the other alternatives.

Personnel from WS would be experienced with managing wildlife damage and would be trained in the employment of methods, which would allow WS' employees to use the WS Decision Model to select the most appropriate methods for taking targeted animals and excluding non-target species. To reduce the likelihood of capturing non-target wildlife, WS would employ the most selective methods for the target species, would employ the use of attractants that are as specific to target species as possible, and

determine placement of methods to avoid exposure to non-targets. SOPs to prevent and reduce any potential adverse impacts on non-targets are discussed in Chapter 3 of this EA. Despite the best efforts to minimize non-target exposure to methods during program activities, the potential for WS to disperse or lethally take non-targets exists when applying both non-lethal and lethal methods to manage damage or reduce threats to safety.

Non-lethal methods have the potential to cause adverse effects to non-targets primarily through exclusion, harassment, and dispersal. Any exclusionary device erected to prevent access of target species also potentially excludes species that were not the primary reason the exclusion was erected; therefore, non-target species excluded from areas may potentially be adversely impacted if the area excluded was large enough. The use of auditory and visual dispersal methods used to reduce damage or threats caused by mammals would also likely disperse non-targets in the immediate area the methods were employed. Therefore, non-targets may be permanently dispersed from an area while employing non-lethal dispersal techniques. However, like target species, the potential impacts on non-target species would expect to be temporary with target and non-target species often returning after the cessation of dispersal methods.

Non-lethal methods that use auditory and visual stimuli to reduce or prevent damage are intended to elicit fright responses in wildlife. When employing those methods to disperse or harass target species, any non-targets near those methods when employed would also likely be dispersed from the area. Similarly, any exclusionary device constructed to prevent access by target species would also exclude access to non-target species. The persistent use of non-lethal methods would likely result in the dispersal or abandonment of those areas where non-lethal methods were employed of both target and non-target species. Therefore, any use of non-lethal methods would be similar to those results on both non-target and target species. Though non-lethal methods do not result in lethal take of non-targets, the use of non-lethal methods can restrict or prevent access of non-targets to beneficial resources. However, non-lethal methods would not be employed over large geographical areas or applied at such intensity that essential resources (*e.g.*, food sources, habitat) would be unavailable for extended durations or over a wide geographical scope that long-term adverse effects would occur to a species' population. Non-lethal methods would generally be regarded as having minimal impacts on overall populations of wildlife since individuals of those species were unharmed. Overall, potential impacts to non-targets from the use of non-lethal methods would not adversely affect populations since those methods are often temporary.

Other non-lethal methods available for use under this alternative include live traps, nets, and repellents. Live traps and nets are considered live-capture methods since wildlife are restrained once captured. Live traps would have the potential to capture non-target species. Trap and net placement in areas where target species were active and the use of target-specific attractants would likely minimize the capture of non-targets. If traps and nets were attended to appropriately, any non-targets captured could be released on site unharmed.

Chemical repellents would also be available to reduce mammal damage. Since FY 2006, WS has not used repellents to reduce mammal damage in the State. However, WS may recommend commercially available repellents when providing technical assistance. Only those repellents registered with the EPA pursuant to the FIFRA and the TDA would be recommended or used by WS under this alternative. The active ingredients in many commercially available repellents are naturally occurring substances (*e.g.*, capsaicin, fish oil, whole egg solids), which are often used in food preparation (EPA 2001). When used according to label instructions, most repellents would be regarded as safe since 1) they are not toxic to animals, if ingested; 2) there is normally little to no contact between animals and the active ingredient, and 3) the active ingredients are found in the environment and degrade quickly (EPA 2001). Only those repellents registered with the EPA pursuant to the FIFRA, and the TDA would be recommended and used by WS under this alternative. Therefore, the use and recommendation of repellents would not have

negative impacts on non-target species when used according to label requirements. Most repellents for mammals pose a very low risk to non-targets when exposed to or when ingested.

Exposure of non-target wildlife to GonaCon™ would occur primarily from secondary hazards associated with wildlife consuming deer that have been injected with GonaCon™. Since GonaCon™ would be applied directly to deer through hand injection after the animal was live-captured and restrained, the risk of directly exposing non-target wildlife to GonaCon™ while being administered to deer would be nearly non-existent. Several factors inherent with GonaCon™ reduce risks to non-target wildlife from direct consumption of deer injected with the vaccine (EPA 2009). The vaccine itself and the antibodies produced by the deer in response to the vaccine are both proteins, which if consumed would be broken down by stomach acids and enzymes (EPA 2009, USDA 2010). The EPA determined that the potential risks to non-target wildlife from the vaccine and the antibodies produced by deer in response to the vaccine "*...are not expected to exceed the Agency's concern levels*" (EPA 2009).

Potential impacts to non-targets from the use of non-lethal methods would be similar to the use of non-lethal methods under any of the alternatives. Non-targets would generally be unharmed from the use of non-lethal methods under any of the alternatives since no lethal take would occur. Non-lethal methods would be available under all the alternatives analyzed; however, the use of Gonacon™ would be restricted to use by the TWRA only under Alternative 2, if registered. WS' involvement in the use of or recommendation of non-lethal methods would ensure the potential impacts to non-targets were considered under WS' Decision Model. Non-lethal methods would not be employed over large geographical areas or applied at such intensity that essential resources (*e.g*., food sources, habitat) would be unavailable for extended durations or over a wide geographical scope that long-term adverse effects would occur to a species' population. Non-lethal methods are generally regarded as having minimal impacts on overall populations of wildlife since individuals of those species are unharmed. Overall, potential impacts to non-targets from the use of non-lethal methods would not adversely affect populations since those methods are often temporary and do not result in lethal take. Potential impacts to non-targets under this alternative from the use of and/or the recommendation of non-lethal methods are likely to be low.

WS could also employ and/or recommend lethal methods under the proposed action alternative to alleviate damage, when those methods were deemed appropriate for use using the WS Decision Model. Lethal methods available for use to manage damage caused by mammals under this alternative would include the recommendation of take during hunting and/or trapping seasons, shooting, body-gripping traps, cable restraints, fumigants, rodenticides, euthanasia chemicals, and euthanasia after live-capture. Available methods and the application of those methods to resolve mammal damage is further discussed in Appendix B.

The use of firearms would essentially be selective for target species since animals would be identified prior to application; therefore, no adverse impacts would be anticipated from use of this method. Similarly, the use of euthanasia methods would not result in non-target take since identification would occur prior to euthanizing an animal.

When using fumigants, burrows and dens would be observed for the presence of non-targets before the use of fumigants. If non-target activity (*e.g*., tracks, scat) were observed, the fumigation of those burrows or dens would not occur. Since non-targets are known to occur in burrows or dens, some risks of unintentional take of non-targets does exist from the use of fumigants. For example, burrows of woodchucks can be used by a variety of non-target species such as the Eastern cottontail, striped skunk, raccoon, red fox, coyote, white-footed mouse, house mouse, and short-tailed shrew (*Blarina brevicauda*) (Hamilton 1934, Grizzell 1955, Dolbeer et al. 1991).

Fumigants would be used in active burrows or dens only, which would minimize risk to non-targets. Dolbeer et al. (1991) found a total of one cottontail rabbit and three mice (*Permyscus* spp.) in three of the 97 woodchuck burrows treated with gas cartridges during the late summer. During 2,064 trap nights at 86 woodchuck burrow entrances targeting small mammals, Swihart and Picone (1995) captured 99 individuals of four small mammal species, which included short-tailed shrews, meadow voles, meadow jumping mouse (*Zapus hudsonius*), and white-footed mice. Risks to non-targets can be minimized by treating only burrows that appear to be active (Dolbeer et al. 1991). There are no secondary poisoning risks involved with the use of gas cartridges as the gas produced dissipates into the atmosphere shortly after activation. Primary risks to non-targets would be minimized by treating only active burrows or dens, by covering entrances of burrows or dens, and by following the pesticide label. Although non-targets could be present in burrows or dens, even after WS' conducts site investigations, the risks would be relatively low and unintentional take from the use of fumigants would be limited.

Zinc phosphide is a toxicant used to kill rodents, lagomorphs, and nutria. Zinc phosphide is two to 15 times more toxic to rodents than to carnivores (Hill and Carpenter 1982). Secondary risks appear to be minimal to predators and scavengers that scavenge carcasses of animals killed with zinc phosphide (Tietjen 1976, Hegdal and Gatz 1977, Hegdal et al. 1980, Hill and Carpenter 1982, Johnson and Fagerstone 1994). Risks would be minimal based on: 1) 90% of the zinc phosphide ingested by rodents is detoxified in the digestive tract (Hegdal et al. 1980), 2) 99% of the zinc phosphide residues occur in the digestive tracts, with none occurring in the muscle, 3) the amount of zinc phosphide required to kill target rodents is not enough to kill most other predatory animals that consume tissue (Johnson and Fagerstone 1994).

In addition, zinc phosphide has a strong emetic action (*i.e.*, causes vomiting) and most non-target animals in research tests regurgitated bait or tissues contaminated with zinc phosphide without succumbing to the toxicant (Hegdal and Gatz 1977, Hegdal et al. 1980, Johnson and Fagerstone 1994). Furthermore, predators tend to eviscerate zinc phosphide-poisoned rodents before eating them or otherwise avoid the digestive tract and generally do not eat the stomach and intestines (Hegdal et al. 1980, Johnson and Fagerstone 1994). Although zinc phosphide baits have a strong, pungent, phosphorous-like odor (garlic like), this characteristic seems to attract rodents, particularly rats, and apparently makes the bait unattractive to some other animals. Many birds appear capable of distinguishing treated from untreated baits and they prefer untreated grain when given a choice (Siegfried 1968, Johnson and Fagerstone 1994). Birds appear particularly susceptible to the emetic effects of zinc phosphide, which would tend to offer an extra degree of protection against bird species dying from the consumption of grain treated with zinc phosphide or, for scavenging bird species, from eating poisoned rodents. Use of rolled oats instead of whole grain also appears to reduce bird acceptance of bait. Uresk et al. (1988) reported on the effects of zinc phosphide on six non-target rodent populations. Uresk et al. (1988) determined that no differences were observed from pretreatment until after treatment in populations of eastern cottontail rabbits and white-tailed jackrabbits (*Lepus townsendii*). However, primary consumption of bait by non-target wildlife could occur and potentially cause mortality. Uresk et al. (1988) reported a 79% reduction in deer mouse populations in areas treated with zinc phosphide; however, the effect was not statistically significant because of high variability in densities and the reduction was not long-term (Deisch et al. 1990).

Ramey et al. (2000) reported that five weeks after treatment, no ring-necked pheasants (*Phasianus colchicus*) had been killed because of zinc phosphide baiting. In addition, Hegdal and Gatz (1977) determined that zinc phosphide did not affect non-target populations and more radio-tracked animals were killed by predators than died from zinc phosphide intoxication (Hegdal and Gatz 1977, Ramey et al. 2000). Tietjen (1976) observed horned larks (*Eremophila alpestris*) and mourning doves (*Zenaida macroura*) on zinc phosphide-treated prairie dog colonies, but observations after treatment did not locate any sick or dead birds, a finding similar to Apa et al. (1991). Uresk et al. (1988) reported that ground-

feeding birds showed no difference in numbers between control and treated sites. Apa et al. (1991) further states that zinc phosphide was not consumed by horned larks because: 1) poisoned grain remaining for their consumption was low (*i.e.*, bait was accepted by prairie dogs before larks could consume it), 2) birds have an aversion to black-colored foods, and 3) birds have a negative sensory response to zinc phosphide.

Reduced impacts on birds have also been reported by Tietjen and Matschke (1982). Deisch et al. (1989) reported on the effect zinc phosphide has on invertebrates. Deisch et al. (1989) determined that zinc phosphide bait reduced ant densities; however, spider mites, crickets, wolf spiders, ground beetles, darkling beetles and dung beetles were not affected. Wolf spiders and ground beetles showed increases after one year on zinc phosphide treated areas (Deisch 1986). Generally, direct long-term impacts from rodenticide treatments were minimal for the population of insects that were sampled (Deisch et al. 1989). Long-term effects were not directly related to rodenticides, but more to habitat changes (Deisch 1986) as vegetative cover and prey diversity increased without prairie dogs grazing and clipping the vegetation (Deisch et al. 1989). In addition, zinc phosphide treated baits would be placed underground or used in bait stations. The application of baits below ground or in bait stations would limit the direct exposure risks by most non-target species.

Use of zinc phosphide on various types of fruit, vegetable, or cereal baits (*e.g.*, apples, carrots, sweet potatoes, oats, barley) has proven to be effective at suppressing target wildlife populations. All chemicals that could be used by WS would be registered under the FIFRA and administered by EPA and TDA. Specific bait applications would be designed to minimize non-target hazards (Evans 1970). WS' personnel that use chemical methods would be certified as pesticide applicators by the TDA and would be required to adhere to all certification requirements set forth in FIFRA and the Tennessee pesticide control laws and regulations. No chemicals would be used on federal or private lands without authorization from the land management agency or property owner/manager.

Anticoagulant rodent baits with warfarin, brodifacoum, or diphacinone as active ingredients could be used in bait stations to target small rodents. WS could utilize locking bait stations to restrict access of non-target species to rodenticides such as anticoagulants. The use and proper placement of bait stations would minimize the likelihood that the bait would be consumed by non-target species. There may be secondary hazards from anticoagulant baits. Those risks are reduced somewhat by the fact that the predator scavenger species would usually need exposure to multiple carcasses over a period of days. Areas where anticoagulants could be used would be monitored and carcasses picked up and disposed of in accordance with label directions.

An issue that has arisen is the potential for low-level flights associated with using firearms from aircraft could potentially disturb wildlife, including T&E species. Aerial operations would be an important method of damage management in Tennessee when used to address damage or threats associated with feral swine and/or coyotes in remote areas where access is limited due to terrain and habitat. Aerial operations would only occur in those areas where a cooperative service agreement allowing the use of aircraft had been signed between WS and the cooperating landowner or manager. Aerial operations would typically be conducted with aircraft between the months of December and April when the foliage has fallen; however, aircraft could be used at any time of year. The amount of time spent conducting aerial operations varies depending on the severity of damage, the size of the area where damage or threats were occurring, and the weather, as low-level aerial activities would be restricted to visual flight rules and would be impractical in high winds or at times when animals were not easily visible.

Aircraft play an important role in the management of various wildlife species for many agencies. Resource management agencies rely on low flying aircraft to monitor the status of many animal populations including large mammals (Lancia et al. 2000), birds of prey (Fuller and Mosher 1987),

waterfowl (Bellrose 1976), and colonial waterbirds (Speich 1986). Low-level flights are also required when aircraft are used to track animal movements by radio telemetry (Gilmer et al. 1981, Samuel and Fuller 1994).

A number of studies have looked at responses of various wildlife species to aircraft overflights. The National Park Service (1995) reviewed the effects of aircraft overflights on wildlife and suggested that adverse effects could occur to certain species. Some species will frequently or at least occasionally show an adverse response to even minor overflights. In general though, it appears that the more serious potential adverse effects occur when overflights are chronic (*i.e.*, they occur daily or more often over long periods). Chronic exposures generally involve areas near commercial airports and military flight training facilities. Aerial operations conducted by WS rarely occur in the same areas on a daily basis and little time is actually spent flying over those particular areas.

The effects on wildlife from military-type aircraft have been studied extensively (Air National Guard 1997), and were found to have no expected adverse effects on wildlife. Examples of species or species groups that have been studied with regard to the issue of aircraft-generated disturbance are as follows:

***Waterbirds and Waterfowl*:** Low-level overflights of two to three minutes in duration by a fixed-wing airplane and a helicopter produced no "*drastic*" disturbance of tree-nesting colonial waterbirds, and, in 90% of the observations, the individual birds either showed no reaction or merely looked up (Kushlan 1979). Belanger and Bedard (1989, 1990) observed responses of greater snow geese (*Chen caerulescens atlantica*) to man-induced disturbance on a sanctuary area and estimated the energetic cost of such disturbance. Belanger and Bedard (1989, 1990) observed that disturbance rates exceeding two per hour reduced goose use of the sanctuary by 50% the following day. They also observed that about 40% of the disturbances caused interruptions in feeding that would require an estimated 32% increase in nighttime feeding to compensate for the energy lost. They concluded that overflights of sanctuary areas should be strictly regulated to avoid adverse effects. Conomy et al. (1998) quantified behavioral responses of wintering American black ducks (*Anas rubripes*), American wigeon (*A. americana*), gadwall (*A. strepera*), and American green-winged teal (*A. crecca carolinensis*) exposed to low-level military aircraft and found that only a small percentage (2%) of the birds reacted to the disturbance. They concluded that such disturbance was not adversely affecting the "*time-activity budgets*" of the species. Aerial operations conducted by WS would not be conducted over federal, state, or other governmental agency property without the concurrence of the managing entity. Those flights, if requested, would be conducted to reduce threats and damages occurring to natural resources and should not result in impacts to bird species. Thus, there is little to no potential for any adverse effects on waterbirds and waterfowl.

***Raptors*:** The Air National Guard analyzed and summarized the effects of overflight studies conducted by numerous federal and state government agencies and private organizations (Air National Guard 1997). Those studies determined that military aircraft noise initially startled raptors, but negative responses were brief and did not have an observed effect on productivity (see Ellis 1981, Fraser et al. 1985, Lamp 1989, USFS 1992 as cited in Air National Guard 1997). A study conducted on the impacts of overflights to bald eagles (*Haliaeetus leucocephalus*) suggested that the eagles were not sensitive to this type of disturbance (Fraser et al. 1985). During the study, observations were made of more than 850 overflights of active eagle nests. Only two eagles rose out of either their incubation or brooding postures. This study also showed that perched adults were flushed only 10% of the time during aircraft overflights. Evidence also suggests that golden eagles (*Aquila chrysaetos*) are not highly sensitive to noise or other aircraft disturbances (Ellis 1981, Holthuijzen et al. 1990). Finally, one other study found that eagles were particularly resistant to being flushed from their nests (see Awbrey and Bowles 1990 as cited in Air National Guard 1997). Therefore, there is considerable evidence that eagles would not be adversely affected by overflights during aerial operations.

Mexican spotted owls (*Strix occidentalis lucida*) (Delaney et al. 1999) did not flush when chain saws and helicopters were greater than 110 yards away; owls flushed to these disturbances at closer distances and were more prone to flush from chain saws than helicopters. Owls returned to their pre-disturbance behavior 10 to 15 minutes following the event and researchers observed no differences in nest or nestling success (Delaney et al. 1999), which indicates that aircraft flights did not result in adverse effects on owl reproduction or survival.

Andersen et al. (1989) conducted low-level helicopter overflights directly at 35 red-tailed hawk (*Buteo jamaicensis*) nests and concluded their observations supported the hypothesis that red-tailed hawks habituate to low level flights during the nesting period; results showed similar nesting success between hawks subjected to overflights and those that were not. White and Thurow (1985) did not evaluate the effects of aircraft overflights, but found that ferruginous hawks (*B. regalis*) were sensitive to certain types of ground-based human disturbance to the point that reproductive success may be adversely affected. However, military jets that flew low over the study area during training exercises did not appear to bother the hawks, nor did the hawks become alarmed when the researchers flew within 100 feet in a small fixed-wing aircraft (White and Thurow 1985). White and Sherrod (1973) suggested that disturbance of raptors by aerial surveys with helicopters may be less than that caused by approaching nests on foot. Ellis (1981) reported that five species of hawks, two falcons (*Falco* spp.), and golden eagles (*Aquila chrysaetos*) were "*incredibly tolerant*" of overflights by military fighter jets, and observed that, although birds frequently exhibited alarm, negative responses were brief and the overflights never limited productivity.

Grubb et al. (2010) evaluated golden eagle response to civilian and military (Apache AH-64) helicopter flights in northern Utah. Study results indicated that golden eagles were not adversely affected when exposed to flights ranging from 100 to 800 meters along, towards and from behind occupied cliff nests. Eagle courtship, nesting, and fledging were not adversely affected, indicating that no special management restrictions were required in the study location.

The above studies indicate raptors were relatively unaffected by aircraft overflights, including those by military aircraft that produce much higher noise levels. Therefore, we conclude that aerial operations would have little or no potential to adversely affect raptors.

***Passerines*:** Reproductive losses have been reported in one study of small territorial passerines ("*perching*" birds that included sparrows, blackbirds) after exposure to low altitude overflights (see Manci et al. 1988 as cited in Air National Guard 1997), but natural mortality rates of both adults and young are high and variable for most species. The research review indicated passerine birds cannot be driven any great distance from a favored food source by a non-specific disturbance, such as military aircraft noise, which indicated quieter noise would have even less effect. Passerines avoid intermittent or unpredictable sources of disturbance more than predictable ones, but return rapidly to feed or roost once the disturbance ceases (Gladwin et al. 1988, USFS 1992). Those studies and reviews indicated there is little or no potential for aerial operations to cause adverse effects on passerine bird species.

***Pronghorn (antelope) and Mule Deer*:** Krausman et al. (2004) found that Sonoran pronghorn (*Antilocapra americana sonoriensis*) were not adversely affected by military fighter jet training flights and other military activity on an area of frequent and intensive military flight training operations. Krausman et al. (1986) reported that only three of 70 observed responses of mule deer (*Odocoileus hemionus*) to small fixed-wing aircraft overflights at 150 to 500 feet AGL resulted in the deer changing habitats. The authors believed that the deer might have been accustomed to overflights because the study area was near an interstate highway that was followed frequently by aircraft. Krausman et al. (2004) also reported that pronghorn and mule deer do not hear noise from military aircraft as well as humans, which potentially indicates why they appeared not to be disturbed as much as previously thought.

***Mountain Sheep*:** Krausman and Hervert (1983) reported that, of 32 observations of the response of mountain sheep to low-level flights by small fixed-wing aircraft, 60% resulted in no disturbance, 81% in no or "*slight*" disturbance, and 19% in "*great*" disturbance. Krausman and Hervert (1983) concluded that flights less than 150 feet AGL could cause mountain sheep to leave an area. When Weisenberger et al. (1996) evaluated the effects of simulated low altitude jet aircraft noise on desert mule deer (*Odocoileus hemionus crooki*) and mountain sheep (*Ovis canadensis mexicana*), they found that heart rates of the ungulates increased according to the dB levels, with lower noise levels prompting lesser increases. When they were elevated, heart rates rapidly returned to pre-disturbance levels suggesting that the animals did not perceive the noise as a threat. Responses to the simulated noise levels were found to decrease with increased exposure.

***Bison*:** Fancy (1982) reported that only two of 59 bison (*Bison bison*) groups showed any visible reaction to small fixed-winged aircraft flying at 200 to 500 feet AGL. The study suggests that bison were relatively tolerant of aircraft overflights.

***Domestic Animals and Small Mammals*:** A number of studies with laboratory animals (*e.g.*, rodents [Borg 1979]) and domestic animals (*e.g.*, sheep [Ames and Arehart 1972]) have shown that these animals can become habituated to noise. Long-term lab studies of small mammals exposed intermittently to high levels of noise demonstrate no changes in longevity. The physiological "*fight or flight*" response, while marked, does not appear to have any long-term health consequences on small mammals (Air National Guard 1997). Small mammals habituate, although with difficulty, to sound levels greater than 100 dbA (USFS 1992).

Although many of those wildlife species discussed above are not present in Tennessee, the information was provided to demonstrate the relative tolerance most wildlife species have of overflights, even those that involve noise at high decibels such as from military aircraft. In general, the greatest potential for impacts to occur would be expected to exist when overflights are frequent such as hourly and over many days that could represent "*chronic*" exposure. Chronic exposure situations generally involve areas near commercial airports and military flight training facilities. Even then, many wildlife species become habituated to overflights that appear to naturally minimize any potential adverse effects where such flights occur on a regular basis. Therefore, it is logical to conclude that the aircraft used in aerial hunting for feral swine and/or coyotes should have far less potential to cause any disturbance to wildlife than military aircraft because the military aircraft produce much louder noise and would be flown over certain training areas many more times per year, and yet were found to have no expected adverse effects on wildlife (Air National Guard 1997).

The fact that WS would only conduct aerial hunting on a very small percentage of the land area of the State indicates that most wildlife would not even be exposed to aerial gunning overflights in the State. Further lessening the potential for any adverse impacts is that such flights occur infrequently throughout the year.

While every precaution would be taken to safeguard against taking non-targets during operational use of methods and techniques for resolving damage and reducing threats caused by mammals, the use of such methods could result in the incidental lethal take of unintended species. The unintentional take and capture of wildlife species during damage management activities conducted under the proposed action alternative would primarily be associated with the use of body-gripping traps and in some situations, with live-capture methods, such as foothold traps, cage traps, and cable restraints. The unintentional lethal take of non-targets by WS from FY 2006 through FY 2011 is shown in Table 4.3 while those non-targets live-captured and released unharmed are shown in Table 4.4.

Table 4.3 – WS' lethal take of non-targets by method in Tennessee, FY 2006 – FY 2011

Species	Method of Lethal Take				Total
	Body Grip	Cage Trap†	Foothold	Neck Snare	
Coyote	0	0	0	1	**1**
Fox Squirrel	0	1	0	0	**1**
Muskrat	21	0	2	0	**23**
Nutria	3	0	0	0	**3**
Opossum	1	4	0	0	**5**
Raccoon	46	0	28	10	**84**
River Otter	125	0	4	11	**140**
Striped Skunk	1	0	0	0	**1**

†These animals were dispatched with firearms to minimize safety concerns of personnel with the release of animals.

The species with the highest level of lethal take were river otters, which were primarily lethally taken during damage management activities targeting beaver. WS has lethally removed 140 river otters as non-targets from FY 2006 through FY 2011, which is an average of 23.3 otter per year. All of those species lethally removed by WS as non-targets are also known to cause damage and could be addressed to alleviate damage or threats of damage by WS when requested under the proposed action alternative.

Table 4.4 – Non-targets live-captured and released by WS in Tennessee, FY 2006 – FY 2011

Species	Method of Live-Capture				Total
	Body Grip†	Cage Trap	Foothold†	Neck Snare†	
Bobcat	0	1	0	0	**1**
Cottontail Rabbit	0	9	0	0	**9**
Feral Cat	0	154	0	0	**154**
Feral Dog	0	6	0	2	**8**
Gray Fox	0	1	0	0	**1**
Gray Squirrel	0	22	0	0	**4**
Woodchuck	0	7	0	0	**7**
Opossum	0	947	0	0	**947**
Raccoons	2	6	4	1	**13**
River Otter	1	0	1	2	**4**
Striped Skunk	0	9	0	0	**9**

†Animals captured in body grip, foothold, or neck snares by the tail or other extremity would be released if they are unharmed and can be released safely.

As shown in Table 4.4, most non-targets captured by WS during damage management activities are live-captured and subsequently released unharmed. The primary species live-captured are opossum and feral cats. Non-targets released have been primarily live-captured during activities targeting raccoons as part of the ORV program (USDA 2005*b*) addressed in Chapter 1 in which WS employs cage traps to live-capture raccoons for sampling. The ORV program and the post-baiting trapping program are further described in the EA addressing those activities (USDA 2005*b*). The capture and limited lethal take that could occur as part of the ORV program and trapping activities are further addressed in the ORV program EA (USDA 2005*b*). Non-targets captured and lethally taken as part of the ORV program are addressed in this EA to ensure a cumulative evaluation of potential effects on non-target populations occurs from those activities that could be conducted under the proposed action alternative.

WS would monitor the take of non-target species to ensure program activities or methodologies used in mammal damage management would not adversely affect non-targets. Methods available to resolve and prevent mammal damage or threats when employed by trained, knowledgeable personnel would be

selective for target species. WS would report to the TWRA any non-target take to ensure take by WS was considered as part of management objectives established for those species by the TWRA. The potential impacts to non-targets would be similar to the other alternatives and would be considered minimal to non-existent.

As discussed previously, the use of non-lethal methods to address damage or threats would generally be regarded as having no impact on a species' population since those individuals addressed using non-lethal methods would be unharmed and no actual reduction in the number of individuals in a species' population occurs. Similarly, the live-capture and release of non-targets is generally regarding as having no adverse effects on a species' population since those individuals are released unharmed and no actual reduction in the number of individuals in a population occurs. Therefore, the live-capture and subsequent releasing of non-targets during damage management activities conducted under the proposed action alternative would not result in declines in the number of individuals in a species' population.

The lethal take of non-targets could result in declines in the number of individuals in a population; however, as shown in Table 4.3, the lethal take of non-targets by WS during damage management activities occurs rarely. A total of 258 non-targets have been lethally taken by WS during damage management activities conducted from FY 2006 through FY 2011, which is an average of 43 non-targets lethally taken annually by WS. The non-targets taken previously by WS are representative of non-targets that could be lethally taken by WS under the proposed action alternative. Although additional species of non-targets could be lethally taken by WS, take of individuals from any species is not likely to increase substantively above the number of non-targets taken annually by WS during previous damage management activities. In addition, all of the species lethally taken or live-captured during those six years are also considered target species in this EA and the level of take analyzed for each species under Issue 1 includes non-target take that could occur by WS. Therefore, the take of those species is evaluated cumulatively under Issue 1, including take that could occur when a species is considered a target or non-target. WS would continue to monitor activities, including non-target take to ensure the annual take of non-targets does not result in adverse effects to a species' population. All the species lethally taken previously can be harvested in the State during annual harvest seasons.

While every precaution would be taken to safeguard against taking non-targets during operational use of methods and techniques for resolving damage and reducing threats caused by mammals, the use of such methods could result in the incidental take of unintended species. Those occurrences would be rare and should not affect the overall populations of any species under the proposed action.

T&E Species Effects

Special efforts would be made to avoid jeopardizing T&E species through biological evaluations of the potential effects and the establishment of special restrictions or minimization measures. SOPs to avoid T&E species effects are described in Chapter 3 of this EA.

Federally Listed Species - The current list of species designated as threatened and endangered in Tennessee as determined by the USFWS was obtained and reviewed during the development of this EA. Appendix C contains the list of species currently listed in the State along with common and scientific names. Based on a review of those T&E species listed in the State during the development of the EA, WS determined that activities conducted pursuant to the proposed action would not likely adversely affect those species listed in the State by the USFWS nor their critical habitats. As part of the development of the EA, WS consulted with the USFWS under Section 7 of the ESA. The USFWS concurred with WS' determination that activities conducted pursuant to the proposed action would not likely adversely affect those species currently listed in the State or their critical habitats (M. Jennings, USFWS pers. comm.

2012). However, WS would consult with the USFWS prior to the initiation of manipulating water levels associated with beaver dams that occur east of the Tennessee River (see Appendix F).

State Listed Species – The current lists of State listed species designated as endangered or threatened as determined by the TWRA and TDEC were obtained and reviewed during the development of the EA (see Appendix D). Based on the review of species listed in the State, WS has determined that the proposed activities would not adversely affect those species currently listed by the State. The TWRA has concurred with WS' determination for State listed species (G. Anderson, TWRA pers. comm. 2012). However, WS would consult with the TWRA prior to the initiation of manipulating water levels associated with beaver dams that occur east of the Tennessee River (see Appendix F).

Alternative 2 – Mammal Damage Management by WS through Technical Assistance Only

Under a technical assistance alternative, WS would have no direct impact on non-target species, including T&E species. Methods recommended or provided through loaning of equipment could be employed by those persons requesting assistance. Recommendations would be based on WS' Decision Model using information provided by the person requesting assistance or through site visits. Recommendations would include methods or techniques to minimize non-target impacts associated with the methods being recommended or loaned. Methods recommended could include non-lethal and lethal methods as deemed appropriate by WS' Decision Model and as permitted by laws and regulations.

The potential impacts to non-targets under this alternative would be variable and based on several factors. If methods were employed, as recommended by WS, the potential impacts to non-targets would likely be similar to the proposed action. If recommended methods and techniques were not followed or if other methods were employed that were not recommended, the potential impacts on non-target species, including T&E species would likely be higher compared to the proposed action.

The potential impacts of harassment and exclusion methods on non-target species would be similar to those described under the proposed action. Harassment and exclusion methods would be easily obtainable and simple to employ. Since identification of targets would occur when employing shooting as a method, the potential impacts to non-target species would likely be low under this alternative.

Those persons experiencing damage from mammals may implement methods and techniques based on the recommendations of WS. The potential for impacts would be based on the knowledge and skill of those persons implementing recommended methods. If those persons experiencing damage do not implement methods or techniques correctly, the potential impacts from providing only technical assistance could be greater than the proposed action. The incorrect implementation of methods or techniques recommended by WS could lead to an increase in non-target take when compared to the non-target take that could occur by WS under the proposed action alternative.

If requestors were provided technical assistance but do not implement any of the recommended actions and take no further action, the potential to take non-targets would be lower when compared to the proposed action. If those persons requesting assistance implement recommended methods appropriately and as instructed or demonstrated, the potential impacts to non-targets would be similar to the proposed action. If WS made recommendations on the use of methods to alleviate damage but those methods were not implemented as recommended by WS or if those methods recommended by WS were used inappropriately, the potential for lethal take of non-targets would likely increase under a technical assistance only alternative. Therefore, the potential impacts to non-targets, including T&E species would be variable under a technical assistance only alternative.

If non-lethal methods recommended by WS under this alternative were deemed ineffective by those requesting assistance, lethal methods could be employed by those experiencing damage. Those persons requesting assistance are those persons likely to use lethal methods since a damage threshold has been met for that individual requestor that has triggered assistance to reduce damage. The potential impacts on non-targets by those persons experiencing damage would be highly variable. People whose mammal damage problems were not effectively resolved by non-lethal control methods would likely resort to other means of legal or illegal lethal control. This could result in less experienced persons implementing control methods and could lead to greater take of non-target wildlife than the proposed action. When those persons experiencing damage caused by wildlife reach a level where assistance does not adequately reduce damage or where no assistance is available, people have resorted to using chemical toxicants that are illegal for use on the intended target species. The illegal use of methods often results in loss of both target and non-target wildlife (White et al. 1989, USFWS 2001, FDA 2003). The use of illegal toxicants by those persons frustrated with the lack of assistance or assistance that inadequately reduces damage to an acceptable level can often result in the indiscriminate take of wildlife species.

The ability to reduce negative impacts caused by mammals to wildlife species and their habitats, including T&E species, would be variable under this alternative. The ability to reduce risks would be based upon the skills and abilities of the person implementing damage management actions. It would be expected that this alternative would have a greater chance of reducing damage than Alternative 3 since WS would be available to provide information and advice on appropriately employing methods and reducing the risk of non-target take.

Alternative 3 – No Mammal Damage Management Conducted by WS

Under this alternative, WS would not be directly involved with damage management activities in the State. Therefore, no direct impacts to non-targets or T&E species would occur by WS under this alternative. Mammals would continue to be taken under permits issued by the TWRA, take would continue to occur during the regulated harvest seasons, and non-native mammal species could continue to be taken without the need for a permit. Risks to non-targets and T&E species would continue to occur from those people who implement damage management activities on their own or through recommendations by the other federal, state, and private entities. Although some risks occur from those people that implement mammal damage management in the absence of any involvement by WS, those risks would likely be low and would be similar to those risks under the other alternatives.

The ability to reduce negative impacts caused by mammals to other wildlife species and their habitats, including T&E species, would be variable based upon the skills and abilities of the person implementing damage management actions under this alternative.

Issue 3 - Effects of Mammal Damage Management Activities on Human Health and Safety

A common concern is the potential adverse effects that methods available could have on human health and safety. The threats to human safety of methods available under the alternatives are evaluated below by each of the alternatives.

Alternative 1 - Continue the Current Adaptive Integrated Mammal Damage Management Program (No Action/Proposed Action)

The cooperator requesting assistance would be made aware through a MOU, cooperative service agreement, or a similar document that those methods agreed upon could potentially be used on property owned or managed by the cooperator. Therefore, the cooperator would be made aware of the possible use of those methods on property they own or manage through the signing of a MOU, cooperative service

agreement, or similar document, which would assist with identifying any risks to human safety associated with the use of those methods.

Under the proposed action, those methods discussed in Appendix B, could be integrated to resolve and prevent damage associated with mammals in the State. WS would use the Decision Model to determine the appropriate method or methods that would effectively resolve the request for assistance. Those methods would be continually evaluated for effectiveness and if necessary, additional methods could be employed. Non-lethal and lethal methods could be used under the proposed action. WS would continue to provide technical assistance and/or direct operational assistance to those persons seeking assistance with managing damage or threats from mammals. Risks to human safety from technical assistance conducted by WS would be similar to those risks addressed under Alternative 2. The use of non-lethal methods as part of an integrated approach to managing damage that could be employed as part of direct operational assistance by WS would be similar to those risks addressed in the other alternatives.

Lethal methods available under the proposed action would include the use of live-capture followed by euthanasia, body-gripping traps, cable restraints, the recommendation of take during hunting and/or trapping seasons, fumigants, rodenticides, and shooting. Those lethal methods available under the proposed action alternative or similar products would also be available under the other alternatives. None of the lethal methods available would be restricted to use by WS only. Euthanasia chemicals would not be available to the public but those mammals live-captured could be killed using other methods.

WS' employees who conduct activities to manage damage caused by mammals would be knowledgeable in the use of methods, wildlife species responsible for causing damage or threats, and WS' directives. That knowledge would be incorporated into the decision-making process inherent with the WS' Decision Model that would be applied when addressing threats and damage caused by mammals. When employing lethal methods, WS' employees consider risks to human safety when employing those methods based on location and method. For example, risks to human safety from the use of methods would likely be lower in rural areas that are less densely populated. Consideration would also be given to the location where damage management activities would be conducted based on property ownership. If locations where methods would be employed occur on private property in rural areas where access to the property is controlled and monitored, the risks to human safety from the use of methods is likely less. If damage management activities occur at parks or near other public use areas, then risks of the public encountering damage management methods and the corresponding risk to human safety increases. Activities would generally be conducted when human activity is minimal (*e.g.*, early mornings, at night) or in areas where human activities are minimal (*e.g.*, in areas closed to the public).

The use of live-capture traps, restraining devices (*e.g.*, foothold traps, some cable restraints), and body-gripping traps have been identified as a potential issue. Live-capture traps available for mammals would typically be walk-in style traps where mammals enter but are unable to exit. Live-traps, restraining devices, and body-gripping traps would typically be set in situations where human activity was minimal to ensure public safety. Those methods rarely cause serious injury and are triggered through direct activation of the device. Therefore, human safety concerns associated with live-traps, restraining devices, and body-gripping traps used to capture wildlife, including mammals, would require direct contact to cause bodily harm. Therefore, if left undisturbed, risks to human safety would be minimal. Signs warning of the use of those tools in the area would be posted for public view at access points to increase awareness that those devices were being used and to avoid the area, especially pet owners.

Other live-capture devices, such as cannon nets, pose minor safety hazards to the public since activation of the device occurs by trained personnel after target species have been observed in the capture area of the net. Lasers also pose minimal risks to the public since application occurs directly to target species by trained personnel, which limits the exposure of the public to misuse of the method.

Safety issues related to the misuse of firearms and the potential human hazards associated with firearms use are issues identified when employed to reduce damage and threats. To help ensure safe use and awareness, WS' employees who use firearms during official duties are required to attend an approved firearm safety-training course and to remain certified for firearm use must attend a safety training course in accordance with WS Directive 2.615. As a condition of employment, WS' employees who carry and use firearms are subject to the Lautenberg Domestic Confiscation Law, which prohibits firearm possession by anyone who has been convicted of a misdemeanor crime of domestic violence (18 USC § 922(g)(9)). A safety assessment based on site evaluations, coordination with cooperating and local agencies (if applicable), and consultation with cooperators would be conducted before firearms were deemed appropriate to alleviate or reduce damage and threats to human safety when conducting activities in the State. WS would work closely with cooperators requesting assistance to ensure all safety issues were considered before firearms would be deemed appropriate for use. The use of all methods, including firearms, would be agreed upon with the cooperator to ensure the safe use of those methods.

The issue of using chemical methods as part of managing damage associated with wildlife relates to the potential for human exposure either through direct contact with the chemical or exposure to the chemical from wildlife that have been exposed. Under the alternatives identified, the use of chemical methods would include immobilizing drugs, euthanasia drugs, reproductive inhibitors, fumigants, rodenticides, and repellents.

The use of immobilizing drugs would only be administered to mammals that have been live-captured using other methods or administered through injection using a projectile (*e.g.*, dart gun). Immobilizing drugs used to sedate wildlife would be used to temporarily handle and transport animals to lessen the distress of the animal from the experience. Drug delivery to immobilize mammals would be likely to occur on site with close monitoring of the animal to ensure proper care of the animal. Immobilizing drugs would be fully reversible with a full recovery of sedated animals occurring. A list and description of immobilizing drugs available for use under the identified alternatives can be found in Appendix B.

Euthanizing drugs would be administered under similar circumstances to immobilizing drugs. Euthanizing drugs would be administered to animals live-captured using other methods. Euthanized animals would be disposed of in accordance with WS Directives; therefore, would not be available for harvest and consumption. If mammals were immobilized for sampling or translocation and released, risks could occur to human safety if harvest and consumption occurred. SOPs employed by WS to reduce risks are discussed in Chapter 3 and in Appendix B.

Drugs used in capturing, handling, and euthanizing wildlife include ketamine, a mixture of ketamine/xylazine, sodium pentobarbital, potassium chloride, and Beuthanasia-D. Meeting the requirements of the AMDUCA should prevent any adverse impacts on human health with regard to this issue (see Section 1.6). SOPs that would be part of the activities conducted would include:

- All drugs used in capturing and handling wildlife would be under the direction and authority of state veterinary authorities, either directly or through procedures agreed upon between those authorities and WS.
- As determined on a state-level basis by those veterinary authorities (as allowed by AMDUCA), wildlife hazard management programs may choose to avoid capture and handling activities that utilize immobilizing drugs within a specified number of days prior to the hunting or trapping season for the target species to avoid release of animals that may be consumed by hunters prior to the end of established withdrawal periods for the particular drugs used. Ear tagging or other marking of animals drugged and released to alert hunters and trappers that they should contact state officials before consuming the animal.

- Most animals administered drugs would be released well before hunting/trapping seasons, which would give the drug time to completely metabolize out of the animals' systems before they might be taken and consumed by humans. In some instances, animals collected for control purposes would be euthanized when they are captured within a certain specified time period prior to the legal hunting or trapping season to avoid the chance that they would be consumed as food while still potentially having immobilizing drugs in their systems.

By following those procedures in accordance with AMDUCA, wildlife management programs would avoid any adverse effects on human health with regard to this issue.

All WS' personnel who handle and administer chemical methods would be properly trained in the use of those methods. Training and adherence to agency directives (see WS Directive 2.430) would ensure the safety of employees applying chemical methods. Mammals euthanized by WS or taken using chemical methods would be disposed of in accordance with WS Directive 2.515. All euthanasia would occur in the absence of the public, whenever possible, which would minimize risks. SOPs are further described in Chapter 3 of this EA.

The recommendation of repellents or the use of those repellents registered for use to disperse mammals in the State could occur under the proposed action as part of an integrated approach to managing mammal damage. Those chemical repellents that would be available to recommend for use or that could be directly used by WS under this alternative would also be available under any of the alternatives. Therefore, risks to human safety from the recommendation of repellents or the direct use of repellents would be similar across all the alternatives. Risks to human safety associated with the use of repellents by WS or the recommendation of repellents by WS is addressed under the technical assistance only alternative (Alternative 2). Risks to human safety would be similar across all the alternatives. WS' involvement, either through recommending the use of repellents or the direct use of repellents, would ensure that label requirements of those repellents were discussed with those persons requesting assistance when recommended through technical assistance or would be specifically adhered to by WS' personnel when using those chemical methods. Therefore, the risks to human safety associated with the recommendation of or direct use of repellents could be lessened through WS' participation.

Gas cartridges would be ignited and placed inside of burrows or dens with the entrance covered by dirt, which traps carbon monoxide inside the burrow. The carbon monoxide would dissipate into the atmosphere and be diluted by the air (EPA 1991). WS would follow label instructions when employing gas cartridges. Therefore, no risks to human safety would occur from the use of gas cartridges.

The recommendation of various rodenticides or the use of those rodenticides registered for use to manage rodents in the State could occur under the proposed action as part of an integrated approach to managing mammal damage. Those rodenticides that would be available for use by WS or could be recommended by WS under this alternative would also be available under any of the alternatives. Therefore, risks to human safety from the recommendation of rodenticides or the direct use of rodenticides would be similar across all the alternatives. WS' involvement, either through recommending the use of rodenticides or their direct use, would ensure that label requirements of these rodenticides are discussed with those persons requesting assistance when recommended through technical assistance or would be specifically adhered to by WS' personnel when using those chemical methods. Therefore, the risks to human safety associated with the recommendation of or direct use of rodenticides could be lessened through WS' participation.

Due to the classification of GonaCon™ as a restricted-use pesticide by the EPA, this product would be restricted to use by federal or state agencies that have successfully completed the requirements of the TDA for the purchase and application of restricted-use pesticides. Risks to human safety would be

limited primarily to the actual applicator due to the necessity to capture and inject GonaCon™ into each animal to be vaccinated. At this time, GonaCon™ has not been registered for use in Tennessee and is therefore not available for use within the State. However, this product could be registered for use in Tennessee and could be administered by TWRA or their agents under any of the alternatives.

Risks to human safety from the use of GonaCon™ would be minimal and would occur primarily to those persons injecting the deer through accidental self-injection or those persons handling syringes. To reduce the risks of accidental exposure through self-injection, the label of GonaCon™ requires the use of long sleeved shirts, long pants, gloves, socks, and shoes. In addition, injection would only occur after deer have been properly restrained to minimize accidental injection during application to the deer. The label also requires that children be absent from the area during application of the vaccine as well as a warning to women that accidental self-injection could cause infertility. WS' employees who are pregnant would not be involved with handling or injecting of the vaccine.

In addition, human exposure could occur through consumption of deer that were treated with GonaCon™. As was discussed previously, the vaccine and the antibodies produced in response to the vaccine are amino acid proteins that if consumed would be broken down by stomach acids and enzymes, posing no risks to human safety. The vaccine would only be used in localized areas where deer populations have exceeded the biological or social carrying capacity. Those areas are likely places where hunting is prohibited or restricted (*e.g.*, in parks); therefore, the consumption of deer is unlikely in those areas where the vaccine would be used since hunting would be prohibited or restricted. Deer injected with the vaccine must also be marked for identification, which would allow for placement of warnings to people that could take and consume a treated deer. Based on the use pattern of GonaCon™ and the chemical make-up of the vaccine and the antibodies, the risks to human safety from the use of the vaccine would be extremely low and would occur primarily to the handler (EPA 2009).

The recommendation by WS that mammals be harvested during the regulated hunting and/or trapping seasons that are established by the TWRA would not increase risks to human safety above those risks already inherent with hunting or trapping those species. Recommendations of allowing hunting and/or trapping on property owned or managed by a cooperator to reduce mammal populations, which could then reduce damage or threats, would not increase risks to human safety. Safety requirements established by the TWRA for the regulated hunting and trapping season would further minimize risks associated with hunting and trapping. Although hunting and trapping accidents do occur, the recommendation of allowing hunting or trapping to reduce localized populations of mammals would not increase those risks.

Consequences of Aerial Wildlife Operations Accidents

Aerial wildlife operations, like any other flying, may result in an accident. WS' pilots and crew members would be trained and experienced to recognize the circumstances which lead to accidents and have thousands of hours of flight time. The national WS Aviation Program has increased its emphasis on safety, including funding for additional training, the establishment of a WS Flight Training Center and annual recurring training for all pilots. Still, accidents may occur and the environmental consequences should be evaluated.

***Major Ground or Wild/Forest Fires*:** Although fires could result from aircraft-related accidents, no such fires have occurred from aircraft incidents previously involving government aircraft and low-level flights.

Fuel Spills and Environmental Hazard from Aviation Accidents: A representative of the National Transportation Safety Board has stated previously that aviation fuel is extremely volatile and will evaporate within a few hours or less to the point that even its odor cannot be detected (USDA 2005*c*). Helicopters used for aerial wildlife operations carry less fuel than fixed-wing aircraft with 30 gallons the

maximum for most helicopters. In some cases, little or none of the fuel would be spilled if an accident occurs. Thus, there should be little environmental hazard from unignited fuel spills.

Oil and Other Fluid Spills: With the size of aircraft used by WS, the quantities of oil (*e.g.*, 3 to 5 quarts in helicopters) capable of being spilled in any accident would be small and insignificant with respect to the potential for environmental damage. The greatest potential amount of oil that could be spilled in one accident would be about eight quarts.

Petroleum products biodegrade through volatilization and bacterial action, particularly when exposed to oxygen (EPA 2000). Thus, small quantity oil spills on surface soils can be expected to biodegrade readily. Even in subsurface contamination situations involving underground storage facilities that would generally be expected to involve larger quantities than would ever be involved in a small aircraft accident, EPA guidelines provide for "*natural attenuation*" or volatilization and biodegradation in some situations to mitigate environmental hazards (EPA 2000). Thus, even where oil spills in small aircraft accidents were not cleaned up, the oil does not persist in the environment or persists in such small quantities that no adverse effects would be expected. Also, WS' accidents generally would occur in remote areas away from human habitation and drinking water supplies. Thus, the risk to drinking water appears to be exceedingly low to nonexistent.

For these reasons, the risk of ground fires or fuel/oil pollution from aviation accidents could be considered low. In addition, based on the history and experience of the program in aircraft accidents, it appears the risk of significant environmental damage from such accidents is exceedingly low.

No adverse effects to human safety have occurred from WS' use of methods to alleviate mammal damage in the State from FY 2006 through FY 2011. The risks to human safety from the use of non-lethal and lethal methods, when used appropriately and by trained personnel, would be considered low.

As stated earlier, the cooperator requesting assistance would be made aware through a MOU, cooperative service agreement, or a similar document that those devices agreed upon could potentially be used on property owned or managed by the cooperator; thereby, making the cooperator aware of the use of those methods on property they own or manage to identify any risks to human safety associated with the use of those methods.

Alternative 2 – Mammal Damage Management by WS through Technical Assistance Only

Under this alternative, WS would be restricted to making recommendations of methods and the demonstration of methods only to resolve damage. WS would only provide technical assistance to those persons requesting assistance with mammal damage and threats. Although hazards to human safety from non-lethal methods exist, those methods are generally regarded as safe when used by trained individuals who are experienced in their use. Risks to human safety associated with non-chemical methods such as resource management methods (*e.g.*, crop selection, limited habitat modification, modification of human behavior), exclusion devices, frightening devices, and cage traps would be considered low based on their use profile for alleviating damage associated with wildlife. Although some risk of fire and bodily harm exists from the use of pyrotechnics and propane cannons, when used appropriately and in consideration of those risks, they can be used with a high degree of safety.

Under a technical assistance only alternative, the use of immobilizing drugs, euthanasia drugs, and Gonacon™ would not be available to the public. However, personnel with the TWRA or their designated agents could use Gonacon™ under this alternative. Drugs used in capturing and handling wildlife could be administered under the direction and authority of state veterinary authorities, either directly or through procedures agreed upon between those authorities and other entities, such as the TWRA. Personnel

employing nets would be present at the site during application to ensure the safety of the public and operators. Although some fire and explosive hazards exist with rocket nets during ignition and storage of the explosive charges, safety precautions associated with the use of the method, when adhered to, would pose minimal risks to human safety and would primarily occur to the handler. Nets would not be employed in areas where public activity was high, which further reduces the risks to the public. Nets would be employed in areas where public access was restricted whenever possible to reduce risks to human safety. Overall, nets would pose minimal risks to the public.

The use of chemical methods that are considered non-lethal would be available under this alternative. Chemical methods available would include repellents. There are few chemical repellents registered for use to manage damage caused by mammals in the State. Most repellents require ingestion of the chemical to achieve the desired affects on target species. Repellents that require ingestion are intended to discourage foraging on vulnerable resources and to disperse mammals from areas where the repellents are applied. Repellents, when used according to label directions, are generally regarded as safe especially when the ingredients are considered naturally occurring. Some risk of exposure to the chemical occurs to the applicator and to others from the potential for drift as the product is applied. Some repellents also have restrictions on whether application can occur on edible plants with some restricting harvest for a designated period after application. All restrictions on harvest and required personal protective equipment would be included on the label and if followed, would minimize risks to human safety associated with the use of those products.

The recommended use of chemical methods that were considered lethal would also be available under this alternative. Lethal chemicals available would consist primarily of those Ready-To-Use toxicants targeting rodents (*e.g.*, D-Con) that were available at local hardware stores for use in managing old world rodents. Those toxicants would require no special certification to use and would generally be considered safe when their use occurred in accordance with label directions. Additional lethal chemicals would be available through WS' recommendation to contact private sector wildlife control operators that have received TDA certification for use of restricted-use pesticides. While those chemicals may not be available to individual landowners, through use of a private sector wildlife control operator, similar chemical use and mammal damage control could be achieved.

The recommendation by WS that mammals be harvested during the regulated hunting and/or trapping season, that would be established by the TWRA, would not increase risks to human safety above those risks already inherent with hunting and trapping mammals. Recommendations of allowing hunting or trapping on property owned or managed by a cooperator to reduce local mammal populations that could then reduce mammal damage or threats would not increase risks to human safety. Safety requirements established by the TWRA for the regulated hunting and trapping season would further minimize risks associated with those activities. Although hunting and trapping accidents do occur, the recommendation of allowing hunting or trapping to reduce localized mammal populations would not increase those risks.

The recommendation of shooting with firearms as a method of direct lethal take could occur under this alternative. Safety issues do arise related to misusing firearms and the potential human hazards associated with firearms use when employed to reduce damage and threats. When used appropriately and with consideration for human safety, risks associated with firearms would be minimal. If firearms were employed inappropriately or without regard to human safety, serious injuries could occur. Under this alternative, recommendations of the use of firearms by WS would include human safety considerations. Since the use of firearms to alleviate mammal damage would be available under any of the alternatives and the use of firearms by those persons experiencing mammal damage could occur whether WS was consulted or contacted, the risks to human safety from the use of firearms would be similar among all the alternatives.

If non-chemical methods were employed according to recommendations and as demonstrated by WS, the potential risks to human safety would be similar to the proposed action. If methods were employed without guidance from WS or applied inappropriately, the risks to human safety could increase. The extent of the increased risk would be unknown and variable. Non-chemical methods inherently pose minimal risks to human safety given the design and the extent of the use of those methods.

The cooperator requesting assistance would also be made aware of threats to human safety associated with the use of those methods. SOPs for methods are discussed in Chapter 3 of this EA. Risks to human safety from activities and methods recommended under this alternative would be similar to the other alternatives since the same methods would be available. If misused or applied inappropriately, any of the methods available to alleviate mammal damage could threaten human safety. However, when used appropriately, methods available to alleviate damage would not threaten human safety.

Alternative 3 – No Mammal Damage Management Conducted by WS

Under the no involvement in damage management by WS alternative, WS would not be involved with any aspect of managing damage associated with mammals in the State, including technical assistance. Due to the lack of involvement in managing damage caused by mammals, no impacts to human safety would occur directly from WS. This alternative would not prevent those entities experiencing threats or damage from mammals from conducting damage management activities in the absence of WS' assistance. The direct burden of implementing permitted methods would be placed on those persons experiencing damage or require those persons to seek assistance from other entities.

Similar to the technical assistance only alternative, Gonacon™, immobilizing drugs, and euthanasia chemicals would not be available under this alternative to the public. However, fumigants, rodenticides, and repellents would continue to be available to those persons with the appropriate pesticide applicators license. Since most methods available to resolve or prevent mammal damage or threats would be available to anyone, the threats to human safety from the use of those methods would be similar between the alternatives. However, methods employed by those persons not experienced in the use of methods or are not trained in their proper use, could increase threats to human safety. Overall, the methods available to the public, when applied correctly and appropriately, pose minimal risks to human safety.

Issue 4 - Effects of Mammal Damage Management Activities on the Socio-cultural Elements of the Human Environment

Another concern often raised is the potential impact the alternatives could have on the aesthetic value that people often regard for mammals. The effects of the alternatives on this issue are analyzed below by alternative.

Alternative 1 - Continue the Current Adaptive Integrated Mammal Damage Management Program (No Action/Proposed Action)

Under the proposed action, methods would be employed that would result in the dispersal, exclusion, or removal of individuals or small groups of mammals to resolve damage and threats. In some instances where mammals were dispersed or removed, the ability of interested persons to observe and enjoy those mammals would likely temporarily decline.

Even the use of exclusionary devices can lead to the dispersal of wildlife if the resource being damaged was acting as an attractant. Thus, once the attractant has been removed or made unavailable, the wildlife would likely disperse to other areas where resources would be more vulnerable.

The use of lethal methods would result in temporary declines in local populations resulting from the removal of mammals to address or prevent damage and threats. The goal under the proposed action would be to respond to requests for assistance and to manage those mammals responsible for the resulting damage. Therefore, the ability to view and enjoy mammals would remain if a reasonable effort were made to locate mammals outside the area in which damage management activities were occurring. The mammals removed by WS would be those mammals that could be removed by the person experiencing damage.

All activities would be conducted where a request for assistance was received and only after the cooperator and WS had signed a cooperative agreement or similar document. Some aesthetic value would be gained by the removal of mammals and the return of a more natural environment, including the return of native wildlife and plant species that may be suppressed or displaced by high mammal densities.

Since those mammals removed by WS under this alternative could be removed with a permit issued by the TWRA, without the need for a permit if the species is unregulated, or during the regulated hunting or trapping seasons, WS' involvement in taking those mammals would not likely be additive to the number of mammals that could be taken in the absence of WS' involvement.

WS' take of mammals from FY 2006 through FY 2011 has been of low magnitude compared to the total mortality and populations of those species. WS' activities would not likely be additive to the mammals that could be taken in the absence of WS' involvement. Although mammals removed by WS would no longer be present for viewing or enjoying, those mammals would likely be taken by the property owner or manager if WS was not involved in the action since take by the property owner or manager could occur under a permit, during the regulated hunting and trapping seasons, or if the mammals were unregulated, take could occur without the need for a permit. Given the limited take proposed by WS under this alternative when compared to the known sources of mortality of mammals and the populations of those species, WS' mammal damage management activities conducted pursuant to the proposed action would not adversely affect the aesthetic value of mammals. The impact on the aesthetic value of mammals and the ability of the public to view and enjoy mammals under the proposed action would be similar to the other alternatives and would likely be low.

Alternative 2 – Mammal Damage Management by WS through Technical Assistance Only

If those persons seeking assistance from WS were those persons likely to conduct damage management activities in the absence of WS' involvement, then technical assistance provided by WS would not adversely affect the aesthetic value of mammals in the State similar to Alternative 1. Mammals could be lethally taken under this alternative by those entities experiencing mammal damage or threats which would result in localized reductions in the presence of mammals at the location where damage was occurring. The presence of mammals where damage was occurring would be reduced where damage management activities were conducted under any of the alternatives. Even the recommendation of non-lethal methods would likely result in the dispersal of mammals from the area if those non-lethal methods recommended by WS were employed by those persons receiving technical assistance. Therefore, technical assistance provided by WS would not prevent the aesthetic enjoyment of mammals since any activities conducted to alleviate mammal damage could occur in the absence of WS' participation in the action, either directly or indirectly.

Under this alternative, the effects on the aesthetic values of mammals would be similar to those addressed in the proposed action. When people seek assistance with managing damage from either WS or another entity, the damage level has often reached an unacceptable economic threshold for that particular person. Therefore, in the case of mammal damage, the social acceptance level of those mammals has reached a level where assistance has been requested and those persons would likely apply methods or seek those

entities that would apply those methods based on recommendations provided by WS or by other entities. Based on those recommendations, methods would likely be employed by the requestor that would result in the dispersal and/or removal of mammals responsible for damage or threatening safety. If those mammals causing damage were dispersed or removed by those persons experiencing damage based on recommendations by WS or other entities, the potential effects on the aesthetic value of those mammals would be similar to the proposed action alternative.

The impacts on aesthetics from a technical assistance program would only be lower than the proposed action if those individuals experiencing damage were not as diligent in employing those methods as WS would be if conducting an operational program. If those persons experiencing damage abandoned the use of those methods, then mammals would likely remain in the area and available for viewing and enjoying for those persons interested in doing so. Similar to the other alternatives, the geographical area in which damage management activities could occur would not be such that mammals would be dispersed or removed from such large areas that opportunities to view and enjoy mammals would be severely limited.

Alternative 3 – No Mammal Damage Management Conducted by WS

Under the no mammal damage management by WS alternative, the actions of WS would have no impact on the aesthetic value of mammals in the State. Those persons experiencing damage or threats from mammals would be responsible for researching, obtaining, and using all methods as permitted by federal, state, and local laws and regulations. Mammals could continue to be dispersed and lethally taken under this alternative in the State. Lethal take could continue to occur when permitted by the TWRA through the issuance of permits, take could occur during the regulated harvest season, and in the case of non-regulated species, take could occur any time without the need for a permit.

Since mammals would continue to be taken under this alternative, despite WS' lack of involvement, the ability to view and enjoy mammals would likely be similar to the other alternatives. The lack of WS' involvement would not lead to a reduction in the number of mammals dispersed or taken since WS' has no authority to regulate take or the harassment of mammals in the State. The TWRA, with management authority over mammals could continue to adjust all take levels based on population objectives for those mammal species in the State. Therefore, the number of mammals lethally taken annually through hunting and under permits would be regulated and adjusted by the TWRA.

Those persons experiencing damage or threats could continue to use those methods they feel appropriate to resolve mammal damage or threats, including lethal take. Therefore, WS' involvement in mammal damage management would not be additive to the mammals that could be taken in the State. The impacts to the aesthetic value of mammals would be similar to the other alternatives.

Issue 5 - Humaneness and Animal Welfare Concerns of Methods

As discussed previously, a common issue often raised is concerns about the humaneness of methods available under the alternatives for resolving mammal damage and threats. The issues of method humaneness relating to the alternatives are discussed below.

Alternative 1 - Continue the Current Adaptive Integrated Mammal Damage Management Program (No Action/Proposed Action)

Under the proposed action, WS would integrate methods using WS' Decision Model as part of technical assistance and direct operational assistance. Methods available under the proposed action could include non-lethal and lethal methods integrated into direct operational assistance conducted by WS. Under this alternative, non-lethal methods would be used by WS that were generally regarded as humane. Non-

lethal methods would include resource management methods (*e.g.*, crop selection, limited habitat modification, modification of human behavior), translocation, exclusion devices, frightening devices, reproductive inhibitors, cage traps, foothold traps, nets, and repellents.

As discussed previously, humaneness, in part, appears to be a person's perception of harm or pain inflicted on an animal. People may perceive the humaneness of an action differently. The challenge in coping with this issue is how to achieve the least amount of animal suffering.

Some individuals believe any use of lethal methods to resolve damage associated with wildlife is inhumane because the resulting fate is the death of the animal. Others believe that certain lethal methods can lead to a humane death. Others believe most non-lethal methods of capturing wildlife to be humane because the animal is generally unharmed and alive. Still others believe that any disruption in the behavior of wildlife is inhumane. With the multitude of attitudes on the meaning of humaneness and the varying perspectives on the most effective way to address damage and threats in a humane manner, agencies are challenged with conducting activities and employing methods that are perceived to be humane while assisting those persons requesting assistance to manage damage and threats associated with wildlife. The goal of WS would be to use methods as humanely as possible to effectively resolve requests for assistance to reduce damage and threats to human safety. WS would continue to evaluate methods and activities to minimize the pain and suffering of methods addressed when attempting to resolve requests for assistance.

Some methods have been stereotyped as "*humane*" or "*inhumane*". However, many "*humane*" methods can be inhumane if not used appropriately. For instance, a cage trap would generally be considered by most members of the public as "*humane*". Yet, without proper care, live-captured wildlife in a cage trap can be treated inhumanely if not attended to appropriately.

Therefore, the goal would be to effectively address requests for assistance using methods in the most humane way possible that minimizes the stress and pain to the animal. Overall, the use of resource management methods, harassment methods, and exclusion devices are regarded as humane when used appropriately. Although some concern arises from the use of live-capture methods, the stress of animals is likely temporary.

Although some issues of humaneness could occur from the use of cage traps, foothold traps, reproductive inhibitors, translocation, immobilizing drugs, nets, and repellents, those methods, when used appropriately and by trained personnel, would not result in the inhumane treatment of wildlife. Concerns from the use of those non-lethal methods would be from injuries to animals while those animals were restrained and from the stress of the animal while being restrained or during the application of the method. Pain and physical restraint can cause stress in animals and the inability of animals to effectively deal with those stressors can lead to distress. Suffering occurs when action is not taken to alleviate conditions that cause pain or distress in animals.

If mammals were to be live-captured by WS, WS' personnel would be present on-site during capture events or capture devices would be checked frequently to ensure mammals captured were addressed in a timely manner and to prevent injury. Although stress could occur from being restrained, timely attention to live-captured wildlife would alleviate suffering. Stress would likely be temporary. Euthanasia of live-captured target mammals by WS under Alternative 1 would occur pursuant to WS Directive 2.505.

Under the proposed action, lethal methods could also be employed to resolve requests for assistance to alleviate or prevent mammal damage and threats. Lethal methods would include shooting, body-gripping traps, cable restraints, fumigants, rodenticides, and the recommendation of take during hunting and/or trapping seasons. In addition, target species live-captured using non-lethal methods could be euthanized

by WS. WS' use of lethal control methods under the proposed action would follow those required by WS' directives (see WS Directive 2.505, WS Directive 2.430).

The euthanasia methods being considered for use under the proposed action for live-captured mammals are carbon dioxide, carbon monoxide, gunshot, and barbiturates or potassium chloride in conjunction with general anesthesia. The use of carbon dioxide, carbon monoxide, barbiturates, and potassium chloride for euthanasia would occur after the animal had been live-captured and away from public view. Although the AVMA guideline also lists gunshot as a conditionally acceptable method of euthanasia for free-ranging wildlife, there is greater potential the method may not consistently produce a humane death (AVMA 2007). WS' personnel that employ firearms to address mammal damage or threats to human safety would be trained in the proper placement of shots to ensure a timely and quick death.

An issue when dealing with aquatic rodent species is the use of foothold traps to create drowning sets and the humaneness of drowning. There is considerable debate and disagreement among animal interest groups, veterinarians, wildlife professionals, fur trappers, and nuisance wildlife control specialists on this issue. The debate centers on an uncertainty as to whether the drowning animals are rendered unconscious by high levels of carbon dioxide (CO_2) and are thus insensitive to distress and pain (Ludders et al. 1999).

The AVMA identifies drowning as an unacceptable method of euthanasia (Beaver et al. 2001, AVMA 2007). Ludders et al. (1999) concluded drowning is not euthanasia based on the animals not dying from CO_2 narcosis, and reported CO_2 narcosis does not occur until 95 millimeters of mercury in arterial blood is exceeded. Ludders et al. (1999) showed death during drowning occurs from hypoxia and anoxia; thus, animals experience hypoxemia. Ludders et al. (1999) also concluded that animals that drown are distressed because of stress related hormones, epinephrine and norepinephrine, and therefore drowning is not euthanasia.

CO_2 causes death in animals by hypoxemia and some animals (cats, rabbits, and swine) are distressed before death (Beaver et al. 2001). Even though those animals are distressed, the AVMA (Beaver et al. 2001) states this death is an acceptable form of euthanasia (AVMA 2007). Thus, the AVMA does not preclude distress or pain in euthanasia.

Death by drowning in the classical sense is caused by the inhalation of fluid into the lungs and is referred to as "*wet*" drowning (Gilbert and Gofton 1982, Noonan 1998). Gilbert and Gofton (1982) reported that all submerged beaver do not die from wet drowning, but die of CO_2-induced narcosis, and the AVMA has stated the use of CO_2 is acceptable (Gilbert and Gofton 1982, Noonan 1998). Gilbert and Gofton (1982) reported that after beaver were trapped and entered the water, they struggled for two to five minutes, followed by a period of reflexive responses. Andrews et al. (1993) stated that with some techniques that induce hypoxia, some animals have reflex motor activity followed by unconsciousness that is not perceived by the animal. Gilbert and Gofton (1982) stated it is unknown how much conscious control actually existed at this stage and they stated anoxia might have removed much of the sensory perception by five to seven minutes post submersion.

However, Gilbert and Gofton (1982) have been criticized because levels of CO_2 in the blood were not reported (Ludders et al. 1999) and there was insufficient evidence that the beaver in their study were under a state of CO_2 narcosis when they died (letter from V. Nettles, D.V.M., Ph.D., Southeastern Cooperative Wildlife Disease Study, to W. MacCallum, MDFW, June 15, 1998). Adding to the controversy, Clausen and Ersland (1970) did measure CO_2 in the blood for submersed restrained beaver; yet, none of the beaver in their study died, so Clausen and Ersland (1970) could not determine if beaver died of CO_2 narcosis. Clausen and Ersland (1970) demonstrated that CO_2 increased in arterial blood while beaver were submersed and CO_2 was retained in the tissues. While Clausen and Ersland (1970) did measure the amounts of CO_2 in the blood of submersed beaver, they did not attempt to measure the

analgesic effect of CO_2 buildup to the beaver (letter from V. Nettles, D.V.M., Ph.D., Southeastern Cooperative Wildlife Disease Study, to W. MacCallum, MDFW, June 15, 1998). When beaver were trapped using foothold traps with intent to "*drown*", the beaver exhibit a flight response. Gracely and Sternberg (1999) reported that there is stress-induced analgesia resulting in reduced pain sensitivity during fight or flight responses. Environmental stressors that animals experience during flight or fight activate the same stress-induced analgesia (Gracely and Sternberg 1999).

The use of drowning trap sets has been a traditional wildlife management technique in trapping aquatic mammals such as beaver and muskrat. Trapper education manuals and other wildlife damage management manuals written by wildlife biologists recommend drowning sets for foothold traps set for beaver (Howard et al. 1980, Randolph 1988, Bromley et al. 1994, Dolbeer et al. 1994, Miller and Yarrow 1994). In some situations, drowning trap sets are the most appropriate and efficient method available to capture beaver and muskrat. For example, a drowning set attachment should be used with foothold traps when capturing beaver to prevent the animals from injuring themselves while restrained, or from escaping (Miller and Yarrow 1994). Animals that drown die relatively quickly (*e.g.*, within minutes) versus the possible stress of being restrained and harassed by people, dogs, and other wildlife before being euthanized. Drowning sets make the captured animal and trap less visible and prevents injury from the trapped animal (*i.e.*, bites and scratches) to people who may otherwise approach a restrained animal. Furthermore, the sight of dead animals may offend some people. Drowning places the dead animal out of public view. Some sites may be unsuitable for body-gripping traps or snares because of unstable banks, deep water, or a marsh with a soft bottom, but those sites would be suitable for foothold traps.

Given the short time period of a drowning event, the possible analgesic effect of CO_2 buildup, the minimal if any pain or distress on drowning animals, the AVMA acceptance of hypoxemia as euthanasia, the AVMA acceptance of a minimum of pain and distress during euthanasia, and the acceptance of catching and drowning muskrats approved by International Humane Trapping Standards (Fur Institute of Canada 2000), WS concludes that drowning, though rarely used by WS, is acceptable. WS recognizes some people would disagree.

Research and development by WS has improved the selectivity and humaneness of management techniques. Research is continuing to bring new findings and products into practical use. Until new findings and products are found practical, a certain amount of animal suffering could occur when some methods were used in situations where non-lethal damage management methods were not practical or effective. Personnel from WS would be experienced and professional in their use of management methods. Consequently, management methods would be implemented in the most humane manner possible. Many of the methods discussed in Appendix B to alleviate mammal damage and/or threats in the State could be used under any of the alternatives by those persons experiencing damage regardless of WS' direct involvement. The only methods that would not be available to those persons experiencing damage associated with mammals would be reproductive inhibitors, immobilizing drugs, and euthanasia drugs. Therefore, the issue of humanness associated with methods would be similar across any of the alternatives since those methods could be employed by other entities in the absence of WS' involvement. Those persons who view a particular method as humane or inhumane would likely continue to view those methods as humane or inhumane under any of the alternatives. SOPs that would be incorporated into WS' activities to ensure methods were used by WS as humanely as possible are listed in Chapter 3.

Alternative 2 – Mammal Damage Management by WS through Technical Assistance Only

The issue of humaneness of methods under this alternative would be similar to the humaneness issues discussed under the proposed action. This perceived similarity is derived from WS' recommendation of methods that some people may consider inhumane. WS would not directly be involved with damage management activities under this alternative. However, the recommendation of the use of methods would

likely result in the requester employing those methods. Therefore, by recommending methods and thus a requester employing those methods, the issue of humaneness would be similar to the proposed action. Under Alternative 2, WS would recommend the use of euthanasia methods pursuant to WS Directive 2.505. However, the person requesting assistance would determine what methods to use to euthanize or kill a live-captured animal under Alternative 2.

WS would instruct and demonstrate the proper use and placement of methodologies to increase effectiveness in capturing target mammal species and to ensure methods were used in such a way as to minimize pain and suffering. However, the efficacy of methods employed by a cooperator would be based on the skill and knowledge of the requestor in resolving the threat to safety or damage situation despite WS' demonstration. Therefore, a lack of understanding of the behavior of mammals or improperly identifying the damage caused by mammals along with inadequate knowledge and skill in using methodologies to resolve the damage or threat could lead to incidents with a greater probability of being perceived as inhumane. In those situations, the pain and suffering would likely be regarded as greater than discussed in the proposed action.

Alternative 3 – No Mammal Damage Management Conducted by WS

Under this alternative, WS would not be involved with any aspect of mammal damage management in Tennessee. Those persons experiencing damage or threats associated with mammals could continue to use those methods legally available. Those methods would likely be considered inhumane by those persons who would consider methods proposed under any alternative as inhumane. The issue of humaneness would likely be directly linked to the methods legally available to the public since methods are often labeled as inhumane by segments of society no matter the entity employing those methods.

The humaneness of methods would be based on the skill and knowledge of the person employing those methods. A lack of understanding of the target species or methods used could lead to an increase in situations perceived as being inhumane to wildlife despite the method used. Despite the lack of involvement by WS under this alternative, those methods perceived as inhumane by certain individuals and groups would still be available to the public to use to resolve damage and threats caused by mammals. Under Alternative 3, euthanasia or killing of live-captured animals would also be determined by those persons employing methods to live-captured wildlife.

Issue 6 - Effects of Mammal Damage Management Activities on the Regulated Harvest of Mammals

The populations of several of the mammal species addressed in this assessment are sufficient to allow for annual harvest seasons that typically occur during the fall. Hunting and trapping seasons are established by the TWRA. Those species addressed in this EA that have established hunting and/or trapping seasons include beaver, muskrat, nutria, woodchuck, cottontail rabbit, black bear, raccoon, long-tailed weasel, mink, striped skunk, spotted skunk, coyote, gray fox, red fox, bobcat, opossum, white-tailed deer, and elk. For many mammal species considered harvestable during hunting and/or trapping seasons, the estimated number of mammals harvested during the season is reported by the TWRA in published reports.

Alternative 1 - Continue the Current Adaptive Integrated Mammal Damage Management Program (No Action/Proposed Action)

The magnitude of lethal take addressed in the proposed action would be low when compared to the mortality of those species from all known sources. When WS' proposed take of mammals was included as part of the known mortality of those species and compared to the estimated populations, the impact on those species' populations was below the level of removal required to lower population levels.

With oversight of mammal populations by the TWRA, the number of mammals permitted by the TWRA to be taken by WS would not limit the ability of those persons interested to harvest those mammal species during the regulated season. All take by WS would be reported to the TWRA annually to ensure take by WS was incorporated into population management objectives established for mammal populations. Based on the limited take proposed by WS and the oversight by the TWRA, WS' take of mammals annually would have no effect on the ability of those persons interested to harvest mammals during the regulated harvest season.

Alternative 2 – Mammal Damage Management by WS through Technical Assistance Only

Under the technical assistance only alternative, WS would have no direct impact on mammal populations in the State. If WS recommended the use of non-lethal methods and those non-lethal methods were employed by those persons experiencing damage, mammals would likely be dispersed from the damage area to areas outside the damage area, which could serve to move those mammals from those less accessible areas to places accessible to hunters. Although lethal methods could be recommend by WS under a technical assistance only alternative, the use of those methods could only occur after the property owner or manager received a permit from the TWRA or when considered a non-regulated species, could be removed at any time using legally available methods. Lethal take could also occurring during the annual hunting and trapping season in areas where those activities were permitted. WS' recommendation of lethal methods could lead to an increase in the use of those methods. However, the number of mammals allowed to be taken under a permit and during the regulated hunting/trapping seasons would be determined by the TWRA. Therefore, WS' recommendation of the use of lethal methods under this alternative would not limit the ability of those persons interested in harvesting mammals during the regulated season since the TWRA determines the number of mammals that may be taken during the hunting/trapping season and under permits.

Alternative 3 – No Mammal Damage Management Conducted by WS

WS would have no impact on the ability to harvest mammals under this alternative. WS would not be involved with any aspect of mammal damage management. The TWRA would continue to regulate populations through adjustments of the allowed take during the regulated harvest season and the continued use of permits.

Issue 7 – Effects of Beaver Removal and Dam Manipulation on the Status of Wetlands in the State

Concern has also been expressed regarding the potential effects of the proposed action and the alternatives on wetland ecosystems associated with activities that could be conducted to address beaver damage or threats. Concerns have been raised that removing and/or modifying beaver dams in an area would result in the loss of wetland habitat and the plant and animal species associated with those wetlands. In addition, concerns are often raised regarding the use of lethal methods to remove beaver to alleviate damage or threats. If beaver were lethally removed from an area and any associated beaver dam was removed or breached, the manipulation of water levels by removing/breaching the dam could prevent the establishment of wetlands in areas where water has been impounded by beaver dams for an extended period.

Over time, the impounding of water associated with beaver dams can establish new wetlands. Because beaver dams may involve waters of the United States, the removal of a beaver dam is regulated under Section 404 of the CWA. The United States Army Corps Of Engineers and the EPA regulatory definition of a wetland (40 CFR 232.2) is: "*Those areas that are inundated or saturated by surface or groundwater at a frequency and duration sufficient to support, and that under normal circumstances do support, a*

prevalence of vegetation typically adapted for life in saturated soil conditions. Wetlands generally include swamps, marshes, bogs, and similar areas."

Although beaver can cause damage to resources, there can be many benefits associated with beaver and beaver activities. Beaver can provide ecological benefits associated with the creation of wetland habitats (Munther 1982, Wright 2002, Rossell et al. 2005, Bergman et al. 2007, Pollock 2007, Fouty 2008*a*, Fouty 2008*b*, Hood and Bayley 2008). Beaver can also provide aesthetic and recreational opportunities for wildlife observation (Wade and Ramsey 1986, Ringleman 1991), improve water quality (Muller-Schwarze and Sun 2003), and provide cultural and economic gains from fur harvest (Hill 1976, McNeely 1995, Lisle 1996, Lisle 2003).

Beaver impoundments can increase surface and groundwater storage, which can help reduce problems with flooding by slowing the downstream movement of water during high-flow events and help to mitigate the adverse impacts of drought (Fouty 2008*a*, Hey and Phillips 1995, Naiman et al. 1988, Wade and Ramsey 1986). Hood and Bayley (2008) determined that the presence of beaver could help reduce the loss of open water wetlands during warm, dry years. The presence of active beaver lodges accounted for over 80% of the variability in the amount of open water present in the mixed-wood boreal region of east-central Alberta (Hood and Bayley 2008). Hood and Bayley (2008) also found temperature and rainfall influenced the amount of open-water wetlands, but to a much lesser extent than the presence of beaver. During wet and dry years, the presence of beaver was associated with a 9-fold increase in open water area over the same areas when beaver were absent. Hood and Bayley (2008) noted that beaver could mitigate some of the adverse impacts of global warming through their ability to create and maintain areas of open water. Beaver ponds and associated wetlands can provide a potential water source for livestock, serve as basins for the entrapment of streambed silt and eroding soil (Hill 1982), and help to filter nutrients from the water; thereby, maintaining the quality of nearby water systems (Arner and Hepp 1989).

Beaver may increase habitat diversity by opening forest habitats via dam building and tree cutting, which can result in a greater mix of plant species, and different-aged plant communities (Hill 1982, Arner and Hepp 1989). The creation of standing water, edge habitat, and plant diversity, all in close proximity, can result in excellent habitat for many wildlife species (Jenkins and Busher 1979, Arner and DuBose 1982, Hill 1982, Arner and Hepp 1989, Medin and Clary 1990, Medin and Clary 1991). The wetland habitat that can be created by beaver ponds can be beneficial to some fish (primarily warm water species), reptiles, amphibians, waterfowl, shorebirds, and furbearers such as muskrats, otter, and mink (Arner and DuBose 1982, Naimen et al. 1986, Miller and Yarrow 1994). For example, in Mississippi, beaver ponds over three years in age were found to have developed plant communities valuable as nesting and brood rearing habitat for wood ducks (Arner and DuBose 1982). Reese and Hair (1976) found that beaver pond habitats were highly attractive to a large number of birds throughout the year and that the value of beaver pond habitat to waterfowl was minor when compared to other species of birds (Novak 1987). Beaver ponds can be beneficial to some T&E species. The USFWS estimates that up to 43% of T&E species rely directly or indirectly on wetlands for their survival (EPA 1995).

Under the proposed action alternative, WS could recommend and/or implement methods to manipulate water levels associated with water impounded by beaver dams to alleviate flooding damage. If the technical assistance alternative was selected, WS could recommend methods to people requesting assistance that could result in the manipulation of water levels associated with water impounded by beaver dams. WS would not be involved with any aspect of activities associated with beaver dams under the no involvement by WS alternative. Methods that would generally be available under all the alternatives would include explosives, exclusion devices, and water flow devices (see Appendix B for additional information). However, the availability to breach or remove beaver dams using explosives would be limited under Alternative 2 and Alternative 3, since the property owner or manager seeking to

remove or breach a dam would be required to locate a person certified to use explosives to conduct the work. In addition, the use of backhoes or other mechanical methods could be employed by property owners or managers to remove or breach beaver dams under any of the alternatives; however, WS would not operationally employ backhoes or other large machinery to remove or breach dams.

Exclusion devices and water control systems have been used for many years to manipulate the level of water impounded by beaver dams with varying degrees of success (United States General Accounting Office 2001). Landowner management objectives play a role in how the efficacy of a level system is perceived (Nolte et al. 2001). Nolte et al. (2001) found that survey respondents classified pond levelers installed to manage wetlands for waterfowl habitat more successful than levelers installed to provide relief from flooding. Langlois and Decker (2004) reported that "*...very few beaver problems...can actually be solved with a water level control device*" with a 4.5% success rate in Massachusetts and a 3% success rate in New York. Nolte et al. (2001) reported only 50% of installed pond levelers in Mississippi met landowner objectives and found that pond levelers placed in sites with high beaver activity more frequently failed if installed without implementing population control measures. Higher success rates have been reported for newer exclusion and water control systems ranging from 87% to 93% (Callahan 2005, Boyles 2006, Simon 2006, Boyles 2007). Lisle (2003) reported the use of water control devices or a combination of a Beaver Deceiver™ and flow management device virtually eliminated the need for maintenance and beaver removal at 20 sites where clogged culverts and flooded roads had previously been a routine issue.

When using exclusion and water control systems, those methods must be specifically designed to meet the needs of each site (Langlois and Decker 2004). Consequently, devices installed by inexperienced individuals may have a higher failure rate than those installed by a professional (Lisle 1996, Callahan 2003, Boyles 2006, Simon 2006, Spock 2006). Higher success rates reported for newer exclusion and water control devices may be indicative of increased understanding of the kinds of situations where those devices work best. For example, Callahan (2005) noted that exclusion and water control systems installed at culvert sites were more successful than similar systems installed at freestanding dams. Callahan (2003) and Callahan (2005) also provided a list of sites that were not well suited to the use of exclusion or water control devices. Boyles (2006) and Boyles (2007) reported some of the highest success rates for newer exclusion and water control systems; however, those devices were only tested at culvert sites.

Beaver build dams to raise water levels to meet their needs for security and access to forage. While pond levelers allow for the retention of some water, if the water level does not meet the needs of the beaver, they may move a short distance upstream or downstream and build a new dam, or abandon the area (Callahan 2003, Langlois and Decker 2004, Clemson University 2006). This may merely result in moving the problem to a new landowner or, depending upon site characteristics, the resulting pond may result in new or increased damage problems for the original landowner. McNeely (1995) reported the most common reasons cited for lack of success of water flow devices were clogging caused by debris or silt and beaver construction of additional dams upstream or downstream of the management device. In a study by Callahan (2005), construction of a new dam upstream or downstream of a pond leveler device was the most common cause of failure for free-standing dams (*e.g*., dams not associated with a culvert or other similar constriction in water flow, 11 of 156 sites). Callahan (2005) also found that insufficient pipe capacity (6 sites), dammed fencing (2 sites), and lack of maintenance (2 sites) were also causes for pond leveler failures. Nolte et al. (2001) also reported the need to address problems with dams upstream or downstream of a device. At culvert sites, Callahan (2005) found a lack of maintenance was the primary cause of failure with culvert exclusion devices (4 of 227 sites). Callahan (2005) also found vandalism resulted in the failure of a culvert device at one of the sites. At two culvert sites, Callahan (2005) found dammed fencing reduced or completely impeded the operation of exclusion devices.

Most pond levelers and exclusion devices require maintenance. The amount of maintenance required can vary considerably among sites, depending on site conditions and the type of device (Nolte et al. 2001, Callahan 2005, Boyles 2006, Spock 2006). Stream flow, leaf fall, floods, and beaver activity can continuously bring debris to the intake of a water control device. Ice damage and damage from debris washed downstream during high water events may also trigger the need for maintenance (*e.g*., cleaning out the intake pipe). Although most exclusion and water control devices generally require some level of maintenance, there are reports of devices that have remained effective for a period of years with no maintenance (Nolte et al. 2001). Nolte et al. (2001) reported that post-installation maintenance had been performed by property owners or managers on 70% of the 20 successfully operating Clemson pond levels installed by WS in Mississippi. The most common action was to adjust the riser on the pipe to manipulate water levels. Other maintenance included removal of vegetation and secondary dams built after the installation of the devices. In a survey of individuals who had received assistance with exclusion and water control devices, Simon (2006) found 18 of 36 survey respondents reported maintaining their devices, while installation program staff monitored an additional 10 devices. Of those survey respondents, Simon (2006) found that 61% reported that routine maintenance took 15 minutes or less while 93% reported that maintenance took a half hour or less. Boyles (2006) reported that time spent in device maintenance ranged from one to 4.75 hours per year.

Installation and upkeep of water control devices vary from site to site. For example, transporting materials over long distances in difficult terrain to install devices in remote locations where road access is not available could increase costs compared to the ability to transport materials for installation at a culvert site along a roadway. Callahan (2005) reported that the average cost for an exclusion fence at a culvert was $750 with an average annual maintenance cost of approximately $200. Flexible leveler pipe systems cost an average of $1,000 to install and $100 per year in maintenance, while the average cost to install a combination fence and leveler was $1,400 with approximately $150 per year in maintenance (Callahan 2005). Over a ten-year period, Callahan (2005) estimated the cost of installation and annual maintenance would range from $200 to $290 per year depending on the device installed. Spock (2006) reported that exclusion and/or water control device installation cost ranged from < $600 to over $3,000 dollars, with slightly more than half the systems (58.2%) ranging between $600 and $1,000 to install. In many cases, Spock (2006) found the cost included the first year of maintenance. The more expensive installations tended to be extensive fence and leveler systems or systems with numerous leveler pipes (Spock 2006). Boyles (2006) reported that device installation cost an average of $1,349 per device and $3,180 per site with subsequent annual maintenance cost averaging $19.75 per site per year (Boyles 2006). However, unlike the study by Callahan (2005) the devices evaluated by Boyles (2006) had only been in place for a relatively short time (average time in place 15 months, range 6 to 22 months versus average time in place 36.6 months, range 3 to 75 months). The cost of maintenance may vary over time as site conditions change.

Alternative 1 - Continue the Current Adaptive Integrated Mammal Damage Management Program (No Action/Proposed Action)

Manipulation of water levels associated with water impoundments caused by beaver dams could be addressed by WS under the proposed action using either dam breaching, dam removal, or the installation of water flow devices, including exclusion devices. Those methods allow dams to be breached or removed to maintain the normal flow of water. Heavy equipment, such as backhoes or bulldozers, would not be used by WS to breach, remove, or install water flow devices; although, heavy machinery could be utilized by a cooperator or their agents. WS may utilize small all terrain or amphibious vehicles and/or watercraft for transporting personnel, equipment, and supplies to worksites.

The breaching or removal of dams could be conducted by hand. Breaching is normally conducted through incremental stages of debris removal from the dam that allows water levels to be gradually

lowered. Breaching of dams would normally occur to limit the potential for flooding downstream by gradually allowing water levels to lower as more of the dam was breached over time. Breaching also minimizes the release of debris and sediment downstream by allowing water to move slowly over or through the dam. Depending on the size of the impoundment, water levels could be slowly lowered over several hours or days when breaching dams. When breaching dams, only that portion of the dam blocking the stream or ditch channel would be altered or breached, with the intent of returning water levels and flow rates to historical levels or to a level that eliminates damage threats or habitat alteration that would be acceptable to the property owner or resource manager. Similar to breaching dams, the removal of the dam removes the debris impounding water and restores the normal flow of water.

Beaver dams would generally be breached or removed by hand with a rake or power tools (*e.g.*, a winch). However, explosives would also be available to remove beaver dams. Explosives could potentially be utilized by WS' personnel specially trained and certified to conduct such activities. Explosives are defined as any chemical mixture or device that serves as a blasting agent or detonator. Explosives would generally be used to remove beaver dams that were too large to remove by hand. Explosives would only be used after beaver were removed from the site.

WS' personnel would only utilize binary explosives (*i.e.*, explosives comprised of two parts that must be mixed at the site before they can be detonated as an explosive material) for beaver dam removal, when requested. Binary explosives consist of ammonium nitrate and nitro-methane; however, those two components separated are not classified as explosives until mixed. Therefore, binary explosives would be subject to fewer regulations and controls. However, once mixed, binary explosives would be considered high explosives and subject to all applicable federal and state regulations. Detonating cord and detonators would also be considered explosives and WS would adhere to all applicable state and federal regulations for storage, transportation, and handling. WS' use of explosives and safety procedures would occur in accordance with WS Directive 2.435.

In addition to dam breaching and removal, water flow devices and exclusion methods would also be available for WS to employ during direct operational assistance or to recommend during technical assistance. Several different designs of water flow devices and exclusion methods would be available; however, the intent of all those methods would be to lower water levels by allowing water to flow through the beaver dam using pipes and wire mesh. After installation, beaver dams would be left intact with water levels maintained at desired levels by adjusting the water flow device. Water flow devices and exclusion methods allow beaver to remain at the site and maintain the beaver dam.

Although dams could be breached/removed manually or with binary explosives, those methods are usually ineffective because beaver can quickly repair or replace the dam if the beaver were not removed prior to breaching or removing the dam (McNeely 1995). Damage may be effectively reduced in some situations by installing exclusion and water control devices. Exclusion and water control devices can be designed so that the level of the beaver-created water impoundment can be managed to eliminate or minimize damage from flooding while retaining the ecological and recreational benefits derived from beaver impounding water over time. For example, WS may recommend modifications to site and culvert design (Jensen et al. 1999) as a non-lethal way of reducing problems with beaver dams at culverts.

Manipulating water levels impounded by beaver dams under the proposed action alternative would generally be conducted to maintain existing stream channels and drainage patterns, and to reduce water levels to alleviate flooding. WS could be requested to assist with manipulation of a beaver dam to alleviate flooding to agricultural crops, timber resources, public property, such as roads and bridges, private property, areas flooded because of beaver dams constructed on adjacent TVA property, and water management structures, such as culverts. The intent of breaching or removing beaver dams would not be to drain established wetlands. With few exceptions, requests for assistance received by WS from public

and private entities would involve breaching or removing dams to return an area to the condition that existed before the dam had been built, or before the impounded water had been affecting the area long enough for wetland characteristics to become established.

Most activities conducted by WS in Tennessee do not have the potential to affect wetlands, since those activities would not be conducted near or in wetlands. Under this alternative, water levels would be manipulated to return streams, channels, dikes, culverts, and canals to their original function. Most requests to alleviate flooding from impounded water would be associated roads, crops, merchantable timber, pastures, and other types of property or resources that were not previously flooded. Most dams removed would have been created because of recent beaver activity. WS' personnel receive most requests for assistance associated with beaver dams soon after affected resource owners discover damage.

Upon receiving a request to manipulate the water levels in impoundments caused by beaver dams, WS would visually inspect the dam and the associated water impoundment to determine if characteristics exist at the site that would meet the definition of a wetland under section 404 of the Clean Water Act (40 CFR 232.2). If wetland conditions were present at the site, the entities requesting assistance from WS would be notified that a permit might be required to manipulate the water levels impounded by the dam and to seek guidance from the TDEC and the United States Corps of Engineers pursuant to State laws and the Clean Water Act. If the area does not already have hydric soils, it usually takes several years for them to develop and a wetland to become established; this often takes greater than 5 years as indicated by the Swampbuster provision of the Food Security Act. Most beaver dam removal by WS would occur under exemptions stated in 33 CFR parts 323 and 330 of Section 404 of the Clean Water Act or parts 3821 and 3822 of the Food Security Act. However, manipulating water levels associated with some beaver dams could trigger certain portions of Section 404 that require landowners to obtain permits from the United States Army Corps of Engineers prior to removing a blockage. WS' personnel would determine the proper course of action upon inspecting a beaver dam impoundment. Appendix E describes the procedures used by WS to assure compliance with the pertinent laws and regulations.

The manipulation of water impoundment levels by WS by dam breaching, dam removal, or installation of water flow devices would typically be associated with dams constructed from recent beaver activity and would not have occurred long enough to take on the qualities of a true wetland (*i.e.* hydric soils, hydrophytic vegetation, and hydrological function). WS' activities associated with beaver dam breaching, beaver dam removal, or the installation of flow control device would only be conducted to restore the normal flow of water through drainages, streams, creeks, canals, and other watercourses where flooding damage was occurring or would occur. Beaver dam breaching or removal would not affect substrate or the natural course of streams.

In the majority of instances, beaver dam removal would be accomplished by manual methods (*i.e.*, hand tools). WS' personnel would not utilize heavy equipment, such as trackhoes or backhoes, for beaver dam removal. Only the portion of the dam blocking the stream or ditch channel would be breached or removed. In some instances, WS would install water flow devices to manage water levels at the site of a breached beaver dam. From FY 2006 through FY 2011, WS breached or removed 2,601 dams (2,352 by hand and 249 by using explosives) during damage management activities associated with beaver. Dams were breached or removed in accordance with exemptions from Section 404 permit requirements established by regulation or as allowed under nationwide permits (NWPs) granted under Section 404 of the Clean Water Act (see Appendix E). The majority of impoundments that WS removed were in existence for only a few months. Those impoundments were not considered wetlands as defined by 40 CFR 232.2; therefore, those impoundments did not possess the same wildlife habitat values as established wetlands.

In those situations where a non-federal cooperator had already made the decision to breach or remove a beaver damage to manipulate water levels with or without WS' assistance, WS' participation in carrying out the action would not affect the environmental status quo.

Additional concern has been raised relating to the lethal removal of beaver by WS or the recommendation of lethal methods to alleviate damage or threats of damage under the proposed action alternative. Beaver lethally removed could be replaced by other beaver requiring additional assistance later. Houston (1995) indicated that beaver tend to reoccupy vacant habitats. The likelihood that a site would be recolonized by beaver varies depending on many factors. For example, removal of beaver and a beaver dam from a relatively uniform section of irrigation canal may resolve the problem for an extended period because the relatively uniform nature of the canal does not predispose a site to repeat problems. Recolonization would also depend on the proximity and density of the beaver population in the surrounding area. Isolated areas or areas with a lower density of beaver would normally take longer for beaver to recolonize than areas with higher beaver densities. Activities conducted under the proposed action would be directed at specific beaver and/or beaver colonies and would not be conducted to suppress the overall beaver population in the State.

In accordance with WS Directive 2.101, preference would be given to non-lethal methods where practical and effective. Although use of exclusion and water control devices could greatly reduce the need for lethal beaver removal, beaver removal may still be needed in some situations even though a flow device or water control system had been installed (Wood et al. 1994, Nolte et al. 2001, Simon 2006, Spock 2006). Callahan (2005) states the trapping of beaver to alleviate damage should occur "*...where a flow device is either not feasible or fails, the water level needs to be drastically lowered, or the landowner wants no beavers or ponds on their property*". Spock (2006) reported that beaver had to be trapped out of one site when an exclusion system was augmented by the installation of a water control device. Lisle (1996) noted that it might be necessary to remove beaver that have learned to dam around exclusion and water control devices. In some instances, trapping during the annual trapping season for beaver continued to occur at or near the area where water control devices were installed but was not prompted by the failure of the devices (Lisle 1996, Simon 2006, Spock 2006).

Exclusion and water control devices may not be the most effective method in specific types of terrain and are not suitable for every site (Wood et al. 1994, Nolte et al. 2001, Langlois and Decker 2004, Callahan 2005). Exclusion devices and water control devices may not be suitable for man-made, uniform channels such as agricultural drainage ditches and irrigation canals; reservoirs; areas where human health, property or safety would be threatened with even minor elevation in water level; and areas where the landowner has expressed zero tolerance for beaver activity on their property (Callahan 2003, Callahan 2005, Simon 2006). Water control devices may be ineffective in beaver ponds in broad, low-lying areas because even a slight increase in water depth can result in a substantial increase in the area flooded (Organ et al. 1996). Exclusion and water control systems would not resolve problems related to beaver construction of bank dens. Depending upon site characteristics, beaver may build bank dens instead of lodges by burrowing into banks, levees, and other earthen impoundments. When bank dens are built in earthen levees or in banks supporting roadways or railroad tracks, they can greatly weaken the earthen structure. Burrowing into embankments can weaken the integrity of impoundments. Burrows allow water to infiltrate embankments, which can allow water to seep through the embankments causing erosion and weakening water impoundments. In those situations, removal of the beaver (either by translocation or by lethal methods) could be the only practical solution to resolve the potential for damage.

Water control devices may also be inappropriate in areas that are managed for aquatic species that need free-flowing water conditions and gravel substrate to survive. The still water and silt that accumulates behind beaver dams can be detrimental to some species. In addition, beaver dams could impede the movement of fish upstream. Avery (2004) found the removal of beaver dams resulted in substantial

increases in the stream area where trout could be found. For example, a 9.8-mile treatment zone on the North Branch of the Pemebonwon River in Wisconsin and an additional 17.9 miles of seven tributaries to the treatment section of the river were maintained free of beaver dams since 1986. In 1982, prior to dam removal, wild brook trout were found in only four of the seven tributaries within the treatment zone and at only four of the 12 survey stations. In the spring of 2000, wild brook trout were present in all seven tributaries and at all 12 survey stations (Avery 2004). In some cases, water control devices could be modified to improve fish passage (Close 2003). Although the presence of beaver dams could be detrimental to some species of fish, some fish species may benefit from the presence of a beaver dam (Rossell et al. 2005, Bergman et al. 2007, Pollock 2007).

Although beaver serve a valuable role in wetland ecology, the presence of beaver dams in intensively managed wetlands could be a concern to property owners or managers. In those areas, man-made water control structures are used to manage the water level in the wetland area in order to maximize habitat value for waterfowl and specific types of wetland-dependent wildlife (USDI 2008). While general elevations or reductions in water levels might conceivably be achieved by installing pipe systems through beaver dams, the devices tend to be more difficult to adjust than the water control structures. More importantly, the primary difficulty comes when drawdowns are used to achieve wetland management objectives. Drawdowns generally involve reducing the water level until large sections of mudflat are exposed. Many plant species valuable to waterfowl and other wetland bird species need exposed mudflats to sprout. Shorebirds use the mudflats to forage for invertebrates (USDI 2008). Once the plants have matured, the water level can be gradually increased until approximately half of the marsh has open water and half has standing plants (USDI 2008). Drawdowns may also be used in fall as a means of eliminating invasive fish (USDI 2008). The extent of the water level reduction conflicts with the beaver's desire for water deep enough to provide protection, and water area of sufficient extent to provide relatively easy access to foraging sites. The extent of the water level reduction during a drawdown would likely increase the risk of new dam creation in other locations that may cause new problems (Callahan 2003).

Alternative 2 – Mammal Damage Management by WS through Technical Assistance Only

The issues regarding the effects on wetlands under this alternative would likely be similar to wetlands issues discussed under the proposed action. This similarity would be based on WS' recommendation of methods to manage damage caused by beaver and the recommendation of methods to manage the water impounded by beaver dams. Based on information provided by the person requesting assistance or based on site visits, WS could recommend that a landowner or manager manipulate beaver dams to reduce flooding damage or threats of damage. WS would not be directly involved with conducting activities associated with the manipulation of beaver dams under this alternative. However, the recommendation of the use of methods would likely result in the requestor employing those methods or employing an agent to employ them. Therefore, by recommending methods and thus a requester employing those methods, the potential for those methods to reduce the presence of impounded water would be similar to the proposed action.

WS could instruct and demonstrate the proper use and placement of flow control and exclusionary devices, as well as recommend the breaching or removal of beaver dams, when appropriate. WS would also assist requestors by providing information on permit requirements and which municipal and state agencies need to be contacted by the requester to obtain appropriate permits to manipulate the levels of water impounded by beaver dams.

The efficacy of methods employed by a cooperator would be based on the skill and knowledge of the requester or their agent despite WS' recommendations or demonstration. Therefore, a lack of understanding of the behavior of beaver along with inadequate knowledge and skill in using methodologies to resolve flooding could lead to incidents with a greater probability of unforeseen impacts

to wetlands. In those situations, the potential for dam manipulation to adversely affect the status of wetlands would likely to be regarded as greater than those discussed in the proposed action.

WS would recommend the landowner or manager seek and obtain the proper permits to manipulate water levels impounded by beaver dams under this alternative; however, WS would not be responsible for ensuring that appropriate permits were obtained, proper methods were implemented for manipulating water levels, or for reviewing sites for the presence of T&E species. Those responsibilities would be incurred by property owner/manager and/or their designated agent who may or may not properly follow WS' recommendations.

Alternative 3 – No Mammal Damage Management Conducted by WS

Under this alternative, WS not be involved with any aspect of managing water levels associated with beaver dam impoundments. Under the no involvement by WS alternative, WS would not be involved with any aspect of managing damage associated with beaver in the State, including technical assistance. Due to the lack of involvement in managing damage caused by beaver, no impacts to wetlands would occur directly from WS. This alternative would not prevent those entities experiencing threats or damage due to flooding from manipulating water levels associated with beaver dams in the absence of WS' assistance. Those methods described previously would be available to other entities to breach or remove dams, including water flow devices. However, the use of explosives to remove dams under this alternative would be limited to those persons trained and licensed to use explosives. A property owner or manager could seek the services of an entity trained and licensed to use explosives to remove beaver dams under this alternative. The direct burden of implementing permitted methods would be placed on those persons experiencing damage.

Since the same methods would be available to resolve or prevent beaver damage or threats related to beaver dams, effects on the status of wetlands in the State from the use of those methods would be similar between the alternatives. However, manipulating water levels by those persons not experienced in identifying wetland characteristics or unaware of the requirement to seek appropriate permits to alter areas considered as a wetland, could increase threats to wetlands and the associated flora and fauna.

4.2 CUMULATIVE IMPACTS OF THE PROPOSED ACTION BY ISSUE

Cumulative impacts, as defined by CEQ (40 CFR 1508.7), are impacts to the environment that result from the incremental impact of the action when added to other past, present, and reasonably foreseeable future actions, regardless of what agency (federal or non-federal) or person undertakes such other actions. Cumulative impacts may result from individually minor, but collectively significant, actions taking place over time.

Under Alternative 1 and Alternative 2, WS would address damage associated with mammals either by providing technical assistance only (Alternative 2) or by providing technical assistance and direct operational assistance (Alternative 1) in the State. WS would be the primary federal agency conducting direct operational mammal damage management in the State under Alternative 1 and Alternative 2. However, other federal, state, and private entities could also be conducting mammal damage management in the State.

WS does not normally conduct direct damage management activities concurrently with such agencies or other entities in the same area, but may conduct damage management activities at adjacent sites within the same period. In addition, commercial companies may conduct damage management activities in the same area. The potential cumulative impacts analyzed below could occur from either WS' damage management program activities over time or from the aggregate effects of those activities combined with

the activities of other agencies and private entities. Through ongoing coordination and collaboration between WS, the TVA, and the TWRA, activities of each agency and the take of mammals would be available. Damage management activities in the State would be monitored to evaluate and analyze activities to ensure they were within the scope of analysis of this EA.

The activities proposed in the alternatives would have a negligible effect on atmospheric conditions including the global climate. Meaningful direct or indirect emissions of greenhouse gases would not occur because of any of the proposed alternatives. Those alternatives would meet the requirements of applicable laws, regulations, and Executive Orders including the Clean Air Act and Executive Order 13514.

Issue 1 - Effects of Mammal Damage Management Activities on Target Mammal Populations

Evaluation of activities relative to target species indicated that program activities would likely have no cumulative adverse effects on mammal populations when targeting those species responsible for damage at the levels addressed in this EA. WS' actions would be occurring simultaneously, over time, with other natural processes and human generated changes that are currently taking place. These activities include, but would not be limited to:

- Natural mortality of mammals
- Mortality through vehicle strikes, aircraft strikes, and illegal harvest
- Human-induced mortality of mammals through private damage management activities
- Human-induced mortality through regulated harvest
- Human and naturally induced alterations of wildlife habitat
- Annual and perennial cycles in wildlife population densities

All those factors play a role in the dynamics of mammal populations. In many circumstances, requests for assistance arise when some or all of those elements have contrived to elevate target species populations or place target species at a juncture to cause damage to resources. The actions taken to minimize or eliminate damage would be constrained as to scope, duration, and intensity for the purpose of minimizing or avoiding impacts to the environment. WS uses the Decision Model to evaluate damage occurring, including other affected elements and the dynamics of the damaging species; to determine appropriate strategies to minimize effects on environmental elements; applies damage management actions; and subsequently monitors and adjusts/ceases damage management actions (Slate et al. 1992). This process allows WS to take into consideration other influences in the environment, such as those listed above, in order to avoid cumulative adverse impacts on target species.

With management authority over statewide mammal populations, the TWRA can adjust take levels, including the take of WS, to ensure population objectives for mammals were achieved. Consultation and reporting of take by WS would ensure the TWRA considers any activities conducted by WS.

WS' take of mammals in Tennessee from FY 2006 through FY 2011 was of a low magnitude when compared to the total known take of those species and the populations of those species. The TWRA considers all known take when determining population objectives for mammals and can adjust the number of mammals that could be taken during the regulated hunting/trapping season and the number of mammals taken for damage management purposes to achieve the population objectives. Any take by WS would occur at the discretion of the TWRA. Any mammal population declines or increases would be the collective objective for mammal populations established by the TWRA through the regulation of take. Therefore, the cumulative take of mammals annually or over time by WS would occur at the desire of the TWRA as part of management objectives for mammals in the State. No cumulative adverse effects on

target and non-target wildlife would be expected from WS' damage management actions based on the following considerations:

Historical outcomes of WS' damage management activities on wildlife

Damage management activities associated with mammals would be conducted by WS only at the request of a cooperator to reduce damage that was occurring or to prevent damage from occurring and only after methods to be used were agreed upon by all parties involved. WS would monitor activities to ensure any potential impacts were identified and addressed. WS would work closely with state and federal resource agencies to ensure damage management activities would not adversely affect mammal populations and that WS' activities were considered as part of management goals established by those agencies. Historically, WS' activities to manage mammals in Tennessee have not reached a magnitude that would cause adverse effects to mammal populations in the State.

SOPs built into the WS program

SOPs are designed to reduce the potential negative effects of WS' actions on mammals, and have been tailored to respond to changes in wildlife populations that could result from unforeseen environmental changes. This would include those changes occurring from sources other than WS. Alterations in programs would be defined through SOPs, and implementation would be insured through monitoring, in accordance with the WS' Decision Model (Slate et al. 1992).

Issue 2 - Effects of Mammal Damage Management Activities on Non-target Wildlife Species Populations, Including T&E Species

Potential effects on non-target species from conducting mammal damage management arise from the use of non-lethal and lethal methods to alleviate or prevent those damages. The use of non-lethal methods during activities to reduce or prevent damage caused by mammals has the potential to exclude, disperse, or capture non-target wildlife. However, the effects of non-lethal methods are often temporary and often do not involve the take (killing) of non-target wildlife species. When using exclusion devices and/or repellents, both target and non-target wildlife can be prevented from accessing the resource being damaged. Since exclusion and repellents do not involve lethal take, cumulative impacts on non-target species from the use of exclusionary or repellent methods would not occur but would likely disperse those individuals to other areas. Exclusionary methods can often be expensive and require constant maintenance to ensure effectiveness. Similarly, repellents can also be expensive to apply and require constant re-application to ensure effectiveness. Therefore, the use of exclusionary devices and repellents would be somewhat limited to small, high-value areas and not used to the extent that non-targets would be excluded from large areas that would cumulatively impact populations from the inability to access a resource, such as potential food sources, denning, or fawning sites. The use of visual and auditory harassment and dispersion methods would generally be temporary with non-target species returning after the cessation of those activities. Dispersal and harassment do not involve the take (killing) of non-target species and similar to exclusionary methods would not be used to the extent or at a constant level that would prevent non-targets from accessing critical resources that would threaten survival of a population.

The use of lethal methods or those methods used to live-capture target species followed by euthanasia also have the potential to affect non-target wildlife through the take (killing) or capture of non-target species. Capture methods used are often methods that would be set to confine or restrain target wildlife after being triggered by a target individual. Capture methods would be employed in such a manner as to minimize the threat to non-target species by placement in those areas frequently used by target wildlife, using baits or lures that are as species specific as possible, and modification of individual methods to exclude non-targets from capture. Most methods described in Appendix B are methods that would be

employed to confine or restrain wildlife that would be subsequently euthanized using humane methods. With all live-capture devices, non-target wildlife captured could be released on site if determined to be able to survive following release. SOPs are intended to ensure take of non-target wildlife is minimal during the use of methods to capture target wildlife.

The use of firearms and euthanasia methods would essentially be selective for target species since identification of an individual would be made prior to the application of the method. Euthanasia methods would be applied through direct application to target wildlife. Therefore, the use of those methods would not affect non-target species.

All chemical methods would be tracked and recorded to ensure proper accounting of used and unused chemicals occurs. All chemicals would be stored and transported according with WS' Directives and relevant federal, state, and local regulations. Chemical methods available for use under the proposed action would include repellents, reproductive inhibitors, rodenticides, fumigants, immobilizing drugs, and euthanasia chemicals, which are described in Appendix B. Except for repellents that are applied directly to the affected resource, all chemical methods would be employed using baits that are highly attractive to target species, used in known burrow/den sites and/or used in areas where exposure to non-targets would be minimal. The use of those methods requires an acclimation period and monitoring of potential bait sites for non-target activity. All chemicals would be used according to product labels, which ensure that proper use would minimize non-target threats. WS' adherence to Directives and SOPs governing the use of chemicals would also ensure non-target hazards would be minimal.

Repellents may be used or recommended by the WS program in Tennessee to manage mammal damage. The active ingredients in numerous commercial repellents are capsaicin, pepper oil, and carnivore urine. Characteristics of these chemicals and potential use patterns indicate that no cumulative impacts related to environmental fate would be expected from their use in WS' programs in Tennessee when used according to label requirements.

When using rodenticides, as required by WS' SOPs and applicable pesticide labels, all potential bait sites would be pre-baited and monitored for non-target use as outlined in the pre-treatment observations section of the label. If non-targets were observed feeding on the pre-bait, the areas would be abandoned and no baiting would occur at those locations. Once sites were baited, sites would be monitored to further observe for non-target feeding activity. If non-targets were observed feeding on bait, those sites would be abandoned. WS would retrieve all dead target species to the extent possible following treatment to minimize any secondary hazards associated with or perceived to be associated with scavengers feeding on target species carcasses. When using rodenticides, appropriate bait stations would be utilized and inspected as required by the applicable label.

The amount of chemicals used or stored by WS would be minimal to ensure human safety. All label requirements of repellents and toxicants would be followed to minimize non-target hazards. Based on this information, WS' use of chemical methods, as part of the proposed action, would not have cumulative impacts on non-targets.

The methods described in Appendix B all have a high level of selectivity and could be employed using SOPs to ensure minimal impacts to non-target species. A total of 258 non-target mammals were lethally taken by WS during mammal damage management activities from FY 2006 through FY 2011, while 1,157 animals were live-captured and released unharmed. All of the species lethally taken as unintentional non-targets are included in take analysis as target species in this EA. The cumulative take of those species, including target and non-target take were evaluated in Chapter 4 of this EA. Based on the methods available to resolve mammal damage and/or threats, WS does not anticipate the number of non-targets taken to reach a magnitude where declines in those species' populations would occur.

Therefore, take under the proposed action of non-targets would not cumulatively affect non-target species. WS' has reviewed the T&E species listed by the TNHP and the USFWS, and has determined that damage management activities proposed by WS would not likely adversely affect T&E species. Cumulative impacts would be minimal on non-targets from any of the alternatives discussed**.**

Issue 3 - Effects of Mammal Damage Management Activities on Human Health and Safety

Non-chemical methods described in Appendix B would be used within a limited period, would not be residual, and do not possess properties capable of inducing cumulative adverse impacts on human health and safety. Non-chemical methods would be used after careful consideration of the safety of those persons employing methods and to the public. Capture methods would be employed where human activity was minimal to ensure the safety of the public. Capture methods also require direct contact to trigger ensuring that those methods, when left undisturbed would have no effect on human safety. All methods would be agreed upon by the requesting entities, which would be made aware of the safety issues of those methods when entering into a MOU, cooperative service agreement, or other comparable document between WS and the cooperating entity. SOPs also ensure the safety of the public from those methods used to capture or take wildlife. Firearms used to alleviate or prevent damage, though hazards do exist, would be employed to ensure the safety of employees and the public.

Personnel employing non-chemical methods would continue to be trained to be proficient in the use of those methods to ensure the safety of the applicator and to the public. Based on the use patterns of non-chemical methods, those methods would not cumulatively affect human safety.

Repellents to disperse mammals from areas of application would be available. Repellents must be registered with the EPA according to the FIFRA and with the TDA. Many of the repellents currently available for use have active ingredients that are naturally occurring and are generally regarded as safe. Although some hazards exist from the use of repellents, hazards occur primarily to the handler and applicator. When repellents are applied according to label requirements, no adverse effects to human safety would be expected. Similarly, fumigants and rodenticides must also be registered for use with the EPA and the TDA. Given the use patterns of repellents, rodenticides, and fumigants, no cumulative adverse effects would occur to human safety.

WS has received no reports or documented any adverse effects to human safety from WS' mammal damage management activities conducted from FY 2006 through FY 2011. No cumulative adverse effects from the use of those methods discussed in Appendix B would be expected given the use patterns of those methods for resolving mammal damage in the State.

Issue 4 - Effects of Mammal Damage Management Activities on the Socio-cultural Elements of the Human Environment

The activities of WS would result in the removal of mammals from those areas where damage or threats were occurring. Therefore, the aesthetic value of mammals in those areas where damage management activities were being conducted would be reduced. However, for some people, the aesthetic value of a more natural environment would be gained by reducing mammal densities, including the return of native species that may be suppressed or dispersed by non-native species.

Some people experience a decrease in aesthetic enjoyment of wildlife because they feel that overabundant species are objectionable and interfere with their enjoyment of wildlife in general. Continued increases in numbers of individuals or the continued presence of mammals may lead to further degradation of some people's enjoyment of any wildlife or the natural environment. The actions of WS could positively affect

the aesthetic enjoyment of wildlife for those people that were being adversely affected by the target species identified in this EA.

Mammal population objectives are established and enforced by the TWRA through the regulation of take during the statewide harvest seasons after consideration of other known mortality factors. Therefore, WS would have no direct impact on the status of mammal populations since all take by WS occurs at the discretion of the TWRA. Since those persons seeking assistance could remove mammals from areas where damage was occurring when permitted by the TWRA, WS' involvement would have no effect on the aesthetic value of mammals in the area where damage was occurring. When damage caused by mammals has occurred, any removal of mammals by the property or resource owner would likely occur whether WS was involved with taking the mammals or not.

In the wild, few animals in the United States have life spans approaching that of humans. Mortality is high among wildlife populations and specific individuals among a species may experience death early in life. This is a natural occurrence and humans who form affectionate bonds with animals experience loss of those animals over time in most instances. A number of professionals in the field of psychology have studied human behavior in response to attachment to pet animals (Gerwolls and Labott 1994, Marks and Koepke 1994, Zasloff 1996, Archer 1999, Ross and Baron-Sorensen 1998, Meyers 2000). Similar observations were probably applicable to close bonds that could exist between people and wild animals. As observed by researchers in human behavior, normal human responses to loss of loved ones proceed through phases of shock or emotional numbness, sense of loss, grief, acceptance of the loss or what cannot be changed, healing, and acceptance and rebuilding which leads to resumption of normal lives (Lefrancois 1999). Those who lose companion animals, or animals for which they may have developed a bond and affection, are observed to proceed through the same phases as with the loss of human companions (Gerwolls and Labott 1994, Boyce 1998, Meyers 2000). However, they usually establish a bond with other individual animals after such losses. Although they may lose the sense of enjoyment and meaning from the association with those animals that die or are no longer accessible, they usually find a similar meaningfulness by establishing an association with new individual animals or through other relational activities (Weisman 1991). Through this process of coping with the loss and establishing new affectionate bonds, people may avoid compounding emotional effects resulting from such losses (Parkes 1979, Lefrancois 1999).

Some mammals with which humans have established affectionate bonds may be removed from some project sites by WS. However, other individuals of the same species would likely continue to be present in the affected area and people would tend to establish new bonds with those remaining animals. In addition, human behavior processes usually result in individuals ultimately returning to normalcy after experiencing the loss of association with a wild animal that might be removed from a specific location. WS' activities would not be expected to have any cumulative adverse effects on this element of the human environment.

Issue 5 - Humaneness and Animal Welfare Concerns of Methods

WS would continue to seek new methods and ways to improve current technology to improve the humaneness of methods used to manage damage caused by wildlife. Cooperation with individuals and organizations involved in animal welfare continues to be an agency priority for the purpose of evaluating strategies and defining research aimed at developing humane methods.

All methods not requiring direct supervision during employment (*e.g.*, live traps) would be checked and monitored to ensure any wildlife confined or restrained were addressed in a timely manner to minimize distress of the animal. All euthanasia methods used for live-captured mammals would be applied

according to WS' directives. Shooting would occur in some situations and personnel would be trained in the proper use of firearms to minimize pain and suffering of mammals taken by this method.

WS employs methods as humanely as possible by applying measures to minimize pain and that allow wildlife captured to be addressed in a timely manner to minimize distress. Through the establishment of SOPs that guide WS in the use of methods to address damage and threats associated with mammals in the State, the cumulative impacts on the issue of method humaneness would be minimal. All methods would be evaluated continually to ensure SOPs were adequate to ensure those methods continue to be used to minimize suffering and that wildlife captured were addressed in a timely manner to minimize distress.

Issue 6 - Effects of Mammal Damage Management Activities on the Regulated Harvest of Mammals

As discussed in this EA, the magnitude of WS' mammal take for damage management purposes from FY 2006 through FY 2011 was low when compared to the total take of mammals and when compared to the estimated statewide population of those species. Since all take of mammals is regulated by the TWRA, the take by WS that would occur annually and cumulatively would occur pursuant to mammal population objectives established in the State. WS' take of mammals (combined take) annually to alleviate damage would be a minor component to the known take that occurs annually during the harvest seasons.

The populations of several mammal species are sufficient to allow for annual harvest seasons that typically occur during the fall. Hunting and trapping seasons are established by the TWRA. Those species addressed in this EA that have established harvest seasons include: beaver, muskrat, nutria, woodchuck, cottontail rabbit, black bear, raccoon, long-tailed weasel, mink, striped skunk, spotted skunk, coyote, gray fox, red fox, bobcat, opossum, white-tailed deer, and elk.

With oversight of mammal take, the TWRA maintains the ability to regulate take by WS to meet management objectives for mammals in the State. Therefore, the cumulative take of mammals would be considered as part of the TWRA objectives for mammal populations in the State.

Issue 7 – Effects of Beaver Removal and Dam Manipulation on the Status of Wetlands in the State

Beaver build dams primarily in smaller riverine streams (intermittent and perennial brooks, streams and small rivers) and in drainage areas with dams consisting of mud, sticks and other vegetative materials. Their dams obstruct the normal flow of water and typically change the pre-existing hydrology from flowing or circulating waters to slower, deeper, more expansive waters that accumulate bottom sediment. The depth of bottom sediment depends on the length of time an area is covered by water and the amount of suspended sediment in the water.

The pre-existing habitat and the altered habitat have different ecological values to the fish and wildlife native to an area. Some species would abound by the addition of a beaver dam, while others would diminish. For example, some fish species require fast moving waters over gravel or cobble beds, which beaver dams can eliminate, thus reducing the habitat's value for these species. In general, it has been found that wildlife habitat values decline around bottomland beaver impoundments because trees are killed from flooding and mast production declines. On the other hand, beaver dams can potentially be beneficial to some species of fish and wildlife such as river otter, neotropical birds, and waterfowl.

If a beaver dam is not breached and water is allowed to stand, hydric soils and hydrophytic vegetation eventually form. This process can take anywhere from several months to years depending on pre-existing conditions. Hydric soils are those soils that are saturated, flooded, or ponded long enough during the growing season to develop anaerobic conditions. In general, hydric soils form much easier where wetlands have preexisted. Hydrophytic vegetation includes those plants that grow in water or on a

substrate that is at least periodically deficient in oxygen because of excessive water content. If these conditions are met, then a wetland has developed that would have different wildlife habitat values than an area that has been more recently impounded by beaver dam activity.

The intent of most dam breaching is not to drain established wetlands. With few exceptions, requests from public and private individuals and entities that WS receives involve dam breaching to return an area back to its pre-existing condition within a few years after the dam was created. If the area does not have hydric soils, it usually takes many years for them to develop and a wetland to become established. This often takes greater than five years as recognized by the Swampbuster provisions. Most beaver dam removal by WS is either exempt from regulation under Section 404 of the CWA as stated in 33 CFR Part 323 or may be authorized under the United States Army Corps of Engineers Nationwide Permit System in 33 CFR Part 330.

However, the breaching of some beaver dams can trigger certain portions of Section 404 that require landowners to obtain permits from the United States Army Corps of Engineers. WS' personnel determine the proper course of action upon inspecting a beaver dam impoundment.

It should also be noted that beaver created wetlands are dynamic and do not remain in one state for indefinite periods. Large beaver ponds may eventually fill with sediment and create a beaver meadow. Beaver may be removed from an area due to natural predation or they may abandon an area due to lack of food. Once a dam is abandoned, it is subject to natural decay and damage due to weather. The dam would eventually fail and the wetland would return to a flowing stream or brook. WS' beaver management activities may accelerate or modify these natural processes by removing beaver and restoring or increasing water flow; however, they are generally processes that would occur naturally over time.

Muskrat management would usually be intended to maintain or protect existing wetlands by reducing threats to natural and man-made wetlands and associated floral, faunal and T&E communities. Wetlands are often created by natural or man-made dams, dikes, levees and berms that contain standing water or control drainage, particularly after precipitation events that could result in flooding. Muskrat burrowing activity can degrade the integrity of these structures by allowing water infiltration or by causing erosion by feeding on vegetation intended to stabilize dirt structures. Muskrats are omnivores and feed on a variety of aquatic and terrestrial plants and aquatic animals. At high population densities, they may disrupt or damage natural wetland floral and faunal communities or they may feed on T&E species. WS activities would be intended to protect existing wetlands from damage caused by muskrats.

Therefore, the activities of WS to manage flooding damage by manipulating beaver dams would not be expected to have any cumulative adverse effects on wetlands in Tennessee when conducted in accordance with the CWA and the Swampbuster provision of the Food Security Act.

CHAPTER 5: LIST OF PREPARERS AND PERSONS CONSULTED

5.1 LIST OF PREPARERS

Brett Dunlap, USDA/APHIS/WS, State Director, Madison, TN
Keith Wehner, USDA/APHIS/WS, Assistant State Director, Madison, TN
Ryan Wimberly, USDA/APHIS/WS, Environmental Management Coordinator, Madison, TN

5.2 LIST OF PERSONS CONSULTED

Gray Anderson, TWRA, Assistant Chief of Wildlife, Nashville, TN

Wesley James, TVA, Senior Specialist – Natural Resources Management, Knoxville, TN
Holly LeGrand, TVA, Biologist/Zoologist, Knoxville, TN
Loretta McNamee, TVA, Contract NEPA Specialist, Knoxville, TN

APPENDIX A
LITERATURE CITED

Agency for Toxic Substances and Disease Registry. 1998. Public Health Assessments and Consultations. http://www.atsdr.cdc.gov/hac/pha/batg/bat_p1.html. Website accesses 2 March 2012.

Air National Guard. 1997. Final environmental impact statement for the Colorado Airspace Initiative, Vol. 1. Impact Analyses. National Guard Bureau, Andrews Air Force Base, Maryland.

ABC. 2011. Domestic cat predation on birds and other wildlife. http://www.abcbirds.org/abcprograms/policy/cats/materials/CatPredation2011.pdf. Accessed July 23, 2012.

American Rivers. 2011. American Rivers homepage. <http://www.americanrivers.org>. Accessed November 7, 2011.

AVMA. 1987. Panel report on the colloquium on recognition and alleviation of animal pain and distress. Journal of the American Veterinary Medical Association 191:1186-1189.

AVMA. 2003. Position statement on abandoned and feral cats. AVMA Membership Directory and Resource Manual. Shaumburg, Illinois, 2003:74.

AVMA. 2004. Animal welfare forum: management of abandoned and feral cats. Journal of the American Veterinary Medical Association. Vol. 225, No. 9, November 1, 2004.

AVMA. 2007. AVMA guidelines on euthanasia. American Veterinary Medical Association. http://www.avma.org/issues/animal_welfare/euthanasia.pdf. Accessed February 2, 2009.

AVMA. 2009. Position on abandoned and feral cats. http://www.avma.org/issues/policy/animal_welfare/feral_cats.asp. Accessed May 9, 2012.

Ames, D. R., and L. A. Arehart. 1972. Physiological response of lambs to auditory stimuli. Journal of Animal Science 34:997-998.

Anderson, E.M. 1987. A critical review and annotated bibliography of literature on the bobcat. Special Report Number 62, Colorado Division of Wildlife, Denver, Colorado.

Andersen, D. E., O. J. Rongstad, and W. R. Mytton. 1989. Response of nesting red-tailed hawks to helicopter overflights. Condor 91:296-299.

Anderson, D.W., J.O. Kieth, G.R. Trapp, F. Gress, and L.A. Moreno. 1989. Introduced small ground predators in California brown pelican colonies. Colonial Waterbirds 12:98-103.

Andrews, E. J., B. T. Bennett, J. D. Clark, K. A. Houpt, P. J. Pascoe, G. W. Robinson, and J. R. Boyce. 1993. Report of the AVMA panel on euthanasia. Journal of the American Veterinary Medical Association 202:229-249.

Apa, A. D., D. W. Uresk, and R. L. Linder. 1991. Impacts of black-tailed prairie dog rodenticides on nontarget passerines. Great Basin Naturalist 51:301-309.

Applegate, R.D. 2010. Results from online game harvest survey 2008-2009. <http://www.tn.gov/twra/pdfs/smallgamereport.pdf.> Accessed November 7, 2011.

Archer, J. 1999. The nature of grief: the evolution and psychology of reactions to loss. Taylor & Francis/Routledge, Florence, Kentucky.

Armitage, K. B. 2003. Marmots (*Marmota monax* and allies). Pp 188-210 *in* G. A. Feldhamer, B. C. Thompson, and J. A. Chapman, eds. Wild mammals of North America: biology, management, and conservation. Second edition. The Johns Hopkins University Press, Baltimore, Maryland.

Arner, D. H. 1964. Research and a practical approach needed in management of beaver and beaver habitat in the Southeastern United States. Transactions of the North American Wildlife Conference 29:150-158.

Arner, D. H., and J. S. DuBose. 1982. The impact of the beaver on the environment and economics in the southeastern United States. International Congress of Game Biologists 14:241-247.

Arner, D. H., and G. R. Hepp. 1989. Beaver pond wetlands: A southern perspective. Pp 117-128 *in* L. M. Smith, R. L. Pederson, and R. M. Kaminski, eds. Habitat management for migrating and wintering waterfowl in North America. Texas Tech University Press, Lubbock, Texas.

Avery, E. L. 1992. Effects of removing beaver dams upon a northern Wisconsin brook trout stream. Wisconsin Department of Natural Resources Final D-J Project Report, F-83R, 185 pp.

Avery, E. L. 2004. A compendium of 58 trout stream habitat development evaluations in Wisconsin – 1985-2000. Wisconsin Department of Natural Resources Research Report 187. 96pp.

Awbrey, F. T., and A. E. Bowles. 1990. The effects of aircraft noise and sonic booms on raptors: a preliminary model and a synthesis of the literature on disturbance. Noise and Sonic Boom Impact Technology, Technical Operating Report 12. Wright-Patterson Air Force Base, Ohio.

Bailey, T. N. 1972. Ecology of bobcats with special reference to social organization. Thesis, University of Idaho, Moscow.

Baker, B. W., and E. P. Hill. 2003. Beaver (*Castor canadensis*). Pages 288-310 *in* G. A. Feldhamer, B. C. Thompson, and J. A. Chapman, editors. Wild mammals of North America: biology, management, and conservation. Second edition. The Johns Hopkins University Press, Baltimore, Maryland.

Baker, R. H., and M. W. Baker. 1975. Montane habitat used by the spotted skunk (*Spilogale putorius*) in Mexico. Journal of Mammalogy, 56: 671-673.

Barrett, R.H. and G.H. Birmingham. 1994. Wild pigs. Pp D65-70 *in* S. E. Hygnstrom, R. M. Timm and G. E. Larson, eds. Prevention and control of wildlife damage. University of Nebraska-Lincoln, Nebraska.

Barrows, P.L. 2004. Professional, ethical, and legal dilemmas of trap-neuter-release. Journal of the American Veterinary Medical Association 225:1365–1369.

Beaver, B. V., W. Reed, S. Leary, B. McKiernan, F. Bain, R. Schultz, B. T. Bennett, P. Pascoe, E. Schull, L. C. Cork, R. Francis-Floyd, K. D. Amass, R. Johnson, R. H. Schmidt, W. Underwood, G. W.

Thorton, and B. Kohn. 2001. 2000 Report of the AVMA panel on euthanasia. Journal of the American Veterinary Medical Association 218:669-696.

Beach, R. 1993. Depredation problems involving feral hogs. Pp. 67-75 *in* C.W. Hanselka and J.F. Cadenhead, eds. Feral swine: a compendium for resource managers. Texas Agric. Ext. Serv., College Station, Texas.

Bekoff, M. 1982. Coyote. Pp 447–459 *in* J. A. Chapman and G. A. Feldhamer, eds. Wild mammals of North America: biology, management, and economics. Johns Hopkins University Press, Baltimore, Maryland.

Belanger, L., and J. Bedard. 1989. Responses of staging greater snow geese to disturbance. Journal of Wildlife Management 53:713-719.

Belanger, L., and J. Bedard. 1990. Energetic cost of man-induced disturbance to staging snow geese. Journal of Wildlife Management. 54:36-41.

Bellrose, F.C. 1976. Ducks, geese and swans of North America. Stackpole, Harrisburg, Pennsylvania.

Beran, G. W. 1994**.** Handbook of zoonoses. CRC Press, Boca Raton, Florida.

Bergman, D., D. Nolte, and J. Taylor. (comp.). 2007. Proceedings of the International Beaver Ecology and Management Workshop. USDA, APHIS, WS, Arizona State Office, 8836 North 23rd Ave. Suite 2, Phoenix AZ 85021.

Berryman, J.H. 1991. Animal damage management: Responsibilities of various agencies and the need for coordination and support. Proc. East. Wildl. Damage Control Conf. 5:12-14.

Berthier, K., M. Langlais, P. Auger, and D. Pontier. 2000. Dynamics of a feline virus with two transmission modes within exponentially growing host populations. Proc. Biol. Sci. 267: 2049–2056.

Bevan, D.J., K.P. Chandroo, and R.D. Moccia. 2002. Predator control in commercial aquaculture in Canada. http://www.aps.uoguelph.ca/aquacentre/files/misc-factsheets/Predator%20Control%20in%20Commercial%20Aquaculture%20in%20Canada.pdf. Accessed March 29, 2012.

Bishop, R. C. 1987. Economic values defined. Pages 24 -33 *in* D. J. Decker and G. R. Goff, editors. Valuing wildlife: economic and social perspectives. Westview Press, Boulder, Colorado.

Blundell, G.M., J.W. Kern, R.T. Bowyer, and L.K. Duffy. 1999. Capturing river otters: a comparison of Hancock and leg-hold traps. Wildlife Society Bulletin 27:184-192.

Boggess, E. K. 1994*a*. Raccoons. Pp C101-C107 *in* S. E. Hygnstrom, R. M. Timm and G. E. Larson, eds. Prevention and control of wildlife damage. University of Nebraska-Lincoln, Lincoln, Nebraska.

Boggess, E. K. 1994*b*. Mink. Pp C89-C92 *in* S. E. Hygnstrom, R. M. Timm and G. E. Larson, eds. Prevention and control of wildlife damage. University of Nebraska-Lincoln, Lincoln, Nebraska.

Bollengier, R. M., Jr. 1994. Woodchucks. Pp B183-B187 *in* S. E. Hygnstrom, R. M. Timm, and G. E. Larson, eds. Prevention and control of wildlife damage. University of Nebraska-Lincoln.

Borg, E. 1979. Physiological aspects of the effects of sound on man and animals. Acta Oto-laryn-gologica, Supplement 360:80-85.

Bounds, D. L. 2000. Nutria: An invasive species of national concern. Wetland Journal 12:9-16.

Boutin, S., and D. E. Birkenholz. 1987. Muskrat and round-tailed muskrat. Pp 282-313 *in* M. Novak, J. A. Baker, M. E. Obbard, and B. Mallock, edis. Wild furbearer management and conservation in North America. Ministry of Natural Resources, Ontario, Canada.

Bouwer, H. 1989. The Bouwer and Rice slug test--an update. Ground Water 27:304-309.

Boyce, P. S. 1998. The social construction of bereavement: an application to pet loss. Thesis, University of New York.

Boyles, S. L. 2006. Report on the efficacy and comparative costs of using flow devices to resolve conflicts with North American beavers along roadways in the Coastal Plain of Virginia. M.S. Thesis, Christopher Newport University, Newport News, Virginia. 48 pp.

Boyles, S., and S. Owens. 2007. North American Beaver (*Castor canadensis*): A technical conservation assessment. [Online]. USDA Forest Service, Rocky Mountain Region. Available: http://www.fs.fed.us/r2/projects/scp/assessments/northamericanbeaver.pdf. Accessed on August 11, 2010.

Brickner, I. 2003. The impact of domestic cat (*Felis catus*) on wildlife welfare and conservation: A literature review. <http://www.tau.ac.il/lifesci/zoology/members/yom-tov/inbal/cats.pdf>. Accessed February 10, 2012.

Bromley, P. T., J. F. Heisterberg, W. T. Sullivan, Jr., P. Sumner, J. C. Turner, R. D. Wickline, and D. K. Woodward. 1994. Wildlife damage management: beavers. North Carolina Cooperative Extension Service.

Brown, S. T., and J. W. Brown. 1999. How to control beaver flooding. and inserts, “Beaver flex pipe and cage”, and “How to protect trees from beavers,” Beavers Wetlands and Wildlife, Dolgeville, New York 13329.

Brown, S., D. Shafer, and S. Anderson. 2001. Control of beaver flooding at restoration projects. WRAP Techical Notes Collection (ERDC TN-WRAP-01-01), U.S. Army Engineer Research and Development Center, Vicksburg MS. www.wes.army.mil/el/wrap.

Buckley, M., T. Souhlas, E. Niemi, E. Warren, and S. Reich. 2011. The economic value of beaver ecosystem services: Escalante River Basin, Utah. ECONorthwest, Eugene, Oregon. 66 pp.

Burt, W. H., and R. P. Grossenheider. 1976. A field guide to the mammals. Houghton Mifflin College, Boston, Massachusetts. 289 pp.

California Department of Fish and Game. 1991. Final environmental document - bear hunting. Title 14 Calif. Code of Regs. Calif. Dept. of Fish and Game, State of California, April 25, 1991. 337 pp.

Callahan, M. 2003. Beaver management study. Association of Massachusetts Wetland Scientists Newsletter 44:12-15.

Callahan, M. 2005. Best management practices for beaver problems. Association of Massachusetts Wetland Scientists Newsletter 53:12-14.

Camenzind, F.J. 1978. Behavioral ecology of coyotes on the National Elk Refuge, Jackson Wyoming. Pp 267-294 *in* M. Bekoff, ed. Coyotes: biology, behavior and management. Academic Press, New York, New York.

Canadian Broadcast Company. 2009. Coyotes kill Toronto singer in Cape Breton. http://www.cbc.ca/news/canada/nova-scotia/story/2009/10/28/ns-coyote-attack-died.html. Accessed November 23, 2011.

Carter, J., and B. P. Leonard. 2002. A review of the literature on the worldwide distribution, spread of, and efforts to eradicate the coypu (*Myocastor coypus*). Wildlife Society Bulletin 30:162-175.

Casey, D., and D. Hein. 1983. Effects of heavy browsing on a bird community in deciduous forest. Journal of Wildlife Management 47:829-36.

Castillo, D., and A. L. Clarke. 2003. Trap/neuter/release methods ineffective in controlling domestic cat "colonies' on public lands. Natural Areas Journal 23:247-253.

CDC. 1990. Compendium of rabies control. Morbidity and mortality weekly report. 39 Number RR-4:6, Atlanta, Georgia.

CDC. 1999. Mass treatment of humans who drank unpasteurized milk from rabid cows – Massachusetts, 1996-1998. CDC – Morbidity and Mortality Weekly Report. 48:228-229.

CDC. 2011. Rabies. http://www.cdc.gov/rabies/index.html. Accessed July 23, 2012.

Chamberlain, M. J., and B. D. Leopold. 2001. Omnivorous furbearers. Pages 278-292 *in* J. G. Dickson editor. Wildlife of southern forests: habitat and management. Hancock House Publishers, Blaine, Washington.

Chapman, J. A., and G. A. Feldhamer. 2003. Wild mammals of North America. John Hopkins University Press, Baltimore, Maryland.

Childs, J.E. 1986. Size dependent predation on rats by house cats in an urban setting. Journal of Mammalogy 67:196-198.

Childs, J.E. 1991. And the cats shall lie down with the rat. Natural History, June 100:16-19.

Churcher, P. B., and J. H. Lawton. 1987. Predation by domestic cats in an English village. Journal of Zoology (London) 212:439-455.

Churcher, P. B., and J. H. Lawton. 1989. Beware of well-fed felines. Natural History 7:40-46.

Clark, F.W. 1972. Influence of jackrabbit density on coyote population change. Journal of Wildlife Management 36:343-356.

Clausen, G., and A. Ersland. 1970. Blood O_2 and acid-base changes in the beaver during submersion. Respiration Physiology 11:104-112.

Clemson University. 2006. Clemson beaver pond leveler Version II. DVD. Clemson University Extension, Clemson, South Carolina.

Close, T. L. 2003. Modifications to the Clemson pond leveler to facilitate brook trout passage. Minnesota Department of Natural Resources Special Publication 158. 9 pp.

Cole, R., and G. O. Batzli. 1979. Nutrition and population dynamics of the prairie vole *Microtus ochrogaster* in central Illinois. Applied Ecology 48:455-470.

Coleman, J. S., and S. A. Temple. 1989. Effects of free ranging cats on wildlife: a progress report. Proceedings of the Eastern Wildlife Damage Control Conference 4:9-12.

Coleman, J. S., S. A. Temple, and S. R. Craven. 1997. Facts on cats and wildlife: a conservation dilemma. Miscellaneous Publications. USDA Cooperative Extension, University of Wisconsin, Madison, Wisconsin.

Coman, B.J., and H.B. Brunner. 1972. Food habits of the feral house cat in Victoria. Journal of Wildlife Management 36:848-853.

Conner, P.F. 1960. The small mammals of Otsego and Schoharie Counties, New York. Bull. New York State Mus. And Sci. Serv. 382:1-84.

Connolly, G.E. 1978. Predator control and coyote populations: A review of simulation models. Pp 327-345 *in* M. Bekoff, ed. Coyotes: Biology, Behavior and Management. Academic Press, New York. 384 pp.

Connolly, G.E. 1992. Coyote damage to livestock and other resources. Pp 161-169 *in* A.H. Boer, ed. Ecology and management of the eastern coyote. Wildlife Research Unit. University of New Brunswick, Fredericton, New Brunswick, Canada.

Connolly, G.E., and W.M. Longhurst. 1975. The effects of control on coyote populations. Univ. Calif., Div. Agric. Sci. Bull. 1872. 37 pp.

Conomy, J. T., J. A. Dubovsky, J. A. Collazo, and W. J. Fleming. 1998. Do black ducks and wood ducks habituate to aircraft disturbance? Journal of Wildlife Management 62:1135-1142.

Conover, M. R. 1982. Comparison of two behavioral techniques to reduce bird damage to blueberries: methiocarb and hawk-kite predator model. Wildlife Society Bulletin 10:211-216.

Conover, M. R. 1997. Monetary and intangible valuation of deer in the United States. Wildlife Society Bulletin 23:298-305.

Conover, M. R., W. C. Pitt, K. K. Kessler, T. J. DuBow, and W. A. Sanborn. 1995. Review of human injuries, illness, and economic losses caused by wildlife in the United States. Wildlife Society Bulletion 23:407-414.

Costanza, R., R. d'Arge, R. de Groot, S. Farber, M. Grasso, B. Hannon, K.Limburg, S. Naeem, R. V. O'Neill, J.Paruelo, R G. Raskin, P. Sutton, and M.van den Belt. 1997. The value of the world's ecosystem services and natural capital. Nature 387:253-260.

Corn, J. L., P. K. Swiderek, B. O. Blackburn, G. A. Erickson, A. B. Thiermann, and V. F. Nettles. 1986. Survey of selected diseases in wild swine in Texas. J. Am. Vet. Med. Assoc. 189:1029-1032.

Courchamp F., J. L. Chapuis, and M. Pascal. 2003. Mammal invaders on islands: impact, control and control impact. Biological Review 78:347-383.

Crabb, W.D. 1941. Food habits of the prairie spotted skunk in southeastern Iowa. Journal of Mammalogy 22:349-364.

Crabb, W. D. 1948. The ecology and management of prairie spotted skunk in Iowa. Ecological Monographs 18:201-232.

Craig, J. R., J. D. Rimstidt, C. A. Bonnaffon, T. K. Collins, and P. F. Scalon. 1999. Surface water transport of lead at a shooting range. Bulletin of Environmental Contamination and Toxicology 63:312-319.

Craven, S. R. 1994. Cottontail rabbits. Pp D75-D80 *in* S. E. Hygnstrom, R. M. Timm and G. E. Larson, eds. Prevention and control of wildlife damage. University of Nebraska-Lincoln, Lincoln, Nebraska.

Craven, S. R., and S. E. Hygnstrom. 1994. Deer. Pp D25-D40 *in* S. E. Hygnstrom, R. M. Timm and G. E. Larson, eds. Prevention and control of wildlife damage. University of Nebraska-Lincoln, Lincoln, Nebraska.

Crawford, B.A., M.R. Pelton, and K.G. Johnson. 1993. Techniques to monitor relative abundance of coyotes in East Tennessee. Proceedings of Annual Conference, Southeastern Association of Fish and Wildlife Agencies 47:62-70.

Crowe, D. M. 1975. A model for exploited bobcat populations in Wyoming. Journal of Wildlife Management 39:408-415.

Crum, J.M. 2003. Non-seasonal mortality white-tailed deer. West Virginia Division of Natural Resources. http://www.wvdnr.gov/Hunting/DeerNSeasMortal.shtm. Accessed April 6, 2012.

Dahl, T.E. 1990. Wetland losses in the United States, 1780s to 1980s. United States Department of Interior, United States Fish and Wildlife Service, Washington, D.C. 21 pp.

Danner, D.A., and N.S. Smith. 1980. Coyote home range, movement, and relative abundance near a cattle feedyard. Journal of Wildlife Management 44:484-487.

Dauphine, N., and R. J. Cooper. 2009. Impacts of free-ranging domestic cats (*Felix catus*) on birds in the United States: A review of recent research with conservation and management recommendations. Proceedings of the fourth international partners in flight conference: tundra to tropics., McAllen, Texas. http://www.abcbirds.org/abcprograms/policy/cats/pdf/impacts_of_free_ranging_domestic_cats.pdf. Accessed July 23, 2012.

Davidson, W. R. 2006. Field manual of wildlife diseases in the southeastern United States. Third edition. The University of Georgia, Athens. 448 pp.

Davidson, W. R. and V. F. Nettles. 1997. Field manual of wildlife diseases in the southeastern United States. 2nd edition. University of Georgia, Athens, Georgia. 417 pp.

DeBenedetti, S. H. 1986. Management of feral pigs at Pinnacles National Monument: why and how. Proceedings of the conference on the conservation and management of rare and endangered plants. California Native Plant Society, Sacramento, California.

deCalesta, D. S. 1997. Deer and ecosystem management. Pp 267-279 *in* W. J. McShea, H. B. Underwood, and J. H. Rappole, eds. The science of overabundance: deer ecology and population management. Smithsonian Institute Press, Washington, D.C.

deCalesta, D. S., and G. W. Witmer. 1994. Elk. Pp D41-D50 *in* S. E. Hygnstrom, R. M. Timm and G. E. Larson, eds. Prevention and control of wildlife damage. University of Nebraska-Lincoln, Lincoln, Nebraska.

Decker, D. J., and G. R. Goff. 1987. Valuing wildlife: economic and social perspectives. Westview Press, Boulder, Colorado.

Decker, D. J., and K. G. Purdy. 1988. Toward a concept of wildlife acceptance capacity in wildlife management. Wildlife Society Bulletin 16:53-57.

Decker, D. J., K. M. Lonconti Lee, and N. A. Connelly. 1990. Incidence and costs of deer-related vehicular accidents in Tompkins County, New York. HDRU Series 89-7, revised Feb 1990. Human Dimensions Research Unit, Department of Natural Resources, New York State College Agriculture and Life Sciences, Cornell University, Ithaca, New York.

Decker, D. J., and L. C. Chase. 1997. Human dimensions of living with wildlife – a management challenge for the 21st century. Wildlife Society Bulletin 28:4-15.

Deisch, M. S. 1986. The effects of three rodenticides on nontarget small mammals and invertebrates. Thesis, South Dakota State University, Brookings, South Dakota.

Deisch, M. S., D. W. Uresk, and R. L. Linder. 1989. Effects of two prairie dog rodenticides on ground-dwelling invertebrates in western South Dakota. Pages 166-170 *in* A. J. Bjugstad, D. W. Uresk, and R. H. Hamre, editors. Ninth Great Plains wildlife damage control workshop proceedings. USDA Forest Service General Technical Report RM-171. Fort Collins, Colorado.

Deisch, M. S., D. W. Uresk, and R. L. Linder. 1990. Effects of prairie dog rodenticides on deer mice in western South Dakota. Great Basin Naturalist 50:347-353.

Delaney, D. K., T. G. Grubb, P. Beier, L. L. Pater, and M. H. Reiser. 1999. Effects of helicopter noise on Mexican spotted owls. Journal of Wildlife Management 63:60-76.

Devault, T.L., J.C. Beasley, L.A. Humberg, B.J. MacGowan, M.I. Retamosa, and O.E. Rhodes, Jr. 2007. Intrafield patterns of wildlife damage to corn and soybean in northern Indiana. Human-Wildlife Conflicts 1:179-187.

Doebel, J.H., and B.S. McGinnes. 1974. Home range and activity of a gray squirrel population. Journal of Wildlife Management 38:860-867.

Dolbeer, R.A. 1998. Population dynamics: the foundation of wildlife damage management for the 21st century. Pp. 2-11 *in* Barker, R. O. and Crabb, A. C., Eds. Eighteenth Vertebrate Pest Conference (March 2-5, 1998, Costa Mesa, California). University of California at Davis, Davis, California.

Dolbeer, R. A. 2000. Birds and aircraft: fighting for airspace in crowded skies. Proceedings of the Vertebrate Pest Conference 19:37-43.

Dolbeer, R.A. 2009. Birds and aircraft: fighting for airspace in ever more crowded skies. Human-Wildlife Conflicts 3:165-166.

Dolbeer, R. A., G. E. Bernhardt, T. W. Seamans and P. P. Woronecki. 1991. Efficacy of two gas cartridge formulations in killing woodchucks in burrows. Wildlife Society Bulletin 19:200-204.

Dolbeer, R. A., N. R. Holler, and D. W. Hawthorne. 1994. Identification and control of wildlife damage. Pp 474-506 *in* T. A. Bookhout, ed. Research and management techniques for wildlife and habitats. The Wildlife Society, Bethesda, Maryland.

Dolbeer, R.A., S.E. Wright, J. Weller, and M.J. Beiger. 2012. Wildlife Strikes to Civil Aircraft in the United States, 1990–2010. U.S. Department of Transportation, Federal Aviation Administration, Office of Airport Safety and Standards, Serial Report No. 17, Washington, D.C.

Drake, D., J. B. Paulin, P. D. Curtis, D. J. Decker, and G. J. San Julian. 2005. Assessment of negative economic impacts from deer in the northeastern United States. Journal of Extension 43(1), Article Number 1RIB5.

Dubey, J.P. 1973. Feline toxoplasmosis and coccidiosis: a survey of domiciled and stray cats. Journal of American Veterinary Medical Association 162:873-877.

Dubey, J.P. , R.M. Weigel, A.M. Siegel, P. Thulliez, U.D. Kitron, M.A. Mitchell, A. Mannelli, N.E. Mateus-Pinilla, S.K. Shen, O.C.H. Kwok, and K.S. Todd. 1995. Sources and reservoirs of *Toxoplasma gondii* infection on 47 swine farms in Illinois. J. Parasitol. 81:723-729.

Eadie, W.R. 1939. A contribution to the biology of *Parascalops breweri*. Journal of Mammalogy 20:150-173.

Eadie, W.R. 1948. C*orpora amylacea* in the prostatic secretion and experiments on the formation of a copulatory plug in some insectivores. Anat. Rec. 102:259-271.

Eagle, T. C., and J. S. Whitman. 1987. Mink. Pp 614-625 *in* M. Novak, J. A. Baker, M. E. Obbard, B. Mallock, eds. Wild furbearer management and conservation in North America. Ministry of Natural Resources, Ontario, Canada.

Eisenberg, J.F. 1989. Mammals of the Neotropics. The Northern Neotropics. The University of Chicago Press, Chicago, Illinois.

Ellis, D. H. 1981. Responses of Raptorial Birds to low level military jets and sonic booms: Results of the 1980-1981 Joint U.S. Air Force-U.S. Fish and Wildlife Service Study. Prepared by the Institute for Raptor Studies for USAF and USFWS. NTIS No. ADA 108-778.

Emmons, L.H., and F. Feer. 1990. Neotropical Rainforest Mammals: A Field Guide. University of Chicago Press, Chicago, Illinois.

EPA. 1986*a*. Quality Criteria for Water 1986. U.S. Environmental Protection Agency, Publication EPA/440/5-86-001. Washington, D.C.

EPA. 1986*b*. Total Exposure Assessment Model (TEAM) Study: Summary and Analysis, Volume I, Final Report. EPA/600/6-87/002a. Washington, D.C.

EPA. 1991. Reregistration eligibility document: Inorganic nitrate/nitrite (sodium and potassium nitrates). List D, Case 4052. Environmental Protection Agency, Office of Pesticide Programs Special Review and Reregistration Division, Washington, D.C.

EPA. 1995. Facts about wetlands. United States Environmental Protection Agency. Office of water, wetlands, oceans, and watersheds (4502F), Environmental Protection Agency 843-F-95-00le.

EPA. 1998. Reregistration Eligibility Decision (RED): Zinc phosphide. United States Environmental Protection Agency, Office of Pesticide Programs Special Review and Reregistration Division, Washington, D.C. 207 pp.

EPA. 2000. Introduction to phytoremediation. EPA/600/R-99/107, Office of Research and Development, Washington, D.C.

EPA. 2001. Selected mammal and bird repellents fact sheet. http://www.epa.gov/oppbppd1/biopesticides/ingredients/factsheets/factsheet_mam-bird-repel.htm. Accessed March 2, 2012.

EPA. 2009. Pesticide fact sheet: Mammalian Gonadotropin releasing hormone (GnRH). United States Environmental Protection Agency, Office of Prevention, Pesticides, and Toxic Substances. Arlington, Virginia.

Erickson, D.W., C.R. McCullough, and W.R. Poranth. 1984. Evaluation of experimental river otter reintroducutions. Final Report of Missouri Department of Conservation. Federal Aid Project. No. W-13-R-38. 47 pp.

Errington, P.L. 1933. Bobwhite winter survival in an area heavily populated with grey fox. Iowa State College Journal of Science 8:127–130.

Errington, P.L. 1943. An analysis of mink predation upon muskrats in north-central United States. Iowa Agricultural Experiment Station Research Bulletin 320:797-924.

Espinoza, H.R., and Rowe, F.P. 1979. Biology and control of the cotton rat (*Sigmodon hipidus*). International Journal of Pest Management 25:251-6.

Evans, J. 1970. About nutria and their control. USDI, Bureau of Sport Fisheries and Wildlife, Resource Publication 86.

Fancy, S.G. 1982. Reaction of bison to aerial surveys in interior Alaska. Canadian Field Naturalist 96:91.

FAA. 2010. FAA National Wildlife Aircraft Strike Database 2010. US Department of Transportation, Federal Aviation Administration, 800 Independence Avenue, SW, Washington, D.C..

Federal Emergency Management Agency. 2005. Dam Owner's Guide to Animal Impacts on Earthen Dams. FEMA L-264.

Fergus, C. 2006. Cottontail rabbit. Wildlife Note – 4, LDR0103. Pennsylvania Game Commission, Bureau of Information and Education, Harrisburg, Pennsylvania.

Fernandez, S. 2008. Ticked off: Deer, Lyme Disease connected? Greenwich Post. September 4, 2008.

Figley, W.K., and L.W. VanDruff. 1982. The Ecology of Urban Mallards. Wildl. Monogr. 81. 40 pp.

Fitzgerald, B. M. 1977. Weasel predation on a cyclic population of the montane vole (*Microtus montanus*) in California. Journal of Animal Ecology 46:367-97.

Fitzgerald, B.M. 1990. House cat. Pp. 330-348 *in* C. M. King, ed. The handbook of New Zealand mammals. Auckland, Oxford University Press.

Fitzgerald, B.M., W.B. Johnson, C.M. King, and P.J. Moors. 1984. Research on Mustelids and cats in New Zealand. WRLG Res. Review No. 3. Wildl. Res. Liaison Group, Wellington. 22 pp.

Fitzwater, W. D. 1994. Feral house cats. Pp C45-50 *in* S. E. Hygnstrom, R. M. Timm and G. E. Larson, eds. Prevention and control of wildlife damage. University of Nebraska-Lincoln, Lincoln, Nebraska.

Flyger, V., and J. E. Gates. 1982. Fox and gray squirrels. Pp 209-229 *in* J. A. Chapman and G. A. Feldhamer, eds. Wild mammals of North America: biology, management, and economics. The John Hopkins University Press, Baltimore, Maryland.

Follmann, E. H. 1973. Comparative ecology and behavior of red and gray fox. Thesis, Southern Illinois University-Carbondale, Carbondale, Illinois.

Food and Drug Administration. 2003. Bird poisoning of federally protected birds. Office of Criminal Investigations. Enforcement Story 2003.

Forrester, D. J. 1991. Parasites and diseases of wild mammals in Florida. University of Florida Press, Gainesville, Florida.

Fouty, S. 2008*a*. Climate change and beaver activity: How restoring nature's engineers can alleviate problems. Beaversprite Spring:4-5, 13.

Fouty, S. C. 2008*b*. Current and historic stream channel response to changes in cattle and elk grazing pressure and beaver activity: southwest Montana and east-central Arizona. Ph.D. Dissertation, Department of Geography, University of Oregon, Corvallis, Oregon.

Fowler, M. E., and R. E. Miller. 1999. Zoo and wild animal medicine. W.B. Saunders Company, Philadelphia, Pennsylvania.

Fowler, J. F., S. Verma, and T. Delphin. 1994. An impact assessment of beaver damage in Louisiana during 1993. Report to the Louisiana State Natural Resources Committee.

Francl, K., and D. Meikle. 2009. A range extension of the hispid cotton rat (*Sigmodon hispidus*) in Virginia. Banisteria 33:54-55.

Fraser, D., J. F. Gardner, G. B. Kolenosky, and S. M. Strathearn. 1982. Estimation of harvest rate of black bears from age and sex data. Wildlife Society Bulletin 10:53-57.

Fraser, J.D., L.D. Frenzel, and J.E. Mathisen. 1985. The impact of human activities on breeding bald eagles in north-central Minnesota. Journal of Wildlife Management 49:585-592.

Fritzell, E. K. 1987. Gray fox and island fox. Pp 408-420 *in* M. Novak, J. A. Baker, M. E. Obbard, B. Mallock, eds. Wild furbearer management and conservation in North America. Ministry of Natural Resources, Ontario, Canada.

Frost, C. C. 1993. Four centuries of changing landscape patterns in the longleaf pine ecosystem. Pp 17-37 *in* S. M. Hermann, ed. The longleaf pine ecosystem: ecology, restoration, and management. Proceedings of the 18th Tall Timbers Fire Ecology Conference, Tallahassee, Florida.

Fuller, M. R., and J. A. Mosher. 1981. Raptor survey techniques. Pages 37-65 *in* B. A. Giron Pendleton, B.A Millsap, K. W. Cline, and D. M. Bird, editors. Raptor management techniques manual. National Wildlife Federation, Washington, D.C.

Fuller, T. K., W. E. Berg, and D. W. Kuehn. 1985. Survival rates and mortality factors of adult bobcats in north-central Minnesota. Journal of Wildlife Management 49:292-296

Fur Institute of Canada. 2000. Traps meeting requirements of Agreement on International Humane Trapping Standards. Press Release.

Gardner, A.L. 1982. Virginia opossum. Pp 3-36 *in* J. A. Chapman and G. A. Feldhamer, eds. Wild mammals of North America: biology, management, and economics. Johns Hopkins University Press, Baltimore, Maryland.

George, W.G. 1974. Domestic cats as predators and factors in winter shortages of raptor prey. Wilson Bulletin 86:384-396.

Gerell, R. 1971. Population studies on mink, *Mustela vison* Schreber, in southern Sweden. Viltrevy 8:83-114.

Gerwolls, M. K., and S. M. Labott. 1994. Adjustment to the death of a companion animal. Anthrozoos 7:172-187.

Gier, H.T. 1948. Rabies in the wild. Journal of Wildlife Management 12:142–153.

Gier, H.T. 1968. Coyotes in Kansas, revised edition. Kansas State College of Agriculture, Experimental Station Bulletin 393, Manhattan, Kansas.

Gilbert, B. 1995. "The 'little armored thing' doesn't get by on looks alone." Smithsonian. 142-151.

Gilbert, F. F., and N. Gofton. 1982. Terminal dives in mink, muskrat, and beaver. Physiology and Behavior 28:835-840.

Gilmer, D. S., L. M. Cowardin, R. L. Duval, L. M. Mechlin, C. W. Shaiffer, and V. B. Kuechle. 1981. Procedures for the use of aircraft in wildlife biotelemetry studies. U.S. Fish and Wildlife Service Resource Publication 140.

Gionfriddo, J.P., J.D. Eisemann, K.J. Sullivan, R.S. Healey, L.A. Miller, K.A. Fagerstone, R.M. Engeman, and C.A. Yoder. 2009. Field test of single-injection gonadotrophin-releasing hormone immunocontraceptive vaccine in female white-tailed deer. Wildlife Research 36:177-184.

Gipson, P.S. 1983. Evaluations of behavior of feral dogs in interior Alaska, with control implications. Vertebrate Pest Control Management Materials 4th Symposium. American Society for Testing Materials 4:285-294.

Gladwin, D. N., D. A. Asherin, and K. M. Manci. 1988. Effects of aircraft noise and sonic booms on fish and wildlife. U.S. Fish and Wildlife Service National Ecology Research Center Report 88/30.

Glover, F.A. 1943. A study of the winter activities of the New York weasel. Pennsylvania Game News 14:8–9.

Glover, F.A. 1942. A population study of weasels in Pennsylvania. Thesis, Pennsylvania State University, University Park, Pennsylvania.

Glueck, T. F., W. R. Clark, and R. D. Andrews. 1988. Raccoon movement and habitat use during the fur harvest season. Wildlife Society Bulletin 16:6-11.

Godbee, J., and T. Price. 1975. Beaver damage survey. Georgia Forestry Commission, Macon, Georgia.

Godfrey, G., and P. Crowcroft. 1960. The life of the mole. London Museum Press, London, England.

Goldburg, R.J., M.S. Elliot, and R.L. Naylor. 2001. Marine Aquaculture in the United States. Prepared for the Pew Oceans Commission. http://www.pewtrusts.org/uploadedFiles/wwwpewtrustsorg/Reports/Protecting_ocean_life/env_pew_oceans_aquaculture.pdf. Accessed March 29, 2012.

Gottschang, J. L. 1965. Range extension and notes on the shrew (*Sorex cinereus*) in Ohio. Ohio Journal of Science 65:46-47.

Gracely, R. H., and W. F. Sternberg. 1999. Athletes: pain and pain inhibition. American Pain Society 9:1-8.

Grasse, J.E., and E.F. Putnam. 1955. Beaver management and ecology in Wyoming. Federal Aid in Wildlife Restoration Project Bulletin Number 6. Wyoming Game and Fish Commission, Cheyenne, Wyoming.

GSMNP. 2011. Elk. USDI, National Park Service. <http://www.nps.gov/grsm/naturescience/elk.htm>. Accessed June 8, 2011.

Green, J. S., and P. S. Gipson. 1994. Feral dogs. Pp C77-C82 *in* S. E. Hygnstrom, R. M. Timm, and G. E. Larson, eds. Prevention and control of wildlife damage. University of Nebraska-Lincoln, Lincoln, Nebraska.

Green, J. S., F. R. Henderson, and M. D. Collinge. 1994. Coyotes. Pp C51-C76 *in* S. E. Hygnstrom, R. M. Timm, and G. E. Larson, eds. Prevention and control of wildlife damage. University of Nebraska-Lincoln, Lincoln, Nebraska.

Greenhall, A. M., and S. C. Frantz. 1994. Bats. Pp D5-24 *in* S. E. Hygnstrom, R. M. Timm, and G. E. Larson, eds. Prevention and control of wildlife damage. University of Nebraska-Lincoln, Lincoln, Nebraska.

Grinnell, J., J. S. Dixon, and J. M. Linsdale. 1937. Fur-bearing mammals of California. University of California Press, Berkeley, California.

Grizzell, Jr., R. A. 1955. A study of the southern woodchuck, *Marmota monax monax*. American Midland Naturalist 53:257-93.

Grubb, T.G., D.K. Delaney, W.W. Bowerman, and M.R. Wierda. 2010. Golden eagle indifference to heli-skiing and military helicopters in Northern Utah. Journal of Wildlife Management 74:1275–1285.

Gurnell, J.C. 1987. The natural history of squirrels. Facts on File, New York.

Hadidian, J., D. Manski, V. Flyger, C. Cox, and G. Hodge, 1987. Urban gray squirrel damage and population management: A case history. Third Eastern Wildlife Damage Control Conference, Paper 19. http://digitalcommons.unl.edu/ewdcc3/19. Accessed July 23, 2012.

Hall, E.R. 1981. The Mammals of North America. 2 volumes. Ronald Press Co.

Hall, E.R., and K.R. Kelson. 1959. The mammals of North America. Volume 2. The Ronald Press Company, New York City, New York. 616 pp.

Hallberg, D.L., and G.R. Trapp. 1984. Gray fox temporal and spatial activity in a riparian–agricultural zone in California's Central Valley. Pp 920-928 *in* R. E. Warner and K. M. Hendrix, eds. Proceedings of the California Riparian Systems Conference. University of California Press, Berkeley, California.

Hallett, J.G. 1978. Mammalian Species. *Parascalops breweri*. by The American Society of Mammalogists. No. 98, pp. 1-4.

Hamilton, D.A. 1982. Ecology of the bobcat in Missouri. Thesis, University of Missouri, Columbia, Missouri.

Hamilton, Jr., W.J. 1933. The weasels of New York. American Midland Naturalist 14:289-337.

Hamilton, Jr., W.J. 1934. The life history of the rufescent woodchuck *Marmota monax rufescens* Howell. Ann. Carnegie Museum 23:85-178.

Hamilton, Jr., W.J. 1939. Activity of Brewer's mole (*Parascalops breweri*). Journal of Mammalogy 20:307–310.

Haroldson, K.J., and E.K. Fritzell. 1984. Home ranges, activity, and habitat use by gray fox in an oak–hickory forest. Journal of Wildlife Management 48:222-227.

Harris, S. 1977. Distribution, habitat utilization and age structure of a suburban fox (*Vulpes vulpes*) population. Mammal Review 7: 25-39.

Harris, S. 1979. Age related fertility and productivity in red fox, *Vulpes vulpes*, in suburban London. Journal of Zoology (London) 187:195-199.

Harris, S, and J.M.V. Rayner. 198*6*. Urban fox *(Vulpes vulpe*s) population estimates and habitat requirements in several British cities. Journal of Animal Ecology 55:575–591.

Hartman, G.D., and J.D. Krenz. 1993. Estimating population density of moles, *Scalopus aquaticus* using assessment lines. Acta Theriologica 38:305–314.

Hasbrouck, J.J., W.R. Clark, and R.D. Andrews. 1992. Factors associated with raccoon mortality in Iowa. Journal of Wildlife Management 56:693-699.

Hatler, D.F. 1976. The coastal mink of Vancouver Island, British Columbia. Ph. D. Thesis. Univ. of British Columbia, Canada.

Hawkins, C.C., W.E. Grant, and M.T. Longnecker. 1999. Effect of subsidized house cats on California birds and rodents. Transactions of the Western Section of The Wildlife Society 35:29-33.

Hefner, J. M., B. O. Wilen, T. E. Dahl, and W. E. Frayer. 1994. Southeast wetlands: status and trends, Mid 1970's to mid 1980's. USDI, Fish and Wildlife Service, Southeast Region, Atlanta, Georgia.

Hegdahl, P. L., T. A. Gatz, and E. C. Fite. 1980. Secondary effects of rodenticides on mammalian predators. Worldwide Furbearer Conference Proceedings, Volume III. August 3-11, 1980, Frostburg, Maryland.

Hegdal, P.O., and T.L. Gatz. 1977. Hazards to seedeating birds and other wildlife associated with surface strychnine baiting for Richardson's ground squirrels. EPA report under Interagency Agreement EPA-IAG-D4-0449.

Henderson, F.R. 1994. Moles. Pp D51-D58 *in* S. E. Hygnstrom, R. M. Timm and G. E. Larson, eds. Prevention and control of wildlife damage. University of Nebraska-Lincoln, Lincoln, Nebraska.

Herrero, S. 1982. Aspects of evolution and adaptation in American black bears (*Ursus americanus Pallas*) and brown and grizzly bears (*U. arctos Linne*) of North America. International Conference on Bear Research and Management 2:221-231.

Herrero, S. 1985. Bear attacks: their causes and avoidance. New Century Publications, Piscataway, New Jersey.

Herrero, S., A. Higgins, J.E. Cardoza, L.I. Hajduk, and T.S. Smith. 2011. Fatal attacks by American black bear on people: 1900–2009. Journal of Wildlife Management 75:596–603.

Hey, D.L., and N.S. Phillips. 1995. Flood reduction through wetland restoration: The Upper Mississippi River Basin as a case study. Restoration Ecology 3:4-17.

Hill, E.P. 1976. Control methods for nuisance beaver in the southeastern United States. Proceedings of the Vertebrate Pest Conference 7:85-98.

Hill, E.P. 1982. Beaver. Pp 256-281 *in* J. A. Chapman and G. A. Feldhamer, eds. Wild mammals of North America: biology, management, and economics. Johns Hopkins University Press, Baltimore, Maryland.

Hill, E.F., and J.W. Carpenter. 1982. Responses of Siberian ferrets to secondary zinc phosphide poisoning. Journal of Wildlife Management 46:678-685.

Hoagland, J.W. 1993. Nuisance beaver damage control proposal. Oklahoma Department of Wildlife Conservation. Internal Document. 20 pp.

Holbrook, H.T., and R.M. Timm. 1986. Moles and their control. NebGuide G86-777. University of Nebraska-Lincoln Cooperative Extension.

Holthuijzen, A. M. A., W. G. Eastland, A. R. Ansell, M. N. Kochert, R. D. Williams, and L. S. Young. 1990. Effects of blasting on behavior and productivity of nesting prairie falcons. Wildlife Society Bulletin 18:270-281.

Honacki, J. H., K. E. Kinman, and J. W. Koeppl, editors. 1982. Mammal species of the world: a taxonomic and geographic reference. Allen Press Incorporated and Associates, Systematics Collections, Lawrence, Kansas.

Hood, G.A., and S.E. Bayley. 2008. Beaver (*Castor Canadensis*) mitigate the effects of climate on the area of open water in boreal wetlands in western Canada. Biological Conservation 141:556-567.

Houston, A.E. 1995. Beaver immigration into a control area. Southern Journal of Applied Forestry 19:127-130.

Howard, W.E., and R.E. Marsh. 1982. Spotted and hog-nosed skunks. Pp 664-673 *in* J. A. Chapman and G. A. Feldhamer, eds. Wild mammals of North America: biology, management, and economics. Johns Hopkins University Press, Baltimore, Maryland.

Howard, R., L. Berchielli, G. Parsons, and M. Brown. 1980. Trapping furbearers: Student manual. New York State Department of Conservation, New York.

Hubalek, Z., F. Treml, Z. Juricova, M. Hundy, J. Halouzka, V. Janik, and D. Bill. 2002. Serological survey of the wild boar (*Sus scrofa*) for tularemia and brucellosis in south Moravia, Czech Republic. Veterinary Medicine (Czech) 47:60-66.

Humphrey, S.R. 1982. Bats. Pp 52-70 *in* J. A. Chapman and G. A. Feldhamer, eds. Wild mammals of North America: biology, management, and economics. Johns Hopkins University Press, Baltimore, Maryland.

Insurance Institute for Highway Safety. 2005. Collisions with deer. Status Report 40(1):4-5.

Iverson, J.B. 1978. The impact of feral cats and dogs on a population of the West Indian rock iguana, *Cyclura carinata*. Biol. Conserv. 24:3-73.

Jackson, H.H.T. 1961. Mammals of Wisconsin. University of Wisconsin Press, Madison, Wisconsin.

Jackson, J.J. 1994*a*. Tree squirrels. Pp B171-B175 *in* S. E. Hygnstrom, R. M. Timm and G. E. Larson, eds. Prevention and control of wildlife damage. University of Nebraska-Lincoln, Lincoln, Nebraska.

Jackson, J.J. 1994*b*. Opossums. Pp D59-D64 *in* S. E. Hygnstrom, R. M. Timm, and G. E. Larson, eds. Prevention and control of wildlife damage. University of Nebraska-Lincoln, Lincoln, Nebraska.

Jackson, W.B. 1951. Food habits of Baltimore, Maryland cats in relation to rat populations. Journal of Mammalogy 32:458-461.

Jackson, S., and T. Decker. 2004. Beavers in Massachusetts natural history, benefits, and ways to resolve conflicts between people and beavers. UMass Extension, USDA and Massachusetts Department of Fish and Wildlife. 14 pp.

Jenkins, S. H., and P. E. Busher. 1979. *Castor canadensis*. Mammalian Species 120:1-8.

Jensen, P.G., P.D. Curtis, and D.L. Hamelin. 1999. Managing nuisance beavers at roadsides: a guide for highway departments. Cornell Cooperative Extension Publicaiton #147BEAV.

Jessup, D.A. 2004. The welfare of feral cats and wildlife. Journal of the American Veterinary Medical Association 225:1377-1382.

Johnson, G.D., and K.A. Fagerstone. 1994. Primary and secondary hazards of zinc phosphide to nontarget wildlife – a review of the literature. Denver Wildlife Research Report No. 11-55-005, U.S. Department of Agriculture, Denver, Colorado.

Johnson, M.R., R.G. McLean, and D. Slate. 2001. Field operations manual for the use of immobilizing and euthanizing drugs. USDA, APHIS, WS Operational Support Staff, Riverdale, Maryland.

Jones, J.K. Jr., R.S. Hoffman, D.W. Rice, C. Jones, R.F. Baker, and M.G. Engstrom. 1992. Revised checklist of North American mammals north of Mexico, 1991. Occasional Paper 146. Texas Technological University Press, Lubbock, Texas.

Jones, S.C., and K.K. Jordan. 2004. Bat Bugs, Extension Fact Sheet. The Ohio State University Extension. http://ohioline.osu.edu/hyg-fact/2000/pdf/2105a.pdf. Accessed March 2, 2012.

Kaller, M.D., and W.E. Kelso. 2003. Effects of feral swine on water quality in a coastal bottomland stream. Proceedings of the Annual Conference of the Southeastern Association of Fish and Wildlife Agencies 57: 291-298.

Kaller, M.D., and W.E. Kelso. 2006. Swine activity alters invertebrate and microbial communities in a coastal watershed. The American Midland Naturalist 156: 165-179.

Kalmbach, E.R. 1943. The Armadillo: Its Relation to Agriculture and Game. Game, Fish and Oyster Commission, Austin, Texas 61 pp.

Kaye, S.V. 1961. Movements of harvest mice tagged with gold-198. Journal of Mammalogy 42:323-327.

Keirn, G., J. Cepek, B. Blackwell, and T. DeVault. 2010. On a quest for safer skies: managing the growing threat of wildlife hazards to aviation. The Wildlife Professional, Summer 2010: 52-55.

Kendall, R. J., T. E. Lacher Jr., C. Bunck, F. B. Daniel, C. Driver, C. E. Grue, F. Leighton, W. Stansley, P. G. Watanabe, and M. Whitworth. 1996. An ecological risk assessment of lead shot exposure in non-waterfowl avian species: upland game birds and raptors. Environmental Toxicology and Chemistry 15:4-20.

Kennedy, M. L., and P. K. Kennedy. 1998. First record of nutria (*Myocastor coypus* (Mammalia: Rodentia), in Tennessee. Brimleyana 25:156-157.

Kennely, J. J., and B. E. Johns. 1976. The estrous cycle of coyotes. Journal of Wildlife Management 40:272-277.

Kern, W. H., Jr. 2002. Raccoons. WEC-34. Florida Cooperative Extension Service, Institute of Food and Agricultural Sciences, University of Florida, Gainesville, Florida.

King, C.M. 1975. The home range of the weasel (*Mustela nivalis*) in an English woodland. Journal of Animal Ecology 44:639-668.

King, C.M. 1989. The advantages and disadvantages of small size to weasels, Mustela species. Pp 302-334 *in* J.L. Gittleman, ed. Carnivore behavior, ecology, and evolution. Cornell University Press, Ithaca, New York.

Kirsch, E. M. 1996. Habitat selection and productivity of least terns on the lower Platte River, Nebraska. Wildlife Monograph 132:1-48.

Knowlton, F.F. 1972. Preliminary interpretations of coyote population mechanics with some management implications. Journal of Wildlife Management 36:369-383.

Knowlton, F.F., and L.C. Stoddart. 1985. Coyote population mechanics: Another look. Pp 93-111 *in* F.L. Bunnell, D.S. Eastman, and J.M. Peck, eds. Symposium on the Natural Regulation of Wildlife Populations. University of Idaho, Moscow, Idaho.

Kolenosky, G. B., and S. M. Strathearn. 1987. Black Bear. Pp 442-454 *in* M. Novak, J. A. Baker, M. E. Obbard, and B. Mallock, eds. Wild furbearer management and conservation in North America. Ministry of Natural Resources, Ontario, Canada.

Krausman, P. R., and J. J. Hervert. 1983. Mountain sheep responses to aerial surveys. Wildlife Society Bulletin 11:372-375.

Krausman, P. R., L. K. Harris, C. L. Blasch, K. K. G. Koenen, and J. Francine. 2004. Effects of military operations on behavior and hearing of endangered Sonoran pronghorn. Wildlife Monographs 157.

Krausman, P. R., B. D. Leopold, and D. L. Scarborough. 1986. Desert mule deer responses to aircraft. Wildlife Society Bulletin 13:71-73.

Krebs, J. W., C.E., Rupprecht, and J.E. Childs. 2000. Rabies surveillance in the United States during 1999. Journal of the American Veterinary Medical Association 217:1799-1811.

Krebs, J. W., J.S. Smith, C.E. Rupprecht, and J.E. Childs. 1998. Rabies surveillance in the United States during 1997. Journal of the American Veterinary Medical Association 213:1713-1672.

Kushlan, J. A. 1979. Effects of helicopter censuses on wading bird colonies. Journal of Wildlife Management 43:756-760.

Laidlaw, M. A., H. W. Mielke, G. M. Filippelli, D. L. Johnson, and C. R. Gonzales. 2005. Seasonality and children's blood lead levels: developing a predictive model using climatic variables and blood lead data from Indianapolis, Indiana, Syracuse, New York, and New Orleans, Louisiana (USA). Environmental Health Perspectives 113:793-800.

Lamp, R. E. 1989. Monitoring of the effect of military air operations at naval air station Fallon on the biota of Nevada. Nevada Department of Wildlife, Reno, Nevada.

Lancia, R. A., C. S. Rosenberry, and M. C. Conner. 2000. Population parameters and their estimation. Pages 64-83 *in* S. Demaris and P. R. Krausman, editors. Ecology and management of large mammals in North America. Prentice-Hall Incorporated, Upper Saddle River, New Jersey.

Langham, N.P.E. 1990. The diet of feral cats (*Felis catus* L.) on Hawke's Bay farmland, New Zealand. New Zealand Journal of Zoology 17:243-255.

Langlois, S. A. and L. A. Decker. 1997. The use of water flow devices and flooding problems caused by beaver in Massachusetts. Massachusetts Division of Fisheries and Wildlife. 13 pp.

Laramie, H. A., Jr., and S. W. Knowles. 1985. Beavers and their control. University of New Hampshire Cooperative Extension Service, Durham, New Hampshire.

Larivière, S., and L.R. Walton. 1997. Mammalian Species, *Lynx rufus*. American Society of Mammalogists 563:1-8.

Latham, R.M. 1960. Bounties are bunk. National Wildlife Federation, Washington, D.C. 10 pp.

Lawhead, D.N. 1984. Bobcat (*Lynx rufus*) home range, density and habitat preference in South-Central Arizona. The Southwestern Naturalist 29:105-113.

Layne, J. N. 2003. Armadillo. Pp 75-97 *in* G. A. Feldhamer, B. C. Thompson, and J. A. Chapman, eds. Wild Mammals of North America: Biology, Management, and Conservation. Second edition. The Johns Hopkins University Press, Baltimore, Maryland and London, United Kingdom.

LeBlanc, D. J. 1994. Nutria. Pp B71-B80 *in* S. E. Hygnstrom, R. M. Timm, and G. E. Larson, eds. Prevention and control of wildlife damage. University of Nebraska-Lincoln, USA.

Lefrancois, G. R. 1999. The Lifespan. Sixth edition. Wadsworth Publishing Company, Belmont, California.

Lembeck, M. 1978. Bobcat study, San Diego County, California. California Department of Fish and Game, Federal Aid Nongame Wildlife Investigation Project. E-W-2 Report.

Levy, J. K., and P. C. Crawford. 2004. Humane strategies for controlling feral cat populations. Journal of the American Veterinary Medical Association 225:1354-60.

Levy, J. K., D. W. Gale, and L. A. Gale. 2003. Evaluation of the effect of a long-term trap-neuter-return and adoption program on a free-roaming cat population. Journal of the American Veterinary Medicine Association 222:42-46.

Lewis, J. W. 1979. Significance of beaver and beaver ponds in the Tombigbee Resource Conservation and Development Area Alabama-1978, Alabama Cooperative Extension Service, Circular CRD-7, Auburn University, Auburn, Alabama.

Liberg, O. 1984. Food habits and prey impact by feral and house based domestic cats in a rural area in southern Sweden. Journal of Mammalogy 65:424-432.

Linnell, M. A., M. R. Conover, and T. J. Ohashi. 1996. Analysis of bird strikes at a tropical airport. Journal of Wildlife Management 60:935-945.

Linscombe, G., N. Kinler, and R. J. Aulerich. 1982. Mink. Pp 629-643 *in* J. A. Chapman and G. A. Feldhamer, eds. Wild mammals of North America: biology, management, and economics. Johns Hopkins University Press, Baltimore, Maryland.

Lipscomb, D. J. 1989. Impacts of feral hogs on longleaf pine regeneration. Southern Journal of Applied Forestry 13:177-81.

Lisle, S. 1996. Beaver Deceivers. Wildlife Control Technology. September-October 42-44.

Lisle, S. 1999. Wildlife Programs at the Penobscot Nation. Transactions of the North American Wildlife and Natural Resource Conference 65:466-477.

Lisle, S. 2003. Use and potential of flow devices in beaver management. Lutra 46:211-216.

Logan, K. A., L. Sweanor, and M. Hornocker. 1996. Cougar population dynamics. Chapter 3 *in* Cougars in the San Andres Mountains, New Mexico (Project No. W-128-R, Final Report). New Mexico Department of Game and Fish, Santa Fe, New Mexico.

Lord, R. D., Jr. 1961. A population study of the gray fox. American Midland Naturalist Journal 66:87–109.

Lowry, D. A. 1978. Domestic dogs as predators on deer. Wildlife Society Bulletin 6:38-39.

Ludders, J. W., R. H. Schmidt, F. J. Dein, and P. N. Klein. 1999. Drowning is not euthanasia. Wildlife Society Bulletin 27:666-670.

MacDonald, D. W., and M. T. Newdick. 1982. The distribution and ecology of fox, *Vulpes vulpes* (L.), in urban areas. Pp 123–135 *in* R. Bornkamm, J. A. Lee, and M. R. D. Seaward, eds. Urban ecology. Blackwell Scientific Publications, Oxford, United Kingdom.

MacKinnon, B., R. Sowden, and S. Dudley. 2001. Sharing the Skies: an Aviation Guide to the Management of Wildlife Hazards. Transport Canada, Aviation Publishing Division, Tower C, 330 Sparks Street, Ottawa, Ontario, K1A 0N8 Canada. 316 pp.

MacLean, S. F., Jr., B. M. Fitzgerald, and F. A. Pitelka. 1974. Population cycles in arctic lemmings: winter reproduction and predation by weasels. Arctic and Alpine Research 6:1-12.

Majumdar, S.K., J.E. Huffman, F.J. Brenner, and A.I. Panah. 2005. Wildlife diseases: landscape epidemiology, spatial distribution and utilization of remote sensing technology. The Pennsylvania Academy of Sciences.

Manci, K. M., D. N. Gladwin, R. Villella, and M. G. Cavendish. 1988. Effects of aircraft noise and sonic booms on domestic animals and wildlife: A literature synthesis. Fort Collins, Colorado/ Kearneysville, West Virginia: U.S. Fish and Wildlife Service and National Ecology Research Center.

Manski, D.M., L.W. VanDruff, and V. Flyger. 1981. Activities of gray squirrels and people in a downtown Washington, D.C. park: management implications. Trans. North Amer. Wild. Nat. Res. Conf. 46:439-454.

Marks, S. G., and J. E. Koepke. 1994. Pet attachment and generativity among young adults. Journal of Psychology 128:641.

Marshall, W. H. 1935. A study of the winter activities of the mink. Journal of Mammalogy 17:382-92.

Massey, B. W. 1971. A breeding study of the California least tern, 1971. Administrative Report 71-9, Wildlife Management Branch, California Department of Fish and Game.

Massey, B. W., and J. L. Atwood. 1981. Second-wave nesting of the California least tern: age composition and reproductive success. Auk 98:596-605.

McCabe, R. A. 1949. Notes on live-trapping mink. Journal of Mammalogy 30:416-423.

McCord, C. M., and J. E. Cardoza. 1982. Pp 728-766 *in* J. A. Chapman and G. A. Feldhamer, eds. Wild mammals of North America: biology, management, and economics. Johns Hopkins University Press, Baltimore, Maryland.

McKnight, T. 1964. Feral livestock in Anglo-America. University of California Publications in Geography Vol. 16. University of California Press, Berkeley, California.

McNeely, R. 1995. Missouri's Beaver: A guide to management, nuisance prevention, and damage control. Missouri Department of Conservation, Jefferson City, Missouri.

Means, D. B. 1999. *Desmognathus auriculatus*. Pages 10-11 *in* Michael Lanoo, editor. Status and Conservation of U. S. Amphibians. Declining Amphibians Task Force Publication No. 1.

Medin, D. E., and W. P. Clary. 1990. Bird populations in and adjacent to a beaver pond ecosystem in Idaho. USDA-Forest Service, Intermountain Research Station, 432.

Medin, D. E., and W. P. Clary. 1991. Small mammals of a beaver pond ecosystem and adjacent riparian habitat in Idaho. USDA-Forest Service, Intermountain Research Station, 445.

Melquist, W. E., and A. E. Dronkert. 1987. River otter. Pp 626-641 *in* M. Novak, J. A. Baker, M. E. Obbard, and B. Mallock, eds. Wild furbearer management and conservation in North America. Ministry of Natural Resources, Ontario, Canada.

Meltzer, M. I. 1996. Assessing the costs and benefits of an oral vaccine for raccoon rabies: a possible model. Emerging Infectious Diseases 2:343-349.

Mengak, M. T. 2005. Nine-banded Armadillo (*Dasypus novemcinctus*). Warnell School of Forest Resources Natural History Series No. 4, May 2005.

Merriam, H.G. 1971. Woodchuck burrow distribution and related movement patterns. Journal of Mammalogy 52:732-46.

Meyers, B. 2000. Anticipatory mourning and the human-animal bond. Pp 537-564 *in* T. A. Rando, ed. Clinical dimensions of anticipatory mourning: theory and practice in working with the dying, their loved ones, and their caregivers. Research Press, Champaign, Illinois.

Millennium Ecosystem Assessment. 2005. Ecosystems and human well-being: Wetlands and water. World Resources Institute, Washington, D.C..

Miller, J. E. 1994. Muskrats. Pp B61-B69 *in* S. E. Hygnstrom, R. M. Timm, and G. E. Larson, eds. Prevention and Control of Wildlife Damage. Univ. Nebr. Coop. Ext., USDA-APHIS-ADC and Great Plains Agric. Council Wildl. Comm., Lincoln, Nebraska.

Miller, L.A. 1995. Immunocontraception as a tool for controlling reproduction in coyotes. Pp 172-176 *in* D. Rollins, C. Richardson, T. Blankenship, K. Canon, and S. Henke, eds. Coyotes in the Southwest: A Compendium of Our Knowledge. Proc. of a Symposium, Dec. 13-15, Texas A&M University, San Angelo, Texas.

Miller, J.E., and G.K. Yarrow. 1994. Beaver. Pp B-1-B-11 *in* S. E. Hygnstrom, R. M. Timm, and G. E. Larson, eds. Prevention and Control of Wildlife Damage. Univ. Nebr. Coop. Ext., USDA-APHIS-ADC and Great Plains Agric. Council Wildl. Comm., Lincoln, Nebraska.

Miller, L. A., B. E. Johns, and G. J. Killian. 2000. Immunocontraception of white-tailed deer with GnRH vaccine. American Journal of Reproductive Immunology 44:266-274.

Mitchell, J. L. 1961. Mink movements and populations on a Montana river. Journal of Wildlife Management 25:48-54.

Mosillo, M., J. E. Heske, and J.D. Thompson. 1999. Survival and movements of translocated raccoons in northcentral Illinois. Journal of Wildlife Management 63:278-286.

Muller, L. I., R. J. Warren, and D. L. Evans. 1997. Theory and Practice of immunocontraception in wild animals. Wildlife Society Bulletin 25:504-514.

Muller-Schwarze, D., and L. Sun. 2003. The beaver: Natural history of a wetlands engineer. Cornell University Press, Ithaca, New York.

Munther, G.L. 1982. Beaver management in grazed riparian ecosystems. Wildilfe Livestock Relationships Symposium 10:234-241.

Naimen, R. J., J. M. Melillo, and J. E. Hobbie. 1986. Ecosystem alteration of boreal forest streams by beaver (*Castor canadensis*). Ecology 67:1254-1269.

NASS. 2002. U.S. Wildlife Damage. USDA, National Agricultural Statistics Service, Washington, D.C..

NASS. 2011. Cattle death loss 2010. Released May 12, 2011. USDA, National Agricultural Statistics Service, Washington, DC. http://www.usda.gov/nass/PUBS/TODAYRPT/catlos11.pdf . Accessed December 22, 2011.

National Audubon Society. 2000. Field guide to North American mammals. J. O. Whitaker, Jr., editor. Alfred A. Knopf Publishing, New York City, New York.

National Park Service. 1995. Report of effects of aircraft overflights on the National Park System. USDI-NPS D-1062, July, 1995.

Nesbitt, W. H. 1975. Ecology of a feral dog pack on a wildlife refuge. Pp 391-396 *in* M. W. Fox, ed. The wild canids. Van Nostrand Reinhold Company, New York City, New York.

Ness, E. 2003. Oh deer: Exploding populations of white-tailed deer are stripping our forests of life. Discover 24(3).

Nicholson, W. S. 1982. An ecological study of the gray fox in east central Alabama. Thesis, Auburn University, Auburn, Alabama.

Nielsen, L. 1988. Definitions, considerations, and guidelines for translocation of wild animals. Pages 12-49 *in* L. Nielsen and R. D. Brown, editors. Translocation of wild animals. Wisconsin Humane Society, Milwaukee, Wisconsin, and Caesar Kleberg Wildlife Research Institute, Kingsville, Texas.

Nielsen, C.K., and A. Woolf. 2001. Spatial organization of bobcats (*Lynx rufus*) in southern Illinois. American Midland Naturalist 146:43–52.

Noah, D.L., M.G. Smith, J.C. Gotthardt, J. W. Krebs, D. Green, and J.E. Childs. 1995. Mass human exposure to rabies in New Hampshire: Exposures, treatment, and cost. Public Health Briefs, National Center for Infectious Diseases, Atlanta, Georgia.

Nolte, D. M., S. R. Swafford, and C. A. Sloan. 2001. Survey of factors affecting the success of Clemson beaver pond levelers installed in Mississippi by Wildlife Services. Pages 120-125 in M C. Brittingham, J. Kays, and R. McPeake editors, Proceedings of the Ninth Wildlife Damage Management Conference. Pennsylvania State University, University Park, Pennsylvania.

Noonan, B. 1998. The Canadian terminal dive study. Wildlife Control Techniques. May-June.

Novak, M. 1987. Beaver. Pp. 282-312 *in* M. Novak, J.A. Baker, M.E. Obbard, and B. Mallock, eds. Wild Furbearer Management and Conservation in North America. Ontario Trappers Assoc., Ontario.

O'Brien, J. M. 1994. Voles. Pp B13-B16 *in* S. E. Hygnstrom, R. M. Timm and G. E. Larson, eds. Prevention and control of wildlife damage. University of Nebraska-Lincoln, Lincoln, Nebraska.

Organ, J. F., T. Decker, J. DiStefano, K. Elowe, P. Rego, and P.G. Mirick. 1996. Trapping and furbearer management: Perspectives from the Northeast. USDI-Fish and Wildlife Service, Hadley, Massachusetts.

Osgood, F.L. 1935. Fluctuations in small mammal populations. Journal of Mammalogy 16:156.

Pagel, M.D., R.M. May, and A.R. Collie. 1991. Ecological aspects of the geographical distribution and diversity of mammalian species. Am. Nat. 137:791-815.

Parker, G. 1995. Eastern coyote: The story of its success. Nimbus Publishing, Halifax, Canada.

Parkes, C. M. 1979. Grief: the painful reaction to the loss of a loved one. Monograph. University of California, San Diego, California.

Parrish, W. F., Jr. 1960. Status of the beaver (*Castor canadensis carolinensis*) in Georgia. Thesis, University of Georgia, Athens, Georgia.

Partington, M. 2002. Preventing beaver dams from blocking culverts. Advantage 3:1-4.

Patterson, D. 1950. Beaver-trout relationships. Wisconsin Conservation Bulletin 15:9-11.

Pearson, O.P. 1964. Carnivore-mouse predation: an example of its intensity and bioenergetics. Journal of Mammalogy 45:177–188.

Pearson, O.P. 1971. Additional measurements of the impact of carnivores on California voles (*Microtus californicus*). Journal of Mammalogy 52:41–49.

Pearson, O. P. 1985. Predation. Pp 535-566 *in* R. A. Tamarin, ed., Biology of new world *Microtus*. American Society of Mammalogy. Shippensburg, Pennsylvania.

Pelton, M. R. 1982. Black Bear. Pp 504-514 *in* J. A. Chapman and G. A. Feldhamer, eds. Wild mammals of North America: biology, management, and economics. Johns Hopkins University Press, Baltimore, Maryland.

Perry, D. 2007. Coexisting with beavers. DVD. Spring Farms Cares, Clinton, New York.

Phillips, R. L. 1996. Evaluation of 3 types of snares for capturing coyotes. Wildlife Society Bulletin 24: 107-110.

Phillips, R. L., and R. H. Schmidt. 1994. Fox. Pp C83-C88 *in* S. E. Hygnstrom, R. M. Timm and G. E. Larson, eds. Prevention and control of wildlife damage. University of Nebraska-Lincoln, Lincoln, Nebraska.

Pollock, M. M., T. M. Beechie, and C. E. Jordan. 2007. Geomorphic changes upstream of beaver dams in Bridge Creek, an incised stream channel in the interior Columbia River basin, eastern Oregon. Earth Surface Processes and Landforms 32:1174-1185.

Pyrah, D. 1984. Social distribution and population estimates of coyotes in north-central Minnesota. Journal of Wildlife Management 48:679-690.

Quick, H.F. 1944. Habits and economics of New York weasel in Michigan. Journal of Wildlife Management 8:71-78.

Quick, H.F. 1951. Notes on the ecology of weasels in Gunnison County, Colorado. Journal of Mammalogy 32:281-290.

Ramey, C. A., J. B. Bourassa, and J. E. Brooks. 2000. Potential risks to ring-necked pheasants in California agricultural areas using zinc phosphide. International Biodeterioration & Biodegradation 45:223-230.

Randolph, J. P. 1988. Virginia trapper's manual. Department of Game and Inland Fisheries, Richmond, Virginia.

Reese, K.P., and J.D. Hair. 1976. Avian species diversity in relation to beaver pond habitats in the Piedmont region of South Carolina. Proceeding of the Annual Conference of the Southeastern Association of Fish and Wildlife Agencies 30:437-447.

Reid, D.G. 1984. Ecological interactions of river otters and beavers in a boreal ecosystem. M.S. Thesis, University of Calgary, Calgary, Alberta, 210 pp.

Richmond, N.D., and H.R. Roslund. 1949. Mammal survey of northwestern Pennsylvania. Pennsylvania Game Commission, Harrisburg, Pennsylvania. 67 pp.

Riley S. P., D. J. Hadidian, and D. A. Manski. 1998. Population density, survival, and rabies in raccoons in an urban national park. Canadian Journal of Zoology 76:1153–1164.

Ringleman, J. K. 1991. Managing beaver to benefit waterfowl. Waterfowl Management Handbook. United States Department of the Interior, Fish and Wildlife Service, Fish and Wildlife Leaflet 13.4.7.

Ritcey, R.W., and R.Y. Edwards. 1956. Live trapping mink in British Columbia. Journal of Mammalogy 37:114-116.

Robinson, M. 1996. The potential for significant financial loss resulting from bird strikes in or around an airport. Proceedings of the Bird Strike Committee (Europe) 23:353-367.

Roblee, K.J. 1983. A wire mesh culvert for use in controlling water levels at nuisance beaver sites. Proc. East. Wildl. Dam. Control Conf. 3:25-33.

Roblee, K. J. 1984. Use of corrugated plastic drainage tubing for controlling water levels at nuisance beaver sites. New York Fish and Game Journal 31:63-80.

Roblee, K.J. 1987. The use of T-culvert guard to protect road culverts from plugging damage by beavers. Proc. East. Wildl. Damage Control Conf. 3:25-33.

Romin, L. A., and J. A. Bissonette. 1996. Deer-vehicle collisions: status of state monitoring activities and mitigation efforts. Wildlife Society Bulletin 24:276-83.

Rosatte, R.C. 1987. Skunks. Pp 599-613 *in* M. Novak, J. A. Baker, M. E. Obbard, and B. Mallock, eds. Wild furbearer management and conservation in North America. Ministry of Natural Resources, Ontario, Canada.

Roseberry, J.L., and A. Woolf. 1998. Habitat-population density relationships for white-tailed deer in Illinois. Wildlife Society Bulletin 26:252-258.

Ross, C. B., and J. Baron-Sorensen. 1998. Pet loss and human emotion: guiding clients through grief. Accelerated Development, Incorporation, Philadelphia, Pennsylvania.

Rossell, R. O. Bozsér, P. Collen, and H. Parker. 2005. Ecological impact of beavers Castor fiber and Castor Canadensis and their ability to modify ecosystems. Mammal Review 35:248-276.

Ruell, E.W., S.P.D. Riley, M.R. Douglas, J.P. Pollinger, and K.R. Crooks. 2009. Estimating bobcat population sizes and densities in a fragmented urban landscape using noninvasive capture-recapture sampling. Journal of Mammalogy 90:129–135.

Saliki, J.T., S.J. Rodgers, and G. Eskew. 1998. Serosurvey of selected viral and bacterial diseases in wild swine in Oklahoma. J. Wildl. Dis. 34:834-838.

Samuel, M. D., and M. R. Fuller. 1994. Wildlife radiotelemetry. Pp 370-417 *in* Research and management techniques for wildlife and habitats, T. A. Bookhout, ed. Allan Press, Inc., Lawrence, Kansas.

Samuel, W. M., M. J. Pybus, and A. A. Kocan, editors. 2001. Parasitic diseases of wild mammals. Iowa State University Press, Ames, Iowa.

Sanderson, G. C. 1987. Raccoons. Pp 486-499 *in* M. Novak, J. A. Baker, M. E. Obbard, B. Mallock, eds. Wild furbearer management and conservation in North America. Ministry of Natural Resources, Ontario, Canada.

Sanderson, G. C., and G. F. Huber, Jr. 1982. Selected demographic characteristics of Illinois (U.S.A) raccoons (*Procyon lotor*). Pages 487-513 *in* J. A. Chapman and D. Pursely, eds. Worldwide furbearer conference proceedings. Maryland Wildlife Administration, Annapolis, Maryland.

Saunders, D.A. 1988. Adirondack Mammals. State University of New York, College of Environmental Science and Forestry. 216 pp.

Schmidt, R. 1989. Wildlife management and animal welfare. Transcripts of the North American Wildlife and Natural Resources Conference 54:468-475.

Schwartz, C. W., and E.R. Schwartz. 2001. The Wild Mammals of Missouri. Second Revised Edition. University of Missouri Press. 365 pp.

Scott, M. D., and K. Causey. 1973. Ecology of feral dogs in Alabama. Journal of Wildlife Management 37:253-265.

Seabrook, W. 1989. Feral cats (*Felis catus*) as predators of hatchling green turtles (*Chelonia mydas*). Journal of Zoology 219:83-88.

Seidensticker, J., M. A. O'Connell, and A. J. T. Johnsingh. 1987. Virginia opossum. Pp 247-261 *in* M. Novak, J. A. Baker, M. E. Obbard, and B. Malloch, eds. Wild furbearer management and conservation in North America. Ministry of Natural Resources, Ontario, Canada.

Seward, N. W., K. C. VerCauteren, G. W. Witmer, and R. M. Engeman. 2004. Feral swine impacts on agriculture and the environment. Sheep and Goat Research Journal 19:34-40.

Sheffield, S.R., and H.H. Thomas. 1997. *Mustela frenata*. Mammalian Species 570:1-9.

Siegfried, W. R. 1968. The reaction of certain birds to rodent baits treated with zinc phosphide. Ostrich 39:197-198.

Simon, L. 2006. Solving beaver flooding problems through the use of water flow control devices. Proceedings of the Vertebrate Pest Conference 22:174-180.

Singer, F. J., W. T. Swank, and E. E. C. Clebsch. 1984. Effects of wild pig rooting in a deciduous forest. Journal of Wildlife Management 48:464-473.

Slate, D. 1980. A study of New Jersey raccoon populations— determination of the densities, dynamics and incidence of disease in raccoon populations in New Jersey. New Jersey Division of Fish, Game and Wildlife, Pittman-Robertson Project W-52-R-8, Final Report.

Slate, D.A., R. Owens, G. Connolly, and G. Simmons. 1992. Decision making for wildlife damage management. Transcripts of the North American Wildlife and Natural Resources 57:51-62.

Slater, M. R. 2004. Understanding issues and solutions for unowned, free-roaming cat populations. Journal of the American Veterinary Medical Association 225:1350-1354.

Smith, R.L. 1996. Ecology and field biology. Haper Collins College Publishers. New York, NY. 5th Ed. 733 pp.

Smithsonian National Museum of Natural History. 2012. North American Mammals. http://www.mnh.si.edu/mna/search_name.cfm. Accessed February 28, 2012.

Speich, S. 1986. Colonial waterbirds. Pages 387-405 *in* A. Y. Cooperrider, R. J. Boyd, and H. R. Stuart, editors. Inventory and monitoring of wildlife habitat. USDI, Bureau of Land Management Service Center, Denver, Colorado.

Spock, M. 2006. Effectiveness of water flow devices as beaver conflict resolution tools: A satisfaction survey of Massachusetts clients. Center for Animals and Public Policy, Tufts University Cummings School of Veterinary Medicine. 50 pp.

Stalling, D. T. 1997. *Reithrodontomys humulis*. Mammalian Species No. 565. American Society of Mammalogy, New York.

Stansley, W., L. Widjeskog, and D. E. Roscoe. 1992. Lead contamination and mobility in surface water at trap and skeet ranges. Bulletin of Environmental Contamination and Toxicology 49:640-647.

State Farm Mutual Automobile Insurance Company. 2011*a*. U.S. deer-vehicle collisions fall 7 percent-Mishaps most likely in November and in West Virginia. http://www.statefarm.com/aboutus/_pressreleases/2011/october/3/us-deer-collisions-fall.asp. Accessed February 9, 2012.

State Farm Mutual Automobile Insurance Company. 2011*b*. Likelihood of collision with deer. http://www.statefarm.com/aboutus/_pdf/deer-chart-2011.pdf. Accessed February 9, 2012.

Stevens, R. L. 1996. The feral hog in Oklahoma. Samuel Roberts Noble Foundation, Ardmore, Oklahoma.

Stewart, C. M., and N. B. Veverka. 2011. The extent of lead fragmentation observed in deer culled by sharpshooting. Journal of Wildlife Management 75:1462-1466.

Stoskopf, M. K., and F. B. Nutter. 2004. Analyzing approaches to feral cat management – one size does not fit all. Journal of the American Veterinary Medical Association 225:1361-1364.

Strole, T. A., and R. C. Anderson. 1992. White-tailed deer browsing: species preferences and implications for central Illinois forests. Natural Areas Journal 12:139-144.

Svendsen, G.E. 1990. Weasels. Pp 613-628 *in* J.A. Chapman, G.A. Feldhamer, eds. Wild Mammals of North America: Biology, Management, and Economics. The Johns Hopkins University Press, Baltimore, Maryland. 1147 pp.

Swihart, R. K. 1992. Home-range attributes and spatial structure of woodchuck populations. Journal of Mammalogy 73:604-18.

Swihart, R. K., and P. M. Picone. 1995. Use of woodchuck burrows by small mammals in agricultural habitats. American Midland Naturalist 133:360-363.

Tennessee Bat Working Group. 2012. Bats of Tennessee. http://www.tnbwg.org/index.html. Accessed September 27, 2012.

TDEC. 2008. Rare Plant List. TDEC Website: <http://tn.gov/environment/na/pdf/plant_list.pdf>. Accessed May 18, 2011.

TDEC. 2009. Rare Animals List. TDEC Website: <http://tn.gov/environment/na/pdf/animal_list.pdf>. Accessed May 18, 2011.

TDEC. 2010. Tennessee 2020: Vision for Parks, People and Landscapes. TDEC Website: <http://tn.gov/environment/recreation/plan>. Accessed May 3, 2011.

TWRA. 2001. Tennessee Furbearer Report 1997/98-1998/99. Technical Report No. 99-14. Tennessee Wildlife Resource Agency, Nashville, Tennessee.

TWRA. 2010. Big Game Harvest Report 2009-2010. Technical Report No. 10-01. Tennessee Wildlife Resource Agency, Nashville, Tennessee.

TWRA. 2011*a*. 2011 Tennessee Hunting and Trapping Guide. Tennessee Wildlife Resources Agency, Nashville, Tennessee.

TWRA. 2011*b*. Elk Reintroduction Questions & Answers. TWRA Website: <http://www.tn.gov/twra/elkquestions.html>. Accessed June 21, 2011.

TWRA. 2012*a*. Tennessee hunting and trapping 2012 guide. Tennessee Wildlife Resources Agency, Nashville, Tennessee.

TWRA. 2012*b*. Deer information, management, and hunting in Tennessee website. http://www.tn.gov/twra/deermain.html. Accessed September 26, 2012.

TVA. 2011*a*. Tennessee Valley Authority: Natural resources plan. https://www.tva.gov/environment/reports/nrp/index.htm. Accessed March 15, 2012.

TVA. 2011*b*. Final Environmental Impact Statement: Natural Rources Plan-Alabama, Georgia, Kentucky, Mississippi, North Carolina, Tennessee, and Virginia. https://www.tva.gov/environment/reports/nrp/index.htm. Accessed March 15, 2012.

Tesky, J.L. 1993. *Castor canadensis*. In: Fire Effects Information System, [Online]. USDA, Forest Service, Rocky Mountain Research Station, Fire Sciences Laboratory (Producer). Available: http://www.fs.fed.us/database/feis/. Accessed April 4, 2011.

Teutsch, S.M., D.D. Juranek, A. Sulzer, J.P. Dubey, and R. K. Sikes. 1979. Epidemic toxoplasmosis associated with infected cats. N. Engl. J. Med. 300:695-699.

The Wildlife Society. 1990. Conservation Policies of the Wildlife Society. The Wildlife Society, Washington., D.C. 20 pp.

Thompson, R. L. 1977. Feral hogs on National Wildlife Refuges. Pages 11-15 *in* G. W. Wood, editor. Reasearch and management of wild hog populations. Belle W. Baruch Forest Science Institute, Clemson University, Georgetown, South Carolina.

Thorpe, J. 1996. Fatalities and destroyed civil aircraft due to bird strikes, 1912-1995. Proceedings of Bird Strike Committee Europe 23:17-32.

Thorpe, J. 1997. The implications of recent serious bird strike accidents and multiple engine ingestions. Bird Strike Committee USA, Boston, Massachusetts.

Tietjen, H. P. 1976. Zinc phosphide – its development as a control agent for black-tailed prairie dogs. U.S. Fish and Wildlife Service, Special Report on Wildlife 195.

Tietjen, H. P., and G. H. Matschke. 1982. Aerial pre-baiting for the management of prairie dogs with zinc phosphide. Journal of Wildlife Management 46:1108-1112.

Tilghman, N.G. 1989. Impacts of white-tailed deer on forest regeneration in northwestern Pennsylvania. Journal of Wildlife Management 53:524-532.

Till J. A. 1992. Behavioral effects of removal of coyote pups from dens. Proc. Vertebr. Pest Conf. 15:396-399.

Timm, R. M. 1994. Norway rats. Pp B105-B120 *in* S. E. Hygnstrom, R. M. Timm and G. E. Larson, ed. Prevention and control of wildlife damage. University of Nebraska-Lincoln, Lincoln, Nebraska.

Timm, R. M., Baker, R. O., Bennett, J. R. and Coolahan, C. C. 2004. Coyote attacks: an increasing urban problem. Proceedings of the 69th North American Wildlife and Natural Resources Conference, March 16-20, 2004, Spokane, Washington.

Trapp, G. R. 1978. Comparative behavioral ecology of the ringtail (*Bassariscus astutus*) and gray fox (*Urocyon cinereoargenteus*) in southwestern Utah. Carnivore 1:3–32.

Truyen U., D. Addie, S. Belák, C. Boucraut-Baralon, H. Egberink, T. Frymus, T. Gruffydd-Jones, K. Hartmann, M.J. Hosie, A. Lloret, H. Lutz, F. Marsilio, M.G. Pennisi, A.D. Radford, E. Thiry, and M.C. Horzinek. 2009. Feline panleukopenia. ABCD guidelines on prevention and management. J. Feline Med. Surg. 11:538-46.

Turner, J. W., J. F. Kirkpatrick, and I. K. M. Liu. 1993. Immunocontraception in white-tailed deer. Pp 147-159 *in* T.J. Kreeger, Technical Coordinator. Contraception in Wildlife Management. USDA/APHIS, Technical Bulletin No. 1853.

Twichell, A. R. 1939. Notes on the southern woodchuck in Missouri. Journal of Mammalogy 20:71-74.

Twichell, A. R., and H. H. Dill. 1949. One hundred raccoons from one hundred and two acres. Journal of Mammalogy 30:130–133.

USDA. 1997. Animal Damage Control Program: Final Environmental Impact Statement (revised). USDA, APHIS, WS-Operational Support Staff, 4700 River Road, Unit 87, Riverdale, Maryland 20737.

USDA. 2004. Wildlife Services cooperative rabies management program: Tennessee 2004. Pages 74-78 *in* C. D. Kostrzewski and K. M. Nelson, eds. Cooperative rabies management program national report 2004. USDA, APHIS, Wildlife Services, Concord, New Hampshire.

USDA. 2005*a*. Environmental Assessment: Mammal Damage Management in the Tennessee Wildlife Services Program. USDA/APHIS/ Wildlife Services, Madison, Tennessee.

USDA. 2005*b*. Supplemental environmental assessment: Oral vaccination to control specific rabies virus variants in raccoons, gray fox, and coyotes in the United States. USDA/APHIS/Wildlife Services, Riverdale, Maryland.

USDA. 2005*c*. Environmental Assessment: Predator Damage Management in Colorado. USDA/APHIS/ Wildlife Services, Denver, Colorado.

USDA. 2007. Wildlife Services cooperative rabies management program: Tennessee 2007. Pages 95-101 *in* K. M. Nelson, ed. Cooperative Rabies Management Program National Report 2007. USDA, APHIS, Wildlife Services, Concord, New Hampshire.

USDA. 2008. Pseudorabies (Aujeszky's disease) and its eradication. United Stated Department of Agriculture, Animal and Plant Health Inspection Service. Technical Bulletin No. 1923.

USDA. 2010. Questions and Answers: GonaconTM-Birth control for deer. United States Department of Agriculture, Animal and Plant Health Inspection Service, Wildlife Services. 3 pp.

USDI. 1979. Mammalian predator damage management for livestock protection in the Western United States. Final Environmental Impact Statement. Washington, D.C. 789 pp.

USDI. 2008. Fish and Wildlife Service, Fergus Falls Wetland Management District - Wetlands. http://www.fws.gov/midwest/FergusFallsWetland/wetland.html. Accessed September 25, 2012.

USFWS. 2001. Inside Region 3: Ohio man to pay more than $11,000 for poisoning migratory birds. Volume 4(2):5.

USFWS. 2011. United States Fish and Wildlife Service, Endangered Species Database. USFWS Website: http://www.fws.gov/endangered. Accessed June 20, 2011.

USFS. 1992. Overview, Report to Congress, Potential Impacts of Aircraft Overflights of National Forest System Wilderness. Report to Congress. Prepared pursuant to Section 5, Public Law 100-91, National Park Overflights Act of 1987.

United States General Accounting Office. 2001. Wildlife Services Program: Information on activities to manage wildlife damage. Report to Congressional Committees. GOA-02-138. 71 pp.

Uresk, D. W., R. M. King, A. D. Apa, M. S. Deisch, and R. L. Linder. 1988. Rodenticidal effects of zinc phosphide and strychnine on notarget species. Pages 57-63 *in* D. W. Uresk, G. L. Schenbeck, and R. Cefkin, tech. coordinators. Eighth Great Plains Wildlife Damage Control Workshop Proceedings, Fort Collins, Colorado, USA. GTR RM-154. USDA Forest Service Rocky Mountain Forest and Range Experiment Station.

VanDruff , L.W. 1971. The ecology of the raccoon and opossum, with emphasis on their role as waterfowl nest predators. Ph.D. Thesis. Cornell University, Ithaca, New York. 140 pp.

Vaughan, M. R. and M. R. Pelton. 2003. Black Bears of North America. National Biological Service, Virginia Cooperative Fish and Wildlife Research Unit. Virginia Polytechnic Institute and State University, Blacksburg, Virginia.

Verts, B.J. 1963. Movements and populations of opossums in a cultivated area. Journal of Wildlife Management 27:127129.

Virchow, D. and D. Hogeland. 1994. Bobcats. Pp C35-C45 *in* S.E. Hygnstrom, R.M. Timm and G.E. Larson, eds. Prevention and control of wildlife damage. University of Nebraska-Lincoln, Lincoln, Nebraska.

Virginia Department of Game and Inland Fisheries. 1999. 1999 Deer management plan. VDGIF Website: http://www.dgif.state.va.us/hunting/va_game_wildlife/management_plans/deer/deer_management_plan.pdf. Accessed 21 June 2011.

Voigt, D. R. 1987. Red fox. Pp 378-392 *in* M. Novak, J. A. Baker, M.E. Obbard, B. Mallock, eds. Wild furbearer management and conservation in North America. Ministry of Natural Resources, Ontario, Canada.

Voigt, D. R., and W. E. Berg. 1987. Coyote. Pp 345-357 *in* M. Novak, J. A. Baker, M. E. Obbard, B. Mallock, eds. Wild furbearer management and conservation in North America. Ministry of Natural Resources, Ontario, Canada.

Voigt, D. R., and B. D. Earle. 1983. Avoidance of coyotes by red fox families. Journal of Wildlife Management 47:852–857.

Voigt, D. R., and D. W. MacDonald. 1984. Variation in the spatial and social behavior of the red fox, *Vulpes vulpes*. Acta Zoologica Fennica 171:261-265.

Voigt, D. R., and R. L. Tinline. 1980. Strategies for analyzing radio tracking data. Pp 387–404 *in* C. J. Amlaner, Jr., and D. W. MacDonald, eds. A handbook on biotelemetry and radio tracking. Pergamon Press, Oxford, United Kingdom.

Wade, D. E., and C. W. Ramsey. 1986. Identifying and managing mammals in Texas: beaver, nutria and muskrat. Texas Agricultural Extension Service and Texas Agriculture Experimental Station. Texas A&M University in cooperation with USDI-USFWS Pub. B-1556, College Station, Texas.

Waller, D. M., and W. S. Alverson. 1997. The white-tailed deer: a keystone herbivore. Wildlife Society Bulletin 25:217-26.

Warren, R. J. 1991. Ecological justification for controlling deer populations in eastern national parks. Transactions of the North American Wildlife and Natural Resources Conference 56:56-66.

Weisenberger, M.E., P.R. Krausman, M.C. Wallace, D.W. De Young, and O.E. Maughan. 1996. Effects of simulated jet aircraft noise on heart rate and behavior of desert ungulates. Journal of Wildlife Management 60:52-61.

Weisman, A. D. 1991. Bereavement and companion animals. Omega: Journal of Death and Dying 22: 241-248.

West, B. C., A. L. Cooper, and J. B. Armstrong. 2009. Managing wild pigs: A technical guide. Human-Wildlife Interactions Monograph 1:1-55.

Whitaker, Jr., J.O., and W.L.J. Hamilton, Jr. 1998. Mammals of the Eastern United States. Cornell University Press, Ithaca, New York. 583 pp.

White, C. M., and S. K. Sherrod. 1973. Advantages and disadvantages of the use of rotor-winged aircraft in raptor surveys. Raptor Res. 7:97-104.

White, C. M., and T. L. Thurow. 1985. Reproduction of Ferruginous Hawks exposed to controlled disturbance. Condor 87:14-22.

White, D.H., L. E. Hayes, and P. B. Bush. 1989. Case histories of wild birds killed intentionally with famphur in Georgia and West Virginia. Journal of Wildlife Diseases 25:144-188.

Wilcove, D.S. 1985. Nest predation in forest tracts and the decline of migratory songbirds. Ecology 66: 1211-1214.

Wilkinson, P. M. 1962. A life history study of the beaver in east-central Alabama. Thesis, Auburn University, Auburn, Alabama.

Williams, D. E., and R. M. Corrigan. 1994. Chipmunks. Pp B13-B16 *in* S. E. Hygnstrom, R. M. Timm and G. E. Larson, eds. Prevention and control of wildlife damage. University of Nebraska-Lincoln, Lincoln, Nebraska.

Williams, E. S., and I. K. Barker, editors. 2001. Infectious diseases of wild mammals. Iowa State University Press, Ames, Iowa.

Windberg, L.A., and F.F. Knowlton. 1988. Management implications of coyote spacing patterns in southern Texas. Journal of Wildlife Management 52:632-640.

Winter. 2004. Trap-neuter-release programs: The reality and the impacts. Journal of the American Veterinary Medical Association 225:1369-1376.

Wiseman G.L., and G.O. Hendrickson. 1950. Notes on the life history of the opossum in southeast Iowa. Journal of Mammalogy 31:331-337.

Wolfe, J. L. 1982. *Oryzomys palustris*. Mammalian Species 176:1-5. American Society of Mammalogy, New York.

Wolfe, J. L., and A. V. Linzey. 1977. *Peromyscus gossypinus*. Mammalian Species 70:1-5. American Society of Mammalogy, New York.

Wood, G. W., and R. H. Barrett. 1979. Status of the wild pig in the United States. Wildlife Society Bulletin 35:237-246.

Wood, G. W., L. A. Woodward, and G. Yarrow. 1994. The Clemson beaver pond leveler. Clemson University Department of Aquaculture, Fisheries and Wildlife Extension Leaflet 1, March 1994. 4 pp.

Woodward, D. K., J. D. Hair, and B. P. Gaffney. 1976. Status of beaver in South Carolina as determined by a postal survey of landowners. Proceedings of the Annual Conference of Southeast Fish and Wildlife Agencies 30:448-54.

Woodward, D. K., R. B. Hazel, and B. P. Gaffney. 1985. Economic and environmental impacts of beavers in North Carolina. Proceedings of the Eastern Wildlife Damage Management Conference 2:89-96.

Woolington, J.D. 1984. Habitat use and movements of river otters at Kelp Bay, Baranof Island, Alaska. M.S. Thesis, University of Alaska, Fairbanks, Alaska.

Wright, J. P., C. G. Jones, and A. S. Flecker. 2002. An ecosystem engineer, the beaver, increases species richness on a landscape scale. Oecologia 132:96-101.

Yeager, L. E., and R. G. Rennels. 1943. Fur yield and autumn foods of the raccoon in Illinois river bottom lands. Journal of Wildlife Management 7:45–60.

Yates, T.L., and D.J. Schmidly. 1978. Mammalian Species. *Scalopus aquaticus*. The American Society of Mammalogists 105:1-4.

Yates, T. L., and R. J. Pederson. 1982. Moles. Pp 37-51 *in* J. A. Chapman and G. A. Feldhamer, eds. Wild mammals of North America: biology, management, and economics. Johns Hopkins University Press, Baltimore, Maryland.

Zasloff, R.L. 1996. Human-animal interactions. Special Issue. Applied Animal Behaviour Science. 47: 43-48.

APPENDIX B

METHODS AVAILABLE FOR RESOLVING OR PREVENTING MAMMAL DAMAGE IN TENNESSEE

The most effective approach to resolving wildlife damage problems would be to integrate the use of several methods, either simultaneously or sequentially. An adaptive plan would integrate and apply practical methods of prevention and reduce damage by wildlife while minimizing harmful effects of damage reduction measures on humans, other species, and the environment. An adaptive plan may incorporate resource management, physical exclusion and deterrents, and population management, or any combination of these, depending on the characteristics of specific damage problems.

In selecting damage management techniques for specific damage situations, consideration would be given to the responsible species and the magnitude, geographic extent, duration and frequency, and likelihood of wildlife damage. Consideration would also be given to the status of target and potential non-target species, local environmental conditions and impacts, social and legal aspects, and relative costs of damage reduction options. The cost of damage reduction may sometimes be a secondary concern because of the overriding environmental, legal, and animal welfare considerations. Those factors would be evaluated in formulating damage management strategies that incorporate the application of one or more techniques.

A variety of methods would potentially be available to the WS program in Tennessee relative to the management or reduction of damage from mammals. Various federal, state, and local statutes and regulations and WS directives would govern WS' use of damage management methods. WS would develop and recommend or implement strategies based on resource management, physical exclusion, and wildlife management approaches. Within each approach there may be available a number of specific methods or techniques. The following methods could be recommended or used by the WS program in Tennessee. Many of the methods described would also be available to other entities in the absence of any involvement by WS.

Non-chemical Wildlife Damage Management Methods

Non-chemical management methods consist primarily of tools or devices used to repel, capture, or kill a particular animal or local population of wildlife to alleviate damage and conflicts. Methods may be non-lethal (*e.g.*, fencing, frightening devices) or lethal (*e.g.*, firearms, body gripping traps). If WS' personnel apply those methods, a MOU, cooperative service agreement, or other similar document must be signed by the landowner or administrator authorizing the use of each damage management method. Non-chemical methods used or recommended by WS could include:

Exclusion pertains to preventing access to resources through fencing or other barriers. Fencing of small critical areas can sometimes prevent animals that cannot climb from entering areas of protected resources. Fencing of culverts, drainpipes, and other water control structures can sometimes prevent beaver from building dams that plug those devices. Fencing installed with an underground skirt can prevent access to areas for many mammal species that dig, including fox, feral cats, and striped skunks. Areas such as airports, yards, or hay meadows may be fenced. Hardware cloth or other metal barriers can sometimes be used to prevent girdling and gnawing of valuable trees and to prevent the entry of mammals into buildings through existing holes or gaps. Construction of concrete spillways may reduce or prevent damage to dams by burrowing aquatic rodent species. Riprap can also be used on dams and levees to deter muskrat, woodchuck, and other burrowing rodents. Exclusion and one-way devices such as netting or nylon window screening can be used to exclude bats from a building or an enclosed structure (Greenhall and Frantz 1994). Electric fences of various

constructions have been used effectively to reduce damage to various crops by deer, raccoons, and other species (Boggess 1994*a*, Craven and Hygnstrom 1994).

Beaver exclusion and the use of water control devices could be recommended or implemented by WS to alleviate flooding damage without removing beaver under the alternatives. Although dams could be breached/removed manually or with binary explosives, those methods are usually ineffective because beaver quickly repair or replace the dam (McNeely 1995). Damage may be effectively reduced in some situations by installing exclusion and water control devices. Exclusion and water control devices can be designed so that the level of the beaver-created pond can be managed to eliminate or minimize damage while retaining the ecological and recreational benefits derived from beaver ponds. WS could also recommend that modifications occur to culvert design (Jensen et al. 1999) as a non-lethal way of reducing problems with beaver dams at culverts.

Beaver exclusion generally involves the placement of fencing to prevent beaver from accessing water intake areas, such as culverts. A variety of exclusion systems could be recommended or implemented by WS, including the Beaver Deceiver™, Beaver Bafflers™, and pre-dams (Lisle 1996, Brown and Brown 1999, Lisle 1999, Brown et al. 2001, Partington 2002, Lisle 2003). The Beaver Deceiver™ is a fencing system that is installed to prevent beaver blockage of culverts by minimizing environmental cues that stimulate beaver to construct dams, and by making culverts less attractive as dam construction sites (Lisle 1996, Lisle 1999, Lisle 2003). Beaver can be deterred from blocking culverts by the installation of a fence on the upstream end of the culvert. Installation of a fence increases the length of the area that must be dammed to impound water, and if beaver build along the fence, may increase the distance between the beaver and the source of the cues that stimulate damming behavior (*e.g.*, water moving through culvert) (Lisle 1996, Lisle 1999, Lisle 2003, Callahan 2005). Beaver prefer to build dams perpendicular to water flow, so fences can be oriented at odd angles to water flow and can be set so that they do not block the stream channel. Fencing can also be used to cover the up and downstream ends of the culverts to prevent beaver from entering the deceiver from the downstream side of the culvert and to prevent any beaver that might make it past the outer fence from plugging the interior of the culvert. Efforts can also be made to reduce the sound of water flowing through the culvert by raising the water level on the down-stream side of the culvert with dam boards or beaver-made dams; by constructing flumes to replace waterfalls, or, in extreme cases, by resetting the culvert (Lisle 1996). To ensure sufficient water flow through the culvert, Beaver Deceivers™ may be used in combination with water control devices (see discussion on Beaver Deceivers™ below).

Cylindrical exclusion devices like the Beaver Bafflers™ can be attached to culvert openings to reduce the likelihood that beaver plug a culvert by spreading the water intake over a larger area (Brown et al. 2001). While cylindrical exclusion devices can be effective in some situations (Partington 2002), in a study of beaver exclusion and water control devices, cylindrical shapes attached in-line with a culvert had a higher failure rate (40%) than trapezoidal shapes (*e.g.*, Beaver Deceivers™; 3% failure rate) and use of the cylindrical devices was discontinued in favor of trapezoidal fences (Callahan 2005).

Unlike Beaver Deceivers™ and cylindrical fences, pre-dam fences (*e.g.*, deep-water fences, diversion dams) (Brown and Brown 1999) can be designed with the specific intention that the beaver build the dam along the fence. Pre-dam fences can be short semicircular or circular fences that are built in an arc around a water inlet. The fence serves as a dam construction platform that allows beaver to build a dam and pond at the site but prevents beaver from plugging the water intake. If the size of the upstream pond created from the impounded water were not a concern, no further modifications of the pre-dam would be needed. However, in most cases, pre-dams would be used in combination with water control devices to manage the size of the upstream pond to alleviate flooding concerns.

Fence mesh size can be selected to minimize risks to beaver and non-target species. Brown et al. (2001) noted that beaver occasionally became stuck in 6-inch mesh and that the risk of beaver entrapment was lower with 5-inch mesh. Lisle (1999) noted that the size of the mesh on the fence of the Beaver Deceivers™ (6-inch mesh) was such that it allowed most species to pass through the fence except beaver and big turtles. In some remote areas where vehicular traffic is infrequent, it may be acceptable for animals that cannot pass through the fence mesh to travel across the road. However, for culverts under busy roads, it may be necessary to design special "*doors*" that allow the passage of beaver, large turtles, and other non-targets through the device. For example, T-joints 30 centimeters in diameter have been used to allow access through Beaver Deceiver™ fences. The T-shape reduces the likelihood that beaver can haul woody debris for dam construction inside the device (Lisle 2003). Fence caps would not be attached to the up and down-stream ends of a culvert when it is necessary to allow passage of species like large turtles and beavers through a culvert.

Water control devices (*e.g.*, pond levelers) are systems that allow the passage of water through a beaver dam. The devices could be used in situations where the presence of a beaver pond is desired but it is necessary to manage the level of water in the pond. Various types of water control devices have been described (Arner 1964, Roblee 1984, Laramie and Knowles 1985, Miller and Yarrow 1994, Wood et al. 1994, Lisle 1996, Organ et al. 1996, Brown and Brown 1999, Lisle 1999, Brown et al. 2001, Close 2003, Lisle 2003, Clemson University 2006, Simon 2006, Spock 2006, Perry 2007). The devices generally involve the use of one or more pipes installed through the beaver dam to increase the flow of water through the dam. Height and placement of pipes can be adjusted to achieve the desired water level in the beaver pond. Beaver generally only check the dam for leaks, so, when site conditions permit, the inlet of the pipe is placed away from the dam to make the source of the water flow more difficult to detect and decrease the likelihood that beaver will attempt to plug the device. To minimize the sound/sensation of water movement and the associated beaver damming behavior, the end of the pipe may be capped with a series of holes or notches cut in the pipe, which allows water to flow into the pipe. Holes and notches may be placed on the underside of the pipe to reduce the sound of water movement. Alternatively, 90-degree elbow joints can be placed facing downward on the upstream end of the pipes to prevent the noise of running water and attracting beaver. A protective cage can be placed around the upstream end of the inlet pipe to prevent beaver from blocking the pipe and to reduce problems with debris blocking the pipe. As noted above, water control systems can be combined with exclusion devices to prevent beaver from blocking culverts while still maintaining a beaver pond at an acceptable level.

Cultural Methods and Habitat Management includes the application of practices that seek to minimize exposure of the protected resource to damaging animals through processes other than exclusion. They may include animal husbandry practices such as employing guard dogs, herders, shed lambing, carcass removal, or pasture selection. Strategies may also include minimizing cover where damaging mammals might hide, manipulating the surrounding environment through barriers or fences to deter animals from entering a protected area, or planting lure crops on fringes of protected crops. Continual destruction of beaver dams and removal of dam construction materials on a daily basis will sometimes cause beavers to move to other locations. Water control devices such as the 3-log drain (Roblee 1983), the T-culvert guard (Roblee 1987), wire mesh culvert (Roblee 1983), and the Clemson beaver pond leveler (Miller and Yarrow 1994) can sometimes be used to control the water in beaver ponds to desirable levels that do not cause damage. Removal of trees from around buildings can sometimes reduce damage associated with tree squirrels and raccoons.

Some mammals that cause damage in urban environments are attracted to homes by the presence of garbage or pet food left outside and unprotected. Removal or sealing of garbage in tight trash receptacles, and elimination of all pet foods from outside areas can reduce the presence of unwanted mammals. If raccoons are a problem, making trash and garbage unavailable, and removing all pet

food from outside during nighttime hours can reduce their presence. Altering how bird feeders are hung and constructing mounting poles for the feeders that cannot be climbed by tree squirrels or chipmunks can reduce the presence of localized populations along with their associated damage.

Beaver dam removal is generally conducted to maintain existing stream channels and drainage patterns, and reduce floodwaters that have affected established silviculture, agriculture, or drainage structures, such as culverts. Beaver dams are made from natural debris such as logs, sticks and mud that beaver take from the immediate area and impound water, creating habitat that they utilize to build lodges and bank dens to raise their young and/or provide protection from predators. The impoundments that WS removes are typically created by recent beaver activity and have not been in place long enough to take on the qualities of a true wetland (*e.g.*, hydric soils, aquatic vegetation, pre-existing function). Unwanted beaver dams can be removed by hand with a rake or power tools (*e.g.*, a winch), or with explosives. Explosives are used only by WS' personnel specially trained and certified to conduct such activities, and only binary explosives are used (*i.e.*, they are comprised of two parts that must be mixed at the site before they can be detonated as an explosive material). Beaver dam removal by hand or with binary explosives does not affect the substrate or the natural course of the stream and returns the area back to its pre-existing condition with similar flows and circulations. Because beaver dams involve waters of the United States, removal is regulated under Section 404 of the Clean Water Act (see Appendix E).

Supplemental feeding is sometimes used to reduce damage by wildlife, such as lure crops. Food is provided so that the animal causing damage would consume it rather than the resource being protected. In feeding programs, target wildlife would be offered an alternative food source with a higher appeal with the intention of luring them from feeding on affected resources.

Animal behavior modification refers to tactics that deter or repel damaging mammals and thus, reduce damage to the protected resource. Those techniques are usually aimed at causing target animals to respond by fleeing from the site or remaining at a distance. They usually employ extreme noise or visual stimuli. Unfortunately, many of these techniques are only effective for a short time before wildlife habituate to them (Conover 1982). Devices used to modify behavior in mammals include electronic guards (siren strobe-light devices), propane exploders, pyrotechnics, laser lights, human effigies, effigies of predators, and the noise associated with the discharge of a firearm.

Live Capture and Translocation can be accomplished using hand capture, hand nets, catch poles, cage traps, suitcase type traps, cable restraints, or with foothold traps to capture some mammal species for the purpose of translocating them for release in other areas. WS could employ those methods in Tennessee when the target animal(s) can legally be translocated or can be captured and handled with relative safety by WS' personnel. Live capture and handling of mammals poses an additional level of human health and safety threat if target animals are aggressive, large, or extremely sensitive to the close proximity of humans. For that reason, WS may limit this method to specific situations and certain species. In addition, moving damage-causing individuals to other locations can typically result in damage at the new location, or the translocated individuals can move from the relocation site to areas where they are unwanted. In addition, translocation can facilitate the spread of diseases from one area to another. The AVMA, the National Association of State Public Health Veterinarians, and the Council of State and Territorial Epidemiologists all oppose the relocation of mammals because of the risk of disease transmission (CDC 1990). Although translocation is not necessarily precluded in all cases, it would be logistically impractical, in most cases, and biologically unwise in Tennessee due to the risk of disease transmission. High population densities of some populations may make this a poor wildlife management strategy for those species. Translocation would be evaluated by WS on a case-by-case basis. Translocation would only occur with the prior authorization of the TWRA.

Trapping can utilize a number of devices, including foothold traps, cage-type traps, and body gripping (Conibear) traps, foot snares, and neck/body snares. Those techniques are implemented by WS' personnel because of the technical training required to use such devices.

Foothold Traps can be effectively used to capture a variety of mammals. Foothold traps are placed either beside or in travel ways being actively used by the target species. Placement of traps is contingent upon the habits of the respective target species, habitat conditions, and presence of non-target animals. Effective trap placement and adjustment, and the use and placement of appropriate baits and lures by trained WS' personnel also contribute to the foothold trap's selectivity. An additional advantage is that foothold traps can allow for the on-site release of non-target animals since animals are captured alive. The use of foothold traps requires more skill than some methods.

Cable Restraints are typically made of wire or cable, and can be set to capture an animal by the neck, body, and foot. Cable restraints may be used as either lethal or live-capture devices depending on how or where they are set. Cable restraints set to capture an animal by the neck are usually lethal but stops can be attached to the cable to increase the probability of a live capture depending on the trap check interval. Snares positioned to capture the animal around the body can be a useful live-capture device, but are more often used as a lethal control technique. Snares can incorporate a breakaway feature to release non-target wildlife and livestock where the target animal is smaller than potential non-targets (Phillips 1996). Snares can be effectively used wherever a target animal moves through a restricted travel lane (*e.g.*, under fences or trails through vegetation). When an animal moves forward into the loop formed by the cable, the noose tightens and the animal is held. Snares must be set in locations where the likelihood of capturing non-target animals is minimized.

The foot or leg snare can be set as a spring-powered non-lethal device, activated when an animal places its foot on the trigger or pan. In some situations, using snares to capture wildlife is impractical due to the behavior or morphology of the animal, or the location of many wildlife conflicts.

Cage traps come in a variety of styles to live-capture animals. The most commonly known cage traps are box traps and corral traps. Box traps are usually rectangular and are made from various materials, including metal, wire mesh, plastic, and wood. These traps are well suited for use in residential areas and work best when baited with foods attractive to the target animal. Box traps are generally portable and easy to set-up.

Corral traps for feral swine are generally large circular traps consisting of panels anchored to the ground using steel posts with a door allowing entrance. Side panels are typically woven metal fencing referred to as hog panels or cow panels. The entrances into the traps generally consist of a door that allow entry into the trap but prevents exit. The doors are often designed to allow swine to continually enter the trap that allows for the possibility of capturing multiple swine.

The disadvantages of using cage traps are: 1) some individual target animals may avoid cage traps; 2) some non-target animals may associate the traps with available food and purposely get captured to eat the bait, making the trap unavailable to catch target animals; 3) cage traps must be checked frequently to ensure that captured animals are not subjected to extreme environmental conditions; and 4) some animals will fight to escape and may become injured; 5) expense of purchasing traps. Disadvantages associated with corral traps include: 1) the expense of purchasing the materials to construct trap, 2) once constructed, corral traps are not moveable until

disassembled and transported, and 3) in remote areas, getting all the required equipment to the location can be difficult.

Trap monitors are devices that send a radio signal to a receiver if a set trap is disturbed and alerts field personnel that an animal may be captured. Trap monitors can be attached directly to the trap or attached to a string or wire and then placed away from the trap in a tree or shrub. When the monitor is hung above the ground, it can be detected from several miles away, depending on the terrain in the area. There are many benefits to using trap monitors, such as saving considerable time when checking traps, decreasing fuel usage, prioritizing trap checks, and decreasing the need for human presence in the area. Trap monitors could be used when using cage traps.

Trap monitoring devices would be employed, when applicable, that indicate when a trap has been activated. Trap monitoring devices would allow personnel to prioritize trap checks and decrease the amount of time required to check traps, which decreases the amount of time captured target or non-targets would be restrained. By reducing the amount of time targets and non-targets are restrained, pain and stress can be minimized and captured wildlife can be addressed in a timely manner, which could allow non-targets to be released unharmed. Trap monitoring devices could be employed where applicable to facilitate monitoring of the status of traps in remote locations to ensure any captured wildlife was removed promptly to minimize distress and to increase the likelihood non-targets could be released unharmed.

Hancock/Bailey Traps (suitcase/basket type cage traps) are designed to live-capture beaver. The trap is constructed of a metal frame that is hinged with springs attached and covered with chain-link fence. The trap's appearance is similar to a large suitcase when closed. When set, the trap is generally baited and opened to allow an animal to enter. When tripped, the panels of the trap close around the animal capturing the animal. One advantage of using the Hancock or Bailey trap is the ease of release of beaver or non-target animals. Beaver caught in Hancock or Bailey traps could also be humanely euthanized. Disadvantages are that those traps are very expensive (>$300 per trap), cumbersome, and difficult to set (Miller and Yarrow 1994). The trap weighs about 25 pounds and is relatively bulky to carry and maneuver. Hancock and Bailey traps can also be dangerous to set (*i.e.*, hardhats are recommended when setting suitcase traps), are less cost and time-efficient than snares, footholds, or body-grip traps, and may cause serious and debilitating injury to river otters (Blundell et al. 1999).

Body-grip Traps are designed to cause the quick death of the animal that activates the trap. Body-grip traps may include snap traps, mole traps, and conibear traps. The conibear trap consists of a pair of rectangular wire frames that close like scissors when triggered, killing the captured animal with a quick body blow. For conibear traps, the traps should be placed so ensure the rotating jaws close on either side of the neck of the animal to ensure a quick death. Conibear traps are lightweight and easily set. Snap traps are common household rat or mouse traps. These traps are often used to collect and identify rodent species that cause damage so that species-specific control tools can be applied, such as identifying the prey base at airports. Spring-powered harpoon traps are used to control damage caused by surface-tunneling moles. Soil is pressed down in an active tunnel and the trap is placed at that point. When the mole reopens the tunnel, it triggers the trap. Two variations of scissor like traps are also used in tunnels for moles. Safety hazards and risks to humans are usually related to setting, placing, checking, or removing the traps. Body-grip traps present a minor risk to non-target animals. Selectivity of body-grip traps can be enhanced by placement, trap size, trigger configurations, and baits. When using body-grip traps, risks of non-target capture can be minimized by using recessed sets (placing trap inside a cubby, cage, or burrow), restricting openings, or by elevating traps. For example, conibear traps set to capture beaver can be placed underwater to minimize risks to non-targets.

Choosing appropriately sized traps for the target species can also exclude non-targets by preventing larger non-targets from entering and triggering the trap. The trigger configurations of traps can be modified to minimize non-target capture. For example, offsetting the trigger can allow non-targets to pass through conibear traps without capture.

Shooting with firearms is very selective for the target species and would be conducted with rifles, handguns, and shotguns. Methods and approaches used by WS may include use of vehicles or aircraft, illuminating devices, bait, firearm suppressors, night vision/thermal equipment, and elevated platforms. Shooting is an effective method in some circumstances, and can often provide immediate relief from the problem. Shooting may at times be one of the only methods available to effectively and efficiently resolve a wildlife problem.

Ground shooting is sometimes used as the primary method to alleviate damage or threats of damage. Shooting is limited to locations where it is legal and safe to discharge a weapon. A shooting program, especially conducted alone, can be expensive because it often requires many staff hours to complete.

Shooting can also be used in conjunction with an illumination device at night, which is especially useful for nocturnal mammals, such as deer or feral swine. Spotlights may or may not be covered with a red lens, which nocturnal animals may not be able to see, making it easier to locate them undisturbed. Night shooting may be conducted in sensitive areas that have high public use or other activity during the day, which would make daytime shooting unsafe. The use of night vision and Forward Looking Infrared (FLIR) devices can also be used to detect and shoot mammals at night, and is often the preferred equipment due to the ability to detect and identify animals in complete darkness. Night vision and FLIR equipment aid in locating wildlife at night when wildlife may be more active. Night vision and FLIR equipment could be used during surveys and in combination with shooting to remove target mammals at night. WS' personnel most often use this technology to target mammals in the act of causing damage or likely responsible for causing damage. Those methods aid in the use of other methods or allow other methods to be applied more selectively and efficiently. Night vision and FLIR equipment allow for the identification of target species during night activities, which reduces the risks to non-targets and reduces human safety risks. Night vision equipment and FLIR devices only aid in the identification of wildlife and are not actual methods of take. The use of FLIR and night vision equipment to remove target mammals would increase the selectivity of direct management activities by targeting those mammals most likely responsible for causing damage or posing threats.

Denning is the practice of locating coyote or fox dens and killing the young, adults or both to stop an ongoing predation problem or prevent future depredation of livestock. Coyote and red fox depredations on livestock often increase in the spring and early summer due to the increased food requirements associated with feeding and rearing litters of pups. Removal of pups will often stop depredations even if the adults are not taken (Till 1992). Pups are typically euthanized in the den using a registered gas fumigant cartridge or by digging out the den and euthanizing the pups with sodium pentobarbital (see discussion of gas cartridges and sodium pentobarbital under *Chemical Wildlife Damage Management Methods*).

Hunting/Trapping: WS sometimes recommends that resource owners consider legal hunting and trapping as an option for reducing mammal damage. Although legal hunting/trapping is impractical and/or prohibited in many urban-suburban areas, it can be used to reduce some populations of mammals.

Aerial Shooting or aerial hunting (*i.e.*, shooting from an aircraft) is a commonly used coyote damage management method; it can be especially effective in removing offending coyotes that have become

"*bait-shy*" to trap sets or are not susceptible to calling and shooting. Aerial shooting is one of the preferred damage management methods for reducing feral swine damage as well, in that local swine populations can quickly be removed when weather and habitat conditions are favorable. Aerial hunting is mostly species-selective (there is a slight potential for misidentification) and can be used for immediate control to reduce livestock and natural resource losses if weather, terrain, and cover conditions are favorable. WS has also used aerial hunting for disease surveillance (*e.g.*, taking deer samples for chronic wasting disease and searching for carcasses in areas where an anthrax outbreak has occurred). Fixed-wing aircraft are most frequently used in flat and gently rolling terrain whereas helicopters with better maneuverability have greater utility and are safer over rugged terrain and timbered areas.

In broken timber or deciduous cover, aerial hunting is more effective in winter when snow cover improves visibility and leaves have fallen. The WS program aircraft-use policy helps ensure that aerial hunting is conducted in a safe and environmentally sound manner, in accordance with federal and state laws. Pilots and aircraft must be certified under established WS program procedures and only properly trained WS' employees are approved as gunners. Ground crews are often used with aerial operations for safety reasons and to assist locating and recovering target animals, as necessary.

Aircraft overflights have created concerns about disturbing wildlife. The National Park Service (1995) reviewed studies on the effects of aircraft overflights on wildlife. Their report revealed that a number of studies documented responses by certain wildlife species that could suggest adverse impacts may occur. Few, if any studies, have proven that aircraft overflights cause significant adverse impacts to wildlife populations, although the report stated it is possible to draw the conclusion that affects to populations could occur. It appears that some species will frequently, or at least occasionally, show adverse responses to even minor overflight occurrences. In general, it appears that the more serious potential impacts occur when overflights are frequent, such as hourly, and over long periods of time, which represents chronic exposure. Chronic exposure situations generally occur in areas near commercial airports and military flight training facilities. WS' aerial hunting operations occur in remote rangeland areas where tree cover is, at most, scattered and allows for visibility of target animals from the air. WS spends relatively little time over any one area.

WS has used fixed-wing aircraft and helicopters for aerial hunting in areas inhabited by wildlife for years. WS conducts aerial activities on areas only under signed agreement and concentrates efforts during certain times of the year and to specific areas. WS' Predator Damage Management Environmental Assessments (*e.g.*, USDA 2005*c*) that have looked at the issue of aerial hunting overflights on wildlife have found that WS has annually flown less than 10 min./mi.^{2} on properties under agreements. WS flies very little over any one property under agreement in any given year. As a result, no known problems to date have occurred with WS' aerial hunting overflights on wildlife, nor are they anticipated in the future.

Aerial Surveying is a commonly used tool for evaluating and monitoring damage and establishing population estimates and locations of various species of wildlife. WS uses aerial surveying throughout the United States to monitor damages and/or populations of coyotes, fox, wolves, feral swine, feral goats, feral dogs, bobcats, mountain lions, white-tailed deer, pronghorn antelope, elk, big-horn sheep, and wild horses but any wildlife species big enough to see from a moving aircraft could be surveyed using this method. As with aerial shooting, the WS program aircraft-use policy helps ensure that aerial surveys are conducted in a safe and environmentally sound manner, in accordance with Federal and State laws. Pilots and aircraft must also be certified under established WS program procedures and policies.

Aerial Telemetry is used in research projects studying the movements of various wildlife species. Biologists will frequently place radio-transmitting collars on selected individuals of a species and then monitor their movements over a specified period. Whenever possible, the biologist attempts to locate the research subject using a hand-held antennae and radio receiver, however, occasionally animals will make large movements that prevent biologists from locating the animal from the ground. In these situations, WS can utilize either fixed wing aircraft or helicopters and elevation to conduct aerial telemetry and locate the specific animal wherever it has moved to. As with any aerial operations, the WS program aircraft-use policy helps ensure that aerial surveys would be conducted in a safe and environmentally sound manner, in accordance with Federal and State laws.

Chemical Wildlife Damage Management Methods

All pesticides used by WS are registered under the FIFRA and administered by the EPA and the TDA. All WS personnel in Tennessee who apply restricted-use pesticides are certified pesticide applicators by TDA and have specific training by WS for pesticide application. The EPA and the TDA require pesticide applicators to adhere to all certification requirements set forth in the FIFRA. Pharmaceutical drugs, including those used in wildlife capture and handling, are administrated by FDA and/or DEA.

Chemicals would not be used by WS on public or private lands without authorization from the land management agency or property owner or manager. The following chemical methods have been proven to be selective and effective in reducing damage by mammals.

GonaCon™ was developed by scientists with the NWRC as a reproductive inhibitor. GonaCon™ is a new single dose immunocontraceptive vaccine. Recent studies have demonstrated the efficacy of this single-shot Gonadotropin-releasing hormone (GnRH) vaccine on California ground squirrels, Norway rats, feral cats and dogs, feral swine, wild horses, and white-tailed deer. Infertility among treated female swine and white-tailed deer has been documented for up to two years without requiring a booster vaccination (Miller et al. 2000). This vaccine overcomes one of the major obstacles of previous two dose vaccines since target wildlife need to be captured only once for vaccination instead of twice. A single-injection vaccine would be much more practical as a field delivery system for use on free-ranging animals.

GonaCon™ was officially registered by the EPA in 2009 for use in reducing fertility in female white-tailed deer under EPA registration number 56228-40. GonaCon™ is registered as a restricted-use pesticide available for use by WS' personnel and personnel of a state wildlife management agency or persons under their authority. Additionally, in order for GonaCon™ to be used in any given state, the product must also be registered with the state and approved for use by the appropriate state agency responsible for managing wildlife. GonaCon™, when injected into the body, elicits an immune response that neutralizes the GnRH hormone being produced naturally by deer. The GnRH hormone in deer stimulates the production of other sexual hormones, which leads to the body reaching a reproductive state. The vaccine neutralizes the GnRH hormone being produced, which then prevents the production of other sexual hormones in the deer vaccinated; thereby, preventing the body of the deer from entering into a reproductive state (USDA 2010).

Ketamine (Ketamine HCl) is a dissociative anesthetic that is used to capture wildlife, primarily mammals, birds, and reptiles. It is used to eliminate pain, calm fears, and allay anxiety. Ketamine is possibly the most versatile drug for chemical capture, and it has a wide safety margin (Fowler and Miller 1999). When used alone, this drug may produce muscle tension, resulting in shaking, staring, increased body heat, and, on occasion, seizures. Usually, ketamine is combined with other drugs such as xylazine. The combination of such drugs is used to control an animal, maximize the reduction of stress and pain, and increase human and animal safety.

Telazol (tiletamine) is another anesthetic used in wildlife capture. It is 2.5 to 5 times more potent than ketamine; therefore, it generally works faster and lasts longer. Currently, tiletamine can only be purchased as Telazol, which is a mixture of two drugs: tiletamine and zolazepam (a tranquilizer). Muscle tension varies with species. Telazol produces extensive muscle tension in dogs, but produces a more relaxed anesthesia in coyotes, wolves, and bears. It is often the drug of choice for those wild species (Fowler and Miller 1999). This drug is sold in a powder form and must be reconstituted with sterile water before use. Once mixed with sterile water, the shelf life is four days at room temperature and 14 days if refrigerated.

Xylazine is a sedative (analgesic) that calms nervousness, irritability, and excitement, usually by depressing the central nervous system. Xylazine is commonly used with ketamine to produce a relaxed anesthesia. It can also be used alone to facilitate physical restraint. Because xylazine is not an anesthetic, sedated animals are usually responsive to stimuli. Therefore, personnel should be even more attentive to minimizing sight, sound, and touch. When using ketamine/xylazine combinations, xylazine will usually overcome the tension produced by ketamine, resulting in a relaxed, anesthetized animal (Fowler and Miller 1999). This reduces heat production from muscle tension, but can lead to lower body temperatures when working in cold conditions.

Sodium Pentobarbital is a barbiturate that rapidly depresses the central nervous system to the point of respiratory arrest. There are DEA restrictions on who can possess and administer this drug. Some states may have additional requirements for personnel training and particular sodium pentobarbital products available for use in wildlife. Certified WS personnel are authorized to use sodium pentobarbital and dilutions for euthanasia in accordance with DEA and state regulations. All animals euthanized using sodium pentobarbital and all of its dilutions (*e.g.* Beuthanasia-D, Fatal-Plus) are disposed of immediately through incineration or deep burial to prevent secondary poisoning of scavenging animals and introduction of these chemicals to non-target animals.

Potassium Chloride used in conjunction with prior general anesthesia is used as a euthanasia agent for animals, and is considered acceptable and humane by the AVMA (2007). Animals that have been euthanized with this chemical experience cardiac arrest followed by death, and are not toxic to predators or scavengers.

Beuthanasia®-D combines pentobarbital with another substance to hasten cardiac arrest. Intravenous (IV) and intracardiac (IC) are the only acceptable routes of injection. As with pure sodium pentobarbital, IC injections with Beuthanasia®-D are only acceptable for animals that are unconscious or deeply anesthetized. With other injection routes, there are concerns that the cardiotoxic properties may cause cardiac arrest before the animal is fully unconscious. It is a Schedule III drug, which means it can be obtained directly from the manufacturer by anyone with a DEA registration. However, Schedule III drugs are subject to the same security and record-keeping requirements as Schedule II drugs.

Fatal-Plus® combines pentobarbital other substances to hasten cardiac arrest. IV is the preferred route of injection; however, IC is acceptable as part of the two-step procedure used by WS. Animals are first anesthetized and sedated using a combination of ketamine/xylazine and once completely unresponsive to stimuli and thoroughly sedated, Fatal-Plus® is administered. Like Beuthanasia®-D, it is a Schedule III drug requiring a DEA registration for purchase and is subject to the security and record-keeping requirements of Schedule II drugs.

Carbon Dioxide is sometimes used to euthanize mammals that are captured in live traps and when relocation is not a feasible option. Live mammals are placed in a sealed chamber. CO_2 gas is

released into the chamber and the animal quickly dies after inhaling the gas. This method is approved as a euthanizing agent by the AVMA. CO_2 gas is a byproduct of animal respiration, is common in the atmosphere, and is required by plants for photosynthesis. It is used to carbonate beverages for human consumption and is the gas released by dry ice. The use of CO_2 by WS for euthanasia purposes is exceedingly minor and inconsequential to the amounts used for other purposes by society.

Zinc phosphide is an inorganic compound used to control rats, mice, voles, ground squirrels, prairie dogs, nutria, muskrats, feral rabbits, and gophers. Zinc phosphide is a heavy, finely ground gray-black powder that is partially insoluble in water and alcohol. When exposed to moisture, it decomposes slowly and releases phosphine gas (PH_3). When zinc phosphide treated bait encounters acids in the stomach, phosphate (PH3) gas is released, which may account in a large part for observed toxicity. Animals that ingest lethal amounts of bait usually succumb overnight with terminal symptoms of convulsions, paralysis, coma, and death from asphyxia. If death is prolonged for several days, intoxication that occurs is similar to intoxication with yellow phosphorous, in which the liver is heavily damaged. Prolonged exposure to phosphine can produce chronic phosphorous poisoning.

Although zinc phosphide baits have a strong, pungent, phosphorous-like odor (garlic like), this characteristic seems to attract rodents, particularly rats, and apparently makes the bait unattractive to some other animals. For many uses of zinc phosphide formulated on grain or grain-based baits, pre-baiting is recommended or necessary for achieving good bait acceptance. Primary toxicity risks to non-target species from the direct consumption of treated bait can be minimized by using bait stations to prevent access by non-target species such as birds.

Because zinc phosphide is not stored in muscle or other tissues of poisoned animals, there is no secondary poisoning with this rodenticide. The bait however, remains toxic up to several days in the gut of the dead rodent. Other animals can be poisoned if they eat enough of the gut content of rodents recently killed with zinc phosphide.

Aluminum phosphide is an inorganic phosphide used to control insects and rodents in a variety of settings. It is mainly used as an indoor fumigant at crop transport, storage or processing facilities (or in shipholds, railcars) for both food and non-food crops. It may also be used as an outdoor fumigant for burrowing rodent and mole control, or in baits for rodent control in crops. Aluminum phosphide is available in pellet and tablet form, and is available in porous blister packs, sachets, or as dusts. As in the case of Phostoxin, it may be formulated as 55% active ingredient along with ammonium carbamate and inert ingredients.

Aluminum phosphide causes acute toxicity with the main routes of exposure occurring through ingestion and inhalation. Dermal absorbtion is not known to occur. The reported rodent oral LD_{50} is 11.5 mg/kg for Phostoxin. Aluminum phosphide ingested orally reacts with water and stomach acids to produce phosphine gas, which may account in a large part for the observed toxicity. Phosphine generated in the gastrointestinal tract is readily absorbed in to the bloodstream, and it is readily absorbed through the lung epithelium.

In chronic toxicity studies, rats fed chow fumigated with aluminum phosphide that averaged 0.51 ppm phosphine residues (approximately 0.43 mg/kg/day) showed no differences from the control animals with respect to blood or urine chemistry and no observable differences in tissue structure. It was reported that workers had probably encountered similar exposures on an intermittent basis (in some cases over as long as a 20-year period) and had yet to show signs of toxicity, which suggests that chronic effects may be minor or have a very long latency period. Inhalation studies were conducted on the effects of phosphine gas on male and female rats exposed at levels of 0.5, 1.5, and

4.5 mg/meters cubed for six hours per day over a 13-week period. Higher exposure groups (7.5 and 15 mg/meters cubed) were added following preliminary acute test results.

Results indicated that 15 mg/m^3 were lethal to 4 out of 10 female rats following 3 days of exposure. Significant treatment-related effects on body weight and decreased food consumption were seen across all treatment groups and sexes, but were reversible. Decreases in red-blood cell counts, hemoglobin, hematocrit, and increased platelet counts were seen in male rats of the 4.5 mg/m^3 group. Dose-related changes in blood urea nitrogen and other clinical parameters were also seen across exposure groups. Post-mortem examination of test animals revealed microscopic lesions in the outer cortex of the kidneys of rats exposed to 15 mg/m^3, but not at lower exposure levels. All of those effects were apparently reversible following a four-week recovery period.

Aluminum phosphide would be used by WS in Tennessee primarily as a fumigant for small field rodents and moles. Products would be used in accordance with label restrictions in a manner defined by application guidelines on the label. Use in Tennessee is infrequent and amounts used would be very small.

Anticoagulant rodent baits with warfarin, brodifacoum, or diphacinone as active ingredients could be used in bait stations to target small rodents. WS would utilize locking bait stations to restrict access of non-target species to rodenticides such as anticoagulants. The use and proper placement of bait stations would minimize the likelihood that the bait would be consumed by non-target species. There may be secondary hazards from anticoagulant baits. Those risks would be reduced somewhat by the fact that the predator scavenger species would usually need exposure to multiple carcasses over a period of days. Areas where anticoagulants are used would be monitored and carcasses picked up and disposed of in accordance with label directions.

Repellents are usually naturally occurring substances or chemicals formulated to be distasteful or to elicit pain or discomfort for target animals when they are smelled, tasted, or contacted. Only a few repellents are commercially available for mammals, and are registered for only a few species. Repellents would not be available for many species that may present damage problems, such as some predators or furbearing species. Repellents are variably effective and depend largely on resource to be protected, time and length of application, and sensitivity of the species causing damage. Again, acceptable levels of damage control would usually not be realized unless repellents were used in conjunction with other techniques.

Gas cartridges (EPA Reg. No. 56228-1, EPA Reg. No. 56228-2) are registered by WS with the TDA and are often used to treat dens or burrows of coyotes, fox, or woodchucks. When ignited, the cartridge burns in the den of an animal and produces large amounts of carbon monoxide, a colorless, odorless, and tasteless, poisonous gas. The combination of oxygen depletion and carbon monoxide exposure kills the animals in the burrow or den. Sodium nitrate is the principle active chemical in gas cartridges and is a naturally occurring substance. Although stable under dry conditions, it is readily soluble in water and likely to be highly mobile in soils. In addition, dissolved nitrate is very mobile, moving quickly through the vadose zone to the underlying water table (Bouwer 1989). However, burning sodium nitrate, as in the use of a gas cartridge as a fumigant in a rodent burrow, is believed to produce mostly simple organic and inorganic gases, using all of the available sodium nitrate. In addition, the human health drinking water tolerance level for this chemical is 10 mg / L, a relatively large amount, according to EPA Quality Criteria for Water (EPA 1986*a*, EPA 1986*b*). The gas along with other components of the cartridge, are likely to form oxides of nitrogen, carbon, phosphorus, and sulfur. Those products are environmentally non-persistent because they are likely to be metabolized by soil microorganisms or enter their respective elemental cycles. In rodent cartridges, sodium nitrate is combined with seven additional ingredients: sulfur, charcoal, red phosphorus, mineral oil, sawdust,

and two inert ingredients. None of the additional ingredients in this formulation is likely to accumulate in soil, based on their degradation into simpler elements by burning the gas cartridge. Sodium nitrate is not expected to accumulate in soils between applications, nor does it accumulate in the tissues of target animals (EPA 1991). The EPA stated sodium nitrates "*...as currently registered for use as pesticides, do not present any unreasonable adverse effects to humans*" (EPA 1991).

Explosives are defined as any chemical mixture or device that serves as a blasting agent or detonator. The procedures and accountability for WS' use of explosive for removing beaver dams and training requirements for explosives certification would adhere to WS Directive 2.435. Explosives are generally used to breach beaver dams that are too large to remove by digging using hand tools. Explosives would be used to remove dams after the beaver were removed using other methods. WS would only use binary explosives to remove beaver dams. Binary explosives consist of two components that are contained separately. The two components of binary explosives are ammonium nitrate and nitromethane and are not classified as explosives until the two components are mixed. Therefore, binary explosives are subject to fewer regulations and controls because they are packaged separately. However, once mixed, binary explosives are considered high explosives and subject to all applicable federal and state requirements. When used to remove beaver dams, the two components would not be mixed until ready for use at the site where the dam was located. Detonating cord and detonators are also considered explosives and WS must adhere to all applicable state and federal regulations for storage, transportation, and handling. All WS' explosive specialists are required to attend extensive explosive safety training and spend time with a certified explosive specialist in the field prior to obtaining certification. All blasting activities are conducted by well-trained, certified employees and closely supervised by professional wildlife biologists in accordance with WS Directive 2.435. Explosive handling and use procedures follow the rules and guidelines set forth by the Institute of Makers of Explosives, which is the safety arm of the commercial explosive industry in the United States and Canada. WS also adheres to transportation and storage regulations from state and federal agencies, such as Occupational Safety and Health Association, Bureau of Alcohol, Tobacco, and Firearms, and the Department of Transportation.

APPENDIX C

FEDERAL LISTED THREATENED AND ENDANGERED SPECIES

TAXA	COMMON NAME	SPECIES	STATUS
Birds	Least tern	*Sterna antillarum*	E
Crustaceans	Nashville crayfish	*Orconectes shoupi*	E
Fishes	Amber darter	*Percina antesella*	E
	Blackside dace	*Phoxinus cumberlandensis*	T
	Blue shiner	*Cyprinella caerulea*	T
	Bluemask darter	*Etheostoma sp.*	E
	Boulder darter	*Etheostoma wapiti*	E
	Chucky madtom	*Noturus crypticus*	E
	Conasauga logperch	*Percina jenkinsi*	E
	Cumberland darter	*Etheostoma susanae*	E
	Duskytail darter	*Etheostoma percnurum*	E
	Goldline darter	*Percina aurolineata*	T
	Laurel dace	*Phoxinus saylori*	E
	Palezone shiner	*Notropis albizonatus*	E
	Pallid Sturgeon	*Scaphirhynchus albus*	E
	Pygmy madtom	*Noturus stanauli*	E
	Slackwater darter	*Etheostoma boschungi*	T
	Slender chub	*Erimystax cahni*	T
	Smoky madtom	*Noturus baileyi*	E
	Snail darter	*Percina tanasi*	T
	Spotfin chub	*Erimonax monachus*	T
	Yellowfin madtom	*Noturus flavipinnis*	T
Insects	American burying beetle	*Nicrophorus americanus*	E
Mammals	Carolina northern flying squirrel	*Glaucomys sabrinus coloratus*	E
	Eastern puma	*Puma concolor couguar*	E
	Florida panther	*Puma concolor coryi*	E
	Gray bat	*Myotis grisescens*	E
	Gray wolf	*Canis lupus*	E
	Indiana bat	*Myotis sodalis*	E
Mussels	Alabama lampmussel	*Lampsilis virescens*	E
	Appalachian elktoe	*Alasmidonta raveneliana*	E
	Appalachian monkeyface	*Quadrula sparsa*	E
	Birdwing pearlymussel	*Conradilla caelata*	E
	Clubshell	*Pleurobema clava*	E
	Coosa moccasinshell	*Medionidus parvulus*	E
	Cracking pearlymussel	*Hemistena lata*	E
	Cumberland bean	*Villosa trabalis*	E
	Cumberland elktoe	*Alasmidonta atropurpurea*	E
	Cumberland monkeyface	*Quadrula intermedia*	E
	Cumberland pigtoe	*Pleurobema gibberum*	E
	Cumberlandian combshell	*Epioblasma brevidens*	E
	Dromedary pearlymussel	*Dromus dromas*	E
	Fanshell	*Cyprogenia stegaria*	E
	Finelined pocketbook	*Lampsilis altilis*	T
	Finerayed pigtoe	*Fusconaia cuneolus*	E

	Georgia pigtoe	*Pleurobema hanleyianum*	E
	Green blossom	*Epioblasma torulosa gubernaculum*	E
Mussels (cont.)	Littlewing pearlymussel	*Pegias fabula*	E
	Orangefoot pimpleback	*Plethobasus cooperianus*	E
	Ovate clubshell	*Pleurobema perovatum*	E
	Oyster mussel	*Epioblasma capsaeformis*	E
	Pale lilliput	*Toxolasma cylindrellus*	E
	Pink mucket	*Lampsilis abrupta*	E
	Pink ring	*Obovaria retusa*	E
	Purple bean	*Villosa perpurpurea*	E
	Purple cat's paw	*Epioblasma obliquata obliquata*	E
	Rough rabbitsfoot	*Quadrula cylindrical strigillata*	E
	Rough pigtoe	*Pleurobema plenum*	E
	Scaleshell mussel	*Leptodea leptodon*	E
	Shiny pigtoe	*Fusconaia cor*	E
	Southern acornshell	*Epioblasma othcaloogensis*	E
	Southern clubshell	*Pleurobema decisum*	E
	Southern pigtoe	*Pleurobema georgianum*	E
	Tan riffleshell	*Epioblasma florentina walkeri* (= *E. walker*)	E
	Triangular kidneyshell	*Ptychobranchus greenii*	E
	Tubercled blossom	*Epioblasma torulosa torulosa*	E
	Turgid blossom	*Epioblasma turgidula*	E
	Upland combshell	*Epioblasma metastriata*	E
	White wartyback	*Plethobasus cicatricosus*	E
	Winged mapleleaf	*Quadrula fragosa*	E
	Yellow blossom	*Epioblasma florentina florentina*	E
Snails	Anthony's riversnail	*Athearnia anthonyi*	E
	Painted snake coiled forest snail	*Anguispira picta*	E
	Royal marstonia	*Pyrgulopsis (= Marstonia) ogmorhaphe*	E
Spiders	Spruce-fir moss spider	*Microhexura montivaga*	E
Vascular Plants	American chaffseed	*Schwalbea americana*	E
	American Hart's-tongue fern	*Asplenium scolopendrium* var. *americanum*	T
	Blue Ridge goldenrod	*Solidago spithamaea*	T
	Braun's rock-cress	*Arabis perstellata*	E
	Cumberland rosemary	*Conradina verticillata*	T
	Cumberland sandwort	*Arenaria cumberlandensis*	E
	Green pitcher-plant	*Sarracenia oreophila*	E
	Guthrie's ground-plum	*Astragalus bibullatus*	E
	Large-flowered skullcap	*Scutellaria montana*	T
	Leafy prairie-clover	*Dalea foliosa*	E
	Morefield's leather flower	*Clematis morefieldii*	E
	Price's potato-bean	*Apios priceana*	T
	Roan Mountain bluet	*Hedyotis purpurea* var. *Montana*	E
	Rock gnome lichen	*Gymnoderma lineare*	E
	Ruth's golden aster	*Pityopsis ruthii*	E
	Small whorled pogonia	*Isotria medeoloides*	T
	Spreading avens	*Geum radiatum*	E
	Spring Creek bladderpod	*Lesquerella perforata*	E
	Tennessee yellow-eyed grass	*Xyris tennesseensis*	E

Virginia spiraea	*Spiraea virginiana*	T

APPENDIX D

STATE LISTED THREATENED AND ENDANGERED SPECIES

TAXA	COMMON NAME	SPECIES	STATUS
Amphibians	Berry cave salamander	*Gyrinophilus gulolineatus*	T
	Tennessee cave salamander	*Gyrinophilus palleucus*	T
Birds	Bachman's sparrow	*Aimophila aestivalis*	E
	Bewick's wren	*Thryomanes bewickii*	E
	Common raven	*Corvus corax*	T
	Golden eagle	*Aquila chrysaetos*	T
	Least tern	*Sterna antillarum athalassos*	E
	Lark sparrow	*Chondestes grammacus*	T
	Northern saw-whet owl	*Aegolius acadius*	T
	Peregrine falcon	*Falco peregrinus*	E
Crustaceans	Big South Fork crayfish	*Cambarus bouchardi*	E
	Brawley's Fork crayfish	*Cambarus williami*	E
	Chickamauga crayfish	*Cambarus extraneus*	T
	Conasauga blue borrower	*Cambarus cymatilis*	E
	Hardin crayfish	*Orconectes wrighti*	E
	Hatchie burrowing crayfish	*Fallicambarus hortoni*	E
	Nashville crayfish	*Orconectes shoupi*	E
	Obey crayfish	*Cambarus obeyensis*	T
	Pristine crayfish	*Cambarus pristinus*	E
	Tennessee cave crayfish	*Orconectes incomptus*	E
	Valley flame crayfish	*Cambarus deweesae*	E
Fishes	Amber darter	*Percina antesella*	E
	Ashy darter	*Etheostoma cinereum*	T
	Barrens darter	*Etheostoma forbesi*	E
	Barrens topminnow	*Fundulus julisia*	E
	Blackside dace	*Phoxinus cumberlandensis*	T
	Blue shiner	*Cyprinella caerulea*	E
	Blue sucker	*Cycleptus elongates*	T
	Bluemask darter	*Etheostoma sp.*	E
	Boulder darter	*Etheostoma wapiti*	E
	Chucky madtom	*Noturus crypticus*	E
	Coldwater darter	*Etheostoma ditrema*	T
	Conasauga logperch	*Percina jenkinsi*	E
	Coppercheek darter	*Etheostoma aquali*	T
	Crown darter	*Etheostoma corona*	E
	Cumberland darter	*Etheostoma susanae*	E
	Duskytail darter	*Etheostoma percnurum*	E
	Egg-mimic darter	*Etheostoma pseudovulatum*	E
	Frecklebelly madtom	*Noturus munitus*	T
	Holiday darter	*Etheostoma brevirostrum*	T
	Lake sturgeon	*Acipenser fulvescens*	E
	Laurel dace	*Phoxinus saylori*	E
	Longhead darter	*Percina macrocephala*	T
	Palezone shiner	*Notropis albizonatus*	E
	Pallid Sturgeon	*Scaphirhynchus albus*	E
	Pygmy madtom	*Noturus stanauli*	E
	Saddled madtom	*Noturus fasciatus*	T

	Sickle darter	*Percina williamsi*	T
	Silverjaw minnow	*Notropis buccatus*	T
Fishes (cont.)	Slackwater darter	*Etheostoma boschungi*	T
	Slender chub	*Erimystax cahni*	T
	Smoky madtom	*Noturus baileyi*	E
	Snail darter	*Percina tanasi*	T
	Spotfin chub	*Erimonax monachus*	T
	Striated darter	*Etheostoma striatulum*	T
	Trispot darter	*Etheostoma trisella*	T
	Tuckasegee darter	*Etheostoma gutselli*	E
	Western sand darter	*Ammocrypta clara*	T
	Yellowfin madtom	*Noturus flavipinnis*	E
Mammals	Carolina northern flying squirrel	*Glaucomys sabrinus coloratus*	E
	Gray bat	*Myotis grisescens*	E
	Indiana bat	*Myotis sodalis*	E
Mussels	Alabama lampmussel	*Lampsilis virescens*	E
	Alabama moccasinshell	*Medionidus acutissimus*	T
	Appalachian elktoe	*Alasmidonta raveneliana*	E
	Appalachian monkeyface	*Quadrula sparsa*	E
	Birdwing pearlymussel	*Conradilla caelata*	E
	Clubshell	*Pleurobema clava*	E
	Coosa moccasinshell	*Medionidus parvulus*	E
	Cracking pearlymussel	*Hemistena lata*	E
	Cumberland bean	*Villosa trabalis*	E
	Cumberland elktoe	*Alasmidonta atropurpurea*	E
	Cumberland monkeyface	*Quadrula intermedia*	E
	Cumberland pigtoe	*Pleurobema gibberum*	E
	Cumberlandian combshell	*Epioblasma brevidens*	E
	Dromedary pearlymussel	*Dromus dromas*	E
	Fanshell	*Cyprogenia stegaria*	E
	Finelined pocketbook	*Lampsilis altilis*	T
	Finerayed pigtoe	*Fusconaia cuneolus*	E
	Littlewing pearlymussel	*Pegias fabula*	E
	Orangefoot pimpleback	*Plethobasus cooperianus*	E
	Ovate clubshell	*Pleurobema perovatum*	E
	Oyster mussel	*Epioblasma capsaeformis*	E
	Pale lilliput	*Toxolasma cylindrellus*	E
	Pink mucket	*Lampsilis abrupta*	E
	Pink ring	*Obovaria retusa*	E
	Purple bean	*Villosa perpurpurea*	E
	Purple cat's paw	*Epioblasma obliquata obliquata*	E
	Rough pigtoe	*Pleurobema plenum*	E
	Rough rabbitsfoot	*Quadrula cylindrical strigillata*	E
	Shiny pigtoe	*Fusconaia cor*	E
	Southern acornshell	*Epioblasma othcaloogensis*	E
	Southern pigtoe	*Pleurobema georgianum*	E
	Tan riffleshell	*Epioblasma florentina walkeri* (= *E. walker*)	E
	Triangular kidneyshell	*Ptychobranchus greenii*	E
	Upland combshell	*Epioblasma metastriata*	E
	White wartyback	*Plethobasus cicatricosus*	E
	Winged mapleleaf	*Quadrula fragosa*	E

Non-Vascular Plants	Alternate leaf archidium moss	*Archidium alternifolium*	T
	American funaria moss	*Funaria americana*	T
	Ammon's tortula	*Ammon's tortula*	E
	Bazzania nudicaulis liverwort	*Bazzania nudicaulis*	T
	Blister ribbon	*Preissia quadrata*	T
	Fragile tortula	*Tortula fragilis*	E
	Funck's rustwort	*Marsupella funckii*	E
	Grandfather Mountain leptodontium	*Leptodontium viticulosoides var. sulphureum*	E
	Gymnomitrion laceratum liverwort	*Gymnomitrion laceratum*	T
	Hot porella	*Porella sp. 1*	E
	Lesser copperwort	*Cephaloziella massalongi*	E
	Lophocolea muricata	*Lophocolea muricata*	T
	Mannia triandra liverwort	*Mannia triandra*	T
	Mount LeConte Moss	*Leptohymenium sharpii*	E
	Ornate cololejeunea	*Cololejeunea ornata*	T
	Palamocladium moss	*Palamocladium leskeoides*	T
	Peak moss	*Brachydontium trichoides*	E
	Pearson's sphenolobopsis	*Sphenolobopsis pearsonii*	E
	Sharp's homaliadelphus	*Homaliadelphus sharpii*	E
	Sharp's Lejeunea	*Lejeunea sharpii*	E
	Sword moss	*Bryoxiphium norvegicum*	T
	Watauga porella	*Porella wataugensis*	T
	Wedge flapwort	*Leptoscyphus cuneifolius*	E
Reptiles	Bog turtle	Glyptemys muhlenbergii	T
	Northern pinesnake	Pituophis melanoleucus melanoleucus	T
	Western pygmy rattlesnake	Sistrurus miliarius streckeri	T
Snails	Anthony's riversnail	*Athearnia anthonyi*	E
	Painted tigersnail	*Anguispira picta*	E
	Royal springsnail	*Pyrgulopsis (=Marstonia) ogmorhaphe*	E
Vascular Plants	Alabama croton	*Croton alabamensis*	E
	Alabama grapefern	*Botrychium jenmanii*	T
	Alabama snow-wreath	*Neviusia alabamensis*	T
	Alderleaf buckthorn	*Rhamnus alnifolia*	E
	American fly-honeysuckle	*Lonicera canadensis*	T
	American water-pennywort	*Hydrocotyle americana*	E
	American wintergreen	*Pyrola americana*	E
	American yew	*Taxus canadensis*	E
	Appalachian bugbane	*Cimicifuga rubifolia*	T
	Appalachian fir clubmoss	*Huperzia appalachiana*	T
	Appalachian waterleaf	*Hydrophyllum virginianum*	T
	Ash-leaved bush-pea	*Thermopsis fraxinifolia*	T
	Barratt's Sedge	*Carex barrattii*	E
	Barrens silky aster	*Symphyotrichium pretense*	E
	Beadle's mountain-mint	*Pycnanthemum beadlei*	E
	Bent avens	*Geum geniculatum*	E
	Blackfoot quillwort	*Isoetes melanopoda*	E
	Blue mud-plantain	*Heteranthera limosa*	T
	Blue Ridge brome sedge	*Carex bromoides ssp. Montana*	T
	Blue Ridge goldenrod	*Solidago spithamaea*	E

	Blue Ridge St. John's-wort	*Hypericum mitchellianum*	T
	Blue-flower coyote-thistle	*Eryngium integrifolium*	T
	Bog fern	*Thelypteris simulate*	E
	Bog-buttons	*Lachnocaulon anceps*	E
	Boykin's milkwort	*Polygala boykinii*	T
	Branched three-awn grass	*Aristida ramosissima*	E
Vascular Plants (cont.)	Branching bur-reed	*Sparganium androcladum*	E
	Braun's rockcress	*Arabis perstellata*	E
	Bristle-fern	*Trichomanes boschianum*	T
	Bristly sedge	*Carex comosa*	T
	Broadleaf bunchflower	*Melanthium latifolium*	E
	Broadleaf goldenrod	*Solidago lancifolia*	E
	Broad-leaved Barbara's-buttons	*Marshallia trinervia*	T
	Broad-leaved tickseed	*Coreopsis latifolia*	E
	Brown bog sedge	*Carex buxbaumii*	T
	Buffalo clover	*Trifolium reflexum*	E
	Buffalo currant	*Ribes odoratum*	T
	Butternut	*Juglans cinerea*	T
	Cain's reedgrass	*Calamagrostis cainii*	E
	Canada anemone	*Anemone canadensis*	E
	Canada burnet	*Sangisorba canadensis*	E
	Canada frostweed	*Helianthum canadense*	E
	Canada lily	*Lilium candense*	T
	Canby's lobelia	*Lobelia canbyi*	T
	Canby's mountain-lover	*Paxistima canbyi*	E
	Capillary hairsedge	*Bulbostylis ciliatifolia* var. *coarctata*	E
	Carolina anemone	*Anemone caroliniana*	E
	Carolina hemlock	*Tsuga Canadensis*	T
	Carolina pink	*Silene caroliniana ssp. Pensylvanica*	T
	Carolina redroot	*Lachnanthes caroliniana*	E
	Carolina saxifrage	*Saxifraga caroliniana*	E
	Chaffseed	*Schwalbea americana*	E
	Chapman's sedge	*Carex chapmanii*	T
	Climbing fumitory	*Adlumia fungosa*	T
	Clingman's hedge-nettle	*Stachys clingmanii*	T
	Coastal false-asphodel	*Triantha racemosa*	E
	Coastal plain yellow-eyed grass	*Xyris ambigua*	E
	Coastal sweet pepper-bush	*Clethra alnifolia*	E
	Compass plant	*Silphium laciniatum*	T
	Copper iris	*Iris fulva*	T
	Cranberry	*Vaccinium macrocarpon*	T
	Creamflower tick-trefoil	*Desmodium ochroleucum*	E
	Creeping St. John's-wort	*Hypericum adpressum*	E
	Crested shield-fern	*Dryopteris cristata*	T
	Cumberland bunchflower	*Minuartia cumberlandensis*	E
	Cumberland featherbells	*Stenanthium diffusum*	E
	Cumberland rose gentian	*Sabatia capitata*	E
	Cumberland rosemary	*Conradina verticillata*	T
	Cumberland rosinweed	*Silphium brachiatum*	E
	Cumberland sandgrass	*Calamovilfa arcuata*	E
	Cutleaf meadow-parsnip	*Thaspium pinnatifidum*	E
	Cutleaf water-milfoil	*Myriophyllum pinnatum*	T
	Death camas	*Zigadenus leimanthoides*	T
	Downy gentian	*Gentiana puberulenta*	E

	Drooping bluegrass	*Poa saltuensis*	T
	Duck River bladderpod	*Lesquerella densipila*	T
	Dwarf filmy-fern	*Trichomanes petersii*	T
	Dwarf huckleberry	*Gaylussacia dumosa*	T
	Dwarf milkwort	*Polygala nana*	E
	Dwarf sundew	*Drosera brevifolia*	T
	Eared goldenrod	*Solidago auriculata*	T
	Earleaved false-foxglove	*Agalinis auriculata*	E
Vascular Plants (cont.)	Eastern turkeybeard	*Xerophyllum asphodeloides*	T
	Eaton's witchgrass	*Dichanthelium acumenatum ssp. spretum*	E
	False dandelion	*Krigia montana*	T
	Fen indian-plantain	*Arnoglossum plantagineum*	T
	Fen orchis	*Liparis loeselii*	T
	Fetter-bush	*Leucothoe racemosa*	T
	Few-flowered beak-rush	*Rhynchospora rariflora*	E
	Fireweed	*Epilobium angustifolium*	T
	Florida hedge-hyssop	*Gratiola floridana*	E
	Fowl bluegrass	*Poa palustris*	E
	Foxtail clubmoss	*Lycopodiella alopecuroides*	T
	Foxtail sedge	*Carex alopecoidea*	E
	Fraser fir	*Abies fraseri*	T
	Fraser's loosestrife	*Lysimachia fraseri*	E
	Fringed black bindweed	*Polygonum cilinode*	T
	Fringed yellow-eyed grass	*Xyris fimbriata*	E
	Gattinger's goldenrod	*Solidago gattingeri*	E
	Giant blue cohosh	*Caulophyllum giganteum*	E
	Giant white-top sedge	*Rhynchospora latifolia*	E
	Glade cleft phlox	*Phlox bifida ssp. Stellaria*	T
	Glade onion	*Allium stellatum*	E
	Globe-fruited false loosestrife	*Ludwigia sphaerocarpa*	T
	Godfrey's stitchwort	*Minuartia godfreyi*	E
	Granite gooseberry	*Ribes curvatum*	T
	Grape honeysuckle	*Lonicera prolifera*	E
	Grassleaf arrowhead	*Sagittaria graminea*	T
	Gray's lily	*Lilium grayi*	E
	Great angelica	*Angelica atropurpurea*	T
	Green pitcher-plant	*Sarracenia oreophila*	E
	Green-and-gold	*Chrysogonum virginianum*	T
	Hairy fimbristylis	*Fimbristylis puberula*	T
	Hairy skullcap	*Scutellaria arguta*	E
	Hairy willow-herb	*Epilobium ciliatum*	T
	Halberd-leaf tearthumb	*Polygonum arifolium*	T
	Harbison's hawthorn	*Crataegus harbisonii*	E
	Harper's fimbristylis	*Fimbristylis perpusilla*	E
	Harper's umbrella-plant	*Eriogonum longifolium var. harperi*	E
	Hart's-tongue fern	*Asplenium scolopenderium var. americanum*	E
	Hay sedge	*Carex argyrantha*	T
	Heartleaf Meehania	*Meehania cordata*	T
	Heart-leaved paper birch	*Betula papyrifera var. cordifolia*	E
	Heart-leaved plantain	*Plantago cordata*	E
	Highland rush	*Juncus trifidus*	E
	Hitchcock's sedge	*Carex hitchcockiana*	T
	Hiwassee quillwort	*Isoetes tennesseensis*	E

	Horned beak-rush	*Rhynchospora capillacea*	E
	Horned bladderwort	*Utricularia cornuta*	E
	Horse-tail spike-rush	*Eleocharis equisetoides*	E
	Kentucky rosinweed	*Silphium wasiotense*	E
	Lake-bank sedge	*Carex lacustris*	T
	Lamance iris	*Iris brevicaulis*	E
	Lambkill	*Kalmia angustifolia var. carolina*	E
	Large purple fringed orchid	*Platanthera grandiflora*	E
	Large round-leaved orchid	*Platanthera orbiculata*	T
	Large-flowered skullcap	*Scutellaria montana*	T
Vascular Plants (cont.)	Large-flowering Barbara's-buttons	*Marshallia grandiflora*	E
	Large-leaf pondweed	*Potamogeton amplifolius*	T
	Larkspur-leaved coreopsis	*Coreopsis delphiniifolia*	E
	Leafy prairie-clover	*Dalea foliosa*	E
	Least trillium	*Trillium pusillum*	E
	Leatherleaf meadowrue	*Thalictrum coriaceum*	T
	Leggett's pinweed	*Lechea pulchella*	E
	Limerock arrow-wood	*Viburnum bracteatum*	E
	Linear-leaved willow-herb	*Epilobium leptophyllum*	T
	Long-bracted green orchis	*Coeloglossum viride var. virescens*	E
	Longleaf stitchwort	*Stellaria longifolia*	E
	Loose-headed beak-rush	*Rhynchospora chalarocephala*	T
	Low frostweed	*Helianthum propinquum*	E
	Lucy Braun's white snakeroot	*Ageratina luciae-brauniae*	T
	Manhart's sedge	*Carex manhartii*	E
	Marsh marigold	*Caltha palustris*	E
	Marsh speedwell	*Veronica scutellata*	E
	Matted spike-rush	*Eleocharis intermedia*	E
	Mayberry	*Vaccinium elliottit*	E
	Menge's fame-flower	*Talinum mengesii*	T
	Michigan lily	*Lilium michiganense*	T
	Missouri primrose	*Oenothera macrocarpa*	T
	Morefield's leather-flower	*Clematis morefieldii*	E
	Moss phlox	*Phlox subulata*	T
	Mountain bittercress	*Cardamine clematitis*	T
	Mountain bush-honeysuckle	*Diervilla sessifolia var. rivularis*	T
	Mountain fetter-bush	*Pieris floribunda*	T
	Mountain sandwort	*Minuartia groenlandica*	E
	Mountain St. John's-wort	*Hypericum graveolens*	E
	Mountain witch-alder	*Fothergilla major*	T
	Muskingum sedge	*Carex muskingumensis*	E
	Narrowleaf bushclover	*Lespedeza angustifolia*	T
	Narrow-leaf ramps	*Allium burdickii*	T
	Narrow-leaved gentian	*Gentiana linearis*	T
	Narrow-leaved meadow-sweet	*Spiraea alba*	E
	Narrow-leaved trillium	*Trillium lancifolium*	E
	Needleleaf witchgrass	*Dichanthelium aciculare*	E
	Nestronia	*Nestronia umbellula*	E
	Nevius's Stonecrop	*Sedum nevii*	E
	Nodding rattlesnake root	*Prenanthes crepidinea*	E
	Northern bush-honeysuckle	*Diervilla lonicera*	T
	Northern starflower	*Trientalis borealis*	T
	Northwest Territory sedge	*Carex utriculata*	E
	Nuttall's milkwort	*Polygala nuttallii*	E
	Obscure beak-rush	*Rhynchospora perplexa*	T
	Oklahoma grass-pink	*Calapogon oklahomensis*	E

	Ovate catchfly	*Silene ovata*	E
	Ozark bunchflower	*Melanthium woodii*	E
	Pale corydalis	*Corydalis sempervirens*	E
	Pale St. John's-wort	*Hypericum ellipticum*	E
	Pale umbrella-wort	*Mirabilis albida*	T
	Pale-purple coneflower	*Echinacea pallida*	T
	Pasture glade-cress	*Leavenworthia exigua var. lutea*	E
	Piedmont Barbara's-buttons	*Marshallia obovata*	T
	Pineland squarehead	*Tetragonotheca helianthoides*	E
	Pink sundew	*Drosera capillaris*	T
	Pinnate-lobed black-eyed susan	*Rudbeckia triloba var. pinnatiloba*	E
	Piratebush	*Buckleya distichophylla*	T
Vascular Plants (cont.)	Pitcher leather-flower	*Clematis pitcheri*	T
	Plains frostweed	*Helianthum bicknellii*	E
	Plains muhly	*Muhlenbergia cuspidata*	E
	Pope's sand-parsley	*Ammoselinum popei*	T
	Porter's reedgrass	*Calamagrostis porteri*	E
	Prairie false-foxglove	*Agalinis heterophylla*	T
	Prairie goldenrod	*Solidago ptarmicoides*	E
	Prairie parsley	*Polytaenia nuttallii*	T
	Prairie rhynchosia	*Rhynchosia latifolia*	E
	Price's potato-bean	*Apios priceana*	E
	Purple gerardia	*Agalinis plukenetii*	E
	Purple giant hyssop	*Agastache scrophulariifolia*	T
	Purple prairie-clover	*Dalea purpurea*	E
	Pursh's rattlebox	*Crotalaria purshii*	E
	Pyne's ground-plum	*Astragalus bibullatus*	E
	Red starvine	*Schisandra glabra*	T
	Ridge-stem false-foxglove	*Agalinis oligophylla*	E
	Roan false goat's-beard	*Astilbe crenatiloba*	E
	Roan Mountain bluet	*Hedyotis purpurea var. montana*	E
	Roan Mountain sedge	*Carex roanensis*	E
	Rock goldenrod	*Solidago rupestris*	E
	Rock skullcap	*Scutellaria saxatilis*	T
	Rockcastle aster	*Eurybia saxicastellii*	E
	Rose pogonia	*Pogonia ophioglossoides*	E
	Rough hawkweed	*Hieracium scabrum*	T
	Rough rattlesnake-root	*Prenanthes aspera*	E
	Roundleaf fame-flower	*Talinum teretifolium*	T
	Roundleaf shadbush	*Amelanchier sanguinea*	T
	Roundleaf sundew	*Drosera rotundifolia*	T
	Royal catchfly	*Silene regia*	E
	Rugel's ragwort	*Rugelia nudicaulis*	E
	Running bittercress	*Cardamine flagellifera*	T
	Running glade clover	*Trifolium calcaricum*	E
	Ruth's golden-aster	*Pityopsis ruthii*	E
	Ruth's sedge	*Carex ruthii*	T
	Sand cherry	*Prunus pumila*	E
	Sand grape	*Vitis rupestris*	E
	Schweinitz's ragwort	*Packera schweinitziana*	T
	Senna seymeria	*Seymeria cassioides*	E
	Sessile water speedwell	*Veronica catenata*	E
	Shadow-witch	*Ponthieva racemosa*	E
	Shaggy false gromwell	*Onosmodium hispidissimum*	E
	Shining ladies-tresses	*Spiranthes lucida*	T
	Short-beaked arrowhead	*Sagittaria brevirostra*	T

	Shortleaf sneezeweed	*Helenium brevifolium*	E
	Short-leaved panicgrass	*Dichanthelium ensifolium ssp. curtifolium*	E
	Short's bladderpod	*Lesquerella globosa*	E
	Showy lady's-slipper	*Cypripedium reginae*	E
	Silverling	*Paronychia argyrocoma*	T
	Skunk-cabbage	*Symplocarpus foetidus*	E
	Slender blazing-star	*Liatris cylindracea*	T
	Slender blue flag	*Iris prismatica*	T
	Small whorled pogonia	*Isotria medeoloides*	E
	Small's stonecrop	*Diamorpha smallii*	E
	Smoky Mountain's mannagrass	*Glyceria nubigena*	T
	Smooth false gromwell	*Onosmodium molle ssp. subsetum*	E
	Snowy orchid	*Platanthera nivea*	E
Vascular Plants (cont.)	Softleaf arrow-wood	*Viburnum molle*	E
	Southern jointweed	*Polygonella americana*	E
	Southern lady's-slipper	*Cypripedium kentuckiense*	E
	Southern lobelia	*Lobelia amoena*	T
	Southern morning-glory	*Stylisma humistrata*	T
	Southern nodding trillium	*Trillium rugelii*	E
	Southern prairie-dock	*Silphium pinnatifidum*	T
	Southern racemose goldenrod	*Solidago arenicola*	T
	Southern twayblade	*Listera australis*	E
	Spinulose shield-fern	*Dryopteris carthusiana*	T
	Spotted coralroot	*Corallorhiza maculata*	T
	Spreading avens	*Geum radiatum*	E
	Spreading rockcress	*Arabis patens*	E
	Spring blue-eyed mary	*Collinsia verna*	E
	Spring Creek bladderpod	*Lesquerella perforata*	E
	Starflower false Solomon's seal	*Maianthemum stellatum*	E
	Sterile sedge	*Carex sterilis*	E
	Stiff clubmoss	*Lycopodium annotinum*	E
	Stones River bladderpod	*Lesquerella stonensis*	E
	Sullivantia	*Sullivantia sullivantii*	E
	Svenson's wild-rye	*Elymus svensonii*	E
	Swamp loosestrife	*Lysimachia terrestris*	E
	Swamp saxifrage	*Saxifraga pensylvanica*	E
	Sweet coneflower	*Rudbeckia subtomentosa*	T
	Sweet pinesap	*Monotropsis odorata*	T
	Sweetbay magnolia	*Magnolia virginiana*	T
	Sweet-fern	*Comptonia peregrina*	E
	Sweetscent ladies-tresses	*Spiranthes odorata*	E
	Sweet-scented indian-plantain	*Hasteola suavolens*	T
	Tall larkspur	*Delphinium exaltatum*	E
	Tawny cotton-grass	*Eriophorum virginicum*	E
	Ten-angle pipewort	*Eriocaulon decangulare*	E
	Tennessee coneflower	*Echinacea tennesseensis*	E
	Tennessee pondweed	*Potamogeton tennesseensis*	T
	Tennessee yellow-eyed grass	*Xyris tennesseensis*	E
	Thicket parsley	*Perideridia americana*	E
	Torrey's dropseed	*Muhlenbergia torreyana*	E
	Trailing stitchwort	*Stellaria alsine*	E
	Trailing trillium	*Trillium decumbens*	E
	Trailing wolfsbane	*Aconitum reclinatum*	E
	Tubercled rein-orchid	*Platanthera flava var. herbiola*	T
	Tufted club-rush	*Trichophorum cespitosum*	E

	Twinflower	*Linnaea borealis*	E
	Velvety sedge	*Carex vestita*	E
	Virginia bunchflower	*Melanthium virginicum*	E
	Virginia mallow	*Sida hermaphrodita*	E
	Virginia spiraea	*Spiraea virginiana*	E
	Water stitchwort	*Stellaria fontinalis*	T
	Water-purslane	*Didiplis diandra*	T
	Wavy-leaf purple coneflower	*Echinacea simulata*	T
	Western false gromwell	*Onosmodium molle ssp. occidentale*	T
	Western hairy rockcress	*Arabis hirsuta*	T
	Western wallflower	*Erysimum capitatum*	E
	White beak-rush	*Rhynchospora alba*	E
	White camas	*Zigadenus glaucus*	E
	White fringeless orchid	*Platanthera integrilabia*	E
Vascular Plants (cont.)	White heather aster	*Symphyotrichium ericoides var. ericoides*	T
	White mandarin	*Steptopus amplexifolius*	T
	White water-buttercup	*Ranunculus aquatilis var. diffusus*	E
	White-bracted thoroughwort	*Eupatorium leucolepis*	E
	White-leaved leather-flower	*Clematis glaucophylla*	E
	White-leaved sunflower	*Helianthus glaucophyllus*	T
	Whorled mountain-mint	*Pycnanthemum verticillatum*	E
	Whorled sunflower	*Helianthus verticillatus*	E
	Wide-leaved yellow-eyed grass	*Xyris laxifolia var. iridifolia*	T
	Willow aster	*Symphyotrichium praealtum*	E
	Wolf spike-rush	*Eleocharis wolfii*	E
	Wood lily	*Lilium philadelphicum*	E
	Woolly sedge	*Carex pellita*	E
	Wooly sandwort	*Arenaria lanuginosa*	E
	Wretched sedge	*Carex misera*	T
	Wright's beak-rush	*Rhynchospora wrightiana*	E
	Yellow fringeless orchid	*Platanthera integra*	E
	Yellow honeysuckle	*Lonicera flava*	T
	Yellow nodding ladies-tresses	*Spiranthes ochroleuca*	E
	Yellow sunnybell	*Schoenolirion croceum*	T
	Yellow water-crowfoot	*Ranunculus flabellaris*	T
	Zigzag bladderwort	*Utricularia subulata*	T

APPENDIX E

CRITERIA FOR BEAVER DAM BREACHING/REMOVAL

Wetlands are recognized by three characteristics: hydric soils, hydrophytic vegetation, and general hydrology. Hydric soils are either entirely composed of, or have a thick surface layer of decomposed plant materials; sandy soils have dark stains or streaks from organic material in the upper layer where plant material has attached to soil particles. In addition, hydric soils may be bluish gray or gray below the surface or brownish black to black and have the smell of rotten eggs. Wetlands also have hydrophytic vegetation such as cattails, bulrushes, willows, sedges, and water plantains. The final indicator is general hydrology which includes standing and flowing water or waterlogged soils during the growing season; high water marks are present on trees and drift lines of small piles of debris are usually present. Beaver dams usually will develop a layer of organic material at the surface because siltation can occur rapidly, but aquatic vegetation and high water marks (a new high water mark is created by the beaver dam) are usually not present. However, cattails and willows can show up rapidly if they are in the vicinity, but most hydrophytic vegetation takes time to establish.

When a dam is removed, debris is discharged into the water. The debris that ends up in the water is considered "*incidental fallback*" or discharge fill. However, in most beaver dam removal operations, the material that is displaced, if considered to be discharge, is exempt from permit requirements under 33 CFR 323 or 330. A permit would be required if the impoundment caused by a beaver dam was considered a true wetland. WS personnel survey the beaver dam site and impoundment and determine whether conditions exist suggesting that the area may be a wetland as defined above. If such conditions exist, the landowner is asked the age of the dam or how long he/she has known of its presence to determine whether Swampbuster, Section 404 permit exemptions or NWPs allow removal of the dam. If not, the landowner is required to obtain a Section 404 permit before the dam will be removed by WS personnel.

The following information explains Section 404 exemptions and conditions that pertain to the removal of beaver dams.

33 CFR 323 – Permits For Discharges of Dredged or Fill Material into Waters of the United States. This regulation provides guidance to determine whether certain activities require permits under Section 404.

> **Part 323.4 Discharges not requiring permits.** This section establishes exemptions for discharging certain types of fill into waters of the United States without a permit. Certain minor drainage activities connected with normal farming, ranching, and silviculture activities where they have been established do not require a permit as long as these drainages do not include the immediate or gradual conversion of a wetland (*i.e.*, beaver ponds greater than 5 years old) to a non-wetland. Specifically part (a)(1)(iii)(C)(i) states, "*…fill material incidental to connecting upland drainage facilities* [*e.g.*, drainage ditches] *to waters of the United States, adequate to effect the removal of excess soil moisture from upland croplands…*". This indicates that beaver dams that block ditches, canals, or other structures designed to drain water from upland crop fields can be removed without a permit.
>
> Moreover, part (a)(1)(iii)(C)(iv) states the following types of activities do not require a permit "*The discharges of dredged or fill materials incidental to the emergency removal of sandbars, gravel bars, or other similar blockages which are formed during flood flow or other events,*

where such blockages close or constrict previously existing drainageways and, if not promptly removed, would result in damage to or loss of existing crops or would impair or prevent the plowing, seeding, harvesting or cultivating of crops in land in established use for crop production. Such removal does not include enlarging or extending the dimensions of, or changing the bottom elevations of, the affected drainageway as it existed prior to the formation of the blockage. Removal must be accomplished within one year of discovery of such blockages in order to be eligible for exemption." This allows the removal of beaver dams in natural streams to restore drainage of agricultural lands within one year of discovery.

Part 323.4(a)(2) allows "*Maintenance, including emergency reconstruction of recently damaged parts, of currently serviceable structures such as dikes, dams, levees, groins, riprap, breakwaters, causeways, bridge abutments or approaches, and transportation structures. Maintenance does not include any modification that changes the character, scope, or size of the original fill design. Emergency reconstruction must occur within a reasonable period of time after damage occurs in order to qualify for this exemption.*" This allows beaver dams to be removed without a permit where they have resulted in damage to roads, culverts, bridges, or levees if it is done in a reasonable amount of time.

Tennessee regulates discharges into the waters of the state through the TDEC and grants exemptions from permitting for discharges based on guidelines and exemptions provided by the Army Corps of Engineers. TDEC does not consider removal of beaver dams by WS in the state to require permits in those situations exempted by the Corps (A. Fritz, TDEC, pers. comm. 2010).

33 CFR 330 – Nationwide Permit Program: The Corps Chief of Engineers is authorized to grant certain dredge and fill activities on a nationwide basis if they have minimal impact on the environment. The NWPs are listed in Appendix A of 33 CFR 330 and permittees must satisfy all terms and conditions established in order to qualify for their use. Individual beaver dam removal activities by WS may be covered by any of the following NWPs if not already exempted from permit requirements by the regulations discussed above. WS complies with all conditions and restrictions placed on NWPs for any instance of beaver dam removal done under a specific NWP.

NWPs can be used **except** in any component of the National Wild and Scenic River System (16 U.S.C. §§ 1271-1287 as amended) such as the designated reaches of the Obed River in Tennessee and any other rivers or reaches and their corridors in Tennessee which have been designated as part of the Wild Rivers system authorized by The Tennessee Scenic Rivers Act. Any beaver dam removal in these designated areas which might be contemplated by WS may require consultation with the Corps and TDEC to obtain permits for any such activities.

NWP 3 authorizes the rehabilitation of those structures, such as culverts, homes, and bridges, destroyed by floods and "*discrete events*" such as beaver dams provided that the activity is commenced within 2 years of the date when the beaver dam was established.

NWP 18 allows minor discharges of dredged and fill material, including the removal of beaver dams, into all waters of the United States provided that the quantity of discharge and the volume of excavated area does not exceed 25 cubic yards below the plane of the ordinary high water mark (this is normally well below the level of the beaver dam) and will not cause the loss of more than $1/10^{th}$ acre of special aquatic site including wetlands. The District Engineer must be "*notified*" (general conditions for notification apply), if the discharge is between 10 and 25 cubic yards for a single project if the project is in a special aquatic site, including wetlands. Beaver dams rarely would exceed 2 or 3 cubic yards of backfill into the waters and probably no more than 5 cubic yards would ever be exceeded. Therefore, this stipulation is not restrictive. Beaver

dams periodically may be removed in a special aquatic area, but in most instances, the aquatic site will be returned to normal. However, if a true wetland exists, and beaver dam removal is not allowed under another permit, then a permit may be obtained from the District Engineer.

NWP 27 provides for the discharge of dredge and fill for activities associated with the restoration of wetland and riparian areas with certain restrictions. On non-federal public and private lands, the owner must have: a binding agreement with USFWS or NRCS to conduct restoration; a voluntary wetland restoration project documented by NRCS; or notified the District Engineer according to "notification' procedures. On Federal lands, including Corps and USFWS, wetland restoration can take place without any contract or notification. This NWP *"…applies to restoration projects that serve the purpose of restoring "natural" wetland hydrology, vegetation, and function to altered and degraded non-tidal wetlands and "natural" functions of riparian areas. This NWP does not authorize the conversion of natural wetlands to another aquatic use…*" If operating under this permit, the removal of a beaver dam would be allowed as long as it was not a true wetland (*i.e*., 5 or more years old), and for non-federal public and private lands the appropriate agreement, project documentation, or notification is in place.

A quick response immediately resulting from permitting requirements can be critical to the success of minimizing or preventing damage. Exemptions contained in the above regulations or NWPs provide for the removal of the majority of beaver dams that WS in Tennessee encounters. The primary determination that must be made by WS personnel is whether a beaver impounded area has become a true wetland or is just a flooded area. The flexibility allowed by these exemptions and NWPs is important for the efficient and effective resolution of many beaver damage problems because damage escalates rapidly in many cases the longer an area remains flooded.

APPENDIX F

CRITERIA FOR BEAVER DAM BREACHING/REMOVAL WHERE THREATENED OR ENDANGERED SPECIES MAY BE PRESENT

Critical habitat is defined in section 3(5)(A) of the ESA as i) the specific areas within the geographic area occupied by a species, at the time it is listed in accordance with the ESA, on which are found those physical or biological features (I) that are essential to the conservation of the species and (II) which may require special management considerations or protection, and ii) specific areas outside the geographic area occupied by a species at the time it is listed, upon a determination that such areas are essential for the conservation of the species. "*Conservation*" means the use of all methods and procedures that are necessary to bring an endangered or a threatened species to the point at which listing under the ESA is no longer necessary. Regulation 50 CFR 424.02(j) defines special management considerations or protection to mean any method or procedure useful in protecting physical and biological features of the environment for the conservation of the listed species.

Critical habitat receives protection under section 7 of the ESA through the prohibition against destruction or adverse modification of critical habitat with regard to actions carried out, funded, or authorized by a Federal agency. Section 7 also requires conferences on federal actions that are likely to result in adverse modification or destruction of proposed critical habitat. Aside from the protection that may be provided under section 7, the ESA does not provide other forms of protection to lands designated as critical habitat.

WS and the USFWS recognize that certain T&E species habitat requirements may not be completely delineated within designated critical habitats. As a result, WS' personnel would evaluate sites proposed for beaver damage management activities based upon the relation of the damage site to currently designated, proposed, and potential critical habitats. Within the State, the Tennessee River (see Figure 1), serves as a dividing line with respect to T&E fish and mussels presence or absence. Currently, no listed T&E fish or mussels are known to exist west of this dividing line. As a result, no additional consultation would be conducted with USFWS on beaver damage management activities in this area. However, at beaver damage sites east of this line, WS' personnel would consult with the USFWS and the TWRA prior to taking action and make determinations about the presence of such fish or mussel populations and the appropriateness of beaver dam removal activities as it relates to the protection of these species. WS would act in accordance with any recommendations provided by USFWS and TWRA.



Figure 1: The Tennessee River divides the State into two areas. No site-specific consultation with USFWS would occur in the shaded area (west of the Tennessee River), while site-specific consultation would occur in the rest of the State prior to beaver dam breaching or removal by WS.

The specific areas that have been deemed critical habitat and the associated federally T&E species found in Tennessee are:

Coosa River Watershed

Conasauga River in Bradley And Polk Counties	Alabama moccasinshell	*Medionidus acutissimus*
	Coosa moccasinshell	*Medionidus parvulus*
	Fine-lined pocketbook	*Lampsilis altilis*
	Georgia pigtoe	*Pleuroblema hanleyianum*
	Ovate clubshell	*Pleuroblema perovatum*
	Southern acornshell	*Epioblasma othcaloogensis*
	Southern clubshell	*Pleuroblema decisum*
	Southern pigtoe	*Pleuroblema georgianum*
	Triangular kidneyshell	*Ptychobranchus greeni*
	Upland combshell	*Epioblasma metastriata*
	Amber darter	*Percina antesella*
	Conasauga logperch	*Percina jenkinsi*

Tennessee River Watershed

Beech Creek, Clinch River, Duck River, Nolichucky River Obed River, and Powell River	Cumberland elktoe	*Alasmidonta atropurpurea*
	Cumberlandian combshell	*Epioblasma brevidens*
	Oyster mussel	*Epioblasma capsaeformis*
	Purple bean	*Villosa perpurpurea*
	Rough rabbitsfoot	*Quadrula cylindrical stigillata*
Buffalo River & tributaries In Lawrence County, and Cypress Creek and Middle Cypress Creek and all Permanent and intermittent Tributaries in Wayne County	Slackwater darter	*Etheostoma boschungi*
Citico Creek in Monroe County	Smoky madtom	*Noturus baileyi*
Clinch River and Powell River in Claiborne and Hancock Counties	Slender chub	*Erimystax (=Hybopsis) cahni*
	Yellowfin madtom	*Noturus flavipinnis*
Clear Creek, Daddy's Creek Emory River and Obed River In Cumberland, Fentress, and Morgan Counties	Spotfin chub	*Cyprinella (=Hybopsis) monacha*

Cumberland River Watershed

Big South Fork & tributaries (Bone Camp Creek, White Oak Creek, North White Oak Creek, New River, Clear Creek, Crooked Creek, Clear Fork, North Prong Clear Fork	Cumberland elktoe	*Alasmidonta atropurpurea*
	Cumberlandian combshell	*Epioblasma brevidens*
	Oyster mussel	*Epioblasma capsaeformis*
	Purple bean	*Villosa perpurpurea*
	Rough rabbitsfoot	*Quadrula cylindrical strigillata*