

BASF Petition (19-317-01p) for Determination of Nonregulated Status of Transgenic GMB151 Soybean Resistant to the Soybean Cyst Nematode and HPPD-inhibiting Herbicides

**OECD Unique Identifier:
BCS-GM151-6**

Draft Environmental Assessment

June 2021

**Agency Contact
Cindy Eck
Biotechnology Regulatory Services
4700 River Road
USDA, APHIS
Riverdale Park, MD 20737**

In accordance with federal civil rights law and U.S. Department of Agriculture (USDA) civil rights regulations and policies, the USDA, its Agencies, offices, and employees, and institutions participating in or administering USDA programs are prohibited from discriminating based on race, color, national origin, religion, sex, gender identity (including gender expression), sexual orientation, disability, age, marital status, family/parental status, income derived from a public assistance program, political beliefs, or reprisal or retaliation for prior civil rights activity, in any program or activity conducted or funded by USDA (not all bases apply to all programs). Remedies and complaint filing deadlines vary by program or incident.

Persons with disabilities who require alternative means of communication for program information (e.g., Braille, large print, audiotope, American Sign Language, etc.) should contact the responsible Agency or USDA's TARGET Center at (202) 720-2600 (voice and TTY) or contact USDA through the Federal Relay Service at (800) 877-8339. Additionally, program information may be made available in languages other than English.

To file a program discrimination complaint, complete the USDA Program Discrimination Complaint Form, AD-3027, found online at [How to File a Program Discrimination Complaint](#) and at any USDA office or write a letter addressed to USDA and provide in the letter all of the information requested in the form. To request a copy of the complaint form, call (866) 632-9992. Submit your completed form or letter to USDA by: (1) mail: U.S. Department of Agriculture, Office of the Assistant Secretary for Civil Rights, 1400 Independence Avenue, SW, Washington, D.C. 20250-9410; (2) fax: (202) 690-7442; or (3) email: program.intake@usda.gov.

Mention of companies or commercial products in this report does not imply recommendation or endorsement by the U.S. Department of Agriculture (USDA) over others not mentioned. USDA neither guarantees nor warrants the standard of any product mentioned. Product names are mentioned solely to report factually on available data and to provide specific information.

TABLE OF CONTENTS

	<u>Page</u>
LIST OF FIGURES.....	- V -
LIST OF TABLES	- VI -
LIST OF ACRONYMS.....	- VII -
1 PURPOSE AND NEED.....	- 1 -
1.1 PURPOSE OF GMB151 SOYBEAN.....	- 1 -
1.2 COORDINATED FRAMEWORK FOR REGULATING BIOTECHNOLOGY..	- 2 -
1.3 PURPOSE AND NEED TO ISSUE A REGULATORY STATUS DETERMINATION.....	- 3 -
2 SCOPING AND PUBLIC INVOLVEMENT	- 5 -
2.1 PUBLIC INVOLVEMENT FOR PETITION 19-317-01p	- 5 -
2.2 ISSUES CONSIDERED	- 6 -
3 ALTERNATIVES	- 7 -
3.1 NO ACTION ALTERNATIVE: CONTINUE REGULATING GMB151 - 7 -- 7 -	
3.3 ALTERNATIVES CONSIDERED BUT EXCLUDED FROM FURTHER DETAILED ANALYSIS IN THIS EA	- 7 -
3.4 COMPARISON OF ALTERNATIVES	- 8 -
4 AFFECTED ENVIRONMENT	- 23 -
4.1 AGRICULTURAL PRODUCTION OF SOYBEANS.....	- 23 -
4.1.1 Acreage and Regional Distribution of Soybean Production.....	- 26 -
4.1.2 Agronomic Practices for Conventional Soybean Production	- 26 -
4.1.3 Agronomic Practices for Organic Soybean Production.....	- 37 -
4.2 PHYSICAL ENVIRONMENT.....	- 38 -
4.2.1 Soil Quality	- 38 -
4.2.2 Water Resources	- 40 -
4.2.3 Air Quality	- 43 -
4.3 BIOLOGICAL RESOURCES	- 44 -
4.3.1 Animal Communities.....	- 44 -
4.3.2 Plant Communities	- 46 -
4.3.3 Gene Flow and Weediness	- 50 -
4.3.4 Microorganisms	- 51 -

4.3.5	Biodiversity	- 53 -
4.4	ANIMAL FEED.....	- 53 -
4.5	HUMAN HEALTH	- 54 -
4.5.1	Public Health	- 54 -
4.5.2	Occupational Health and Worker Safety	- 55 -
4.6	SOCIOECONOMIC ISSUES.....	- 56 -
4.6.1	Domestic Economic Environment.....	- 56 -
4.6.2	International Trade Environment.....	- 57 -
5	ENVIRONMENTAL CONSEQUENCES	- 60 -
5.1	ENVIRONMENTAL ANALYSIS.....	- 60 -
5.2	SCOPE OF THE ENVIRONMENTAL ANALYSIS	- 60 -
5.3	AGRICULTURAL PRODUCTION OF SOYBEAN	- 61 -
5.3.1	Locations and Acreage of Soybean Production: No Action Alternative.....	- 61 -
5.3.2	Locations and Acreage of Soybean Production: Preferred Alternative.....	- 61 -
5.3.3	Agronomic Practices: No Action Alternative.....	- 62 -
5.3.4	Agronomic Practices: Preferred Alternative.....	- 62 -
5.3.5	Soybean Seed Production: No Action Alternative	- 63 -
5.3.6	Soybean Seed Production: Preferred Alternative	- 63 -
5.3.7	Organic and Conventional Soybean Production: No Action Alternative.....	- 64 -
5.3.8	Organic and Conventional Soybean Production: Preferred Alternative.....	- 64 -
5.4	PHYSICAL ENVIRONMENT.....	- 64 -
5.4.1	Soil Quality: No Action Alternative	- 64 -
5.4.2	Soil Quality: Preferred Alternative.....	- 65 -
5.4.3	Water Resources: No Action Alternative	- 65 -
5.4.4	Water Resources: Preferred Alternative	- 66 -
5.4.5	Air Quality: No Action Alternative	- 66 -
5.4.6	Air Quality: Preferred Alternative.....	- 67 -
5.5	BIOLOGICAL RESOURCES	- 67 -
5.5.1	Animal Communities: No Action Alternative.....	- 67 -
5.5.2	Animal Communities: Preferred Alternative.....	- 67 -
5.5.3	Plant Communities: No Action Alternative.....	- 68 -
5.5.4	Plant Communities: Preferred Action Alternative.....	- 68 -

5.5.5	Gene Flow and Weediness: No Action Alternative.....	- 69 -
5.5.6	Gene Flow and Weediness: Preferred Alternative	- 69 -
5.5.7	Microorganisms: No Action Alternative	- 70 -
5.5.8	Microorganisms: Preferred Alternative	- 70 -
5.5.9	Biodiversity: No Action Alternative.....	- 71 -
5.5.10	Biodiversity: Preferred Alternative	- 71 -
5.6	ANIMAL HEALTH.....	- 72 -
5.6.1	Animal Feed—No Action Alternative.....	- 72 -
5.6.2	Animal Feed—Preferred Alternative.....	- 73 -
5.7	HUMAN HEALTH	- 73 -
5.7.1	Public Health: No Action Alternative	- 73 -
5.7.2	Public Health: Preferred Alternative	- 74 -
5.7.3	Worker Safety: No Action Alternative	- 74 -
5.7.4	Worker Safety: Preferred Alternative.....	- 75 -
5.8	SOCIOECONOMIC IMPACTS.....	- 76 -
5.8.1	Domestic Economic Environment: No Action Alternative	- 76 -
5.8.2	Domestic Economic Environment: Preferred Alternative.....	- 76 -
5.8.3	International Trade Economic Environment: No Action Alternative	- 76 -
5.8.4	International Trade Economic Environment: Preferred Alternative	- 77 -
6	CUMULATIVE IMPACTS	- 78 -
6.1	METHODOLOGY AND ASSUMPTIONS.....	- 78 -
6.2	CUMULATIVE IMPACTS: SOYBEAN AGRICULTURAL PRODUCTION ..	- 79 -
6.3	CUMULATIVE IMPACTS: PHYSICAL ENVIRONMENT	- 80 -
6.4	CUMULATIVE IMPACTS: BIOLOGICAL RESOURCES	- 80 -
6.5	CUMULATIVE IMPACTS: ANIMAL FEED AND HUMAN HEALTH	- 81 -
6.6	CUMULATIVE IMPACTS: SOCIOECONOMICS.....	- 82 -
7	THREATENED AND ENDANGERED SPECIES.....	- 85 -
7.1	ESA REQUIREMENTS FOR FEDERAL AGENCIES.....	- 85 -
7.2	POTENTIAL EFFECTS OF GMB151 SOYBEAN ON T&E SPECIES AND CRITICAL HABITAT	- 87 -
7.2.1	Threatened and Endangered Plant Species and Critical Habitat	- 88 -
7.2.2	Threatened and Endangered Animal Species and Critical Habitat.....	- 89 -
7.3	SUMMARY.....	- 93 -

8	CONSIDERATION OF EXECUTIVE ORDERS, STANDARDS, AND TREATIES RELATING TO ENVIRONMENTAL IMPACTS	- 94 -
8.1	FEDERAL LAWS AND REGULATIONS	- 94 -
8.1.1	National Environmental Policy Act (NEPA).....	- 94 -
8.1.2	Clean Water Act, Safe Drinking Water Act, and Clean Air Act	- 94 -
8.1.3	National Historic Preservation Act (NHPA) of 1966 as Amended	- 94 -
8.2	EXECUTIVE ORDERS WITH DOMESTIC IMPLICATIONS	- 95 -
8.3	INTERNATIONAL IMPLICATIONS	- 97 -
8.4	IMPACTS ON UNIQUE CHARACTERISTICS OF GEOGRAPHIC AREAS	- 99 -
9	LIST OF PREPARERS	- 100 -
10	REFERENCES.....	- 101 -

LIST OF FIGURES

	<u>Page</u>
Figure 1. Soybean Planted Acres by County for Selected States.....	- 24 -
Figure 2. U. S. Soybean Acreage: 2011-2020.....	- 25 -
Figure 3. U.S. Soybean Yield: 1991-2019.....	- 26 -
Figure 4. Schematic Diagram of the Development of Herbicide Resistance.	- 48 -
Figure 5. U.S. Soybean Export Volume and Percent Exported.....	- 58 -
Figure 6. U.S. Soybean Export Volume and Percent of Global Market.	- 58 -

LIST OF TABLES

	<u>Page</u>
Table 1. Summary of Potential Impacts and Consequences of Alternatives.....	- 9 -
Table 2. Biotech Soybeans as a Percentage of the Total U.S. Soybean Crop.	- 27 -
Table 3. Nutrient Removal Rates for Commonly Grown U.S. Grain Crops.....	- 30 -
Table 4. Insecticides Most Commonly Applied to Soybean Acreage in 2017.....	- 32 -
Table 5. Five Most Used Herbicides to Treat U.S. Soybean Acreage in 2017.	- 34 -
Table 6. Examples of Estimated Annual Fuel Used for Different Tillage Methods.	- 43 -
Table 7. Summary of World-wide HR Weeds by Herbicide Group	- 49 -
Table 8. World Soybean Production.....	- 59 -
Table 9. Proteins Produced by GMB151 Soybean that are Novel in Soybean.....	- 88 -

ACRONYMS AND ABBREVIATIONS

AOSCA	Association of Official Seed Certifying Agencies
APHIS	Animal and Plant Health Inspection Service
C	carbon
CAA	Clean Air Act
CFR	Code of Federal Regulations
CH ₄	methane
CO	carbon monoxide
CO ₂	carbon dioxide
CWA	Clean Water Act
DNA	deoxyribonucleic acid
EA	environmental assessment
EO	Executive Order
EPA	Environmental Protection Agency
ESA	Endangered Species Act
FDA	Food and Drug Administration
FFDCA	Federal Food, Drug, and Cosmetic Act
FIFRA	Federal Insecticide, Fungicide, and Rodenticide Act
FQPA	Food Quality Protection Act
FR	Federal Register
GR	glyphosate-resistant
HPPD-4	4-hydroxyphenylpyruvate dioxygenase
HR	herbicide-resistant
HRAC	Herbicide Resistance Action Committee
IP	identity preservation
IPPC	International Plant Protection Convention
lb/A	pounds per acre
lbs	pounds
LMO	living modified organisms
MGs	maturity groups
MOA	mode of action

MT	metric tons
N ₂ O	nitrous oxide
NAAQS	National Ambient Air Quality Standards
NAPPO	North American Plant Protection Organization
NEPA	National Environmental Policy Act
NHPA	National Historic Preservation Act
NMFS	National Marine Fisheries Service
NO ₂	nitrogen dioxide
NOP	National Organic Program
O ₃	ozone
Pb	lead
PDP	Pesticide Data Program
PIPs	plant-incorporated protectants
PM	coarse particulate matter
PM _{2.5}	fine particles less than 2.5 micrometers in diameter
PM ₁₀	particles greater than 2.5 micrometers and less than 10 micrometers in diameter
PPRA	Plant Pest Risk Analysis
SDWA	Safe Drinking Water Act of 1974
SO ₂	sulfur dioxide
SOM	soil organic matter
SSA	sole source aquifer
T&E	threatened and endangered species
TMDL	total maximum daily loads
TSCA	Toxic Substances Control Act
U.S.	United States
U.S.C.	United States Code
USDA	United States Department of Agriculture
USFWS	United States Fish and Wildlife Service
WPS	Worker Protection Standard
WSSA	Weed Science Society of America

1 PURPOSE AND NEED

BASF Corporation, Research Triangle Park, North Carolina (referred to as BASF in this document) submitted a petition (19-317-01p) to the U.S. Department of Agriculture (USDA), Animal and Plant Health Inspection Service (APHIS) on January 28, 2020 (BASF 2020). The petitioner requested that APHIS make a regulatory determination for Soybean Event IND 00410-5 (referred to as GMB151 Soybean in this document), which was developed using genetic engineering. The petitioner asserted that GMB151 Soybean should no longer be regulated under Title 7 of the Code of Federal Regulations part 340 (7 CFR 340) because it does not pose a plant pest risk. As described in more detail elsewhere in this chapter, APHIS regulations provide that any person or entity with legal standing may submit a petition to the Agency with a request that an organism not be regulated because it is unlikely to pose a plant pest risk.

This Draft Environmental Assessment (EA) was prepared by APHIS to evaluate the potential of effects from its regulatory decision for GMB151 Soybean to cause significant impacts on the human environment¹. One purpose of a NEPA analysis is to ensure that agencies assess possible effects of their actions for significant impacts, and consider them in their decision-making process. This Draft EA reports results of the Agency's thorough analysis of both a decision to continue regulating or to no longer regulate GMB151 Soybean as a plant pest. It includes a review of the Agency's findings about the potential for significant impacts from the effects, both beneficial and adverse, from either decision, and inform the public of these findings.

This EA has been prepared in compliance with the National Environmental Policy Act (NEPA, 42 U.S.C. § 4321 et seq.), the Council of Environmental Quality's (CEQ) NEPA-implementing regulations (40 CFR parts 1500-1508), and USDA and APHIS NEPA-implementing regulations (7 CFR part 1b, and 7 CFR part 372).

1.1 PURPOSE OF GMB151 SOYBEAN

GMB151 Soybean produces a modified 4-hydroxyphenylpyruvate dioxygenase (HPPD-4) enzyme, which is also produced naturally by the bacterium, *Pseudomonas fluorescens*. HPPD-4 confers resistance to herbicides that are HPPD inhibitors such as isoxaflutole. The source organism of the HPPD-4 protein, *P. fluorescens*, is a non-pathogenic bacterium that is ubiquitous in nature and has a history of safe use. HPPD proteins are also ubiquitous in nature in nearly all aerobic organisms, (e.g., bacteria, fungi, plants, and animals including mammals).

GMB151 Soybean also expresses a crystalline protein, Cry14Ab-1, that is an endotoxin naturally produced by a strain of the bacterium, *Bacillus thuringiensis*. Expression of Cry14Ab-1 in GMB151 Soybean confers resistance to the soybean cyst nematode (SCN), *Heterodera glycines* Ichinohe.

BASF has indicated that the method of Hoekema et al. (1983), which involves the disarmed bacterium, *Agrobacterium tumefaciens* strain LBA4404, was used to mediate gene transfer. The *Agrobacterium*-mediated transformation vector, pSZ8832, containing the *cry14Ab-1.b* and

¹Human environment includes the natural and physical environment and the relationship of people with that environment. When economic or social and natural or physical environmental effects are interrelated, the NEPA analysis may also address these potential impacts (40 CFR §1508.14).

hppdPf-4Pa gene cassettes, was used to insert the *cry14Ab-1.b* gene from *B. thuringiensis* and the *hppdPf-4Pa* gene from *P. fluorescens* into the GMB151 Soybean genome (BASF 2020).

SCN is a major pest of soybeans worldwide; it is an invasive pest occurring in most of the soybean growing regions of the United States (NCSRP No Date). Losses to soybean growers from SCN vary depending on location, genetic background of the variety, and the cropping system used. Susceptible varieties grown in the United States suffer yield reductions estimated at 15-30% or more (NCSRP No Date).

Planting soybean varieties resistant to SCN and rotating plantings with other crops that are not hosts of SCN are the most effective ways of managing SCN infestations. To date, more than one thousand SCN-resistant soybean varieties are available to farmers in the United States (Tylka and Mullaney 2018). If GMB151 Soybean is not regulated, it will likely be crossed with other commercially available SCN-resistant varieties, thereby creating lines with multiple resistance to extend the durability of both GMB151 Soybean and other SCN-resistant soybean varieties.

In addition to controlling invertebrate pests, effective weed management in soybean cropping systems is critical for maintaining high-yield production. Establishing good weed control is especially important during early vegetative growth, and also during the early reproductive growth stages (Van Acker and Swanton 1993). Herbicides are essential for weed management of many crops because they are a cost-effective method for weed control that helps make farming a profitable venture. Inclusion of resistance to HPPD-inhibiting herbicides in GMB151 Soybean will provide growers with an additional mode of action for the control of herbicide-resistant (HR) weeds, which provides more options to growers, when choosing weed control products.

1.2 COORDINATED FRAMEWORK FOR REGULATING BIOTECHNOLOGY

On June 26, 1986, the White House Office of Science and Technology Policy issued the “Coordinated Framework for the Regulation of Biotechnology” (referred to as the Coordinated Framework in this document), which outlined federal regulatory policy for ensuring the safety of biotechnology products (51 FR 23302 1986). The primary federal agencies responsible for oversight of biotechnology products are the U.S. Department of Agriculture (USDA), the U.S. Environmental Protection Agency (EPA), and the U.S. Food and Drug Administration (FDA).

In 2015, the Executive Office of the President (EOP) issued a memorandum directing the USDA, EPA, and FDA to clarify current roles and responsibilities in the regulation of biotechnology products; develop a long-term strategy to ensure that the federal biotechnology regulatory system is prepared for the future products of biotechnology; and commission an independent, expert analysis of the future landscape of biotechnology products. In 2016, the Emerging Technologies Interagency Policy Coordination Committee’s Biotechnology Working Group (BTW) published the “National Strategy for Modernizing the Regulatory System for Biotechnology Products” (ETIPCC 2016). One recommendation was to modernize the Coordinated Framework to make the policy consistent with technological changes that occurred after its initial development in 1986. In response to the BTW recommendations, the policy was updated and published on January 4, 2017 (81 FR 65414 2017).

USDA-APHIS is responsible for protecting animal and plant health. USDA-APHIS regulates products of biotechnology that may pose a risk to agricultural plants and agriculturally important natural resources under the authorities provided by the plant pest provisions of the PPA, as

amended (7 U.S. Code (U.S.C.) 7701–7772), and implementing regulations at 7 CFR 340.

The purpose of EPA oversight is to protect human and environmental health. EPA regulates pesticides under the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA) (7 U.S.C. 136 et seq.). EPA regulatory oversight of pesticides includes those expressed by organisms produced using genetic engineering, which are referred to as plant-incorporated protectants (PIPs). EPA has oversight authorized under the Federal Food, Drug, and Cosmetic Act (FFDCA; 21 U.S.C. 301 et seq.) to either set maximum residue limits, commonly referred to as tolerances, for pesticide residues that may remain on or in food and animal feed, or establish an exemption from the requirement for a tolerance.

Both FDA and USDA monitor food and animal feed for pesticide residues to enforce tolerances to ensure protection of human health. USDA collects data on pesticide residues as part of the Pesticide Data Program (PDP). PDP activities include sampling, testing, and reporting residues detected in meat and dairy products in the U.S. food supply. The program is implemented through cooperation with state agriculture departments and other federal agencies.

EPA uses PDP data to prepare pesticide dietary exposure assessments. EPA also regulates certain transgenic microorganisms (agricultural uses other than pesticides) under the Toxic Substances Control Act (15 U.S.C. 53 et seq.).

The purpose of FDA oversight is to ensure that human and animal foods and drugs are safe and sanitary. The FDA regulates a wide variety of products, including human and animal foods, cosmetics, human and veterinary drugs, and human biological products under the authority of the FFDCA and Food Safety Modernization Act.

1.3 PURPOSE AND NEED TO ISSUE A REGULATORY STATUS DETERMINATION

Under the authority of the plant pest provisions of the Plant Protection Act (7 U.S.C. 7701 *et seq.*), the regulations in 7 CFR 340 for movement of organisms modified or produced using genetic engineering regulate importation, interstate movement, or release into the environment of such organisms that are plant pests or pose a plausible plant pest risk.²

APHIS recently revised 7 CFR 340 and issued a final rule, published in the *Federal Register* on May 18, 2020 (85 FR 29790-29838, Docket No. APHIS-2018-0034).³ However, the final rule is being implemented in phases. APHIS' new Regulatory Status Review (RSR) process, which replaces the petition process for determination of nonregulated status process, became effective April 5, 2021 for corn, soybean, cotton, potato, tomato, and alfalfa. The RSR process is effective for all crops as of October 1, 2021. However, for crops for which an RSR process is still under development APHIS will continue to receive and consider petitions for determination of

²Genetic engineering in the context of 7 CFR 340 refers to biotechnology-based techniques that use recombinant, synthesized, or amplified nucleic acids to modify or create a genome. Various terms are used in the lay and scientific peer-reviewed literature in reference to new plant varieties that have been developed using modern molecular biology tools: “agricultural biotechnology,” “genetically engineered,” and “genetically modified.” In this EA, the terms “genetic engineering” and “biotechnology” may be used interchangeably. The term “transgenic” may also be used when discussing or referring to a transgene introduced into the genome of a plant. The USDA does not regulate plants developed through by traditional breeding techniques, including biolistics-, chemical- and radiation-based mutagenesis if they are not plant pests or developed using plant pests.

³To view the final rule, go to www.regulations.gov and enter “APHIS-2018-0034” in the Search field.

nonregulated status in accordance with the legacy regulations in § 340.6 of 7 CFR 340 (85 FR 29815).

The petition for a determination of nonregulated status that is the subject of this Draft EA (BASF 2020) is being evaluated in accordance with the regulations at 7 CFR 340.6 (2020), as it was received by APHIS in January 2020. Pursuant to the terms set forth in the final rule, any person or entity can submit a petition to APHIS seeking a determination that an organism should not be regulated under 7 CFR 340. APHIS must respond to petitioners with a decision to approve or deny the regulatory action requested by a petitioner. An organism produced using genetic engineering is no longer subject to the requirements of 7 CFR 340 or the plant pest provisions of the PPA if APHIS determines, through the conduct of a Plant Pest Risk Assessment (PPRA), that it is unlikely to pose a plant pest risk.

Consistent with APHIS regulations (7 CFR 340), importation into, interstate movement within, and field trials of GMB151 Soybean in the United States require permits issued by APHIS or notifications acknowledged by the Agency. Since 2013, field trials of GMB151 Soybean have been conducted by BASF in diverse growing regions within the United States. Results from these field trials are reported in the GMB151 Soybean petition (BASF 2020), and analyzed for plant pest risk in a Draft APHIS Plant Pest Risk Assessment (PPRA) (USDA-APHIS 2020).

In its petition, BASF provided evidence that GMB151 Soybean does not pose a plant pest risk or weediness potential, so it should not be regulated by APHIS (BASF 2020). If the Agency makes a determination of nonregulated status, it would pertain to GMB151 Soybean itself, and any progeny derived from crossing it with other soybean varieties that are not regulated under 7 CFR 340 (i.e., conventional soybean varieties, or other soybean varieties produced using genetic engineering that APHIS has determined are not regulated as plant pests). APHIS prepared this Environmental Assessment (EA), as required by the National Environmental Policy Act (NEPA), to determine if its regulatory decision (either to continue regulating or no longer regulate GMB151 Soybean) could have any significant impacts on the human environment. As part of the process required for NEPA compliance, this EA also informs the public about the environmental analysis the Agency made. It is intended to promote public participation and provide input to assist the APHIS decision maker in determining the regulatory status of GMB151 Soybean.

As required by 7 CFR 340.6, APHIS must respond to petitioners that request a determination of the regulated status of organisms produced using genetic engineering, including plants such as GMB151 Soybean. When a petition for nonregulated status is submitted, APHIS must determine if the organism is unlikely to pose a plant pest risk. The petitioner is required under 7 CFR 340 (§340.6(c)(4)) to provide information related to plant pest risk that the Agency may use to compare the plant pest risk of the regulated article to that of the corresponding (unmodified) organism. An organism produced using genetic engineering is no longer subject to the regulatory requirements of 7 CFR 340 or the plant pest provisions of the PPA, when APHIS determines that it is unlikely to pose a plant pest risk.

APHIS must respond to the January 2020 petition (19-317-01p) from BASF (BASF 2020) with a regulatory status decision for GMB151 Soybean consistent with 7 CFR 340. APHIS prepared this EA to document possible environmental effects of its decision, and evaluate their potential to cause significant impacts consistent with the Council of Environmental Quality's (CEQ) NEPA regulations (40 CFR parts 1500-1508), and the USDA departmental and APHIS NEPA-implementing regulations and procedures (7 CFR part 1b, and 7 CFR part 372).

2 SCOPING AND PUBLIC INVOLVEMENT

APHIS seeks public comment on draft EAs through notices published in the Federal Register. On March 6, 2012, APHIS published in the *Federal Register* (77 FR 13258-13260, Docket No. APHIS-2011-0129) a notice describing our public review process for soliciting public comments and information when considering petitions for determinations of nonregulated status for organisms developed using genetic engineering.⁴

2.1 PUBLIC INVOLVEMENT FOR PETITION 19-317-01p

APHIS made the BASF petition requesting non-regulated status for GMB151 Soybean available for public review in a notification⁵ in the *Federal Register* (85 FR 32004 2020) on May 28, 2020. The 60-day public comment period closed on July 27, 2020. APHIS received nine comments. The petition and comments are available⁶ for public review on regulations.gov, the U.S. federal government web site that serves as an internet portal and document repository for U.S. government documents (Docket No. APHIS-2020-0023).

Two comments were supportive of removing regulatory constraints on the GMB151 Soybean variety. Another comment addressed the pesticide registration issue about the Cry14Ab-1 protein PIP, urging EPA to make it available as a new active ingredient. Two other comments, while generally supportive of the development of crops produced using genetic engineering, expressed concerns about the risks and liabilities from possible disruptive effects on U.S. exports if residues from a deregulated biotech soybean enter the supply chain in cases where the soybean has not been approved in export markets. They emphasized the need for careful vetting of biotech crops and the need for stewardship measures if commercialized. BASF emphasized in its petition, a commitment to stewardship to meet applicable regulatory requirements for GMB151 Soybean in the country of intended production and for key import countries to ensure compliance, maintain product integrity, and assist in minimizing the potential for trade disruptions (BASF 2020).

Four comments expressed opposition to a determination of nonregulated status for GMB151 Soybean based on general opposition to the use of organisms produced using genetic engineering, but did not cite or provide documentation specific to why the GMB151 Soybean variety should continue to be regulated under 7 CFR 340. All comments were considered, carefully analyzed for relevancy, and addressed in this EA according to NEPA regulatory requirements.

APHIS determined from its initial review of the petition for GMB151 Soybean that the review process for the draft PPRA and draft NEPA documents (EA and FONSI) should follow

⁴*Federal Register*, Vol. 77, No. 44, Tuesday, March 6, 2012, p.13258 – Biotechnology Regulatory Services; Changes Regarding the Solicitation of Public Comment for Petitions for Determinations of Nonregulated Status for Genetically Engineered Organisms. This notice can be accessed at: <http://www.gpo.gov/fdsys/pkg/FR-2012-03-06/pdf/2012-5364.pdf>

⁵This notice can be accessed at: <https://www.federalregister.gov/documents/2020/05/28/2020-11492/basf-corporation-petition-for-a-determination-of-nonregulated-status-for-plant-parasitic>

⁶The docket can be accessed at: <https://www.regulations.gov/docket/APHIS-2020-0023>

The petition alone is also accessible on the APHIS web site:

<https://www.aphis.usda.gov/aphis/ourfocus/biotechnology/permits-notifications-petitions/petitions/petition-status>

Approach 2, as described in the Agency's 2012 revisions (77 FR 13258 2012) to the procedures it follows to promote public participation in its decision making relevant to the regulation of organisms produced using genetic engineering. This decision was made because APHIS has not previously analyzed plant pest risk for any soybean varieties that express the Cry14Ab-1 protein.

2.2 ISSUES CONSIDERED

The issues addressed in this Draft EA were developed by considering similar ones identified and addressed in prior NEPA documents, those identified in public comments for BASF's petition and other petitions for organisms produced using genetic engineering, information in the scientific literature on agricultural biotechnology, and issues identified by APHIS as specific to soybean. These issues were addressed in this Draft EA under the following subject categories:

Agricultural Production:

- Areas and Acreage of Soybean Production
- Agronomic Practices
- Soybean Seed Production
- Organic Soybean Production

Environmental Resources:

- Soil Quality
- Water Resources
- Air Quality
- Animal Communities
- Plant Communities
- Soil Microorganisms
- Biological Diversity
- Gene Movement

Animal Health:

- Animal Feed Quality
- Livestock Health

Human Health:

- Public Health
- Worker Health and Safety

Socioeconomics:

- Domestic Economic Environment
- Trade Economic Environment

Cumulative Impacts

Threatened and Endangered Species

Other U.S. Regulatory Approvals and Compliance with Other Laws

3 ALTERNATIVES

NEPA implementing regulations (40 C.F.R. § 1502.14) require the evaluation of all alternatives that appear reasonable and appropriate to the purpose and need of an agency's action. For this USDA APHIS action, a regulatory determination for BASF GMB151 Soybean, two alternatives were evaluated in this EA: (1) No Action Alternative, which would continue the current regulated status of GMB151 Soybean if selected; (2) Preferred Alternative, which would result in nonregulated status for GMB151 Soybean if selected.

3.1 NO ACTION ALTERNATIVE: CONTINUE REGULATING GMB151 SOYBEAN

Under the No Action Alternative, APHIS would deny the petition request by BASF (BASF 2020), so there would be no change in the regulatory status of GMB151 Soybean; it and any soybean varieties derived from it would continue to be regulated articles under 7 CFR 340. APHIS would continue to require permits for introductions and movement of GMB151 Soybean grown in the United States. Because APHIS has concluded from its draft PPRA (USDA-APHIS 2020) that GMB151 Soybean is unlikely to pose a plant pest risk, choosing this alternative would not be an appropriate response to the petition for nonregulated status because it would not satisfactorily meet the purpose and need for making a science-based regulatory status decision pursuant to the requirements of 7 CFR 340.

3.2 PREFERRED ALTERNATIVE: NONREGULATED STATUS FOR GMB151 SOYBEAN

Under the Preferred Alternative, GMB151 Soybean and any varieties derived from crosses between it and other soybean varieties that are not regulated would no longer be regulated under 7 CFR 340. APHIS has determined that GMB151 Soybean is unlikely to pose a plant pest risk based on available scientific evidence (USDA-APHIS 2020), therefore, if this alternative is selected, permits or notifications acknowledged by APHIS would no longer be required to grow GMB151 Soybean or progeny derived from it in the United States. This alternative best meets the purpose and need to respond appropriately to the petition for nonregulated status of IND00410-5 Soybean based on the requirements in 7 CFR 340 and the Agency's authority under the plant pest provisions of the PPA.

3.3 ALTERNATIVES CONSIDERED BUT EXCLUDED FROM FURTHER DETAILED ANALYSIS IN THIS EA

APHIS considered several other alternatives for this EA. These included: approve the petition only in part as provided for in § 340.6(d)(3)(i) of the regulations (e.g., allow nonregulated status for GMB151 Soybean crops grown in limited regions of the United States); establish mandatory rules for isolation or geographic separation of biotech and non-biotech cropping systems; require testing for the presence of biotech crop plant material in non-biotech crops and commodities.

Based on the draft PPRA (USDA-APHIS 2020) for GMB151 Soybean and the Agency's past experience with regulating biotech soybean varieties under 7 CFR 340, APHIS concluded that it is unlikely to pose a plant pest risk. Therefore, the imposition of testing, release, and/or isolation

requirements on GMB151 Soybean would be inconsistent with the Agency's statutory authority under the plant pest provisions of the PPA, implementing regulations at 7 CFR 340, and the federal regulatory policies of the Coordinated Framework. Because it would neither be reasonable nor appropriate for APHIS to evaluate alternatives for actions that exceed its statutory authority, the alternatives summarized above were excluded from further detailed analysis in this EA.

3.4 COMPARISON OF ALTERNATIVES

Table 1 includes a summary and comparison of possible impacts associated with selection of each of the alternatives evaluated in this EA. Details about the impact assessment for each are reported in Chapter 4 (Environmental Consequences).

Table 1. Summary of Potential Impacts and Consequences of Alternatives.

Attribute/Measure	Alternative A: No Action	Alternative B: Determination of Nonregulated Status
Meets Purpose and Need, and Objectives:	No	Yes
Agricultural Production		
Areas and Acreage of Soybean Production:	Current trends in acreage and areas of production are likely to continue to be driven by market conditions and federal policies that influence demand for U.S. soybeans (e.g., demand for animal feed, biodiesel and exports). U.S. 2020 soybean planted acreage (83.8 million) was up 10% from 2019 (USDA NASS 2020a), and is projected to remain level through 2028 (USDA-OCE 2018); selection of the No Action Alternative would not be expected to change this estimate, so would not increase or decrease soybean acreage.	If GMB151 Soybean were no longer regulated it would only be expected to be planted as an alternative to other varieties in the United States, so soybean acreage under the Preferred Alternative would be about the same as for the No Action Alternative.

Attribute/Measure	Alternative A: No Action	Alternative B: Determination of Nonregulated Status
Agronomic Practices:	Soybean management practices and methods that increase yield such as fertilization, crop rotation, irrigation, pest management, and plant residue management would be expected to continue as currently practiced. Some conservation tillage practices may be replaced by conventional tillage, where this is the only alternative to control increasing HR weed problems.	The agronomic characteristics and cultivation practices used for the production of GMB151 Soybean are the same as those used for the cultivation of other commercially available soybean varieties, so they would remain unchanged from the No Action Alternative.
Pesticide Use:	The EPA approves and labels uses of pesticides on soybeans. Commercial soybean growers would continue to use the same pesticides for soybean insect pests and weeds as are currently used.	The EPA regulatory oversight of pesticides would not change. Most nematicides for SCN are prescribed as seed treatments to be used in conjunction with resistant varieties. With the exception of SCN, GMB151 Soybean is susceptible to the same insect and other invertebrate pests and pathogens that affect most other commercially available conventional and biotech soybean varieties, so pest management practices would not change from the No Action Alternative. Growers with weeds resistant to herbicides with other modes of action may choose this HPPD-resistant variety for weed management.

Attribute/Measure	Alternative A: No Action	Alternative B: Determination of Nonregulated Status
Organic Soybean Production:	Methods currently used for certified seed production to maintain soybean seed identity and meet National Organic Standards would continue unchanged. The availability of biotech soybean is unrelated to the market share proportion of organic soybeans.	Measures used by organic soybean producers to manage, identify, and preserve organic production systems would not change. Similar to other commercially available biotech soybean varieties, GMB151 Soybean does not present any new or different issues or impacts for organic soybean producers or consumers. Other HR soybean varieties that are not regulated are currently planted by growers. GMB151 Soybean would only replace these as another HR alternative.
Soybean Seed Production:	Quality control methods, such as those of the Association of Official Seed Certifying Agencies (https://www.aosca.org/) for certifying seed to ensure varietal purity would continue to be available.	Practices to ensure varietal purity would remain the same as for the No Action Alternative. Tests would be available to determine the presence of genes that convey SCN and HPPD resistance traits in GMB151 Soybean.
Physical Environment		

Attribute/Measure	Alternative A: No Action	Alternative B: Determination of Nonregulated Status
Water Resources:	<p>Agronomic practices that could impact water resources (e.g., irrigation, tillage practices, and the application of pesticides and fertilizers) would be expected to continue. The use of EPA-registered pesticides for soybean production in accordance with label directions would continue to prevent unacceptable risks to water quality. Historic trends of increased soybean yield on existing cropland would continue unchanged, so any current impacts on water resources from soybean production would not change significantly.</p>	<p>Except for replacing herbicides with other modes of action with HPPD-based herbicides, the production of GMB151 Soybean is not expected to change current agronomic practices, acreage, or the range of production areas, so current effects from runoff on water resources would not change. Use of HPPD-based herbicides likely offsets the need to change tillage practices to control HR weeds resistant to currently available herbicides, so soil erosion impacts on water quality from soybean production may be reduced or would not change. Other HPPD HR soybean varieties that APHIS assessed previously (USDA-APHIS 2014, 2013) are not regulated and are currently available to growers. If it is not regulated, GMB151 Soybean will only be another HPPD-resistant alternative to growers, so herbicide use will not change.</p>
Air Quality:	<p>Current soybean agronomic practices that impact air quality, such as tillage, application of farm chemicals, and use of exhaust-emitting mechanized equipment would not change, so current environmental impacts would not change significantly.</p>	<p>Except for replacing herbicides with other modes of action with HPPD-based herbicides, the production of GMB151 Soybean is not expected to differ significantly from the No Action Alternative. Use of HPPD herbicides would likely offset the need to change tillage practices to control HR weeds resistant to currently available herbicides, so soil erosion impacts on air quality from soybean production may be reduced or would not change significantly from that of the No Action alternative. HPPD use is not expected to increase relative to the no action alternative.</p>

Attribute/Measure	Alternative A: No Action	Alternative B: Determination of Nonregulated Status
Soil Quality:	Most cropping practices that impact soil such as tillage, contouring, cover crops, agricultural chemical management, and crop rotation would continue unchanged, but some tillage practices (e.g., conservation), may change to conventional where this is the only alternative to control increasing HR weed problems.	Production of GMB151 Soybean would not be expected to change cropping practices. Use of HPPD herbicides would likely offset the need to change tillage practices to control HR weeds resistant to currently available herbicides, which would prevent or reduce soil quality losses from erosion. HPPD use is not expected to increase relative to the no action alternative.
Biological Resources		
Animal Communities:	Non-biotech and biotech soybeans that are not regulated have been shown to have no allergenic or toxic effects on animal communities. Soybean agronomic practices such as tillage, cultivation, farm chemical applications, and the use of mechanized agricultural equipment would continue to impact animal communities unchanged.	Field trials demonstrated that growth and disease characteristics of GMB151 Soybean are not significantly different from other soybean varieties that are not regulated, so no changes to soybean agronomic practices potentially impacting animal communities would occur other than the use of HPPD herbicide applications, where HR weeds resistant to other modes of action are a problem. HPPD use is not expected to increase relative to the no action alternative

Attribute/Measure	Alternative A: No Action	Alternative B: Determination of Nonregulated Status
<p>Plant Communities:</p>	<p>Most commercial soybean acreage is planted with transgenic varieties, and this would continue unchanged. Most agronomic practices would not change except where the continuing increasing problem of HR weeds forces growers to modify methods (e.g., tillage; alternative herbicide choices) to control weeds. Herbicide use in accordance with the EPA registration requirements would continue to ensure that no unacceptable risks to non-target plants and plant communities would occur.</p>	<p>Field trials and laboratory analyses show no differences between GMB151 Soybean and other soybean varieties (conventional and transgenic) in growth, reproduction, and susceptibility to pathogens and other pests except the target species (SCN). Except for the option to substitute HPPD herbicides with other herbicides currently used, agronomic practices to cultivate GMB151 Soybean would not differ from the No Action Alternative. HPPD use is not expected to increase relative to the no action alternative. .</p>

Attribute/Measure	Alternative A: No Action	Alternative B: Determination of Nonregulated Status
<p>Gene Movement:</p>	<p>GMB151 Soybean would continue to be cultivated only under regulated conditions. The availability of biotech, conventional, and organic soybeans would not change as a result of the continued regulation of GMB151 Soybean. Because there are no wild soybean relatives in the United States, and soybeans are mostly self-pollinated at rates that decrease significantly with distance, gene flow and introgression from soybean to wild or weedy species are highly unlikely. Any risk is further limited because soybeans are not frost tolerant, do not reproduce vegetatively, exhibit poor seed dispersal, and any volunteers that persist in warmer U.S. climates can be easily controlled with common agronomic practices.</p>	<p>Field and laboratory test results show that there are no significant differences among the traits in GMB151 Soybean that influence gene flow or weediness, when compared to soybean varieties that are not regulated. Traits for SCN resistance and HPPD herbicide resistance would not change gene movement characteristics, so there would be no significant impacts compared to the No Action Alternative.</p>

Attribute/Measure	Alternative A: No Action	Alternative B: Determination of Nonregulated Status
Soil Microorganisms:	<p>Agronomic practices used for soybean production, such as soil inoculation, tillage and the application of agricultural chemicals (pesticides and fertilizers) that potentially impact microorganisms would continue unchanged.</p>	<p>Field and greenhouse tests show no significant differences from other nonregulated soybean varieties in the parameters measured to assess the symbiotic relationship of GMB151 Soybean with its <i>Rhizobium</i> spp. symbionts. GMB151 Soybean would not result in any significant changes to current soybean cropping practices that may impact microorganisms except that HPPD herbicides may be substituted for herbicides with other modes of action, where HR weeds are a problem. Other HPPD HR transgenic soybean varieties that APHIS assessed previously (USDA-APHIS 2014, 2013) are not regulated and are currently available to growers. If it is not regulated, GMB151 Soybean will only be another HPPD-resistant alternative to growers, so herbicide use would not be expected to change.</p>

Attribute/Measure	Alternative A: No Action	Alternative B: Determination of Nonregulated Status
Biological Diversity:	Agronomic practices used for soybean production and yield optimization, such as tillage, the application of agricultural chemicals (pesticides and fertilizers), timing of planting, and row spacing, would be expected to continue unchanged. Agronomic practices that benefit biodiversity both on cropland (e.g., intercropping, agroforestry, crop rotations, cover crops, and no-tillage) and on adjacent non-cropland (e.g., woodlots, fencerows, hedgerows, and wetlands) would remain the same.	GMB151 Soybean would not change current soybean cropping practices that may impact biodiversity because field and laboratory testing demonstrate its growth, reproduction, and interactions with pests and diseases are the same as or not significantly different from other nonregulated varieties other than its resistance to SCN. GMB151 Soybean poses no potential for naturally occurring, pollen-mediated gene flow and transgene introgression, so is not expected to affect genetic diversity. Testing has confirmed that the Cry14Ab-1e protein expressed by GMB151 Soybean does not have unacceptable risks to or impacts on non-target organisms (BASF 2020).
Public Health		
Farm Worker Safety and Health:	Farm workers are exposed to potential allergens from soybean plants, hazards from farm equipment used to grow and harvest soybeans, and pesticides applied to soybeans. Hazards to farm workers would not change from selection of the No Action Alternative.	EPA Worker Protection Standards (WPS) implement protections for agricultural workers, handlers, and their families (40 CFR 170). If the Preferred Alternative were selected, GMB151 Soybean would not change current soybean cropping practices, so hazards would be the same as under the No Action Alternative.

Attribute/Measure	Alternative A: No Action	Alternative B: Determination of Nonregulated Status
<p>Human Health:</p>	<p>Compositional and nutritional characteristics of nonregulated biotech soybean varieties have been determined to pose no risk to human health. EPA-approved pesticides would continue to be used for pest management in both biotech and conventional soybean cultivation. Use of registered pesticides in accordance with EPA-approved labels protect human health and worker safety. EPA also establishes tolerances for pesticide residue that give a reasonable certainty of no harm to the general population and any subgroup from the use of pesticides at the approved levels and methods of application.</p>	<p>Laboratory and field testing demonstrated that there are no biologically meaningful differences for compositional and nutritional characteristics between conventional and GMB151 Soybean. Safety testing of the GMB151 Soybean Cry14Ab-1 and HPPD proteins showed that they are degraded rapidly and completely in simulated gastric fluid. Testing also showed that the GMB151 Soybean Cry14Ab-1 and HPPD proteins have no similarities to known allergens, and are not toxic to mammals.</p> <p>On January 28, 2019, BASF initiated a consultation (BNF 172) with FDA that included molecular, compositional, nutritional data, and other food and feed safety assessment data related to GMB151 Soybean (BASF 2020). EPA has established a permanent exemption (82FR57137) from the requirement for a tolerance for the HPPD-4 protein expressed in all food commodities when used as an inert ingredient. In addition, EPA concluded on June 8, 2020 that <i>B. thuringiensis</i> Cry14Ab-1 protein residue in or on soybean food and feed commodities are exempt from the requirement of a tolerance when expressed as a PIP in soybean plants (85 FR 35008; 40 CFR 174.540).</p>

Attribute/Measure	Alternative A: No Action	Alternative B: Determination of Nonregulated Status
Animal Feed:	<p>GMB151 Soybean would remain regulated and not be allowed for distribution to the animal feed market. Soybean-based animal feed would still be available from currently cultivated soybean crops, including both biotech and conventional soybean varieties. Nonregulated biotech soybean varieties used as animal feed have been previously determined not to pose any risk to animal health.</p>	<p>Laboratory and field testing demonstrated that there are no biologically meaningful differences for compositional and nutritional characteristics between conventional and GMB151 Soybean. Safety testing of the GMB151 Soybean Cry14Ab-1 and HPPD proteins showed that they have no toxic potential to mammals, and are degraded rapidly and completely in simulated gastric fluid, when present in animal feed. On January 28, 2019, BASF initiated a consultation (BNF 172) with the FDA that included molecular, composition, and nutrition data, and other food and feed safety assessment data related to GMB151 soybean (BASF 2020). In addition, EPA concluded on June 8, 2020 (40 CFR 174.540) that the Cry14Ab-1 protein is exempt from a food and feed tolerance, when it is expressed in soybean plants.</p>
Socioeconomic Environment		

Attribute/Measure	Alternative A: No Action	Alternative B: Determination of Nonregulated Status
<p>Domestic Economic Environment:</p>	<p>GMB151 Soybean would remain regulated by APHIS. Domestic growers would continue to utilize biotech and conventional soybean varieties based upon availability and market demand. U.S. soybeans would likely continue to be used domestically for animal feed with lesser amounts and byproducts used for oil or fresh consumption. Agronomic practices and conventional breeding techniques using herbicide- and pest-resistant varieties currently used to optimize yield and reduce production costs would be expected to continue unchanged. Average soybean yield is expected to continue to increase without expansion of soybean acreage while grower net returns are estimated to increase.</p>	<p>Field tests show the performance and composition of GMB151 Soybean is not substantially different from that of other conventional soybean reference varieties and although yield potential is increased, it would be similar to other commercially available conventional and biotech soybean varieties and subject to the same variables affecting agronomic practices and yields as other varieties. GMB151 Soybean would likely only replace other varieties of biotech soybean on existing cropland and not impact organic soybean production or markets. Since biotech soybeans represent over 90% of soybeans produced, the addition of GMB151 Soybean will have little incremental impact on the biotech sensitive market. Because losses from SCN would be reduced, soybean growers would likely experience improved profits under Alternative B.</p>

Attribute/Measure	Alternative A: No Action	Alternative B: Determination of Nonregulated Status
Trade Economic Environment:	<p>If GMB151 Soybean remains regulated by APHIS, U.S. soybean plantings will not be affected and are projected to rebound and remain relatively steady over the course of the next decade. U.S. soybeans will continue to be a major component of global production, and as a source of supply in the international market (USDA 2020). Although U.S. exports are expected to increase overall, increasing competition and tariffs on U.S. soybean exports are expected to reduce the U.S. export share (Hubbs 2018).</p>	<p>A determination of nonregulated status for GMB151 Soybean is not expected to have an effect on current trends affecting the trade economic environment. GMB151 Soybean is similar to other transgenic varieties. If it becomes commercially available as a non-regulated variety, it would only be substituted to replace other varieties where SCN- and/or HPPD-resistant varieties are required for pest management. If the Preferred Alternative is selected, there would not be any difference from choosing the No Action Alternative.</p> <p>BASF emphasized in its petition a commitment to stewardship to meet applicable regulatory requirements for GMB151 Soybean in the country of intended production and for key import countries to ensure compliance, maintain product integrity, and assist in minimizing the potential for trade disruptions (BASF 2020).</p>
Other Regulatory Approvals		
U.S. Agencies:	<p>Existing approvals for other nonregulated transgenic soybeans would not change.</p>	<p>EPA has concluded (40 CFR 174.540) that the Cry14Ab-1 protein is exempt from a food and feed tolerance, when it is expressed in soybean plants.</p>

Attribute/Measure	Alternative A: No Action	Alternative B: Determination of Nonregulated Status
Other countries	The existing status of other transgenic soybeans regulated in other countries would not change.	No Change from the No Action Alternative. BASF emphasized in its petition a commitment to stewardship to meet applicable regulatory requirements for GMB151 Soybean in the country of intended production and for key import countries to ensure compliance, maintain product integrity, and assist in minimizing the potential for trade disruptions (BASF 2020).
Compliance with Other Laws		
CAA, CWA, EOs:	Fully compliant	Fully compliant
	Fully compliant	Fully compliant

4 AFFECTED ENVIRONMENT

This chapter includes a review of the current status of the human environment as defined in the CEQ regulations for NEPA (40 CFR §1508.14). The components of the human environment that may be affected by a regulatory determination for GMB151 Soybean by APHIS under 7 CFR 340 were listed (see “Issues Considered”) in Chapter 1. A detailed description for each component follows. This information was the basis for comparison to identify effects that may result from a regulatory determination for GMB151 Soybean, as the Agency performed its analysis to assess the potential for significant impacts from it decision.

4.1 AGRICULTURAL PRODUCTION OF SOYBEANS

Soybean (*Glycine max* (L.) Merr.) is an economically important leguminous crop that is a source of vegetable oil and protein. Soybeans are grown for their seed, which is processed to yield oil and meal. Among oil seed crops, soybeans are ranked first in the world as a source of oil production (Chung and Singh 2008). In the United States, soybeans are also a major source of livestock animal feed and biodiesel fuel (USB 2012).

The genus *Glycine*, includes two subgenera and more than 25 species (Sherman-Broyles et al. 2014b). The subgenus *soja* consists of only two species, the cultivated *G. max* and its annual wild soybean progenitor, *G. soja* Sieb. & Zucc. The subgenus *glycine* includes at least 26 species. Most are perennials native to Australia and its surrounding islands. The domestication of *G. max* from its wild progenitor soybean (*G. soja* Sieb. & Zucc.) occurred in China or Southeast Asia between 3,000 and 9,000 years ago (Hymowitz 1970; Hymowitz and Newell 1981; Sedivy, Wu, and Hanzawa 2017).

Soybean is a self-pollinating species, propagated commercially by seed (OECD 2000). Soybean seeds contain about 18% oil and 38% protein (Hartman, West, and Herman 2011). Nearly all soybean meal (98%) is used for livestock or aquaculture feed (Hartman, West, and Herman 2011). Soybeans are grown worldwide. Leading soybean-producing countries include Argentina, Brazil, China, India and the United States (USDA-FAS 2019a, 2019b).

U.S. soybeans are grown mostly in the Midwest (Figure 1). Acreage increased rapidly after World War II until the late 1970s as a result of increased vegetable oil demand and higher meat consumption (USDA-ERS 2006). U.S. soybean acreage stabilized in the 1980s mostly because of farm programs that encouraged planting other crops.

In the 1990s, changes in farm programs, overseas demand, and lower production costs associated with herbicide-resistant (HR) crops, resulted in an increase in soybean acreage (USDA-ERS 2006). From 1992 to 2012, U.S. soybean acreage increased 31% from about 59.1 to 77.2 million acres (USDA-NASS 2012e, 2012d). About 90 million acres were planted in 2017 (Figure 2), followed by a decline in acreage that rebounded to about 83 million planted and harvested acres by 2020 (USDA-NASS 2020).

Soybean acreage in the major producing states (Figure 1) is commonly rotated with corn. Total soybean production in the United States (Figures 2 and 3) has increased in recent years because of an increase in both the area under cultivation and yield per unit area (USDA-NASS 2018b, 2017, 2017). A significant factor contributing to these increases in recent years is that soybean cultivation has expanded into the northern and western parts of the country because new

improved short-season soybean varieties have been developed that are better adapted to the climate of that region, and provide better profits (USDA-ERS 2010b) than wheat or older soybean varieties.




Figure 1. Soybean Planted Acres by County for Selected States.

Source: (USDA-NASS 2016b, 2020)

Soybean production has increased 35.6%, from nearly 2.2 billion bushels or 59.88 million metric tons (MT) in 1992 to approximately 3.0 billion bushels (81.7 million MT) in 2012. From 1991 to 2011, average yield increased approximately 17.6% from 34.2 bushels per acre to 41.5 bushels, but declined nationally in 2012 to 39.3 bushels per acre compared to 2011 average yields. By 2020, the harvest was 53.3 bushels per acre (USDA NASS 2020b).




Figure 2. U. S. Soybean Acreage: 2011-2020.

Source (USDA-NASS 2020)

USDA projects an estimated 3.6 billion bushels of soybeans (97.99 million MT) will be produced by the end of the 2021/2022 growing season. About 2.1 billion bushels (57.16 million MT) of this production will be used for domestic consumption and 1.6 billion bushels (43.55 million MT) will be exported (USDA-OCE 2012).

Improvements in soybean yield are challenged by both biotic and abiotic stress factors. Some typical abiotic stress factors include salinity, non-optimal temperatures, drought, flooding, and poor soil quality (Chung and Singh 2008). One objective of soybean breeding programs is to develop varieties that maintain yield under a broad array of environmental conditions.

Varieties have for many years been developed using conventional plant breeding methods. Combined with improved agronomic practices, these varieties have resulted in improved yields. The multigene components influencing soybean yield are complex. Genetic selection to develop soybean varieties adapted to lower yielding areas, and the need to develop regional soybean varieties for specific environments limits the availability and identification of those traits that can provide yield improvements across the entire spectrum of soybean production environments.




Figure 3. U.S. Soybean Yield: 1991-2019

Source: (USDA NASS 2020b)

4.1.1 Acreage and Regional Distribution of Soybean Production

Field testing of biotech crops began in the 1980s, (Fernandez-Cornejo and Caswell 2006), and biotech soybeans became commercially available in 1996 (USDA-ERS 2011a; Fernandez-Cornejo and Caswell 2006). By 2017, a biotech variety was grown on 94% of U.S. soybean acreage (Table 2) (USDA-ERS 2020a).

4.1.2 Agronomic Practices for Conventional Soybean Production

Soybeans are an herbaceous annual that grows as an erect bush (OECD 2000). It is a short-day plant, so flowers when days have fewer daylight hours (OECD 2000). As a result, photoperiod and temperature responses are important in determining areas of specific varietal adaptation. Soybean varieties are identified based on geographic bands of adaptation that run east-west, determined by latitude and day length. In North America, there are 13 described maturity groups (MGs), ranging from MG 000 in the north (45° latitude) to MG X near the equator. Within each maturity group, varieties are described as early, medium, or late maturing (OECD 2000).

Soybean seeds germinate at about 50°F (10°C). Under favorable conditions, seedlings emerge in 5-7 days. Inoculation of soybean fields, that were previously used to grow another crop, with

Bradyrhizobium japonicum, a nitrogen fixing bacterium that develops a symbiotic relationship with soybeans, dramatically increases plant production (OMAFRA 2011; Pedersen 2007; Missouri University of Science and Technology No Date). Inoculation is necessary for optimum efficiency of the nodules that form on soybean root systems (Pedersen 2007; Berglund and Helms 2003a). In the 1990s, the row spacing recommendation for soybeans was narrowed to seven inches to achieve greater yields. This was later changed to 15 inches to promote greater air circulation, which reduces diseases that impact yields (USDA-ERS 2010b).

Table 2. Biotech Soybeans as a Percentage of the Total U.S. Soybean Crop.

STATE	Transgenic Percentage of U.S. Soybean Crops Grown in:	
	2016	2017
Arkansas	96	97
Illinois	94	93
Indiana	92	92
Iowa	97	94
Kansas	95	94
Michigan	95	94
Minnesota	96	96
Mississippi	99	99
Missouri	89	87
Nebraska	96	94
North Dakota	95	95
Ohio	91	91
South Dakota	96	96
Wisconsin	94	92
Other States*	94	94
UNITED STATES	94	94
*All other states in the U.S. estimating program		

Source: (USDA-NASS 2018d)

Soybeans require more moisture to germinate than corn, and seed-to-soil contact is important for good early season soybean growth. An adequate water supply is especially important at planting, during pod-filling, and seed filling (Hoeft et al. 2000). Soybeans require approximately 20-25 inches of water during the growing season. In 2008, only 9% of harvested soybean acreage (about 12 million acres) was irrigated. States with the most irrigated soybean acreage are Nebraska, Arkansas, Mississippi, Missouri, and Kansas (USDA-NASS 2010). There was no substantial change in irrigated U.S. soybean acreage between 2008 and 2015, Irwin, et al. 2017).

Soybeans tolerate a broad spectrum of growing environments, but maximum yields require optimum soil conditions that include a pH range of 6.0-7.0 (NSRL No Date), and adequate levels of phosphorus, potassium, calcium, and magnesium, plus other minor nutrients. Because soybeans develop a symbiotic relationship with *B. japonicum* that promotes nitrogen fixation from atmospheric nitrogen, fertilizer nitrogen is not always required for optimum soybean production. In areas with increased amounts of salt or carbonates, or those that have no past history of soybean production, nitrogen amendments prior to or at the time of planting have been shown to increase yield if soil tests reveal levels that are not adequate (Franzen 1999; Berglund and Helms 2003a).

When grown in rotation with corn, a common practice is to fertilize the preceding corn crop with enough phosphorus and potassium to provide sufficient carry over for the subsequent soybean crop, so no supplemental fertilizer is needed (Ebelhar et al. 2004b; Franzen 1999; Berglund and Helms 2003a; Franzen 2019). Adequate amounts of calcium and magnesium are normally present if soil pH is at or near the optimum pH or has been recently treated with dolomitic limestone to achieve an optimum pH (Frank 2000; Harris 2011).

Crop Rotation

Crop rotation is a sustainable agriculture practice of growing a series of different crops in the same field in succession, usually according to a planned cycle of plantings. The primary goal of crop rotation is to achieve maximum short term (annual or seasonal) crop yields in a system that sustains the long-term productivity of the fields. It is a strategy designed to prevent long-term profit loss from depletion of resources by maintaining them at a level that supports profitable crop productivity (Hoeft et al. 2000). When applied effectively, rotating crops can improve soil quality and fertility. Since the roots of soybean plants share a symbiotic relationship with *B. japonicum* that fixes atmospheric nitrogen, this may decrease the requirements for fertilizer inputs for following crops, such as corn or wheat. Crop rotation also tends to reduce the incidence of plant diseases, insect pests and weed competition (Berglund and Helms 2003a; USDA-ERS 1997). Crop rotation may also include fallow periods in which no crop is grown for a season, or seeding of fields with a cover crop that prevents soil erosion and can provide livestock forage (USDA-NRCS 2010a; Hoeft et al. 2000).

Maximizing economic returns results from rotating crops in a sequence that efficiently produces the most net profit. Many factors at the individual farm level influence crop rotation choices, including soil type, anticipated commodity prices, farm labor requirements, fuel, fertilizer and seed costs, and regional climatic conditions (Duffy 2011; Hoeft et al. 2000; Langemeier 1997).

Soybeans are commonly rotated with corn, winter wheat, spring cereals, and dry beans (OECD 2000). Cropland used for soybean and corn production is nearly identical in many areas, such as Illinois, where more than 90% of the cropland is planted in a two-year corn-soybean rotation (Hoeft et al. 2000). Approximately 95% of U.S. soybean acreage is in a rotation system.

Soybeans may also be a cover crop in short rotations for its fixed nitrogen contribution to soil (Hoorman, Islam, and Sundermeier 2009). Where continuous soybean production is undertaken, yields may be reduced in the second or later years because pest and disease incidence may increase (Monsanto 2010; Pedersen et al. 2001; Whitaker 2017). In the Midwest, the crops planted most often in rotation with soybean include corn and wheat. Those soybean plantings in the Southeast that are grown in a rotation are most frequently followed by corn and cotton. Corn is most often the crop of choice for rotation with soybeans grown in the coastal states of the eastern United States. Double-cropping soybeans is also an option to increase returns. Soybeans are frequently planted in winter wheat stubble to produce a second crop in the same growing season. Double-cropping maximizes profits if high commodity prices can support it, but careful management to achieve uniform stands to sustain high yields and profitability is needed. These requirements include selection of appropriate varieties, a higher seeding rate, closer row spacing, and adequate moisture for germination (McMahon 2011).

Tillage

Soybean growers till soil to prepare seedbeds, dislodge compaction, incorporate fertilizers and herbicides, manage drainage within and outside fields, and control weeds (Heatherly et al. 2009). Tillage systems include conventional, reduced, conservation (including mulch-till, strip-till, ridge-till, and no-till), and deep. The primary purpose of conservation tillage is to reduce soil erosion (Heatherly et al. 2009).

In conventional tillage, after harvest crop residue is plowed into the soil to prepare a clean seedbed for planting and to reduce the growth of weeds, leaving less than 15% of crop residue on the surface (Heatherly et al. 2009; Towery and Werblow 2010). Conservation tillage uses tools that disturb soil less and leave more crop residue on the surface (at least 30%). No-till farming only disturbs the soil for planting seed (Towery and Werblow 2010; USDA-NRCS 2005). Crop residue includes materials left in an agricultural field after the crop has been harvested, including stalks and stubble (stems), leaves and seed pods (USDA-NRCS 2005). Residue aids in conserving soil moisture and reducing wind and water-induced soil erosion (Heatherly et al. 2009; USDA-ERS 1997; USDA-NRCS 2005). No-till systems are not meant to control weeds or dislodge soil compaction, so other strategies such as herbicide applications and track management of heavy machinery must be used in no-till fields for these problems (Heatherly 2012).

Since 1996, the use of a no-till has increased more than any other reduced tillage system. Nearly all of this shift is attributable to reliance on HR crop varieties (e.g., soybean, corn, cotton, canola) (Fawcett and Towery 2002). A 1997 survey revealed that farmers using no-till practices were more likely to adopt HR soybeans as an effective weed control practice, but the same study also found that the reliance on HR soybean varieties did not encourage adoption of no-till practices. However, later surveys revealed a corresponding increase in the use of no-till production practices (Sankula 2006; Carpenter et al. 2002) with the increase in availability of GR soybean varieties. From the introduction of GR soybeans in 1996 until 2004, no-till practices increased by 64% (Sankula 2006). Use of conservation tillage practices by U.S. soybean growers increased by 12 million acres (4.9 million hectares) from 51% in 1996 to 63% in 2008 (NRC 2010).

No-till soybean production is not suitable for all producers or areas. For example, no-till soybean production is less successful in heavier, cooler soils more typical of northern latitudes (NRC

2010; Kok, Fjell, and Kilgore 1997) where the potential for increased weed and insect pests and disease requires careful management (Pedersen et al. 2001; Peterson 1997; Shoup 2016; Peterson 2016).

Agronomic Inputs

Agronomic inputs, including water, soil and foliar nutrients, inoculants, fungicides, pesticides, and herbicides, are used in soybean production to maximize yields (Clevenger 2010; Hoeft et al. 2000; OECD 2000; OMAFRA 2011). Soil and foliar macronutrient applications to soybean primarily include nitrogen, phosphorous (phosphate), potassium (potash), calcium, and sulfur, with other micronutrient supplements such as zinc, iron, and magnesium applied as needed. Irrigation provides essential water for growth where rainfall is insufficient or erratic (see sections on Water Resources in this chapter, and Soil Quality in Chapter 4 for more details).

Nutrients. Fertilizers and other nutrients may be applied to the soil or sprayed on foliage in soybean production. Soil fertilizers have differential availability to plants based upon soil characteristics and moisture. For example, in a drought year, potassium may become fixed between clay layers until water moves through the soil again (Corn and Soybean Digest 2012). Fertilizers such as nitrogen, potassium and phosphorous may be incorporated into the soil at soybean planting by tillage or drilling. Fertilizer may be purposefully concentrated in bands at varying depths in the soil to enhance nutrient availability at different growth stages (Vitosh, Johnson, and Mengel 2007; Fernandez and White 2012). In conservation tillage, phosphorous and potassium may become vertically stratified from use of surface broadcast fertilizers that minimize soil disturbance. Therefore, there is a trend among farmers to enhance nutrient availability to sustain higher yields (Fernandez and White 2012).

On average, soybean removes 0.85 lbs of phosphate (phosphorous) and 1.2 lbs of potash (potassium) per bushel of seed produced (CAST 2009). Table 3 includes a summary of removal rates of nitrogen, phosphate and potassium for soybean, corn and wheat that are commonly rotated with soybean (Silva 2011). The data show that soybeans remove more nitrogen, potassium and phosphorous than either corn or wheat.

Table 3. Nutrient Removal Rates for Commonly Grown U.S. Grain Crops.

Crop	Pounds of Fertilizer Removed/Bushel Produced/Acre:		
	Nitrogen	Potash	Phosphate
Corn	0.9	0.37	0.27
Soybean	3.8	0.8	1.4
Wheat	1.2	0.63	0.37

Source: Silva (2011)

Research summarized by the Council for Agricultural and Science Technology indicates that adding nitrogen displaces rather than supplements natural cost-free nitrogen production in soybean cultivation, as the size, weight, and number of nitrogen-fixing nodules formed on soybean roots are actually reduced (CAST 2009). Application of nitrogen under drought conditions in acid subsoil conditions, in soils having low residual nitrogen, in a high-yield environment, or in late or double crop plantings has raised soybean yields, but not enough to offset the added cost. Potassium may change considerably from one testing time to the next, so it should also be regularly monitored to ensure optimum yields (CAST, 2009).

Phosphorous should be applied at least at the crop removal rate determined by regular soil testing. There is some concern that phosphorous for crop fertilizer is depleting phosphate rock deposits, which are a finite resource. However, finite does not necessarily mean that world reserves are being depleted. Price changes for phosphorous fertilizer are not caused by phosphate rock depletion; they're caused by many market factors (Heckenmueller, Narita, and Klepper 2014).

Soybeans are often grown in rotation with corn, and soil nutrient supplements applied to corn are often adequate to support soybean crops the following year without additional supplementation (Bender et al. 2013), making it more economical to apply nutrients such as nitrogen, potassium and phosphorous ahead of the corn crop in two-year corn-soybean rotations (CAST 2009). Other research has found that annual supplementation of potassium and phosphorous is most beneficial in the South where soybean to soybean rotation is more common (Heatherly 2012). Corn and soybean take up nutrients and both concentrate different nutrients in different parts of the plants (Mallarino et al. 2011). In plants, potassium is located mainly in the cytoplasm of cells and cell vacuoles where it activates enzymes, regulates stomata functions, and assists in transfer of compounds across membranes. In contrast, most phosphorous is located in cell membranes and nucleic acids, which is incorporated into plant organic matter, and is a major component of the metabolic, energy-rich compounds that drive plant metabolism.

Compared to potassium much more phosphorous is absorbed from the soil by soybeans. Some portion of these nutrients absorbed by the crop may be returned to the soil by leaving plant residue such as soybean foliage in the field (Mallarino et al. 2011).

Data for average chemical fertilizer application rates (USDA-NASS 2018e) for 2017 for USDA program states showed that nitrogen was applied at 18 lb/A, and phosphate and potash were applied at an average annual rate of 52 and 91 lb/A respectively. These supplements were applied on average only once per crop year. The relatively low rate of soybean nutrient supplementation likely results because most soybeans are rotated after corn crops that leave sufficient nutrients to sustain the subsequent soybean crop.

Inoculants. When added to soil as an inoculant, the bacterial symbiont, *B. japonicum* can increase soybean yields by about one bushel per acre (Conley and Christmas 2005). Historically, a nonsterile peat powder was applied to seed at planting as an inoculant carrier to the field. Improvements have since been made in inoculant manufacturing, such as using sterile carriers, adhesives to stick the inoculant to seeds, liquid carriers, concentrated frozen products, new organism strains, pre-inoculants, and inoculants containing extended biofertilizer and biopesticidal properties (Conley and Christmas 2005).

Pesticides. Feeding on soybean foliage, seed pods and roots by several different types of insects can reduce yield (Lorenz et al. 2006; Whitworth, Michaud, and Davis 2011). Nematodes are also

serious pests, especially the soybean cyst nematode, because effective pesticides are not available (Nelson and Bradley 2003). A combination of crop rotation to a non-susceptible host and the use of resistant varieties are used to manage the problem (Nelson and Bradley 2003). However, these resistant soybean varieties often provide lower yields than other commercially available varieties.

Insecticides and Nematicides: Economic thresholds for soybean insect infestations are used to decide if and when to apply integrated pest management (IPM) control measures (Higgins 1997; Whitworth 2016). Thresholds are typically based on field survey data for the number of pests present and/or extent of defoliation, such as those developed for management strategic plans of the National Information System of Regional IPM Centers (USDA 2011; IPM 2019). Data summarizing USDA NASS chemical insecticide a.i. (active ingredient) usage in 2017 for U.S. soybeans are included in Table 4. Based on total soybean acreage treated, the three most commonly applied insecticides were lambda-cyhalothrin, bifenthrin, and chlorpyrifos, which were applied to 8%, 5% and 3% of soybean acreage, respectively (USDA-NASS 2018a). Some growers may use other methods to control insect infestations including crop rotation, tillage, and biological control (i.e., beneficial organisms [predators and parasites of pest species]).

Table 4. Insecticides Most Commonly Applied to Soybean Acreage in 2017.

Insecticide (Active Ingredient: a.i.)	Percent of Soybean Acreage Treated	Average Application Rate (lbs a.i./acre)	Total Applied (lbs a.i.)
Lambda-cyhalothrin	8	0.03	215,000
Bifenthrin	5	0.06	247,000
Chlorpyrifos	3	0.34	876,000
Imidacloprid	2	0.08	109,000
Zeta-cypermethrin	2	0.02	25,000
Acephate	1	0.69	861,000
Chlorantraniliprole	1	0.06	55,000
Thiamethoxam	1	0.04	44,000
Beta-Cyfluthrin	1	0.04	41,000
Source: (USDA-NASS 2018a)			

There are only a limited number of nematicides registered for use on soybeans to control SCN and other nematode pests because applying chemical methods for their control is not cost effective compared with other management alternatives. Most nematicides for SCN are prescribed as seed treatments to be used in conjunction with resistant varieties ((University of

Arkansas Cooperative Extension Service 2021) (Tylka and Mullaney 2018; Program 2014; Nelson and Bradley 2003) (USDA-NASS 2018b; Shoup 2016).

Fungicides: Most plant diseases are caused by fungi, bacteria, or viruses. Planting resistant varieties is the most commonly used management method to control those plant pathogens that reduce soybean yields. In contrast, chemical pesticides are used less often, but among those used to treat soybean diseases, those used most frequently are fungicides.

Diseases that infect soybeans are caused by bacteria, fungi and viruses (Jardine 1997; Bradley 2018). The most serious soybean diseases include *Cercospora* foliar blight, purple seed stain, aerial blight, soybean rust, pod and stem blight, and anthracnose (Benedict 2011). Besides selecting varieties with resistance to the diseases prevalent in a growing region (Jardine 1997; Bradley 2018), growers plant sterilized disease-free seed (Jardine 1997), and use other best management practices (BMPs) such as rotating crops to prevent buildup of disease organisms in fields, and providing adequate nutrients and water for growth (Kandel 2019; Nelson 2011). Seed treatments with various chemicals such as fungicides, promote successful seed germination (Jardine 1997; Bradley 2018).

When other management measures fail to control diseases, soybean growers have chemical treatment options, but most are only effective on diseases caused by fungi (Benedict 2011). Fungicides most commonly applied to soybeans in 2017 were pyraclostrobin, fluxapyroxad, azoxystrobin, propiconazole, trifloxystrobin, and picoxystrobin (applied to 5%, 5%, 4%, 3%, 2%, and 1% of U.S. soybean acreage respectively (USDA-NASS 2018a).

Herbicides: Weed management for soybeans has been done primarily with herbicides since the mid-1960s, and will continue to be an important practice for the foreseeable future. One review of aggregate data for crop yield losses and herbicide use estimated a \$16 billion (20%) U.S. crop production loss in value if herbicides were not used (Gianessi and Reigner 2007), even if additional tillage and hand weeding labor replaced herbicides.

The herbicides most commonly applied to soybeans in 2017 are listed in Table 5. Growers consider several factors when selecting a weed control program including cost, potential adverse effects on the crop, residual effects that limit following crop choices in a rotation cycle, and control efficacy. All herbicides and other pesticides can only be applied legally in strict accordance with their EPA registration labels. What is allowed by the label is a primary consideration for growers.

Management of Weed Herbicide Resistance

There are currently 262 species of weeds (152 dicots and 110 monocots) and 514 unique cases of HR weeds reported from throughout the world. Weeds have evolved resistance to 23 of the 26 known herbicide modes of action (MOAs) and to 167 different herbicides. HR weeds have been reported in 93 crops in 70 countries (Heap 2021).

For many years, growers were able to effectively control or suppress virtually all weeds in soybean with glyphosate. A number of weed species eventually developed resistance to glyphosate however, and the number of acres infested with resistant biotypes has been increasing. The Weed Science Society of America (WSSA) currently lists 50 weed species worldwide that are resistant to one or more herbicide MOAs and are associated with soybean cultivation. Seventeen are found in the United States (Heap 2021).

Table 5. Five Most Used Herbicides to Treat U.S. Soybean Acreage in 2017.

Herbicide (Active Ingredient: a.i.)	Percent of Acreage Treated	Average Application Rate (lbs a.i./acre)	Total Applied (million lbs a.i.)
Glyphosate isopropylamine salt	46	1.145*	44.2*
Glyphosate potassium salt	30	1.590*	40.3*
Sulfentrazone	22	0.179	3.3
Fomesafen sodium	19	0.240*	3.9*
Metribuzin	18	0.256	3.7
Source: (USDA-NASS 2018a)		*Expressed as acid equivalent	

Herbicide usage trends since the adoption of biotech crops are the subject of much interest, research and debate. The initial assessments indicated a decline in herbicide use in the early years of HR crop production (Carpenter et al. 2002). Some argue that this was followed by an increase in the volume of herbicide usage as the HR varieties became increasingly popular (Benbrook 2009).

Others report a continuing decline in herbicide use with the adoption of biotech crops (Fernandez-Cornejo and Caswell 2006), or little or no change in the amount of herbicide active ingredients applied to soybeans (Brookes and Barfoot 2010). The contradictory findings have been attributed to the different measurement approaches used by researchers, the way data were adjusted for the effects of factors affecting pesticide use such as weather or cropping patterns, and the statistical procedures used to analyze the data (NRC 2010).

Herbicide applications and other weed control practices exert selection pressures on weed communities. This can change weed community structure (i.e., change the types of weeds present) that favors weeds that don't respond to the herbicide or other control methods being used (Owen 2008). Herbicide resistance is primarily caused by natural selection for individuals with HR traits within a population by repeated sub-lethal exposure to one or a limited number of herbicides (Duke 2005; Durgan and Gunsolus 2003). Dispersal of the HR weed seed and further selection by herbicide treatment contributes to the rapid spread of HR weeds to new locations.

Both the increased selection pressure resulting from the exclusive or extensive and widespread use of glyphosate herbicides on GR crops without other types of herbicides, and changes in weed management practices (i.e., conservation tillage or no-till) have resulted in weed population shifts and increased glyphosate resistance among some weed populations (Duke 2005; Owen 2008).

GR crops, themselves, do not influence weeds any more than non-transgenic crops. HR weed biotypes result from natural selection for those biotypes under the current weed control methods, rather than gene transfer from the crop to the weed. It is the prolonged use of the same weed

control tactics by growers that causes long-term selection pressures that change weed communities and contribute to the induction of HR weeds (Owen 2008). More details about HR weeds are reviewed in the Plant Communities section of this chapter.

The management of GR weeds has become a substantial challenge for U.S. agriculture especially soybean production because good alternative options are limited (Owen 2011a; Powles 2008a; Powles 2008b). Some strategies proposed to manage GR weeds (Boerboom 1999; Beckie 2006; Sammons et al. 2007; Frisvold, Hurley, and Mitchell 2009), include:

- Rotating different herbicides that have different modes of action
- Site specific herbicide applications
- Use of highest labeled application rate allowed by the label (prevents sublethal dosing)
- Crop rotation
- Use of tillage for supplemental weed control
- Cleaning equipment between fields
- Controlling weed escapes
- Controlling weeds early
- Scouting for weeds before and after herbicide applications

Volunteer soybeans are not a widespread management problem, and occur most often in parts of the southern United States where winters are mild. In production systems where soybeans are rotated with other crops, soybeans can be a volunteer weed (Owen and Zelaya 2005a), but this is not considered difficult to manage because soybean seeds rarely remain viable the following season and any interference they may pose to subsequent crops is minimal. Furthermore, herbicides, such as atrazine and metolachlor that are usually used for weed control in corn, the crop most often rotated with soybeans, are also effective at controlling volunteer soybean (Owen and Zelaya 2005a). Conversely, volunteer GR corn in soybean is a greater concern (Owen and Zelaya 2005a). With the widespread use of both GR corn and soybean, glyphosate is no longer effective to control corn volunteers. Growers must often include graminicides (herbicides that control weedy grasses) as part of their weed management strategy (Owen and Zelaya 2005a).

Soybean Yield Increases

Because of recent trends in farm production and land area, soybean growers will have the future challenge of expanding agricultural output by raising productivity on a stable or reduced land area (OECD-FAO 2020). This implies that most of the projected expansion in soybean production is expected to come from increasing yield, not increasing crop acreage (OECD-FAO 2020).

Egli (2008) reviewed historical trends in U.S. corn and soybean back to the first available data in 1924 to document soybean yield increase. Improved management practices such as mechanization, narrow-row planting, earlier planting, adoption of conservation tillage, increased weed control, and decreased harvest loss contributed to increased yields (Egli 2008). Agricultural biotechnology has further enhanced crop yields. For example, De Bruin and Pedersen (2009) estimate SCN-resistant soybean varieties increased yields by 17--19% compared to non-resistant varieties.

USDA projections through 2021/2022 show an average annual rate of increased average yields of 0.45 bushels per acre for the period 2012/2013 to 2021/2022, which results in an average U.S. yield of 46.05 bushels per acre for the period (USDA-OCE 2018). While USDA projects

increasing yields, the projected rate of increase is lower than the past rate. Current and future factors that negatively affect yield increases are the expansion of soybean production into northern and western parts of the country, where yields are typically lower than in the Midwest.

Soybean Seed Production

Growers may plant certified soybean seed, uncertified seed, and soybean seed grown and stored on individual farms (Oplinger and Amberson 1986). Seed production differs from grain production because of additional biological, technical, and quality control factors required to maintain varietal purity. Genetic purity in the production of commercial soybean seed is regulated through a system of seed certification which ensures the desired traits in that particular seed remain within purity standards (Bradford 2006).

The production and certification of foundation, registered, certified, or quality assurance seeds are administered by state and regional crop improvement associations, several of which are chartered under the laws of the state(s) they serve (e.g., see Virginia Crop Improvement Association No Date; SSCA No Date-a; Illinois Crop Improvement Association 2013; Mississippi Crop Improvement Association 2008, 2015; Illinois Crop Improvement Association 2019). These agencies certify varietal purity and identity, while issues concerning germination and mechanical purity are governed under state and federal seed laws.

Seed quality includes a variety of attributes, including genetic purity, vigor, weed seed content, seed borne diseases, and the presence of foreign material such as dirt or chaff (Bradford 2006). The genetic purity of the seed must be maintained to maximize the value of the new variety (Sundstrom et al. 2002). Some general examples of seed production practices include certification of origin and class, documentation of field cropping history, isolation from weeds and soybean grain crops, decontamination of cultivation, transportation, and storage equipment, and inspection and laboratory analysis of harvested seeds from approved fields (Virginia Crop Improvement Association No Date; South Dakota Crop Improvement Association 2011; SSCA No Date-a; Illinois Crop Improvement Association 2013; Mississippi Crop Improvement Association 2008). There are also crop specific field, inspection, isolation, and harvested seed purity standards (e.g., percentage of pure seed, inert matter, weed seeds, other crop seeds, other variety seeds, and germination) (South Dakota Crop Improvement Association 2011; Virginia Crop Improvement Association 2013; SSCA No Date-b).

The U.S. Federal Seed Act of 1939 recognizes seed certification and official certifying agencies. Implementing regulations further recognize land history, field isolation, and varietal purity standards for seed. States have developed laws to regulate the quality of seed available to farmers (Bradford 2006). Most of the laws are similar in nature and have general guidelines for providing information on the label for the following:

- Commonly accepted names of agricultural seeds
- Approximate total percentage by weight of purity
- Approximate total percentage of weight of weed seeds
- Name and approximate number per pound of each kind of noxious weed seeds
- Approximate percentage of germination of the seed
- Month and year the seed was tested

Various seed associations have standards to help maintain the quality of soybean seed. The AOSCA (AOSCA 2012c, 2013, 2019) defines the classes of seed as follows:

- *Breeder*: developed and used by plant breeders
- *Foundation*: progeny of Breeder or Foundation maintained to preserve specific genetic identity and purity
- *Registered*: progeny of Breeder or Foundation maintained for satisfactory genetic identity and purity
- *Certified*: progeny of Breeder, Foundation, or Registered handled to maintain satisfactory genetic identity and purity

Seed certification systems differ from Identity Preservation (IP) systems for certain agricultural commodities. IP refers to a system of production, handling, and marketing practices used to maintain the integrity and purity of crop products throughout the food supply chain (Sundstrom et al. 2002). IP systems are used to meet the demands for specialized grain products, including those from crops with output-specific traits (e.g., high oleic oil), without specific traits or attributes (e.g., conventional crops), grown under specific production methods (e.g., organic crops), and requiring rigorous safeguards and confinements practices (e.g., pharmaceutical and industrial crops) (Elbehri 2007).

Soybean is self-pollinated and propagated commercially from seed (OECD 2000; Hoeft et al. 2000). In the United States, there are no *Glycine* spp. found outside of cultivation, so the potential for outcrossing is minimal (OECD 2000). Minimum Land, Isolation, Field, and Seed Standards (7 CFR part 201.76) specify isolation distances for the production of Foundation, Registered and Certified soybean seeds to prevent mechanical mixing from potential contaminating sources.

4.1.3 Agronomic Practices for Organic Soybean Production

In the United States, only products produced using specific methods and certified under the USDA's Agricultural Marketing Service National Organic Program (NOP) can be labeled as "USDA Organic" (USDA-AMS 2008, 2020). Organic certification is a process for validation of production practices, not certification of the end product. USDA organic certification requires that specific production methods be documented by the producer and certified by an independent auditor.

An accredited organic certifying auditor conducts an annual review of a producer's organic system plan and practices documented in records maintained on site. The auditor also makes on-site inspections to confirm accuracy of recordkeeping. Organic growers must maintain records to show that production and handling procedures comply with USDA organic standards. The NOP regulations (7 CFR § 205.2) specifically exclude certain methods that cannot be used for the production of products labeled "USDA Organic."

Common practices organic growers may use to exclude biotech products include planting only organic seed, staggering planting earlier or later than neighboring farmers who may be using biotech crops, so that the crops will flower at different times to establish adequate isolation between organic fields and neighboring fields, where non-organic crops are grown to minimize the possibility of cross-pollination (NCAT 2012).

Although the NOP standards prohibit the use of excluded methods, they do not require testing of inputs or products for the presence of excluded methods. The presence of a detectable residue of a product of excluded methods alone does not necessarily constitute a violation of the NOP standards (USDA-AMS 2008, 2020). The current NOP regulations do not specify an acceptable

threshold level for the adventitious presence of biotech materials in an organic-labeled product. The unintentional presence of the products of excluded methods will not affect the status of an organic product or operation when the operation has not used excluded methods and has taken reasonable steps to avoid contact with the products of excluded methods as detailed in their approved organic system plan” (USDA-AMS 2008, 2020; Ronald and Fouche 2006).

4.2 PHYSICAL ENVIRONMENT

Components of the physical environment affected by soybean production in the United States are reviewed in this section. These include soil, water, and air quality.

4.2.1 Soil Quality

Soil consists of solids (minerals and organic matter), liquids, and gases. Inorganic and organic matter harbor a wide variety of fungi, bacteria, and arthropods, as well as the growth medium for terrestrial plant life (USDA-NRCS 2004). Soil is characterized by its layers (USDA-NRCS 1999b), and is further distinguished by its ability to support rooted plants in a natural environment. Soil establishes the capacity of a site’s biomass vigor and production in terms of air, water, temperature moderation, protection from toxins, and nutrient availability. Soils also determine a site’s susceptibility to erosion by wind and water, and its flood attenuation capacity.

Important soil properties include temperature, pH, soluble salts, amount of organic matter, the carbon-nitrogen ratio, numbers of microorganisms and soil fauna, and all vary seasonally, and over extended periods of time (USDA-NRCS 1999b). Soil texture and organic matter levels directly influence its shear strength, nutrient holding capacity, and permeability. Soils are differentiated based on characteristics such as particle size, texture, and organic matter content (USDA-NRCS 2010b).

Soybeans are normally grown in agricultural fields managed for crop production and are best suited to fertile, well-drained medium-textured loam soils, but can be produced in a wide range of soil types (Berglund and Helms 2003a; NSRL No Date). Soybeans need a variety of macronutrients, such as nitrogen, phosphorus, potassium, calcium, magnesium, and sulfur, at various levels. They also require smaller amounts of micronutrients such as iron, zinc, copper, boron, manganese, molybdenum, cobalt, and chlorine. These micronutrients may be deficient in poor, weathered soils, sandy soils, alkaline soils, or soils excessively high in organic matter.

As with proper nutrient levels, soil pH is critical for soybean development. Soybeans grow best in soil that is slightly acidic (pH range: 6.0 - 7.0) (NSRL No Date). Soil with a pH that is too high (7.3 or greater) negatively affects yield (NSRL No Date; Cox et al. 2003). Similarly, soils that are high in clay and low in humus may impede plant emergence and development. Soils with some clay content may increase moisture availability during periods of low precipitation (Cox et al. 2003).

Soybean yield is highly dependent upon soil and climatic conditions. In the United States, the soil and climatic requirements for growing soybean are very similar to corn. The soils and climate in the Midwest, portions of the Great Plains and eastern regions of the United States provide sufficient water under normal climatic conditions to produce a soybean crop. Soil texture and structure are key components in determining water availability in soils. Medium-textured

soils hold more water, allowing soybean roots to penetrate deeper in medium-textured soils than in clay soils (Berglund and Helms 2003a; Cox et al. 2003).

Land management practices for soybean cultivation can affect soil quality. While practices such as tillage, fertilization, the use of pesticides and other management tools can improve soil health, they can also cause substantial damage if not properly used. Several concerns relating to agricultural practices include increased erosion, soil compaction, degradation of soil structure, nutrient loss, increased salinity, change in pH, and reduced biological activity (USDA-NRCS 2001).

Conventional and conservation tillage may be used for the cultivation of soybean. Reducing excessive tillage through practices such as conservation tillage minimizes the loss of organic matter and protects the soil surface by leaving plant residue on the surface. Management of crop residue is one of the most effective conservation methods to reduce wind and water erosion. It also benefits air and water quality and wildlife (USDA-NRCS 2006a). Residue management that uses intensive tillage and leaves low amounts of crop residue on the surface results in greater losses of soil organic matter (SOM).

Intensive tillage turns the soil over and buries the majority of the residue, stimulating microbial activity and increasing the rate of residue breakdown (USDA-NRCS 1996). The residues left after conservation tillage increase organic matter and improve infiltration, soil stability and structure, and soil microorganism habitat (Fawcett and Caruana 2001; USDA-NRCS 2006b)). Organic matter is probably the most vital component in maintaining quality soil. It is instrumental in maintaining soil stability and structure, reduces the potential for erosion, provides energy for microorganisms, improves infiltration and water holding capacity, and is important in nutrient cycling, cation exchange⁷ capacity, and the degradation of pesticides (USDA-NRCS 1996).

The residue left from conservation tillage practices increases SOM in the top three inches of the soil and protects the surface from erosion, while maintaining water-conducting pores. Soil aggregates in conservation tillage systems are more stable than that of conventional tillage because the products of SOM decomposition, and the presence of fungal hyphae (filamentous structures that compose the main growth) and soil bacteria bind aggregates and soil particles together (USDA-NRCS 1996). Although soil erosion rates are dependent on numerous local conditions such as soil texture and crops grown, a comparison of 39 studies contrasting conventional and no-till practices showed that, on average, no-till practices reduce erosion by a factor of 488 times compared to conventional tillage (Montgomery 2007).

From 1982 through 2003, erosion on U.S. cropland dropped from 3.1 billion tons per year to 1.7 billion tons per year (USDA-NRCS 2006a). This can partially be attributed to the increased effectiveness of weed control through the use of herbicides and the corresponding reduction in the need for mechanical weed control (Carpenter et al. 2002). Conservation tillage also minimizes soil compaction because it reduces, but does not eliminate the number of times a field is tilled.

⁷Cation Exchange Capacity is the ability of soil anions (negatively charged clay, organic matter and inorganic minerals such as phosphate, sulfate, and nitrate) to adsorb and store soil cation nutrients (positively charged ions such as potassium, calcium, and ammonium).

Other methods to improve soil quality include careful management of fertilizers and pesticides, the use of cover crops to increase plant diversity and limit the time soil is exposed to wind and rain, and the use of buffer strips, contour strips, wind breaks, crop rotations, and varying tillage practices (USDA-NRCS 2006b). Planting cover crops is another management practice that has become recognized as a way to increase plant diversity, reduce compaction, suppress disease, control weeds, and enhance soil nutrients (NWF 2012; USDA-NASS 2012c; Lee et al. No Date; SARE 2012; Corn and Soybean Digest 2013; MDA 2012; USDA-NRCS 2011a; Hoorman et al. 2009; University of Georgia Soybean Team 2019) in addition to suppressing erosion by limiting the time soil is exposed to wind and rain effects.

Although conservation tillage benefits soil quality in several ways, it can also have negative effects. For example, under no-till practices, soil compaction may become a problem because tillage disrupts compacted areas (USDA-NRCS 1996). Another concern is that not all soils (such as wet and heavy clay soils) are suited for no-till. No-till practices may also increase pest abundance compared to conventional tillage (NRC 2010).

Numerous kinds of organisms that live in soils, ranging from microorganisms to larger macroinvertebrates, such as worms and insects, affect soil quality. The microorganisms that make up the soil community include bacteria, fungi, protozoa, and nematodes. Decomposers, such as bacteria, and saprophytic fungi, degrade plant and animal remains, organic materials, and some pesticides (USDA-NRCS 2004). Other organisms, such as protozoans, mites and nematodes, consume the decomposer microbes and release macro- and micronutrients, making them available for plant uptake.

Mutualists are another important group of soil microorganisms. These are the mycorrhizal fungi, nitrogen-fixing bacteria, and some free-living microbes that have coevolved with plants, and supply nutrients to and obtain food from their plant hosts (USDA-NRCS 2004). The bacterium, *B. japonicum*, associated with soybeans fixes nitrogen in root nodules on the plants (Franzen 1999). If a field has not been planted recently with soybeans (3-5 years), either the seed or seed zone must be inoculated with *B. japonicum* prior to soybean planting (Elmore 1984; Pedersen 2007).

Pesticide use has the potential to affect soil quality because it can impact the soil microbial community (see the section on Microorganisms in this chapter for more details). The length of persistence of herbicides in the environment is dependent on the concentration and rate of degradation by biotic and abiotic processes (Carpenter et al. 2002). Persistence is measured by the half-life, which equates to the length of time needed for the herbicide residue to degrade to half of its original concentration. The degradation of pesticides may be dependent on mineralization by microbes in soil, photodegradation in water, and leaching (US-EPA 2005). In soil, pesticide persistence may be strongly influenced by moisture, temperature, organic matter content and pH (FAO 1997; Senseman 2007).

4.2.2 Water Resources

Surface water quality is determined by the natural, physical, and chemical properties of the land that surrounds the water body. Topography, soil type, vegetative cover, minerals, and climate all influence water quality. Surface runoff is affected by meteorological factors such as rainfall intensity and duration, and physical factors such as vegetation, soil type, and topography. When land use affects one or more of these natural physical characteristics of the land, water quality is

almost always impacted to some extent. These impacts may be positive or negative, depending on the type, duration and extent of land use.

Agricultural practices have the potential to substantively impact water quality because of the vast amount of acreage devoted to farming nationwide and the physical and chemical demands that agricultural use has on the land. The most common types of agricultural pollutants include excess sediment, fertilizers, animal manure, and pesticides. Agricultural non-point source pollution is the leading source of impacts on rivers and lakes, the third largest source of impairment to estuaries, and a major source of impairment to groundwater and wetlands (USDA-NRCS 2011b).

The principal law regulating pollution of the nation's water resources is the Federal Water Pollution Control Act of 1972, which is commonly referred to as the Clean Water Act (CWA). The EPA sets water quality standards, permitting requirements, and monitors water quality. The EPA sets the standards for water pollution abatement for all waters of the United States under CWA programs, but in most cases, gives qualified states the authority to issue and enforce permits. The CWA provides the authority to establish water quality standards, control discharges into surface and subsurface waters (including groundwater), develop waste treatment management plans and practices, and issue permits for discharges under the National Pollutant Discharge Elimination System. Section 303(d) of the CWA established a process for states to identify those waters within its boundaries that do not meet minimum water quality standards. Waters that do not meet clean water standards are classified under the CWA as "Impaired Waters." Impaired Waters cannot support one or more designated uses (e.g., swimming, propagation of aquatic life, drinking, and agricultural or industrial supply). Common pollutants evaluated include sediment, chemicals, fuels, biological contaminants and pathogens, and characteristics such as oxygen availability, water temperature, and water clarity.

Once a water body or stream segment is listed as impaired, the state must complete a plan to address the issue causing the impairment. States then develop total maximum daily loads (TMDLs) for priority waters that identify the amount of a specific pollutant from various sources that may be discharged to a water body, but still ensure that water quality standards are met for that body of water. Completion of the plan is generally all that is required to remove the stream segment from the 303(d) impaired water list and does not mean that water quality has changed. Once the TMDL is completed and approved by EPA (US-EPA 2012), the stream segment is placed on the 305(b) list of impaired streams with a completed TMDL.

Groundwater is water that flows underground and is stored in natural geologic formations called aquifers. It is ecologically important because it sustains ecosystems by releasing a constant supply of water into wetlands and contributes a sizeable amount of flow to permanent streams and rivers. Currently, the largest use of groundwater in the United States is irrigation, representing approximately 67.2% of all the groundwater pumped (NGWA 2017).

Approximately 9% of the planted acres of soybeans in the United States are irrigated (USDA-NASS 2010; USDA-ERS 2011b; USDA-NASS 2011). A majority (approximately 73%) of U.S. irrigated soybean farms occur in the Missouri and Lower Mississippi Water Resource Regions with soybean farms in the states of Nebraska, Arkansas, Mississippi, Missouri, and Kansas accounting for 85% of all irrigated soybean acres (USDA-NASS 2010).

In the United States, approximately 47% of the population depends on groundwater for its drinking water supply (NGWA 2017). Drinking water is protected under the Safe Drinking

Water Act of 1974 (SDWA) (Public Law 93-523, 42 U.S.C. 300 *et seq.*). SDWA and subsequent amendments authorize the EPA to set national health-based standards for drinking water from source water to the tap to protect against both naturally-occurring and man-made contaminants that may be found. In an effort to protect source water, the Sole Source Aquifer (SSA) Program was developed to protect drinking water supplies in areas where there are few or no alternative sources to the groundwater resource for drinking water and other needs. EPA defines an SSA as an aquifer that supplies at least 50% of the drinking water consumed in the area overlying the aquifer. There are 77 designated SSAs in the United States and its territories (US-EPA 2011c). The designation protects an area's groundwater resource by requiring EPA to review certain proposed projects receiving federal funds or approval within the designated area to ensure that they do not endanger the water source.

Use of pesticides can introduce chemicals to water through spray drift, cleaning of pesticide equipment, soil erosion, and filtration through soil to groundwater. Solubility (whether it readily dissolves in water), its adsorptive qualities (how tightly it binds to clay and humus particles in the soil), and its degradation (how fast it breaks down into harmless components) are some of the factors that influence the degree to which herbicide residue can infiltrate ground or surface water.

Planting HR soybean varieties enables growers to treat post-emergent soybeans for weed control, reducing or eliminating the need for cultivation. Approximately 94% of the soybean acreage in the United States is planted with HR soybean varieties (USDA-ERS 2017). Therefore growers who plant HR soybean varieties are more likely to use conservation tillage and no-till practices than growers of non-HR soybeans (Givens et al. 2009; Dill, CaJacob, and Padgett 2008). This trend has resulted in reduced surface water runoff and soil erosion (Locke, Zablotowicz, and Reddy 2008). Reduced tillage agricultural practices result in improved soil quality having high organic material that binds nutrients within the soil (see the section on Soil Quality in this chapter for more details). An increased amount of plant residue on the soil surface reduces the effects of pesticide usage on water resources by forming a physical barrier to erosion and runoff, allowing more time for absorption into the soil, and slowing soil moisture evaporation (Locke, Zablotowicz, and Reddy 2008). The use of HR soybean varieties has also promoted a shift to herbicides that have lower environmental impact, such as glyphosate (Fernandez-Cornejo and McBride 2002).

Nutrient applications to soybeans primarily include nitrogen, phosphorus (phosphate), potassium (potash), calcium, and sulfur, with other micronutrient supplements such as zinc, iron, and magnesium applied as needed. Runoff from cropland areas receiving manure or fertilizers contributes to increased phosphorous and nitrogen delivery to streams and lakes. This causes eutrophication⁸ primarily from phosphorus, which is the limiting nutrient in freshwater ecosystems. Ammonium runoff into surface waters can result in the poisoning of aquatic organisms. Nitrate in runoff from fields is carried into rivers and lakes. Elevated nitrate levels in the Gulf of Mexico contribute to the hypoxia zone, an area depleted of oxygen and marine life.

⁸Eutrophication is the process by which a body of water becomes enriched in dissolved nutrients (such as phosphates) that stimulate the growth of aquatic plant life usually resulting in the depletion of dissolved oxygen.

Conservation tillage and other management practices are used to trap and control sediment and nutrient runoff. Water quality conservation practices benefit agricultural producers by lowering input costs and enhancing the productivity of working lands.

4.2.3 Air Quality

The Clean Air Act (CAA) requires the EPA to set National Ambient Air Quality Standards (NAAQS) for certain common and widespread pollutants. The NAAQS, developed by the EPA to protect public health, sets limits for six criteria pollutants: ozone (O₃), nitrogen dioxide (NO₂), carbon monoxide (CO), sulfur dioxide (SO₂), lead (Pb), and inhalable particulates (particulates greater than 2.5 micrometers and less than 10 micrometers in diameter are defined as coarse particulate matter [PM₁₀], and those less than 2.5 micrometers in diameter are classified as fine particulate matter [PM_{2.5}]). The CAA requires states to achieve and maintain the NAAQS within their jurisdiction.

Each state may adopt requirements stricter than those of the national standard and each is also required by EPA to prepare an implementation plan with strategies to achieve and maintain the national standard for air quality within the state. Areas that violate air quality standards are designated as non-attainment areas for the criteria pollutant(s), whereas areas that comply with air quality standards are designated as attainment areas.

Primary sources of emissions associated with crop production include exhaust from motorized equipment such as tractors and irrigation equipment, soil particulates from tillage and wind-induced erosion, particulates from burning of fields, and aerosols from herbicide and pesticide applications.

Because they reduce the need to till for weed control, HR soybeans have promoted the use of no-till or conservation tillage for soybean production. Decreased tillage reduces the use of emission-producing equipment (Table 6) and also causes less dust from particulates, potentially lowering rates of wind erosion, which benefits air quality (Towery and Werblow 2010).

Table 6. Examples of Estimated Annual Fuel Used for Different Tillage Methods.

Estimated Use/1,000 Acres of Soybeans in Urbana, Illinois	Tillage Method			
	Conventional	Mulch-till	Ridge-till	No-till
Total fuel used*	5,239	4,369	3,460	2,330
Estimated fuel saved compared to that used for conventional tillage	--	870	1,779	2,909
Percent estimated savings	--	17%	34%	56%
*Diesel fuel in gallons				

Source: USDA-NRCS (2013a)

Volatilization of fertilizers, herbicides and pesticides from soil and plant surfaces introduces these chemicals into the air. One study in the Chesapeake Bay region (USDA-ARS, 2011) determined that volatilization is highly dependent upon exposure of disturbed unconsolidated soils and variability in measured compound levels is correlated with temperature and wind conditions. Another study of volatilization of certain herbicides after application to fields found moisture in dew and soils in higher temperature regimes significantly increases volatilization rates (USDA-ARS, 2011).

Prescribed burning is a land treatment used under controlled conditions to accomplish resource management objectives. Open combustion produces particles of widely ranging size, depending to some extent on the rate of energy release of the fire (US-EPA 2011a). The extent to which agricultural and other prescribed burning may occur is regulated by individual state implementation plans to achieve compliance with the NAAQS. Prescribed burning of fields would likely occur only as a pre-planting option for soybean production based on individual farm characteristics.

Pesticide and herbicide spraying may impact air quality from drift and diffusion. Drift is defined by EPA as “the movement of pesticide through air at the time of application or soon thereafter, to any site other than that intended for application” (US-EPA 2000). Diffusion is gaseous transformation into the atmosphere (FOCUS 2008). Factors affecting drift and diffusion include application equipment and method, weather conditions, topography, and the type of crop being sprayed (US-EPA 2000).

Other conservation practices, as required by USDA to qualify for crop insurance and beneficial federal loans and programs, effectively reduce crop production impacts to air quality through the use of windbreaks, shelterbelts, reduced tillage, and cover crops that promote soil protection on highly erodible lands.

4.3 BIOLOGICAL RESOURCES

Biological resources include animal, plant and microbiological organisms, and their assemblages that form living community structures in the environment.

4.3.1 Animal Communities

Agriculture dominates human uses of land (Robertson and Swinton 2005). In 2011, 917 million acres (approximately 47%) of the conterminous 48 states were dedicated to farming, including: crop production, pasture, rangeland, Conservation Reserve Program and Wetlands Reserve Program lands, or other government program uses. About 10% (88.8 million) of farmed acreage was for soybean production (Senseman 2007; USDA-NASS 2012b).

Intensely cultivated lands, such as those for commercial soybeans, provide less suitable habitat for wildlife than natural areas. A wide array of wildlife species occurs within the 31 major soybean-producing U.S. states. The types and numbers of animal species found in and near soybean fields is less diverse as compared to unmanaged natural habitats. How these lands are maintained influences the function and integrity of the wildlife populations they support and the ecosystem services they provide. Animal communities considered in this EA include wildlife species and their habitats. Wildlife refers to both native and introduced species of mammals, birds, amphibians, reptiles, invertebrates, and fin and shellfish.

Birds and Mammals

Soybean fields can provide both food and cover for a variety of birds, and small and large mammals. During the spring and summer months, soybean fields provide browse for rabbits, deer, rodents, other mammals, and birds such as upland gamebirds (Palmer, Bromley, and Anderson No Date). During the winter months, leftover and unharvested soybeans provide a food-source for wildlife; however, soybeans are poorly suited for meeting nutrient needs of wildlife, such as waterfowl, which require a high-energy diet (Krapu, Brandt, and Cox 2004).

A shift from conventional agricultural practices to conservation tillage and no-till practices has occurred on farms planting HR soybean varieties (Givens et al. 2009; Dill, CaJacob, and Padgett 2008). This increased use of conservation tillage practices has benefitted wildlife through improved water quality, availability of waste grain, retention of cover in fields, and increased populations of invertebrates (Sharpe 2010; Brady 2007). Conservation tillage practices that leave greater amounts of crop residue serve to increase the diversity and density of birds and mammals (USDA-NRCS 1999a).

Invertebrates

Increased residue from the shift to conservation tillage also provides habitat for insects and other arthropods, consequently increasing this food source for insect predators. Insects are important during the spring and summer brood rearing season for many upland game birds and other birds, as they provide a protein-rich diet source to fast growing young, and a nutrient-rich diet for migratory birds (USDA-NRCS 2016).

Insects, nematodes and other invertebrates can be beneficial to soybean production by cycling nutrients and preying on plant pests. Conversely, there are some invertebrates that are detrimental to soybean crops, including: soybean cyst nematode (*Heterodera glycines*); root knot nematodes (*Meloidogyne* spp.); bean leaf beetle (*Cerotoma trifurcata*); beet armyworm (*Spodoptera exigua*); blister beetle (*Epicauta* spp.); soybean podworm (*Helicoverpa zea*); shorthorned grasshoppers (*Acrididae* spp.); green cloverworm (*Hypena scabra*); seed corn beetle (*Stenolophus lecontei*); seed corn maggot (*Delia platura*); soybean aphid (*Aphis glycines*); soybean looper (*Pseudoplusia includens*); soybean stem borer (*Dectes texanus*); spider mites (*Tetranychus urticae*); stink bugs (green [*Acrosternum hiliare*] and brown [*Euschistus* spp.]); and velvetbean caterpillar (*Anticarsia gemmatilis*) (Palmer 2012; Whitworth 2016; Whitworth, Michaud, and Davis 2012).

While insects and nematodes are considered less problematic than weeds in U.S. soybean production, injury can impact yield, plant maturity, and seed quality. Consequently, these pests are managed during the growth and development of soybean to enhance soybean yield (Higley and Boethel 1994; Aref and Pike 1998).

Under FIFRA, all pesticides, including herbicides, insecticides and nematicides, that are sold or distributed in the United States must be registered by the EPA (US-EPA 2005). Registration decisions are based on scientific studies that assess the chemical's potential toxicity and environmental impact. To be registered, a pesticide must be able to be used without posing unreasonable risks to the environment, including wildlife.

All pesticides registered prior to November 1, 1984 must also be reregistered to ensure that they meet the current, more stringent standards. EPA must find during its registration process (US-EPA, 2018) that a pesticide does not cause unreasonable adverse effects to the environment if

used in accordance with the EPA-approved label instructions. Growers must adhere to EPA label use restrictions for herbicides and pesticides. These measures help to minimize potential impacts of their use on non-target wildlife species.

4.3.2 Plant Communities

Although U.S. soybeans are grown commercially in 31 states, most (95%) are produced in 18 states in the Midwest and Southeast (Figure 1), encompassing a wide range of physiographic regions, ecosystems, and climatic zones (USDA-NASS 2019). The types of vegetation, including the variety of weeds within and adjacent to soybean fields can vary greatly, depending on their geographic location.

In general, plant diversity surrounding crop fields is an important component of a sustainable agricultural system (Scherr and McNeely 2008; CBD 2020). Hedgerows, woodlands, fields, and other assemblages of plants in habitats surrounding crop fields serve as important reservoirs for a range of organisms—both beneficials and pests.

Soybean fields and surrounding edges for example, are habitats for weeds that adversely impact crop production directly through interference and resource competition. They serve as seed sources for weeds that invade soybean fields and also support insect pests and plant pathogens that impact soybeans. However, both the weedy and non-weedy plant species in habitats around fields provide valuable ecosystem services. Examples include habitat for pollen and nectar resources, and harborage for beneficial arthropods like biological control agents of soybean pests (e.g., lady beetles, spiders, and parasitic wasps) (Nichols and Altieri 2012; Scherr and McNeely 2008). Although soybeans are self-pollinated, pollen and nectar resources are indirectly important to soybeans by supporting some of the plant and animal species beneficial to soybean production. Surrounding plant communities for example can help regulate run-off, reduce soil erosion, and improve water quality, so when effectively managed, they provide benefits for crop production (Altieri and Letourneau 1982; Nichols and Altieri 2012).

Non-crop vegetation in soybean fields is limited by the extensive cultivation and weed control programs practiced by soybean producers. Plant communities bordering soybean fields can range from forests and woodlands to grasslands, aquatic habitats, and residential areas. Adjacent crops frequently include other soybean varieties, corn, cotton, or other field crops.

Weeds are classified as annuals, biennials, or perennials. Annuals and biennials are plants that complete their lifecycle within one year or two years respectively. Perennials are plants that live for more than two years. Weeds are also classified as broadleaf (dicots) or grass (monocots). Weeds can reproduce by seeds, rhizomes (underground creeping stems), or other underground parts. Annual grass and broadleaf weeds are considered the most common weed problems in soybeans (Krausz et al. 2001). However, with increased rates of conservation tillage, increases in perennial, biennial, and winter annual weed species are being observed (Durgan and Gunsolus 2003; Green and Martin 1996). Winter perennials are particularly competitive and difficult to control, as these weeds re-grow every year from rhizomes or root systems.

At least 55 weed species have been identified as commonly occurring in soybean production. Among the most common are: common lambsquarter (*Chenopodium album*), morning glory species (*Ipomoea* spp.), velvetleaf (*Abutilon theophrasti*), pigweed, (*Amaranthus* spp.), common cocklebur (*Xanthium strumarium*), foxtail (*Setaria* spp.), ragweed species (*Ambrosia* spp.),

crabgrass (*Digitaria* spp.), barynyard grass (*Echinochloa crus-galli*), Johnsongrass (*Sorghum halepense*), and thistles (*Cirsium* spp.) (Heap 2021; Shoup 2016).

An important concept in weed control is the seed bank, which is the reservoir of seeds that are in the soil and have the potential to germinate. Agricultural soils contain reservoirs of weed seeds ranging from 4,100 to 137,700 seeds per square meter of soil (May and Wilson 2006). Climate, soil characteristics, cultivation, crop selection, and weed management practices affect the seed bank composition and size (May and Wilson 2006).

Herbicide resistance is described by the Weed Science Society of America as the “inherited ability of a plant to survive and reproduce following exposure to a dose of herbicide normally lethal to the wild type” (WSSA 2013). The first reports of weed resistance to herbicides were in the 1950s (WSSA 2011). Individual plants within a species can exhibit different responses to the same herbicide rate. Initially, herbicide rates are set to work effectively on the majority of the weed population under normal growing conditions.

Genetic variability, including herbicide resistance, is exhibited naturally in normal weed populations, although at very low frequencies. When only a single herbicide is continuously relied upon as the primary means of weed control, the number of weeds resistant to that herbicide compared to those susceptible to the herbicide may change as the surviving resistant weeds reproduce (Figure 4). With no change in weed control strategies, in time, the weed population may be composed of more and more resistant weeds (WSSA 2011).

The adoption of GR crops, including soybeans, resulted in growers changing historical weed management strategies and relying on a single herbicide, glyphosate, to control weeds in the field (Owen et al. 2011; Weirich et al. 2011). Reliance on a single management technique for weed control resulted in the selection for weeds resistant to that technique (Owen et al. 2011; Weirich et al. 2011). The development of GR weeds has necessitated a diversification of weed management strategies by growers. GR weeds have forced growers to respond to the problem by applying herbicides with different modes of action, using tank mixes, increasing the frequency of applications, and returning to tillage and other cultivation techniques to physically control HR species, when a specific herbicide proves to be ineffective (CAST 2012). Integrated weed management programs that use herbicides from different groups, vary cropping systems, rotate crops, and that use mechanical as well as chemical weed control methods, delay or prevent the selection of HR weed populations (Sellers, Ferrell, and MacDonald 2011; Gunsolus 2002).

There are 495 unique HR biotypes (Table 7) with herbicide resistance in 23 HRAC herbicide groups (Heap 2021). Strategies for managing and avoiding the development of HR weed populations are well developed. In most instances, crop producers are advised to and use IWM practices to address HR weed concerns (e.g., (Shaw et al. 2011; Vencill et al. 2012; Wilson et al. 2009). IWM consists of integrating multiple practices, including mechanical, cultural, chemical, and biological weed control tactics, into a weed management program to optimize control of a particular weed problem. IWM can include specifically timed applications of herbicides, the use of herbicides with multiple modes of action, crop rotation, cover crops, various tillage practices, weed surveillance, and hand-pulling or hoeing (CLI 2015; Garrison et al. 2014; Owen 2011b).

Developers of HR varieties provide stewardship and IWM guidance to crop producers in accordance with and responsive to EPA requirements and WSSA recommendations. In 2017, EPA issued PR Notice 2017-2, *Guidance for Herbicide-Resistance Management, Labeling, Education, Training and Stewardship* (US-EPA 2017b). Through PRN 2017-2, EPA provides

HR weed management guidance for herbicides undergoing registration review and for label registration (i.e., new herbicide active ingredients, and new uses proposed for HR crops and other case-specific registration actions). To assist growers in managing weeds, individual states track the prevalent weeds in crops in their area and provide the most effective means for their management, typically through state agricultural extension services and regional IPM Centers (e.g., see (IPM 2015) that work with USDA to develop crop profiles and timelines.




Figure 4. Schematic Diagram of the Development of Herbicide Resistance.

Source: Adapted from (Tharayil-Santhakumar 2003)

Table 7. Summary of World-wide HR Weeds by Herbicide Group

Herbicide Group	HRAC Group	Herbicide Example	Dicots	Monocots	Total
ALS inhibitors	B	Chlorsulfuron	98	62	160
Photosystem II inhibitors	C1	Atrazine	51	23	74
ACCase inhibitors	A	Sethoxydim	0	48	48
EPSP synthase inhibitors	G	Glyphosate	22	20	42
Synthetic Auxins	O	2,4-D	30	8	38
PSI Electron Diverter	D	Paraquat	22	10	32
PSII inhibitors	C2	Chlorotoluron	11	18	29
PPO inhibitors	E	Oxyfluorfen	10	3	13
Microtubule inhibitors	K1	Trifluralin	2	10	12
Lipid Inhibitors	N	Triallate	0	10	10
Carotenoid biosynthesis (unknown target)	F3	Amitrole	1	5	6
Long chain fatty acid inhibitors	K3	Butachlor	0	5	5
PSII inhibitors (Nitriles)	C3	Bromoxynil	3	1	4
Carotenoid biosynthesis inhibitors	F1	Diflufenican	3	1	4
Glutamine synthase inhibitors	H	Glufosinate-ammonium	0	4	4
Cellulose inhibitors	L	Dichlobenil	0	3	3
Antimicrotubule mitotic disrupter	Z	Flamprop-methyl	0	3	3
HPPD inhibitors	F2	Isoxaflutole	2	0	2
DOXP inhibitors	F4	Clomazone	0	2	2
Mitosis inhibitors	K2	Propham	0	1	1
Unknown	Z	Endothall	0	1	1
Cell elongation inhibitors	Z	Difenzoquat	0	1	1
Total Number of Unique HR Biotypes:			256	239	495

*HRAC: Herbicide Resistance Action Committee

Source: (Heap 2021)

Runoff, spray drift, and volatilization of herbicides have the potential to impact non-target plant communities in proximity to fields where herbicides are used. The extent of damage to non-target plants exposed to herbicides is determined by their overall vigor, the amount and type of herbicide to which the plant is exposed, and the growing conditions after contact.

The total rainfall the first few days after herbicide applications can influence the amounts of leaching and runoff. However, it has been estimated that even after heavy rains, herbicide losses to runoff generally do not exceed 5-10% of the total applied. Planted vegetation, such as grass buffer strips, or crop residues can effectively reduce runoff. Volatilization typically occurs during application, but herbicide deposits on plants or soil can also volatilize (Tu, Hurd, and Randall 2001; USDA-FS 2009).

Spray drift is a concern for non-target effects on susceptible plants growing adjacent to fields when herbicides are used in the production of soybeans. This potential impact results from off-target herbicide drift (US-EPA 2010b). Damage from spray drift typically occurs at field edges or at shelterbelts (i.e., windbreaks), but highly volatile herbicides may drift farther into a field. The risk of off-target herbicide drift is recognized by the EPA, which has incorporated both equipment and management restrictions to address drift on the EPA-approved herbicide labels. These EPA label restrictions include requirements that the grower manage droplet size, spray boom height above the crop canopy, restrict applications to specified wind speeds and environmental conditions, and use drift control agents (US-EPA 2010b).

4.3.3 Gene Flow and Weediness

Gene flow to and from agroecosystems can occur on both spatial and temporal scales. In general, plant pollen is an important way that genes are transmitted. The rate and success of pollen-mediated gene flow is dependent on numerous external factors in addition to the donor and recipient plant. General external factors related to pollen-mediated gene flow include the presence, abundance, and distance of sexually-compatible plant species; overlap of flowering phenology between populations; the mechanism of pollination; the biology and amount of pollen produced; and weather conditions, including temperature, wind, and humidity (Zapiola et al. 2008). Seed-mediated gene flow also depends on many factors, including the presence, and magnitude of seed dormancy, contribution and participation in various dispersal pathways, and environmental conditions and events.

Soybean (*G. max*) is native to Asia. It does not have any feral or weedy relatives in the United States. Soybean is considered a highly self-pollinated species, propagated by seed (OECD 2000). Pollination typically takes place on the day the flower opens. The soybean flower stigma is receptive to pollen approximately 24 hours before anthesis (i.e., the period in which a flower is fully open and functional) and remains receptive for 48 hours after anthesis. Anthesis normally occurs in late morning, depending on the environmental conditions. The pollen usually remains viable for two to four hours, and no viable pollen can be detected by late afternoon. Natural or artificial cross-pollination can only take place during the short time when the pollen is viable, and soybean's reproductive characteristics (e.g., flower orientation that reduces its exposure to wind, internal anthers, and clumping and stickiness of the pollen) decrease the dispersion ability of pollen (Yoshimura 2011).

As a highly self-pollinated species, cross-pollination of soybean plants to adjacent plants of other soybean varieties occurs at a very low (0-6.3%) frequency (Ray et al. 2003; USDA-APHIS 2011; Caviness 1966; Yoshimura, Matsuo, and Yasuda 2006). A study of soybeans grown in Arkansas

found that cross-pollination of soybeans in adjacent rows averaged between 0.1% and 1.6%, but may be as high as 2.5% (Ahrent and Caviness 1994). Abud et al. (2007) illustrated that as distance is increased from the soybean pollination source, the chance of cross-pollination is decreased. This study found that at a distance of 1 meter, outcrossing averaged about 0.5%, at 2 meters about 0.1%, at 4 meters about 0.05%, and at 10 meters less than 0.01%.

Gene flow by seed is usually dependent on natural dispersal mechanisms, such as water, wind, or animals, or by human actions, and is favored by characteristics such as small and lightweight seed size, prolific production, seed longevity and dormancy, and long distance seed transport (Mallory-Smith and Zapiola 2008). Soybean seeds do not possess the characteristics for efficient seed-mediated gene flow. Soybean seeds are heavy and, therefore, are not readily or naturally dispersed by wind or water (Mallory-Smith and Zapiola 2008). Similarly, soybean seeds and seedpods do not have physical characteristics that encourage animal transport (OECD 2000). Soybeans also lack dormancy, a characteristic that allows dispersal in time by maintaining seeds and their genes within the soil for several years (OECD 2000; Mallory-Smith and Zapiola 2008).

Horizontal gene transfer and expression of DNA from a plant species to bacteria is unlikely to occur (Keese 2008). Many bacteria that are closely associated with plants have been described by sequencing them genetically, including *Agrobacterium* and *Rhizobium* (Kaneko et al. 2000; Kaneko et al. 2002; Wood et al. 2001). In cases where a review of sequence data implied that horizontal gene transfer occurred, evidence indicated that these events occurred over an evolutionary time scale (i.e., over millions of years) (Brown 2003; Koonin, Makarova, and Aravind 2001). The FDA has also evaluated horizontal gene transfer from the use of antibiotic resistance marker genes, and concluded that the likelihood of transfer of antibiotic resistance genes from plant genomes to microorganisms in the gastrointestinal tract of humans or animals, or in the environment, is remote (US-FDA 1998).

4.3.4 Microorganisms

Soil microorganisms significantly influence soil structure formation, decomposition of organic matter, toxin removal, nutrient cycling, and most biochemical soil processes (Garbeva, van Veen, and van Elsas 2004). They also suppress soil-borne plant diseases and promote plant growth (Doran, Sarrantonio, and Liebig 1996). One estimated range of the number of bacterial species in a gram of soil is between 6,400 and 3838 thousand (Curtis, Sloan, and Scannell 2002). The soil microbial community includes nitrogen-fixing microbes such as the soybean mutualist *B. japonicum*, mycorrhizal fungi, and free-living bacteria⁹; saprophytic fungi responsible for decomposition, denitrifying bacteria and fungi, phosphorus-solubilizing bacteria and fungi, and pathogenic microbes (USDA-NRCS 2004).

The main factors affecting microbial population size and diversity include soil type (texture, structure, organic matter, aggregate stability, pH, and nutrient content), plant type (providers of specific carbon and energy sources into the soil), agricultural management practices (crop rotation, tillage, herbicide and fertilizer application, and irrigation) and cropping history (Garbeva, van Veen, and van Elsas 2004; Garbeva, van Elsas, and van Veen 2008).

⁹Organisms that are able to obtain food without the need for a host organism.

Some types of soil microorganisms share metabolic pathways with plants that may be affected by herbicides. Tillage disrupts multicellular relationships among microorganisms, and crop rotation changes soil conditions in ways that favor different microbial communities. Plant roots, including those of soybean, release a variety of compounds into the soil creating a unique environment for microorganisms in the rhizosphere (root zone). Microbial diversity in the rhizosphere may be extensive and differs from the microbial community in the bulk soil (Garbeva, van Veen, and van Elsas 2004). More information about how soybean microbes, biotech crops, and herbicide use may affect soil microbial communities follows.

Soybean Microbes

An important group of soil microorganisms associated with legumes, including soybean, are the mutualists. These include mycorrhizal fungi, nitrogen-fixing bacteria, and some free-living microbes that have co-evolved with plants that supply nutrients to and obtain food from their plant hosts (USDA-NRCS 2004). Legumes have developed symbiotic relationships with specific nitrogen-fixing bacteria in the family *Rhizobiaceae* that induce the formation of root nodules where bacteria reduce atmospheric nitrogen to ammonia that is usable by plants (Gage 2004). *B. japonicum* is the bacterium specifically associated with soybeans (Franzen 2019). If a field has not been planted with soybean within 3-5 years, either the seed or seed zone must be inoculated with *B. japonicum* prior to soybean planting (Berglund and Helms 2003a; Pedersen 2007).

In addition to beneficial microorganisms, there are also several microbial pathogens that cause disease in soybean and vary somewhat depending on the region. These include fungal pathogens such as rhizoctonia stem rot (*Rhizoctonia solani*), brown stem rot (*Phialophora gregata*), sudden death syndrome (*Fusarium solani* race A), charcoal root rot (*Macrophomina phaseolina*); bacterial pathogens such as bacterial blight (*Pseudomonas syringae*) and bacterial pustule (*Xanthomonas campestris*), and viral pathogens such as soybean mosaic virus and tobacco ringspot virus (Ruhl 2007; SSDW No Date). The soybean cyst nematode (*Heterodera glycines*) is a microscopic parasite that infects the roots of soybeans. Management to control disease outbreaks varies by region, and pathogen/parasite, but include common practices such as crop rotation, weed control, planting resistant varieties, and proper planting and tillage practices.

Biotech Crop Impacts on Microbes

All soils, including agricultural soils are complex, dynamic ecosystems. Changes in agricultural practices and natural variations in season, weather, plant development stages, geographic location, soil type, and plant species or varieties can impact the microbial community (Kowalchuk, Bruinsma, and van Veen 2003; US-EPA 2009). Direct impacts may include changes to the structure (species richness and diversity) and function of the microbial community in the rhizosphere caused by the biological activity of the inserted gene(s). Indirect impacts may result from changes in the composition of root exudates, plant litter, or agricultural practices (Kowalchuk, Bruinsma, and van Veen 2003; US-EPA 2009). Several reviews of the investigations into the impact of transgenic plants on microbial soil communities found that most of them concluded there was either minor or no detectable non-target effects (Kowalchuk, Bruinsma, and van Veen 2003; US-EPA 2009; Hart 2006).

4.3.5 Biodiversity

Biodiversity refers to all plants, animals, and microorganisms interacting in an ecosystem (Wilson 1988). Biodiversity provides valuable genetic resources for crop improvement (Harlan 1975) and also provides other functions beyond food, fiber, fuel, and income. These include pollination, genetic introgression, biological control, nutrient recycling, competition against natural enemies, soil structure, soil and water conservation, disease suppression, control of local microclimate, control of local hydrological processes, and detoxification of noxious chemicals (Altieri 1999). Loss of biodiversity can result in more costly management practices to provide these functions to the crop (Altieri 1999).

The degree of biodiversity in an agroecosystem depends on four primary characteristics: (1) diversity of vegetation within and around the agroecosystem; (2) permanence of various crops within the system; (3) intensity of management; and (4) extent of isolation of the agroecosystem from natural vegetation (Altieri 1999). Agricultural land subject to intensive farming practices, such as that used in crop production, generally has low levels of biodiversity compared with adjacent natural areas. Tillage, seed bed preparation, planting of a monoculture crop, pesticide use, fertilizer use, and harvesting limit the diversity of plants and animals (Lovett, Price, and Lovett 2003).

Biodiversity can be maintained or reintroduced into agroecosystems through the use of woodlots, fencerows, hedgerows, and wetlands. Agronomic practices that may be used to support biodiversity include intercropping (the planting of two or more crops simultaneously to the same field), agroforestry, crop rotations, cover crops, no-tillage, composting, green manuring (growing a crop specifically for the purpose of incorporating it into the soil in order to provide nutrients and organic matter), addition of organic matter (compost, green manure, animal manure, etc.), and hedgerows and windbreaks (Altieri 1999). Integrated pest management strategies include several practices that increase biodiversity such as retaining small, diverse natural plant refuges and minimal management of field borders.

A variety of federally supported programs, such as the USDA funded Sustainable Agriculture Research and Education Program, and partnership programs between EPA and the agricultural community, support sustainable agricultural practices that are intended to protect the environment, conserve natural resources, and promote cropland biodiversity (i.e.,(USDA-NIFA 2017; US-EPA 2017a).

4.4 ANIMAL FEED

Animal agriculture consumes 98% of the U.S. soybean meal produced and 70% of soybeans produced worldwide (USB 2011d). Poultry consume more than 48% of domestic soybean meal or 11.92 million MT of the U.S. soybean crop with soy oil increasingly replacing animal fats and oils in broiler diets (ASA 2012b; USB 2011c). Soybean can be the dominant component of livestock diets, such as in poultry, where upwards of 66% of their protein intake is derived from soy. Other animals fed domestic soybean include swine (26%), beef cattle (12%), dairy cattle (9%), other (e.g., farm-raised fish 3%), and household pets (3%) (USB 2011a; ASA 2010a; USB 2011b).

Although the soybean market is dominated by seed production, soybeans have a long history in the United States as a nutritious grazing forage, hay, and silage crop for livestock (Blount et al.

2009). Soybean may be harvested for hay or grazed from the flowering stage to near maturity. The best soybean for forage is in the beginning pod stage (Johnson, Dunphy, and Poore 2007; Gonzalez and Burch 2019). For silage, it should be harvested at maturity before leaf loss, and mixed with a carbohydrate source, such as corn, for optimal fermentation characteristics (Blount et al. 2009). Varieties of soybean have been developed specifically for grazing and hay, but use of the standard varieties are recommended by some because of the whole plant feeding value.

Similar to the regulatory oversight for direct human consumption of soybean under the FFDCA, it is the responsibility of feed manufacturers to ensure that the products they market are safe and properly labeled. Feed derived from transgenic soybean must comply with all applicable legal and regulatory requirements, which in turn protects human health. To help ensure compliance, biotech organisms used for feed may undergo a voluntary consultation process with the FDA before release onto the market, which provides the applicant with any needed direction regarding the need for additional data or analysis, and allows for interagency discussions regarding possible issues. A developer who intends to commercialize a food derived from a biotech source consults with FDA to identify and discuss relevant safety, nutritional, or other regulatory issues regarding food derived from such crops, and then submits a summary of its scientific and regulatory assessment of the food to FDA. FDA evaluates the submission and responds to the developer by letter.

Growers must adhere to EPA label use restrictions for pesticides used to produce a soybean crop before using it as forage, hay, or silage. Under Section 408 of FFDCA, EPA regulates the levels of pesticide residues that can remain on feed from pesticide applications (US-EPA 2010a). These tolerances are the maximum amount of pesticide residue that can legally be present in food or feed, and if pesticide residues in food or feed are found to exceed the tolerance value, it is considered adulterated and subject to seizure.

4.5 HUMAN HEALTH

This section provides a summary of the human health concerns for public health related to the human consumption of products derived from transgenic soybeans, and those related to occupational health and health and safety from potential exposure to agricultural hazards.

4.5.1 Public Health

Human health concerns surrounding biotech soybeans focus primarily on human and animal consumption. Soybeans yield both solid (meal) and liquid (oil) products. Soybean meal is high in protein and is used for products such as tofu, soymilk, meat replacements, and protein powder. It also provides a natural source of dietary fiber (USB 2018). Nearly 98% of soybean meal produced in the United States is used as animal feed, while less than 2% is used to produce soy flour and proteins for food use. Soybean liquids are used to produce salad and cooking oils, baking and frying fat, and margarine. Soybean oil is low in saturated fats, high in polyunsaturated and monounsaturated fats, and contains essential omega-3 fatty acids. Soybean oil comprises nearly 70% of the oils consumed in U.S. households (ASA 2010b).

Soybean varieties developed for conventional use or for use in organic production systems, are not routinely required to be evaluated by any regulatory agency in the United States for human food or animal feed safety prior to market release. Food and feed manufacturers are responsible under FFDCA rules for ensuring that the products they market are safe and properly labeled.

Biotech organisms for food and feed may undergo a voluntary consultation process with the FDA prior to release onto the market. In a consultation, a developer who intends to commercialize a food derived from a biotech source meets with the FDA representatives to identify and discuss relevant safety, nutritional, or other regulatory issues. It then submits to the FDA a summary of its scientific and regulatory assessment of the food. This process includes: (1) an evaluation of the amino acid sequence introduced into the food crop to confirm whether the protein is related to known toxins and allergens; (2) an assessment of the protein's potential for digestion; and (3) an evaluation of the history of safe use in food (Hammond and Jez 2011). The FDA evaluates the submission and responds to the developer by letter with any concerns it may have or additional information it may require. Several international agencies also review food safety associated with food derived from biotech sources including the European Food Safety Agency and the Australia and New Zealand Food Standards Agency.

Foods derived from biotech sources undergo a comprehensive safety evaluation before entering the market, including reviews under the Codex Alimentarius, the European Food Safety Agency, and the World Health Organization (FAO 2009; Hammond and Jez 2011). Food safety reviews frequently will compare the compositional characteristics of the biotech crop with non-transgenic, conventional varieties of that crop. This comparison also evaluates the composition of the modified crop under actual agronomic conditions, including various agronomic inputs (FAO 2009; Aumaitre et al. 2002). Composition characteristics evaluated in these comparative tests include moisture, protein, fat, carbohydrates, ash, minerals, dietary fiber, essential and non-essential amino acids, fatty acids, vitamins, and antinutrients.

There are multiple ways in which organisms can be genetically modified through human intervention (e.g., traditional cross breeding, chemical or radiation-mediated mutagenesis, and genetic engineering using the methods of biotechnology). Unexpected and unintended compositional changes can arise with all forms of genetic modification, including both conventional hybridizing and genetic engineering (NRC 2004), however, no adverse health effects from genetic engineering have been documented in the human population. Reviews on the nutritional quality of foods have generally concluded that there are no significant nutritional differences in food and animal feeds derived from conventional versus biotech plants (Faust 2002; Flachowsky, Chesson, and Aulrich 2005).

Before a pesticide can be used on a food crop, the EPA must establish a tolerance, the maximum residue level of a pesticide that can remain on a crop or in foods processed from a crop, or establish an exemption for a tolerance (US-EPA 2010a). Both the FDA and USDA monitor foods for pesticide residues and enforce tolerances (USDA-AMS 2011). If pesticide residues in excess of a tolerance are detected on food, the food is considered adulterated and is subject to seizure. The USDA has implemented the Pesticide Data Program (PDP) to collect data on pesticides residues on food (USDA-AMS 2016). The EPA uses PDP data to prepare pesticide dietary exposure assessments pursuant to the FQPA. Pesticide tolerances have been established for most commodities, including soybeans, and have been published in the *Federal Register*, 40 CFR part 180, and the *Indexes to Part 180 Tolerance Information for Pesticide Chemicals in Food and Feed Commodities* (US-EPA 2011b).

4.5.2 Occupational Health and Worker Safety

Agriculture is one of the most hazardous U.S. work environments. Pesticides, particularly herbicides, are used on most soybean acreage in the United States. To protect all workers, the

National Institute of Occupational Safety and Health has been authorized by Congress to establish and enforce safety standards as part of a program to address high-risk issues in the work place. In response to the specific risks of poisoning and injuries among agricultural workers from pesticide exposure, the EPA has also established safeguards under its Worker Protection Standard (WPS) (40 CFR part 170) (US-EPA 2017c). The WPS establishes protections for more than 2.5 million agricultural workers in the United States who handle pesticides at more than 560,000 workplace sites on farms, forests, nurseries, and greenhouses.

The Occupational Safety and Health Administration requires all employers to protect their employees from hazards associated with pesticides and herbicides. The EPA WPS, updated in 2015 (US-EPA, 2017d), establishes specific safety procedures that must be followed by employers who hire workers who handle pesticides. The WPS requires pesticide safety training, notification of pesticide applications, use of personal protective equipment, restricted entry intervals (reentry times) following pesticide application, decontamination supplies and practices, and access to emergency medical assistance. The EPA pesticide registration process also includes protections for worker health. Under FIFRA, all pesticides sold or distributed in the United States must be registered by the EPA (US-EPA 2018). Registration decisions are based on scientific studies that assess the chemical's potential toxicity and environmental impact. All pesticides registered prior to November 1, 1984 must also be reregistered to ensure that they meet the current, more stringent standards. During the registration decision, the EPA must find that a pesticide does not cause unreasonable adverse effects to human health or the environment if used in accordance with its approved label instructions (OSTP 2001).

EPA labels for pesticides include use restrictions and safety measures to mitigate exposure risks. Pesticide applicators are required to use registered pesticides consistent with the instructions issued by the EPA that are listed on the label for each registered pesticide product. Worker safety precautions and use restrictions are included on pesticide registration labels. These include instructions for the levels of personal protection required for agricultural workers to safely handle and apply pesticides. Further details to achieve compliance are provided in the EPA WPS (US-EPA 2017c). When used in accordance with the EPA registration label, pesticides do not cause any unacceptable health risk to workers.

4.6 SOCIOECONOMIC ISSUES

Socioeconomic issues that are affected by soybean production in the United States are reviewed in this section. These issues are separated into two categories here, domestic economic environment and international trade environment.

4.6.1 Domestic Economic Environment

In 2020, soybeans accounted for 22% of all crops (excluding horticultural ones) grown in the United States. The 2020 crop represented a 10% increase in total acreage from 2019 with 88.8 million acres planted in the United States. HR varieties produced using genetic engineering were planted on 94% of this acreage (USDA-NASS 2020). Fourteen states (Arkansas, Illinois, Indiana, Iowa, Kansas, Michigan, Minnesota, Mississippi, Missouri, Nebraska, North Dakota, Ohio, and South Dakota and Wisconsin) accounted for most of the U.S. soybean production.

Soybean production cost data are compiled every 4-8 years by USDA-ERS as part of its Agricultural Resource Management Survey. For soybeans in 2019, typical operating costs per

planted acre included purchased seed (\$56.10), fertilizer and soil amendments (\$24.48), other chemicals (\$26.05), and irrigation water (\$0.07). Total operating costs were \$159.27 per planted soybean acre (USDA-ERS 2020b).

There is consistent evidence that farmers obtain substantial financial and non-financial benefits as a result of adoption of biotech crops. These benefits include increased income from off-farm labor, increased flexibility and simplicity in the application of pesticides, an ability to adopt more farming practices that have less environmental impact; increased consistency of weed control; increased human safety; equipment savings; and labor savings (Fernandez-Cornejo and McBride 2000; Duke and Powles 2009; Fernandez-Cornejo, C. Hendricks, and Mishra 2005; Fernandez-Cornejo and McBride 2002; Hurley, Mitchell, and Frisvold 2009; Marra, Piggott, and Carlson 2004; USDA-ERS 2020a).

Most of the soybean crop is crushed to produce oil and meal. In the United States, almost all (98%) of the soybean meal is used for animal feed. Most of the oil produced is used for human consumption; the balance is used for industrial products. Soybean oil represents almost 70% of the oils consumed by U.S. households. A noteworthy ongoing shift affecting soybean demand is an increased interest in using soybeans for biofuel production. From 1999 to 2009, the consumption of soybean biodiesel has increased from 0.5 to 1,070 million gallons (ASA 2012a).

Organic production methods account for only about 0.14% of the total U.S. soybean crop value. Although only a small proportion (about 0.09%) of the U.S. soybean crop is grown organically (USDA-ERS 2020b), it is more profitable per unit (bushel) of production than conventional systems that use non-biotech or biotech soybean varieties because the premium prices paid for soybeans certified as organically grown more than offset higher production costs (McBride and Greene 2009). Another factor contributing to enhanced profitability of organic soybeans is consumer demand for organically certified food in general, which has been experiencing a double-digit growth rate for more than a decade (USDA-ERS 2020b). The incentive for growers to choose conventional production systems is that organic methods are much more labor intensive, so organic farms tend to be much smaller than those that use conventional, highly mechanized systems. As a result, the latter yield higher overall profits per grower because the volume of production more than offsets premium prices paid for organically certified soybeans (McBride and Greene 2009).

4.6.2 International Trade Environment

Processed soybeans are the world's largest source of animal protein feed and the second largest source of vegetable oil. The United States is one of the world's leading soybean producers and the second-leading exporter. Soybeans comprise about 90% of U.S. oilseed production, while other oilseeds, including peanuts, sunflower seed, canola, and flax, make up the remainder (USDA-ERS 2017).

The total value of U.S. agricultural exports was \$135 billion in 2016 (USDA-ERS 2017). Of this total, \$23 billion was from soybean exports, ranking them first among all U.S. agricultural commodity exports. Since 2005, the percentage of U.S. soybean production that has been exported has increased from about 30% to nearly 50% (Figure 5). Despite the long-term trend in increasing export volume of U.S. soybean production, the U.S. share of the global export market has been declining (Figure 6) since 1980.




Figure 5. U.S. Soybean Export Volume and Percent Exported.

Source: (USDA-NASS 2018f)




Figure 6. U.S. Soybean Export Volume and Percent of Global Market.

Source: (USDA-NASS 2018f)

The United States, Brazil, Argentina, Paraguay, and Canada, account for 96.1% of the bulk soybean exported, while Argentina, Brazil, the United States, India, and Paraguay account for 90.4% of the soybean meal exported (Table 8). Argentina, the European Union (EU), and Brazil are the dominant countries in terms of soybean oil exports accounting for 75.4%. China, the EU, Mexico, and Japan are the major importers of world bulk soybean, accounting for 82.9% of total imports, whereas the EU, Indonesia, Thailand, Japan and Vietnam are the largest importers of soybean meal with a world share of 55.0% (USDA-FAS 2013). China and India are the major importers of soybean oil with a world share of 35.8% (USDA-FAS 2013). Between 1996 and 2011, 28 countries, including the United States, adopted the use of biotech crops, the largest being U.S., Brazil, Argentina, India, and Canada (Clive 2011). Prior to exporting GMB151 Soybean, BASF is expected to seek biotechnology regulatory approvals in all major import countries that have a functioning regulatory system to assure global compliance and support of international trade.

Table 8. World Soybean Production* (metric tons) in 2018.

Location	Soybean
Argentina	37.80
Brazil	122.00
Canada	7.72
China	15.20
European Union	2.54
India	8.35
Mexico	0.43
Paraguay	10.3
United States	341.54
Total Foreign	221.48

*Major Producers

Source: (USDA-NASS 2018f)

5 ENVIRONMENTAL CONSEQUENCES

Possible environmental impacts from selecting either the No Action Alternative or the Preferred Alternative as part of regulatory decision making by APHIS for GMB151 Soybean were considered in this chapter. Details about how APHIS evaluated environmental impacts, results of the analyses it performed to assess whether or not they caused impacts, and the Agency's conclusions about the significance of impacts it identified are presented in this chapter. Pursuant to CEQ regulations APHIS considers the direct impacts of both alternatives.

5.1 ENVIRONMENTAL ANALYSIS

For this chapter, those impacts that were categorized as direct were evaluated. A direct impact was considered to be one solely caused by an Agency action without any intervening intermediate steps. An example is conversion of land use from non-agricultural to agricultural in response to an action that increases demand for a crop.

5.2 SCOPE OF THE ENVIRONMENTAL ANALYSIS

Those resource areas listed in Chapter 1 (see Issues Considered) that may be affected by selecting either the No Action Alternative or the Preferred Alternative were considered for this EA. Impacts were defined as those effects likely to result in permanent changes to the environment. Impacts were evaluated for significance by analyzing the positive or negative changes from the existing (baseline) conditions described in Chapter 3 (Affected Environment). Wherever possible, APHIS used data that supported a quantitative analysis of the impacts of selecting either the No Action Alternative or the Preferred Alternative. When data were not available or were insufficient to support a quantitative assessment, APHIS made qualitative assessments of the impacts of an Agency regulatory decision for GMB151 Soybean.

APHIS limited its environmental analyses to the geographic areas that currently support U.S. soybean production. These analyses were also made under the assumption that most U.S. farmers who produce soybeans rely on widely accepted best management practices (BMPs). It was also assumed that if GMB151 Soybean was no longer regulated by APHIS and became widely planted, farmers would use the same BMPs that are currently used for soybean production in the United States.

The Cry14Ab-1 protein expressed by GMB151 Soybean confers resistance to the soybean cyst nematode (SCN), *Heterodera glycines*, the. GMB151 also produces a modified 4-hydroxyphenylpyruvate dioxygenase (HPPD-4) enzyme that confers resistance to HPPD-inhibitor herbicides such as isoxaflutole (BASF 2020). Production of the HPPD-4 enzyme in transgenic crops has been evaluated previously in EAs by APHIS (USDA-APHIS 2018, 2013, 2014) and those assessments are incorporated into this EA by reference. In one of these evaluations (USDA-APHIS 2013), the organism (*Pseudomonas fluorescens* strain 32) that was the source of the gene that produces the HPPD-4 protein is the same one used to develop GMB151 Soybean (BASF 2020). Since the previous APHIS assessments did not identify any significant impacts associated with the HPPD-4 enzyme or the gene that produces it, no further analysis of the HPPD-4 enzyme was performed for this EA.

APHIS emphasizes that it has regulatory authority over GMB151 Soybean plants, but EPA has regulatory authority over herbicides that are applied to the crop. The scope of this EA covers the possible direct impacts that would result primarily from the cultivation and use of the plant. EPA is considering impacts from the use of HPPD-inhibitor herbicides on GMB151 plants as part of its registration process. USDA is relying on EPA's authoritative assessments and will not duplicate the assessment prepared by EPA. This EA will provide informative assessments, but not the determinative document for any impacts of herbicide usage, since that analysis will be completed by EPA under its regulatory authority.

5.3 AGRICULTURAL PRODUCTION OF SOYBEAN

U.S. soybean acreage is concentrated primarily in the Midwest (Figure 1), where yields are highest (USDA-NASS 2016a). Soybean acreage has expanded recently to the northern and western parts of the country. This has resulted because of the availability of newer improved soybean varieties better adapted to provide higher yields under the short-season climatic conditions of those areas (USDA-ERS 2010b). This has been a major factor contributing to a 31% increase in recent years to a total U.S. soybean acreage of about 83 million acres (USDA-NASS 2019). Current trends in U.S. biotech soybean production are expected to continue unchanged in the major soybean producing states listed previously in Chapter 3 (Table 2).

5.3.1 Locations and Acreage of Soybean Production: No Action Alternative

Under the No Action Alternative, GMB151 Soybean would remain regulated and would not be commercially available for production. Transgenic soybeans were introduced in the United States in 1996 (USDA-ERS 2011a; Fernandez-Cornejo and Caswell 2006). By 2017, 94% of U.S. soybean acreage was planted in a variety produced using genetic engineering (USDA-NASS 2018c). Most of this shift by growers to biotech soybeans resulted because of the cost-effective benefits gained from improved weed control with HR varieties.

The trend of planting primarily biotech soybeans in the United States will likely continue under the No Action Alternative as new varieties are developed with new traits or that combine different traits desired by growers and consumers. For example, during the past decade, APHIS has considered petitions for nonregulated status for biotech soybean varieties that combine resistance to multiple herbicides, provide insect resistance, or modify nutritional properties of the oil derived from soybeans. Although these current trends for development of new biotech soybean varieties are expected to continue, U.S. soybean acreage and production is not expected to change in the foreseeable future, and selection of the No Action Alternative is unlikely to alter this projection.

5.3.2 Locations and Acreage of Soybean Production: Preferred Alternative

BASF conducted phenotypic, agronomic, and environmental interaction trials with GMB151 Soybean and its non-transgenic soybean control parental line having a genetic background similar to GMB151 Soybean. The results of the combined trials demonstrated that there were no substantial agronomic or phenotypic differences between GMB151 Soybean and its comparator control or other commercial soybean varieties. Other than resistance to SCN, GMB151 Soybean confers no novel agronomic benefit compared to other soybean varieties. It would only be expected to replace currently planted soybean varieties, where SCN has been a pest problem, so soybean production trends and U.S. soybean acreage are unlikely to change.

Because GMB151 Soybean is anticipated to increase yields, it might be expected to replace other varieties of soybean currently grown. Since the middle of the last century, changes in soybean varieties have contributed to increased yields, as have improved management practices. From 1991 to 2011, average soybean yield increased approximately 17.6% from 34.2 bushels per acre in 1991 to 41.5 bushels per acre in 2011, then declined slightly in 2012 to 39.3. Since 1991, U.S. soybean production acreage has increased 31% (USDA-NASS 2018c).

As described for the No Action Alternative, the USDA has projected that soybean acreage will remain relatively steady during the next decade (USDA-OCE, 2018). Based upon its phenotypic and agronomic similarity to other soybean varieties, GMB151 Soybean is also subject to the same variables affecting yield in other soybean varieties, such as management practices and weather (see “Agronomic Inputs” in Chapter 3 for more details). It is unlikely that a significant change in U.S. soybean production acreage would result from a determination of nonregulated status of GMB151 Soybean. Therefore, effects on soybean production acreage under the Preferred Alternative would be the same as for the No Action Alternative and would not differ from current baseline conditions reviewed in Chapter 3, so no significant impacts are anticipated.

5.3.3 Agronomic Practices: No Action Alternative

Under the No Action Alternative, GMB151 Soybean would continue to be subject to the regulatory requirements of 7 CFR 340 and plant pest provisions of the PPA. However, growers would still have access to conventional and nonregulated biotech soybean varieties currently available. The potential environmental impacts associated with the agronomic practices and inputs used for the production of biotech and conventional soybean varieties such as conventional and conservation tillage, soil and foliar fertilization, crop rotation, irrigation, pest (insects and weeds) and disease management with herbicides, pesticides, and fungicides, and crop residue management would be unaffected by continued regulation of GMB151 Soybean.

A variety of herbicide choices are available to growers, including those used for preplant only, pre-emergent herbicides often with residual activity, those used as post-emergent herbicides, and combinations of both. EPA approves and labels uses of pesticides on soybeans. Under the No-Action Alternative, commercial soybean growers would continue to use the same pesticides for soybean insect pests and weeds as are currently used.

5.3.4 Agronomic Practices: Preferred Alternative

A determination of nonregulated status of GMB151 Soybean is not expected to result in changes to current soybean cropping practices. BASF’s studies (BASF 2020) demonstrated that except for its capacity to control SCN, GMB151 Soybean is essentially the same as other commercial soybean varieties in terms of agronomic characteristics and cultivation practices.

Soybeans are grown mostly in rotation with corn. In two-year rotations, a common practice is to fertilize the previous year’s corn crop with enough phosphorus and potassium to allow for the subsequent soybean crop to be grown with no supplemental fertilizer as it is more economical than two separate applications (Franzen 2019; Berglund and Helms 2003b; Ebelhar et al. 2004a). About two-thirds of U.S. soybean is grown in rotation with corn. However, annual supplementation of nutrients is common in soybeans that are not grown in rotation with another crop. Regular testing of soil fertility levels and supplementation if needed is already widely recommended in soybean production for achieving optimal yields. Current practices in soybean include 41% of soybean acreage annually supplemented with phosphate (phosphorous) and 42%

annually supplemented with potash (potassium) (USDA-NASS 2018b). These and other agronomic baseline conditions as reviewed in Chapter 3 would be the same as for the No-Action Alternative, so no significant impacts on agronomic practices are anticipated if the Preferred Alternative is selected.

5.3.5 Soybean Seed Production: No Action Alternative

Under the No Action Alternative, current soybean seed production practices are not expected to change. Several factors influence optimal planting rate for soybean such as row spacing, seed germination rate, soil conditions, climate, disease and pest pressure, past tillage practices and crop rotation (Robinson and Conley 2007). Seeding rate is also determined by the plant population desired by the grower. Growers may plant certified soybean seed, uncertified seed, and seed that is grown and stored on individual farms (Oplinger and Amberson 1986).

The production of the soybean seed crop for foundation, registered, certified, or quality control seed requires biological, technical, and quality control practices to maintain varietal purity greater than that for soybean grain production. The production and certification of soybean seed is regulated by state or regional crop improvement agencies that are chartered under the laws of the state(s) they serve (e.g., see Virginia Crop Improvement Association No Date; SSCA No Date-a; Illinois Crop Improvement Association 2013; Mississippi Crop Improvement Association 2008, 2015; Illinois Crop Improvement Association 2019). The procedures followed by certified seed producers to ensure varietal purity and identity during the cultivation, harvest, storage, and transportation of soybean seed are not expected to change under the No Action Alternative.

Seed genetic purity is maintained to maximize the value of a new variety (Sundstrom et al. 2002), of which a seed certification process ensures that the desired traits remain within purity standards (Bradford 2006) (for more details see Chapter 3: Soybean Seed Production). Seed producers routinely submit applications to the AOSCA National Variety Review Boards for review and recommendation for inclusion into seed certification programs. For example, in September 2012, AOSCA recommended the inclusion of 60 varieties of soybean expressing high yield traits by three seed producing companies for certification (AOSCA 2012b, 2012a). It is expected that soybean seed producers would continue to implement measures to preserve the identity of their seed varieties if the No Action Alternative is selected.

5.3.6 Soybean Seed Production: Preferred Alternative

Field trials conducted by BASF have not demonstrated any agronomic or phenotypic differences between GMB151 Soybean and conventional soybean varieties that would require changes to soybean seed production practices (BASF 2020). Based on the data provided by BASF, APHIS has concluded that the availability of GMB151 Soybean under the Preferred Alternative would not alter the agronomic practices, cultivation locations, seed production practices or quality characteristics of conventional and biotech soybean seed production. BASF has also indicated GMB151 Soybean will be adopted into existing maturation groups to match the area in which it would be cultivated. Therefore, its adoption would not alter planting practices of soybean grown for seed. Based on this information, soybean seed production associated with the Preferred Alternative would not be any different than practices under the No Action Alternative so no significant impacts on soybean seed production are anticipated if the Preferred Alternative is selected.

5.3.7 Organic and Conventional Soybean Production: No Action Alternative

Under the No Action Alternative, GMB151 Soybean would remain subject to the regulatory requirements of 7 CFR 340 and the plant pest provisions of the PPA. Biotech, conventional and organic soybean production would not change as a result of the continued regulation of GMB151 Soybean. Organic and conventional soybean growers would continue to use the same methods they currently use to manage crop identity, preserve the integrity of their production systems, and maintain organic certification. As described in Chapter 3 (Affected Environment: Organic Soybean Production), organic and conventional soybean production is a small portion of the soybean market (USDA NASS 2017).

5.3.8 Organic and Conventional Soybean Production: Preferred Alternative

Biotech soybeans are already extensively used by farmers, while organic (less than 1.0%) and conventional (less than 10.0%) soybean production represents a small percentage of the total U.S. soybean acreage. Organic and -conventional soybean acreage is unlikely to change significantly, regardless of whether new conventional or biotech soybean varieties, such as GMB151 Soybean, become available for commercial production.

When compared to other biotech varieties of soybean, GMB151 Soybean does not present any new or different issues or potential impacts for organic and other specialty soybean producers and consumers. Organic producers employ a variety of measures to manage, identify and preserve the integrity of organic production systems. Agronomic tests conducted by BASF found that GMB151 Soybean is substantially equivalent to conventional soybeans (BASF 2020). Pollination characteristics are similar to other soybean varieties currently available to growers. Since soybeans exhibit limited pollen movement and are mostly self-pollinating (Abud et al. 2007; Caviness 1966; OECD 2000; Ray et al. 2003; Yoshimura 2011), there is no indication that organic and conventional soybean crops will be affected by a determination of nonregulated status for GMB151 Soybean if they continue to be produced in accordance with current agronomic practices to meet organic standards such as those of the NOP.

The trend in the cultivation of biotech, conventional, and organic soybean varieties, and the corresponding production systems to maintain varietal integrity are likely to remain the same as those for the No Action Alternative. Therefore, impacts on organic and conventional soybean growers if a determination of nonregulated status for GMB151 Soybean is made (i.e., selection of the Preferred Alternative) would be the same as or similar to the No Action Alternative, so no significant impacts on organic and conventional soybean production are anticipated if the Preferred Alternative is selected.

5.4 PHYSICAL ENVIRONMENT

5.4.1 Soil Quality: No Action Alternative

Under the No Action Alternative, current soybean soil management practices that affect soil quality, including the use of cover crops to limit the time soil is exposed to wind and rain, tillage methods to reduce erosion and compaction, control weeds, and enhance nutrients, careful management of fertilizers and pesticides, crop rotation, establishing windbreaks and contour plowing (for more details see Chapter 3: Agronomic Practices) would be expected to continue unchanged. Growers would continue to choose methods based on weed, insect, and disease pressure, as well as the costs of seed and other inputs consistent with BMPs they currently use.

5.4.2 Soil Quality: Preferred Alternative

A determination of nonregulated status for GMB151 Soybean is not expected to result in changes to current soybean cropping practices that may impact soil quality. Studies conducted by BASF demonstrated that GMB151 Soybean is essentially indistinguishable from conventional and other biotech soybean varieties in terms of agronomic characteristics and cultivation practices (BASF 2020).

The Cry14Ab-1 protein expressed by GMB151 Soybean is not expected to have any effects on the physicochemical characteristics of soil (BASF 2020). In particular, mutual symbiotic relationships between soybean and the Rhizobiaceae and Bradyrhizobiaceae are unlikely to be negatively affected. Many Cry proteins derived from *B. thuringiensis* are rapidly degraded in a variety of soil types and these proteins do not accumulate (Head et al. 2002; Mendelsohn et al. 2003; Dubelman et al. 2005). For a few Cry proteins, residual amounts of Bt proteins may persist for extended periods, but the levels detected are not biologically significant (Feng et al. 2011). However, EPA has concluded in a risk assessment that available data indicates that there is short term accumulation of Cry1F and Cry1Ac proteins in agricultural soil. It concluded that Cry proteins have a short half-life, and are unlikely to affect soil invertebrates or significantly impact soil microbiota (US-EPA 2014).

Soil quality may be impacted by a soybean crop through direct interaction with soil fauna at the root system and by the degradation of remaining plant tissue after harvest. However, compositional analysis of GMB151 Soybean forage tissue (i.e., stems and leaves) revealed no significant or consistent differences between it and the conventional control variety (BASF 2020). There also were no differences between GMB151 Soybean and the conventional control variety with respect to plant-environment interactions or plant-symbiont interactions. Because of the compositional similarities between GMB151 Soybean and conventional soybeans, and the examined safety of the GMB151 Soybean gene products, it is not anticipated that GMB151 Soybean interactions with soil fauna or the impact of degradation of its stubble remaining in fields following harvest would be significantly different from that of conventional soybean. Based on the Agency's analyses of this information, overall impacts to soil under the Preferred Alternative are not expected to differ from those of the No Action Alternative, so no significant impacts on soil are anticipated if the Preferred Alternative is selected.

5.4.3 Water Resources: No Action Alternative

Current soybean management practices, including irrigation, and pesticide and fertilizer applications, would be expected to continue unchanged under the No Action Alternative. Under the authority of FIFRA, environmental risks of pesticide use are assessed during the registration processes of the EPA, and are regularly reevaluated to ensure that registered uses continue to pose no unreasonable risks to humans or the environment, including risks to water resources.

The trend towards conservation tillage or no-tillage practices since the adoption of HR soybean varieties is expected to continue, resulting in reduced surface water run-off and soil erosion (Dill, CaJacob, and Padgett 2008; Givens et al. 2009). Conservation tillage and other management practices are used to trap and control sediment and nutrient runoff. Water quality conservation practices benefit agricultural producers by lowering input costs and enhancing the productivity of working lands.

As of 2018, nitrogen, potassium, and phosphorous were applied to 18%, 25%, and 23% respectively of soybean acreage in 19 states surveyed (USDA-NASS 2018e). Production practices for any soybean variety remove more nutrients from soils (Pedersen 2008) than less intensive methods. Regular testing of soil fertility is required and applications of nutrients are not uncommon in soybean production (USDA-NASS 2018a). Nitrogen is not usually applied to soybeans because they fix nitrogen in the soil through their symbiotic relationship with rhizomatous bacteria (CAST 2009).

5.4.4 Water Resources: Preferred Alternative

No differences in morphological characteristics and agronomic requirements were found between GMB151 Soybean and a non-transgenic comparator (BASF 2020). Therefore, cultivation of GMB151 Soybean would not necessitate changes in current agronomic practices for soybean production, so current impacts of soybean cultivation on water quality would not change if GMB151 Soybean were no longer regulated under 7 CFR 340. BASF evaluated on a site-specific basis abiotic stressors such as drought and flood, and found no difference between GMB151 Soybean and its comparator. As described previously, if GMB151 Soybean is no longer regulated under 7 CFR 340, neither total U.S. soybean acreage nor its locations would change, so there would be no shifts in how or where water quality impacts related to soybean cultivation would occur in the United States. For these reasons, a determination of nonregulated status of GMB151 Soybean is unlikely to change the current irrigation practices in commercial soybean production.

Adoption of HR crops is associated with increased use of no-till and reduced till practices that benefit water quality by reducing runoff loads from soil erosion (Givens et al. 2009; Dill, CaJacob, and Padgett 2008). The adoption rate of HR soybeans has steadily increased since their introduction in 1996. Today, more than 94% of U.S. commercially grown soybeans are herbicide resistant (USDA-NASS 2018c). This trend is unlikely to change if GMB151 Soybean were to become commercially available.

Runoff from cropland areas receiving manure or fertilizer contributes to increased phosphorous and nitrogen in streams and lakes. In fresh water systems, phosphorus is the limiting factor causing eutrophication (see Chapter 3, Water Resources, for more details). Up to 41% of soybean acreage has been annually supplemented with phosphorous (USDA-NASS 2018a). Since GMB151 Soybean is unlikely to change total U.S. soybean acreage or where soybeans are grown in the United States, impacts to water resources from fertilization are not expected to differ from those of the No Action Alternative, so no significant impacts on water resources are anticipated if the Preferred Alternative is selected.

5.4.5 Air Quality: No Action Alternative

Under the No Action Alternative, current air quality impacts from the soybean agronomic practices described in Chapter 3 such as tillage, cultivation, and agrochemical applications would continue unchanged. Applications of EPA-registered pesticides would continue unchanged, as would any associated environmental impacts because as part of its reregistration process, the EPA regularly reevaluates registered pesticides to ensure that they continue to pose no unacceptable risks. Of particular relevance to air quality, this process includes identifying methods to reduce pesticide drift, which are included on pesticide labels and approved by the EPA. Under the No Action Alternative use of pesticides according to EPA-approved labels would not pose unreasonable risk to air quality. The trend towards conservation tillage and no-

till practices associated with cultivating HR soybean varieties, which reduces exhaust emissions from agricultural equipment and airborne dust from soil disturbance is also likely to continue (Dill, CaJacob, and Padgett 2008; Givens et al. 2009).

5.4.6 Air Quality: Preferred Alternative

No differences in morphological characteristics and agronomic requirements were found between GMB151 Soybean and a conventional comparator (BASF 2020). Therefore, if the Preferred Alternative is selected, cultivation of GMB151 Soybean would not result in changes to most current soybean agronomic practices (e.g., fertilizer and pesticide applications). The addition of HPPD-herbicide resistance provides another option (e.g., isoxaflutole) for resistant weed control, which when combined with SCN resistance should contribute to maintaining the current level of usage of no-till practices. Therefore, no changes to emission sources (i.e., tillage and fossil-fuel-burning equipment) would be expected. As described previously, commercial use of GMB151 Soybean would neither increase the total U.S. soybean acreage nor modify the existing U.S. soybean production range. Since no changes to agronomic practices for the cultivation of GMB151 Soybean, and no increase in area or acreage are expected if the Preferred Alternative is selected, impacts to air quality are expected to be the same as the No Action Alternative, so no significant impacts on air quality are anticipated if the Preferred Alternative is selected.

5.5 BIOLOGICAL RESOURCES

5.5.1 Animal Communities: No Action Alternative

Under the No Action Alternative, animal species would continue to be affected by agronomic practices associated with soybean production, such as tillage, cultivation, pesticide and fertilizer applications, and the use of agricultural equipment (Sharpe 2010; Brady 2007; Palmer, Bromley, and Anderson No Date; USDA-NRCS 1999a) no differently than they are currently. Some of these current practices have potential to impact animal communities. For example, if tillage rates were to increase as a means of weed suppression, it could possibly diminish benefits to wildlife from conservation tillage practices. Some pesticides for weed, insect, and disease control may also impact animal communities. However, environmental risks of pesticides to wildlife and their habitats are assessed by the EPA in its registration process and are regularly reevaluated to establish uses that have a reasonable certainty of not causing harm to non-target animals and their habitats.

5.5.2 Animal Communities: Preferred Alternative

A determination of nonregulated status of GMB151 Soybean is not expected to result in changes to current soybean cropping practices. BASF's studies demonstrated GMB151 Soybean is the same as other soybean varieties in terms of agronomic characteristics and cultivation practices (BASF 2020). Therefore, replacement of other biotech and conventional soybean varieties with GMB151 Soybean will not alter agronomic practices currently used (e.g., crop rotation; weed management; cultivation), so no changes in effects from those of current soybean cultivation practices on wildlife that use soybean fields for cover and forage (as described in Chapter 3 in "Animal Communities") are likely.

Field trials showed that GMB151 Soybean does not confer any biologically significant differences to susceptibility or tolerance to invertebrate pests other than SCN. This indicates that

there would be no changes to agronomic practices, such as increased use of pesticides, that could impact wildlife.

Results of testing of non-target organisms presented by BASF indicated that the Cry14Ab-1 protein expressed by GMB151 Soybean is unlikely to be toxic when consumed by animals other than the target species (SCN). On January 28, 2019, BASF initiated a consultation (BNF 172) with the FDA that included molecular, composition, and nutrition data, and other food and feed safety assessment data related to GMB151 Soybean (BASF 2020). In addition, EPA concluded on June 8, 2020 (40 CFR 174.540) that the Cry14Ab-1 protein is exempt from a food and feed tolerance, when it is expressed in soybean plants. Based on the above information, there are no expected hazards associated with the Cry14Ab-1 protein expressed by GMB151 Soybean from exposure to or consumption by animals that reside in or near soybean fields.

The source organism of the HPPD-4 protein, *P. fluorescens* strain A32, is a non-pathogenic bacterium that is ubiquitous in nature and has a history of safe use. HPPD proteins are also ubiquitous in nature in nearly all aerobic organisms, (e.g., bacteria, fungi, plants, and animals including mammals). The HPPD-4 protein has no significant amino acid sequence similarity to known allergens or toxins, is rapidly degraded in simulated gastric fluid, and exhibited no effects in an acute oral mouse toxicity test. Also, EPA has established a permanent exemption (82FR57137) from the requirement for a tolerance for the HPPD-4 protein expressed in all food commodities when used as an inert ingredient.

Based on available evidence, GMB151 Soybean is unlikely to pose a hazard to wildlife species. As a result, APHIS concluded, effects, if any on wildlife species, would be the same as those for the No Action Alternative, so no significant impacts on animal communities are anticipated if the Preferred Alternative is selected.

5.5.3 Plant Communities: No Action Alternative

Under the No Action Alternative, GMB151 Soybean would remain under APHIS regulation. Current soybean production would likely continue unchanged. Growers would continue to select the agronomic practices such as tillage, irrigation, row spacing, timing of planting, and weed management that optimize soybean yield and efficiency that they currently use.

Plant species that typically compete with soybean production would be managed through the use of mechanical, cultural, and chemical control methods. Multiple herbicides would continue to be used for weed control in soybean fields. Runoff, spray drift, and volatilization of herbicides have the potential to impact non-target plant communities growing in proximity to fields in which herbicides are used. The environmental effects of pesticide use are assessed by the EPA in the pesticide registration process and are regularly reevaluated by the EPA in its reregistration process under FIFRA. In this process, where appropriate, steps to reduce pesticide drift and volatilization are included on a pesticide's label approved by the EPA to minimize off-target effects.

5.5.4 Plant Communities: Preferred Action Alternative

A determination of nonregulated status of GMB151 Soybean is not expected to result in changes to current soybean cropping practices. Field trials and laboratory analyses conducted by BASF showed no evidence of differences between GMB151 Soybean and other biotech and conventional soybean in growth, reproduction, or interactions with pests and diseases except for resistance to SCN (BASF 2020). The expression of the Cry14Ab-1 protein by GMB151 Soybean

will provide a PIP that will suppress SCN and reduce or eliminate soybean crop damage from it. Is not expected to cause plant disease or increase susceptibility of GMB151 Soybean or other soybean varieties derived from it to diseases or other pests (USDA-APHIS 2020).

Similar to the No Action Alternative, weeds within fields of GMB151 Soybean could be managed using mechanical, cultural, and chemical control. There are no differences expected in the use of herbicides or other pesticides in the production of GMB151 Soybean, when compared to other biotech and conventional soybean varieties (USDA-APHIS 2020). Except for the option to substitute isoxaflutole for other herbicides used, agronomic practices to cultivate GMB151 Soybean would not differ from the No Action Alternative. Other isoxaflutole-resistant transgenic soybean varieties that are not regulated are currently available to growers. GMB151 Soybean would only replace these as another alternative to growers, so isoxaflutole use is not expected to change. If a determination of nonregulated status is made for GMB151 Soybean, the risks to wild plants and agricultural productivity from weedy GMB151 Soybean populations are negligible because volunteer soybean populations can be easily managed and there are no feral or weedy relatives in the United States (Carpenter et al. 2002).

Based on the information reviewed, APHIS has determined that the effects on other vegetation in and around soybean fields from a determination of nonregulated status for GMB151 Soybean are identical to those under the No Action Alternative so no significant impacts on plant communities are anticipated if the Preferred Alternative is selected.

5.5.5 Gene Flow and Weediness: No Action Alternative

Under the No Action Alternative, GMB151 Soybean would remain under APHIS regulation. The availability of biotech, conventional and organic soybeans would not change as a result of the continued regulation of GMB151 Soybean. Because soybean is mostly self-pollinated, its cross-pollination rate significantly decreases with distance, and there are no wild and weedy relatives in the US, introgression of soybean pollen to wild or weedy species is virtually impossible. In addition, volunteer soybeans are typically not a major problem in agroecosystems, and regionally where volunteer soybean populations can develop, the volunteer plants are manageable and do not represent a serious weed threat (York, Beam, and Culpepper 2005).

5.5.6 Gene Flow and Weediness: Preferred Alternative

A determination of nonregulated status of GMB151 Soybean is not expected to pose greater pollen- or seed-mediated gene flow, or increased potential for weediness than that of currently cultivated soybean varieties. There were no differences between GMB151 Soybean and conventional soybean varieties tested that would increase the potential for gene flow from GMB151 Soybean or otherwise increase its weediness (BASF 2020).

APHIS evaluated information in its PPRA (USDA-APHIS 2020) on the inserted genetic material, the potential for vertical and horizontal gene transfer, and weedy characteristics of GMB151 Soybean and concluded it would not represent any plant pest risk. Field trials and laboratory data for GMB151 Soybean indicate no plant pathogenic properties or weediness characteristics (BASF 2020). Based on agronomic data and compositional analyses, GMB151 Soybean was found to be substantially equivalent to conventional soybeans and would no more likely become a plant pest than conventional soybeans. The reproductive characteristics of GMB151 Soybean are essentially equivalent to other biotech and conventional soybean varieties (BASF 2020). GMB151 Soybean would not persist in unmanaged environments and does not

demonstrate a competitive advantage compared to conventional soybeans. Neither of the transgenic traits in GMB151 Soybean (SCN and resistance to HPPD-inhibiting herbicides such as isoxaflutole) present a risk of increased weediness. The reproductive mechanism in soybeans also makes the potential for cross-pollination of GMB151 Soybean with other soybean varieties highly unlikely (BASF 2020).

In reference to interspecific gene transmission, studies have indicated that horizontal gene transfer and expression of DNA from a plant species to bacteria is unlikely to occur (Keese 2008). Furthermore, there is no evidence that bacteria closely associated with plants and/or their constituent parts contain genes derived from plants (Kaneko et al. 2000; Kaneko et al. 2002; Wood et al. 2001). When horizontal gene transfer has been observed, it has been on an evolutionary time scale of millions of years (Brown 2003; Koonin, Makarova, and Aravind 2001). Based on this information, APHIS has concluded that horizontal gene flow from GMB151 Soybean to other unrelated organisms would be highly unlikely.

If a determination of nonregulated status is made for GMB151 Soybean, the risks to wild plants and agricultural productivity from weedy GMB151 Soybean populations are negligible, as volunteer soybean populations can be easily managed and there are no feral or weedy relatives in the United States (Carpenter et al. 2002). If present as volunteer soybean, GMB151 Soybean would not be considered difficult to control, as soybean seeds rarely remain viable the following season and are easily managed with cultivation, hand weeding, or the application of herbicides. In addition, since no feral or weedy species of soybean exist in the United States (Ellstrand, Prentice, and Hancock 1999; OECD 2000), GMB151 Soybean poses no potential for either naturally occurring, pollen-mediated gene flow or transgene introgression. Based on the information reviewed, APHIS has determined that the effects on other vegetation in and around soybean fields from a determination of nonregulated status for GMB151 Soybean are indistinguishable from those under the No Action Alternative. Therefore, no significant impacts associated with gene flow or weediness are anticipated if the Preferred Alternative is selected.

5.5.7 Microorganisms: No Action Alternative

Under the No Action Alternative, GMB151 Soybean would remain under APHIS regulation. The availability of biotech, conventional, and organic soybeans would not change as a result of the continued regulation of GMB151 Soybean. Agronomic practices used for soybean production, such as soil inoculation, tillage, and the application of agricultural chemicals (pesticides and fertilizers) that potentially impact microorganisms, would continue unchanged.

5.5.8 Microorganisms: Preferred Alternative

A determination of nonregulated status of GMB151 Soybean is not expected to change current soybean cropping practices that may affect microorganisms. Possible impacts of GMB151 Soybean on microbial communities would be identical or similar to those for conventional soybeans and other biotech varieties. GMB151 Soybean could have some effect on the structure of the soil microbial community in which it is planted, which could include nitrogen-fixing bacteria and mycorrhizal fungi; bacteria, actinomycetes (filamentous bacteria), and saprophytic fungi responsible for decomposition; denitrifying bacteria and fungi; phosphorus-solubilizing bacteria and fungi; as well as pathogenic and parasitic microbes (USDA-NRCS 2004). Testing by BASF revealed no significant differences found in the parameters measured to assess the relationship of the legume and its associated symbiont between GMB151 Soybean and the conventional comparators (BASF 2020).

As with other biotech soybean varieties in high yield production systems, the cultivation of GMB151 Soybean may remove more nutrients, particularly phosphorus and potassium, than lower yielding conventional varieties, necessitating testing and possibly increased soil nutrient amendments. Soil organisms require varying amounts of both macronutrients, including phosphorus, and micronutrients (USDA-NRCS 2004). Several studies have demonstrated *B. japonicum* activity, root nodulation, and nitrogen fixation are positively correlated with phosphorus levels (Beck and Munns 1984; Cassman, Whitney, and Stockinger 1980; Israel 1987; Mullen, Israel, and Wollum 1988; Sa and Israel 1991; Tsvetkova and Georgiev 2003). Likewise, potassium is necessary for nodule formation and bacteria-mediated nitrogen fixation in soybean and other nitrogen-fixing legumes (IPNI 1998; Mengel, Haghparast, and Kock 1974). Applications of these nutrients to soybean is not an uncommon practice and is widely recommended to sustain the yields of all soybean varieties.

Field and greenhouse tests show no significant differences from other nonregulated conventional and biotech soybean varieties in the parameters measured to assess the relationship of GMB151 Soybean with its symbionts. GMB151 Soybean would not result in any significant changes to current soybean cropping practices except that it may be grown where SCN is a problem and/or where isoxaflutole may be substituted for herbicides with other modes of action that are ineffective against HR weeds. Other isoxaflutole-resistant biotech soybean varieties that are not regulated are currently available to growers. GMB151 Soybean would only replace these as another alternative to growers, so isoxaflutole usage would not be expected to change.

Based on the above information, overall impacts to microorganisms under the Preferred Alternative would be indistinguishable from those under the No Action Alternative. Therefore, there are no indications that the effects of commercially growing GMB151 Soybean would cause significant impacts to microorganisms if the Preferred Alternative were selected.

5.5.9 Biodiversity: No Action Alternative

Under the No Action Alternative, GMB151 Soybean would remain under APHIS regulation. The availability of biotech, conventional and organic soybeans would not change. Agronomic practices used for soybean production and yield optimization, such as tillage, the application of agricultural chemicals (pesticides and fertilizers), timing of planting, row spacing, and scouting for pest infestations would be expected to continue unchanged. Agronomic practices that benefit biodiversity both on cropland (e.g., intercropping, agroforestry, crop rotations, cover crops, and no-tillage) and on adjacent non-cropland (e.g., woodlots, fencerows, hedgerows, and wetlands) would also remain the same.

5.5.10 Biodiversity: Preferred Alternative

A determination of nonregulated status of GMB151 Soybean is not expected to result in changes in current soybean cropping practices that may impact biodiversity. Trials conducted by BASF showed no differences between GMB151 Soybean and other biotech and conventional soybean in growth, reproduction, or interactions with pests and diseases that might impact biodiversity. Similar to the No Action Alternative, weeds within fields of GMB151 Soybean could be managed using mechanical, cultural, and chemical control. Growers would determine the best method necessary to manage pests based on individual needs. The environmental risks of pesticide use are assessed by the EPA in the pesticide registration process and are regularly reevaluated by the EPA as part of its reregistration process under FIFRA. Pesticide use in accordance with label instructions established by the EPA would not result in unreasonable risks

to the environment. Under the Preferred Alternative, potential impacts to biodiversity from runoff, spray drift, and volatilization of agricultural chemicals such as pesticides, herbicides, and fungicides are not expected to be substantially different from those associated with the No Action Alternative.

Possible risks to biodiversity from the production of biotech crops include the disturbance of biosystems, including the agroecosystem, and permanent loss or changes in species diversity or the genetic diversity within a species (Snow et al. 2005). The intensive farming practices associated with agricultural lands limit the diversity of plants and animals (Lovett, Price, and Lovett 2003).

Diversity in adjacent natural areas, and those areas established to promote biodiversity (e.g., woodlots, fencerows, hedgerows, and wetlands) tend to have greater biodiversity than does cropland. Agronomic practices for the production of GMB151 Soybean are not expected to change from those currently used for other commercially available biotech and conventional soybean varieties. Because of cost effectiveness, control measures for SCN rely mostly on planting resistant varieties rather than making nematicide applications to soybeans. Most nematicides are used for soybean seed treatments, which would not change if the Preferred Alternative is selected. Therefore, impacts on species diversity would be the same as or similar to those of the No Action Alternative. Agronomic practices commonly used to increase farm-scale biodiversity are also unlikely to change. As described previously for gene flow and weediness, GMB151 Soybean has no potential to produce naturally occurring, pollen-mediated gene flow and transgene introgression, so is not expected to affect genetic diversity, which could have an impact on biodiversity.

For GMB151 Soybean, the most important concern related to biodiversity is the possibility that expression of the Cry14Ab-1 protein might adversely impact non-target organisms. Residues of certain Cry proteins are known to persist in soil (Feng et al. 2011), but at levels so low as to be negligible in terms of their capacity to impact non-target organisms. EPA assesses risks of biopesticides, which include possible impacts of Cry proteins that are PIPs (US-EPA 2014), and has concluded that risks associated with the GMB151 Soybean Cry14Ab-1 protein are negligible, so does not require a food or feed tolerance (40 CFR 174.540) for it. Test results reported by BASF for off-target effect of the Cry14Ab-1 protein expressed by GMB151 Soybean indicate that there are no unacceptable risks to or impacts on non-target organisms (BASF 2020).

Based on the information summarized above, overall effects on biodiversity under the Preferred Alternative are expected to be the same as or so similar to the No Action Alternative, that they would be indistinguishable. Therefore, there are no indications that the effects of commercially growing GMB151 Soybean would cause significant impacts to biodiversity if the Preferred Alternative were selected.

5.6 ANIMAL HEALTH

5.6.1 Animal Feed—No Action Alternative

Under the No Action Alternative, GMB151 Soybean would continue to be regulated. Soybean-based animal feed derived from both conventional and those biotech varieties that are not regulated under 7 CFR 340 would continue to be available. Nonregulated biotech soybeans used as animal feed have previously been determined to not pose any risk to animal health.

5.6.2 Animal Feed—Preferred Alternative

Results of studies conducted by BASF confirmed that there are no differences in the quality of animal feed produced from GMB151 Soybean compared to feed derived from both conventional and those biotech varieties that are not regulated under 7 CFR 340 (BASF 2020). APHIS critically reviewed data provided by and information in the scientific literature cited by (BASF 2020), and concluded that a determination of nonregulated status of GMB151 Soybean would not alter the nutritional quality of animal feed derived from it.

Possible effects to livestock from a determination of nonregulated status for GMB151 Soybean are related to concerns about the potential health impacts on animals from consuming soybean products containing residues of the Cry14Ab-1 protein. Safety evaluations conducted by BASF followed Codex Alimentarius Commission procedures recommended to assess potential adverse impacts to animals and humans.

Safety studies included: (1) characterization of the physicochemical and functional properties of the Cry14Ab-1 protein; (2) quantification of the Cry14Ab-1 protein levels in plant tissues; (3) comparison of the amino acid sequence of the Cry14Ab-1 protein in GMB151 Soybean to known allergens, gliadins, glutenins, toxins, and other biologically-active proteins known to have adverse effects on mammals; (4) evaluation of the digestibility of the Cry14Ab-1 protein in simulated gastric and intestinal fluids; (5) documentation of the presence of related proteins in several plant species currently consumed; and (6) investigation of the potential mammalian toxicity through an oral gavage assay. The Cry14Ab-1 protein was determined to have no amino acid sequence similar to known allergens, lacked toxic potential to mammals, and was degraded rapidly and completely in gastric fluid (85 FR 35008).

As part of its regulatory compliance process, BASF submitted supporting data and EPA concluded that the Cry14Ab-1 is not toxic or allergenic to mammals, so it does not require a food or feed tolerance (40 CFR 174.540) for Cry14Ab-1 protein when it is a plant-incorporated protectant (US-EPA 2020). Pesticide residue tolerances for pesticides listed in 40 CFR § 180 establish residue limits for soybean forage, hay, hulls, and seed (US-EPA, 2010a) that are protective of livestock and human health.

Based on the above information, there are no expected hazards associated with the consumption of GMB151 Soybean by animals, so it is unlikely to pose a hazard to any livestock species. The results of studies conducted by BASF confirmed that the crops containing these proteins can be safely used as animal feed (BASF 2020). There are no differences in feed safety between the GMB151 Soybean and other varieties currently available under the No Action Alternative. Therefore, APHIS has concluded that a determination of nonregulated status of GMB151 Soybean would not have significant impacts on animal feed or the health of livestock that consume it. Overall impacts of selecting the Preferred alternative would be the same as or similar to those of the No Action Alternative.

5.7 HUMAN HEALTH

5.7.1 Public Health: No Action Alternative

Under the No Action Alternative, GMB151 Soybean would remain under APHIS regulation. Human exposure to existing biotech and conventional soybean varieties would not change under this alternative. The same EPA-registered pesticides would continue to be used for pest

management in conjunction with both biotech and conventional soybean cultivation. The environmental risks of pesticide use are assessed by the EPA in its pesticide registration process and are regularly reevaluated under its reregistration process to ensure that pesticides do not cause unreasonable adverse effects on human health or the environment.

The EPA also establishes maximum residue limits for pesticides that are referred to as tolerances (US-EPA 2010a). Tolerances represent the maximum amount of pesticide residues that can remain on or in food or feed. These levels have been carefully determined using scientific data to establish exposure levels that will not cause adverse health effects. The EPA sets tolerances for pesticides to meet FQPA safety standards for the U.S. population and designated sensitive populations (i.e., infants and children) to ensure that there is a reasonable certainty of no harm to the general population and any subgroup. Food or feed may not be distributed for consumption if it contains residues of one or more pesticides that exceed a tolerance. Food and feed with pesticide residues that exceed a tolerance are considered adulterated and may be seized. The FDA and USDA monitor foods for pesticide residues and enforce tolerances (USDA-AMS, 2018). For more details about tolerances, go to the EPA web site at:

<http://www.epa.gov/pesticides/bluebook/chapter11.html>

5.7.2 Public Health: Preferred Alternative

BASF conducted safety evaluations using Codex Alimentarius Commission procedures to assess any potential adverse impacts to humans or animals resulting from environmental releases and consumption of GMB151 Soybean (BASF 2020). The gene regulating the expression of the Cry14Ab-1 protein, and the Cry14Ab-1 protein itself were determined to have no amino acid sequences similar to known allergens and lacked toxic potential to mammals. As part of its regulatory compliance process, BASF submitted supporting data, and EPA concluded that risks associated with the GMB151 Soybean are negligible, so it does not require a food or feed tolerance (40 CFR 174.540) for Cry14Ab-1 protein when it is a plant-incorporated protectant. Pesticide residue tolerances for pesticides listed in 40 CFR § 180 establish residue limits for soybean forage, hay, hulls, and seed (US-EPA, 2010a) that are protective of livestock and human health.

Based on this information, including field and laboratory data and scientific literature provided by BASF (2017, 2018) and safety data for other biotech soybeans, APHIS concluded that there would not be any adverse human health effects from a determination of nonregulated status of GMB151 Soybean. Human consumption of food products derived from GMB151 Soybean would not be different from those derived from conventional or biotech soybean varieties that are not regulated under 7 CFR 340. Likewise, human consumption of food products derived from livestock feed derived from GMB151 Soybean would not be different from products from livestock feed from conventional or biotech soybean varieties that are not regulated under 7 CFR 340 because no significant impacts on animal health were identified for GMB151 Soybean (see the preceding section for details). Impacts from choosing the Preferred Alternative would be the same as those for the No Action Alternative, so would not have significant impacts on human health.

5.7.3 Worker Safety: No Action Alternative

The availability of biotech, conventional, and organic soybeans would not change as a result of the continued regulation of GMB151 Soybean. Because of cost effectiveness, control measures for SCN rely mostly on planting resistant varieties rather than making nematicide applications to

soybeans. Most nematicides are used for soybean seed treatments, which would not change if the Preferred Alternative is selected. Therefore, agronomic practices used for soybean production, such as the application of agricultural chemicals (pesticides and fertilizers), would be expected to continue unchanged. Growers will continue to choose agronomic practices based on weed, insect and disease pressures, cost of seed and other inputs, technology fees, human safety, potential for crop injury, and ease and flexibility of the production system (Farnham 2001; Heiniger 2000; University of Arkansas 2006). Worker safety is taken into consideration by the EPA in the pesticide registration and reregistration processes. Pesticides are regularly reevaluated by the EPA for each pesticide to maintain its registered status under FIFRA. Occupational Safety and Health Administration requires all employers to protect their employees from hazards associated with pesticides and herbicides. When used according to label directions, pesticides can be used with a reasonable certainty of no harm to human health and no unreasonable risks to the environment.

The EPA Worker Protection Standards (WPS) (40 CFR Part 170) implement protections for agricultural workers, handlers, and their families. These WPS requirements were revised in 2015 to implement even stronger standards that became effective on January 2, 2017, with further revisions implemented on January 2, 2018. The EPA has also issued guidance for farm managers about how to implement the new standards (US-EPA 2017c).

5.7.4 Worker Safety: Preferred Alternative

A determination of nonregulated status of GMB151 Soybean is not expected to result in changes in current soybean cropping practices. Similar to the No Action Alternative, it is expected that EPA-registered pesticides, fertilizers, and other chemicals that are currently used for soybean production would continue to be used by growers. The EPA's core pesticide risk assessment and regulatory processes ensure that each registered pesticide continues to meet the highest standards of safety including all populations of non-target species and humans, and if used in accordance with the label, can be demonstrated to pose a reasonable certainty of no harm to humans, including those employed in agricultural and farm-related occupations, and no unreasonable adverse effects to the environment. The EPA WPS (40 CFR Part 170) would be the same as that described for the No Action Alternative. Growers are required to use pesticides in accordance with the application instructions provided on the EPA registration label for each pesticide product label, and follow the additional guidance (US-EPA 2017c) issued by the EPA to ensure farm worker safety. These label restrictions are legally enforceable and are enforced by EPA and the states (Federal Insecticide, Fungicide, and Rodenticide Act 7 USC 136j (a)(2)(G) Unlawful Acts).

Exposure to GMB151 Soybean under the Preferred Alternative is not expected to pose any changes to existing human health risks. Based on the above information, occupational health and safety risks under the Preferred Alternative are expected to be the same as or similar to those associated with the No Action Alternative.

5.8 SOCIOECONOMIC IMPACTS

5.8.1 Domestic Economic Environment: No Action Alternative

Under the No Action Alternative, GMB151 Soybean would remain under APHIS regulation. Growers and other parties who are involved in production, handling, processing, or consumption of soybeans would continue to have access to other nonregulated biotech and conventional soybean varieties. Domestic growers would continue to utilize biotech and conventional soybean varieties based upon availability and market demand. Current production practices, using biotech varieties to optimize yield and reduce production costs would not change if the No Action Alternative is selected. Grower net returns are estimated to increase approximately 24% from \$303 to \$375 per acre by the end of the period, 2013/2014 to 2021/2022, despite an estimated 3% rise in seed and residual costs, and 10.3% rise in overall per acre cost of production.

5.8.2 Domestic Economic Environment: Preferred Alternative

In field tests conducted by BASF, the performance and composition of GMB151 Soybean was determined not to be substantially different from that of other soybean varieties. If no longer regulated, GMB151 Soybean would be subject to the same variables that affect yield of other biotech and conventional soybean varieties such as weather, timing and density of planting, and soil nutrients. Growers are familiar with yield improvements using increased yield varieties obtained through traditional breeding techniques, and more recently, increased yields from better weed control and disease resistance in biotech soybean varieties. As noted previously, soybean yields have increased steadily since 1924 (USDA-NASS 2012d).

GMB151 Soybean would be expected to be adopted by growers who are already growing biotech soybeans. The rate of adoption would depend on SCN distribution and this equates to increased yield and ultimately profitability after production costs. It is unlikely the availability of GMB151 Soybean would significantly impact the domestic economic environment. Past and recent increases in U.S. soybean acreage have occurred as growers replaced other crops with soybeans; not by bringing new lands into production. U.S. total cropland has remained relatively stable since the mid-20th century. Since 94% of U.S. soybean acreage is planted with biotech soybean varieties (USDA-NASS 2018c), it is likely that GMB151 Soybean would only replace other varieties of biotech soybean grown on existing cropland. Historically, soybean yields have been increasing for decades. In more recent times, this has resulted from conventional cross-breeding of high yielding varieties with biotech varieties, and applying improved management practices. Growers would have to make an independent assessment as to whether the benefits of GMB151 Soybean would offset higher seed cost.

Based upon the preceding information, the potential domestic economic impacts from a determination of nonregulated status for GMB151 Soybean would be similar to or no different than those under the No Action Alternative, so there would not be any significant impacts from selecting the Preferred Alternative.

5.8.3 International Trade Economic Environment: No Action Alternative

If the No Action Alternative is selected, GMB151 Soybean would continue to be regulated by APHIS. It is unlikely the current soybean market trade trends would change if GMB151 Soybean remained a regulated article. U.S. soybeans will continue to be a major contributor to global soybean production, and the United States will continue to be major exporter and supplier in the international market.

5.8.4 International Trade Economic Environment: Preferred Alternative

There are several factors that influence worldwide prices for oilseed, including soybean and its products. These include energy costs, fluctuations in currency exchange rates, government policies, national population size, per capita income, global market conditions, and trends and practices in market trading and speculation (Trostle 2008b; Trostle 2008a; Irwin and Good 2009). These factors influence the value derived from soybeans. If this value increases, it gets distributed between consumers in the form of lower product prices and growers and distributors as increased profits.

Projections from current trends in U.S. production indicate that it is unlikely that U.S. soybean acreage will increase significantly, so if it became commercially available, GMB151 Soybean, would likely only replace other biotech soybean varieties, where SCN is a pest problem. Any impact on soybean market prices from the potential increased yield from GMB151 Soybean production would likely be negligible because GMB151 would be used primarily to prevent potential losses and maintain current yield levels. Therefore, it would not alter the value currently derived from U.S. soybean production, so would not have any significant impact on the international trade environment for U.S. exports of soybean products.

USDA projects that from 2013/2014 to 2021/2022, the national annual average of U.S. soybean yield is expected to increase approximately 8% without expanding acreage, but the U.S. average farm price per bushel of soybean is predicted to vary only between \$10.30 and \$11.35. Grower annual net returns per acre are estimated to increase on average approximately 24% over the same period, despite an estimated approximately 3% rise in seed and residual costs, and 10.3% rise in overall per acre cost of production (USDA-OCE 2012). Adoption of GMB151 Soybean would likely be gradual at a pace equal to the extent growers find value in another higher than average yielding soybean variety.

It is not expected that if available, GMB151 Soybean would affect world attitudes towards biotech crops. The adventitious presence of biotech products in other food or feed continues to be a concern of some international trade partners, but if available, GMB 151 Soybean would not be expected to change the acceptability of U.S. soybean exports because most U.S. soybeans are already produced from biotech varieties. In its petition, BASF has asserted its commitment to stewardship to meet applicable regulatory requirements for GMB151 Soybean in the country of intended production and for key import countries to ensure compliance, maintain product integrity, and assist in minimizing the potential for trade disruptions (BASF 2020).

Based on these considerations, the potential impacts to the trade economic environment from a determination of nonregulated status of GMB151 Soybean would be similar to or no different than those currently observed for the No Action Alternative. Therefore, a determination of nonregulated status of GMB151 Soybean is not expected to have any significant impact on total annual U.S. soybean production, and no significant impacts on the international trade economic environment affecting U.S. soybean exports.

6 CUMULATIVE IMPACTS

The CEQ regulations define a cumulative impact (40 CFR part 1508, Section 1508.7 as follows: *Cumulative impact* is the impact on the environment which results from the incremental impact of the action when added to other past, present, and reasonably foreseeable future actions regardless of what agency (Federal or non-Federal) or person undertakes such other actions. Cumulative impacts can result from individually minor but collectively significant actions taking place over a period of time.

In Chapter 4, APHIS analyzed individually the environmental consequences that may result from choosing either the No Action or Preferred Alternative. As part of that analysis, APHIS considered the potential direct impacts on those aspects of the human environment related to the petition, and any subsequent commercial production of GMB151 Soybean. In this chapter, potential cumulative impacts of a decision about the regulatory status of GMB151 Soybean are reviewed.

6.1 METHODOLOGY AND ASSUMPTIONS

For its analysis, APHIS assumed that if no direct impacts on a resource area were identified as part of its analyses of impacts from a regulatory decision for GMB151 Soybean under Environmental Consequences (Chapter 4), then there cannot be any cumulative impacts on that resource area. When possible, effects were quantified for the analysis to measure the potential to cause significant impacts; otherwise qualitative assessments were made.

APHIS limited its cumulative impacts analysis to the areas in the United States where soybeans are commercially grown. The potential for significant impacts from effects identified by the Agency as being reasonably foreseeable cumulative impacts were analyzed under the assumption that farmers, who grow transgenic soybeans conventional or organic soybeans would continue to use the same BMPs they currently use if GMB151 Soybean were no longer regulated.

Biotech soybeans grown in the United States are frequently produced from varieties that have multiple biotech traits. Such varieties are referred to as “stacked” hybrids, and some have been developed using the same recombinant DNA techniques used to produce single-trait biotech varieties. These are subject to APHIS regulation under 7 CFR 340 until a determination of nonregulated status is made. However, stacked hybrids can also be developed using traditional cross breeding to combine biotech traits from different biotech varieties, including those that have previously been evaluated individually by APHIS and have been determined to have nonregulated status. Therefore, if APHIS makes a determination of nonregulated status for GMB151 Soybean, it is likely that the traits from GMB151 will be bred into other soybean varieties that are not regulated by the Agency.

If it is no longer regulated, traditional plant breeding methods could be used to develop stacked trait hybrids between GMB151 Soybean and other biotech soybean varieties that have previously been determined to have nonregulated status. These include, for example, varieties that are resistant to herbicides and certain insect pests, and those expressing modified nutritional profiles. If GMB151 Soybean were no longer regulated, cross breeding it with other biotech soybean varieties to produce stacked trait varieties is a reasonably foreseeable action.

6.2 CUMULATIVE IMPACTS: SOYBEAN AGRICULTURAL PRODUCTION

Except for its resistance to SCN and isoxaflutole, GMB151 Soybean is agronomically and compositionally similar to conventional and biotech soybean varieties that are not regulated by APHIS (BASF 2020).

Neither the No Action Alternative nor the Preferred Alternative are expected to change total U.S. soybean acreage or cause any shift in the regions where soybean crops are currently grown for grain or seed. Total U.S. cropland has remained relatively steady since the middle of the last century. Increases in soybean acreage have occurred during this period, but this is the result of replacing other crops on existing cropland (USDA-ERS 2011c). Future increases in soybean production will likely be from improved soybean varieties and production methods that increase yield rather than expand production area (OECD-FAO 2008). Most soybeans currently grown in the United States are biotech varieties (USDA-ERS 2012a). Long-term projections indicate that soybean acreage will remain level until 2028 (USDA-OCE 2018). If it were no longer regulated, it is expected that GMB151 Soybean would replace other similar biotech soybean varieties and would not increase current total U.S. acreage or change the areas where soybeans are grown. Therefore, there would be no difference in the environmental impacts of selecting either the Preferred Alternative or the No Action Alternative on total U.S. soybean acreage or the locations where soybeans are grown for seed or grain, so there would not be any associated cumulative impacts.

Based upon past and current trends, the addition of another biotech soybean variety would not have any impacts on the ability of organic soybean producers to maintain their current market share. U.S. organic soybean production acreage has fluctuated somewhat from year to year between 82,143 and 126,000 acres during the period, 1997-2011 (USDA-ERS 2010a; USDA-NASS 2012a). This represented about 0.09% of total U.S. soybean acreage in 2011 (USDA-NASS 2012a). The most recent data puts U.S. organic soybean acreage at 124,591 in 2016, compared to 94,841 in 2015 (USDA-AMS 2017), which indicates little fluctuation from the previously reported trends. Availability of another biotech soybean variety, such as GMB151 Soybean would not be expected to alter any impacts that biotech soybeans currently have on organic soybean production, so no cumulative impacts will be associated with selecting either the No Action Alternative or the Preferred Alternative.

Studies by BASF demonstrated that, in terms of agronomic characteristics and cultivation practices such as tillage, fertilization, irrigation, pest and disease control measures, crop rotation, and irrigation, GMB151 Soybean is similar to other biotech currently grown. Therefore, GMB151 Soybean production is likely to require the same fertilizer inputs as other high yield soybean systems utilizing conventional or biotech soybean varieties. Supplementing soybean crops with nutrients is not uncommon (USDA-NASS 2018e), and BMPs that include soil fertility testing and supplementation recommendations to optimize nutrient replacement and maximize yield potential are widely used (Pedersen 2008; CAST 2009; Mallarino et al. 2011; Silva 2011; Snyder 2000; Specht et al. 2006).

If GMB151 Soybean were no longer regulated, it would be grown in rotation with other crops such as corn or wheat, no differently than any other high yield soybean varieties. In two-year corn-soybean rotations, enough potassium and phosphorus amendments are commonly applied to the corn crop to sustain the soybean crop the following year without additional supplementation (Bender et al. 2013). However, recent research has shown higher yielding corn varieties may

remove more phosphorous than is applied on average, and soil fertility testing prior to soybean planting is recommended (Bender et al. 2013). Potassium and phosphorus are commonly applied annually where soybeans are not rotated, which is the predominant practice in the South (Heatherly 2012). Testing soil fertility and supplementing nutrients is widely recommended and used in soybean production to achieve optimal yield potential (Pedersen 2008; CAST 2009; Mallarino et al. 2011; Silva 2011; Snyder 2000; Specht et al. 2006). There is no evidence that cultivation of GMB151 Soybean would require changes to any of these fertilization practices in soybean production.

Since the agronomic requirements and cultivation practices for GMB151 Soybean are the same as those for other high yield conventional and biotech soybean varieties currently grown in the United States, any environmental impacts from current soybean production in the United States would not be altered if GMB151 Soybean were no longer regulated by APHIS. Because there would be no changes in impacts, APHIS concluded that there would not be any cumulative impacts associated with selecting either the No Action Alternative or the Preferred Alternative.

6.3 CUMULATIVE IMPACTS: PHYSICAL ENVIRONMENT

Current agronomic practices for soybeans are important sources of impacts on the physical environment. Agronomic practices that have the potential to impact soil, water, and air quality, such as tillage, agricultural inputs (fertilizers and pesticides), and irrigation would not change following a determination of nonregulated status of GMB151 Soybean because GMB151 Soybean is agronomically and morphologically similar to other biotech and conventional soybeans. Other practices that benefit these resources, such as contouring, use of cover crops to limit the time soil is exposed to wind and rain, crop rotation, and windbreaks would also remain the same under both Alternatives.

Because of its similarity to other commercially available biotech soybean varieties, and the likelihood that GMB151 Soybean would only replace other similar varieties, it would not change the acreage or locations of current U.S. soybean production. Therefore, any existing impacts on water, soil, and air quality from current U.S. soybean production practices would not change if GMB151 Soybean were no longer regulated by APHIS. As a result, there would be no difference in impacts from choosing either the Preferred Alternative or the No Action Alternative, and APHIS concluded that selection of either the No Action Alternative or the Preferred Alternative would not result in any cumulative impacts on the physical environment.

6.4 CUMULATIVE IMPACTS: BIOLOGICAL RESOURCES

Commercial cultivation of GMB151 Soybean, or progeny derived from it would not be expected to contribute in a cumulative manner to impacts on biological resources any differently than that of cultivation of current soybean varieties. GMB151 Soybean is both agronomically and compositionally similar to other conventional and biotech soybean varieties that are not regulated (BASF 2020). Therefore, if it were no longer regulated, GMB151 Soybean would not alter current U.S. soybean agronomic practices, so the impacts of those practices on animal and plant communities, microorganisms, and biodiversity would not change.

The traits for protection from SCN and resistance to isoxaflutole in GMB151 Soybean do not

exert any influence on its weediness, so they do not represent any weediness risks that differ from other currently available soybean varieties. If present as a volunteer in crops rotated with soybeans, GMB151 Soybean would not be difficult to control because soybean seeds rarely remain viable the following season and are easily managed by hand weeding, cultivation, or herbicide applications.

The reproductive characteristics of GMB151 Soybean are also equivalent to other biotech and conventional soybean varieties (BASF 2020). Since soybean plants are mostly self-pollinating and have limited ability to disperse pollen, there is little or no potential for cross pollination of GMB151 Soybean with other soybean varieties. Since no feral or weedy species of soybean exist in the United States (Ellstrand, Prentice, and Hancock 1999; OECD 2000), GMB151 Soybean poses no potential for either naturally occurring, pollen-mediated gene flow or transgene introgression. The risk of gene flow and weediness of GMB151 Soybean is no greater than that of other conventional and nonregulated biotech soybean varieties.

The maximum amount of herbicide active ingredient applied to varieties of GMB151 Soybean stacked with additional HR traits would be limited by the EPA registration for the product used. Other isoxaflutole-resistant biotech soybean varieties that are not regulated are currently available to growers. GMB151 Soybean would only replace these as another alternative to growers, so isoxaflutole usage would not change. As with other herbicides used for soybean cultivation, isoxaflutole used in accordance with EPA registration requirements would continue to ensure that there are no unacceptable risks to non-target organisms or the environment. Since there is no anticipated increase in U.S. soybean acreage in the foreseeable future, and no anticipated change in the acreage of isoxaflutole-resistant soybeans if GMB151 Soybean were no longer regulated, total isoxaflutole usage is unlikely to change because current EPA-labeled uses of isoxaflutole are expected to remain the same. Possible impacts on biological resources from the application of pesticides to stacked GMB151 Soybean varieties would not be any different from those resulting from biotech soybeans, when used in accordance with label instructions.

Results of the Agency's analysis, which is summarized above, support the conclusion that there would be no impacts on biological resources if GMB151 Soybean were no longer regulated. In addition, existing impacts on biological resources associated with current soybean cultivation in the United States would not be altered. Because potential direct impacts on biological resources do not significantly differ between the No Action and Preferred Alternatives, there are no reasonably foreseeable cumulative impacts that would derive from the commercial cultivation of GMB151 Soybean or its progeny.

6.5 CUMULATIVE IMPACTS: ANIMAL FEED AND HUMAN HEALTH

Food and feed derived from biotech soybeans must be in compliance with all applicable legal and regulatory requirements. Results of previous analyses presented in Chapter 4 in the Human Health and Animal Feed sections described how the Agency considers the EPA pesticide registration process in APHIS EAs for biotech organisms that affect how pesticides are used. Under the authorizations of FIFRA, the EPA assesses environmental risks of pesticides, and once they are registered for use, regularly reevaluates them. As part of the registration process, the EPA considers human health impacts from the use of pesticides and must determine that the pesticide will not cause unreasonable adverse effects on human health.

Worker safety is considered by the EPA in the pesticide registration and reregistration processes. If GMB151 Soybean is determined to have nonregulated status and is subsequently stacked in biotech soybean varieties with other HR traits, the total amount of herbicides that may be applied would be limited to the per application and per year rates established by the EPA. When used in compliance with EPA registration label specifications, pesticides present minimal risk to human health and worker safety. Pesticide residue tolerances for pesticides listed in 40 CFR § 180 establish residue limits for soybean forage, hay, hulls, and seed (US-EPA 2010a) that are protective of livestock and human health. EPA has also concluded that risks associated with the GMB151 Soybean are negligible, so it does not require a food or feed tolerance (40 CFR 174.540) for the Cry14Ab-1 protein when it is a plant-incorporated protectant (CFR 174.522).

APHIS has determined that there are no past, present, or reasonably foreseeable actions that would aggregate with the effects of a determination of nonregulated status for GMB151 Soybean that would adversely impact human health or animal feed. Based on its review of available information, APHIS has concluded that there is no evidence that any impacts that may result from a determination of nonregulated status for GMB151 Soybean would compound to cause significant cumulative impacts to human health or animal feed. Therefore, selection of either the No Action Alternative or the Preferred Alternative would not result in any cumulative impacts on human health and animal feed.

6.6 CUMULATIVE IMPACTS: SOCIOECONOMICS

The increase in U.S. soybean acreage during the past few decades has been associated with an increase in double cropping and the replacement of other crops with soybeans, not by bringing new lands into production (USDA-ERS 2011d). If it were no longer regulated, GMB151 Soybean would likely replace conventional or other biotech soybean varieties on existing cropland. Most (94%) U.S. soybean acreage is currently planted with biotech soybean varieties (USDA-NASS 2018c), and combined trials have confirmed that GMB151 Soybean is phenotypically and agronomically similar to other soybean conventional and biotech varieties that are not regulated by APHIS (BASF 2020), so adding a new soybean variety would not impact the domestic economic environment. Since impacts to the domestic economic environment would not change, there would not be any cumulative impacts on the domestic economic environment associated with selecting either the No Action Alternative or the Preferred Alternative.

Soybean yields have been increasing for decades. During the past few decades this has resulted from the development of conventionally bred and biotech varieties with high yielding traits, and improved management practices (Pedersen 2008; Specht et al. 2006). U.S. soybean acreage is projected to remain level at least until 2028, but with an anticipated 8% per acre yield gain. Despite potential increased production, prices for soybeans per bushel are not expected to change appreciably (remaining between \$10.30 and \$11.35 per bushel), but annual production net value is expected to increase (USDA-OCE 2012). Soybean supply is a function of the amount of acreage planted and crop yield. While domestic soybean yield has recently increased primarily without increasing production acreage, demand for soybean products has also increased, offsetting any downward pressure on farm soybean prices from any potential over supply (NRC 2010).

Nonregulated GMB151 Soybean is not expected to adversely impact the current trends affecting the seed, feed, or food trade and may have a negligible impact from increased yields. Apart from SCN and isoxaflutole resistance, GMB151 Soybean is essentially indistinguishable from other soybean varieties in terms of agronomic, morphologic, and compositional characteristics (BASF 2020). Increased farm productivity if GMB151 Soybean were no longer regulated may increase U.S. competitiveness in the global economy, although many other factors affect worldwide prices for soybean, including energy costs, monetary exchange rates, government policies, population size and growth rate, per capita income, global market conditions, and trends and practices in market trading and speculation (Trostle 2008b; Trostle 2008a; Irwin and Good 2009). How any value derived from GMB151 Soybean is distributed between consumers in the form of reduced prices and growers as increased profits would be subject to these factors. Based upon the above information, any impact to soybean market prices from production of GMB151 Soybean would be negligible.

Another consideration is that since GMB151 Soybean is agronomically and compositionally similar to other commercially available soybean varieties, there would be no major changes to agronomic inputs or practices if it were determined to have nonregulated status. Like any other high-yield soybean variety, GMB151 Soybean has been shown to deplete potassium and phosphorous in soil more than other varieties. But as described above, supplementation of these nutrients in soybean production is not uncommon, and soil fertility testing and supplementation as indicated by tests and known crop soil nutrient removal rates is widely recommended in soybean production to achieve yield potential (Pedersen 2008; CAST 2009; Mallarino et al. 2011; Silva 2011; Snyder 2000; Specht et al. 2006). Advances in soybean yield have been attributed to development of conventionally bred higher yield varieties that also have transgenic herbicide and or other resistance traits.

Another possible way that the soybean socioeconomic environment might be impacted by GMB151 Soybean would be if it were stacked with other biotech soybean traits that altered production costs of the agronomic practices used to produce soybeans. Although conservation tillage is used in conjunction with soybean production, there is an increasing trend to use strip tillage to support adequate soil fertility (Fernandez and White 2012).

More than half of the U.S. soybean crop acreage is in a two-year rotation with corn or wheat. Fertilization of a preceding corn crop is usually made at a level that supports the following soybean crop. However, recent research has shown that high yielding transgenic HR and IR corn varieties may remove more phosphorous than is applied, so soil testing prior to planting of any soybean variety is recommended (Bender et al. 2013). In the South, where soybean crops are not rotated, fertilizer is applied annually (Heatherly 2012). Where crop rotation is practiced (e.g., corn-soybean-wheat rotation) phosphorous applications to replace nutrients removed and balance nutrient inputs are recommended (PPI 2003).

If GMB151 Soybean were no longer regulated, it would not be expected to change the choices of production systems soybean growers currently use (i.e., biotech, conventional, or organic). Organic soybean growers in particular supply a niche market that is a small portion of the U.S. market. As mentioned above regarding cumulative impacts on organic soybean production, adding biotech varieties to the domestic market is not related to the ability of organic production systems to maintain their market share.

It is possible that GMB151 Soybean would not be approved for import into other countries. Because the United States and other countries already have access to other soybean varieties, and GMB151 Soybean would only replace other biotech varieties already in the marketplace, its availability only to U.S. producers would not likely impact the economic trade environment. In 2011/2012, 42% of domestically produced U.S. soybean was dedicated to the export market (USDA-ERS 2012b, 2016). If GMB151 Soybean were not approved for import by other countries, but were not regulated in the United States, it would not likely affect the supply of U.S. soybean eligible for export to other countries. In contrast, if it were approved in the United States and for import by other countries, because of its similarity to other soybean varieties, the likelihood is that it would replace other such varieties, so would not increase the acreage or locations of soybean production in the United States. In its petition, BASF has also asserted its commitment to stewardship to meet applicable regulatory requirements for GMB151 Soybean in the country of intended production and for key import countries to ensure compliance, maintain product integrity, and assist in minimizing the potential for trade disruptions (BASF 2020). Therefore, it's unlikely GMB151 Soybean would impact the supply of U.S. soybean available for export, there would be no potential cumulative impacts related to past and present actions if either the Preferred Alternative or the No Action Alternative is selected.

7 THREATENED AND ENDANGERED SPECIES

The Endangered Species Act (ESA) of 1973, as amended, is a far-reaching wildlife conservation law. Congress passed the ESA to prevent extinctions facing many species of fish, wildlife and plants. The purpose of the ESA is to conserve threatened and endangered species (T&E) and the habitats on which they depend as key components of America's natural heritage. The U.S. Fish & Wildlife Service (USFWS) and the National Marine Fisheries Service (NMFS) together comprise "the Services" and implement the ESA by working in coordination with other federal, state, and local agencies, Tribes, non-governmental organizations, and private citizens.

Before a plant or animal species can receive protection under the ESA, it must be added to the federal list of T&E animal and plant species. T&E species are those plants and animals at risk of becoming extinct throughout all or part of their geographic range (endangered species) or those at risk of becoming endangered in the foreseeable future throughout all or a significant portion of their range (threatened species).

The Services add a species to the list when they determine the species to be endangered or threatened by any of the following factors:

- The present or threatened destruction, modification, or curtailment of its habitat or range;
- Overutilization for commercial, recreational, scientific, or educational purposes;
- Disease or predation;
- Inadequacy of existing regulatory mechanisms; and
- Natural or manmade factors affecting its survival.

Once a species is added to the list, in accordance with the ESA, protective measures apply to the species and its habitat. These measures include protection from adverse effects of federal activities.

7.1 ESA REQUIREMENTS FOR FEDERAL AGENCIES

Section 7(a)(2) of the ESA requires consultation with USFWS and/or the NMFS. Federal agencies must consult to ensure that any action they authorize, fund, or carry out "... is not likely to jeopardize the continued existence of any endangered species or threatened species or result in the destruction or adverse modification of habitat of such species . . .". It is the responsibility of the federal agency taking the action to assess the effects of the action and consult with the USFWS and NMFS if it is determined that the action "may affect" listed species or designated critical habitat (a process known as a Section 7 Consultation).

To facilitate the development of its ESA consultation requirements, APHIS met with the USFWS from 1999 to 2003 to discuss factors relevant to APHIS' regulatory authority and effects analysis practices for petitions for nonregulated status of biotech crop lines. By working with USFWS, APHIS developed a process for conducting an effects determination consistent with the Plant Protection Act (PPA) of 2000 (Title IV of Public Law 106-224). APHIS uses this process to fulfill its obligations and responsibilities under Section 7 of the ESA for biotechnology regulatory actions.

As described in Chapters 1 and 2, the APHIS regulatory authority over biotech organisms under the PPA is limited to those instances when there are reasons to believe a biotech organism could

pose a plant pest risk, or when the Agency does not have sufficient information to determine that a biotech organism is unlikely to pose a plant pest risk. BASF has requested that APHIS determine that GMB151 Soybean is not a plant pest as defined by the PPA (BASF 2020; USDA-APHIS 2018). If APHIS concludes from its PPRA that GMB151 Soybean does not pose a plant pest risk, then it is not subject to the plant pest provisions of the PPA or the regulations of 7 CFR part 340, and the Agency has no authority to regulate it. For this EA, APHIS analyzed the potential effects of GMB151 Soybean on listed T&E species, their critical habitats, and proposed T&E species and their proposed critical habitats. For the analysis as part of this EA, APHIS thoroughly reviewed data related to the transgene/transgenic cultivar, GMB151 Soybean, and supporting data to determine possible effects on listed and proposed T&E species and critical habitat.

For each transgene/transgenic plant petition, APHIS considers the following:

- A review of the biology, taxonomy, and weediness potential of the crop plant and its sexually compatible relatives
- Characterization of each transgene with respect to its structure and function and the nature of the organism from which it was obtained
- A determination of where the new transgene and its products (if any) are produced in the plant and their quantity
- A review of the agronomic performance of the plant including disease and pest susceptibilities, weediness potential, and agronomic and environmental impact
- Determination of the concentrations of known plant toxicants (if any are known in the plant)
- Analysis to determine if the transgenic plant is sexually compatible with any T&E species or a host of any T&E species
- Any other information about the potential for an organism to pose a plant pest risk

APHIS met with USFWS officials on June 15, 2011, to discuss and clarify whether APHIS has any obligations under the ESA regarding analyzing the effects on T&E species that may occur from the use of pesticides associated with biotech crops. As a result of these joint discussions, USFWS and APHIS have agreed that it is not necessary for APHIS to perform an ESA effects analysis on pesticide use associated with biotech crops because EPA has both regulatory authority over the labeling of pesticides under FIFRA, and the necessary technical expertise to assess pesticide effects on the environment. APHIS has no statutory authority to authorize or regulate the use of pesticides by soybean growers. Transgenic plants that elicit plant-incorporated protectants (PIPs) are regulated by EPA under FIFRA. Under APHIS regulations (7 CFR 340), the Agency only has authority to regulate GMB151 Soybean and other biotech organisms if they pose a plant pest risk (7 CFR § 340.1). APHIS does not have jurisdiction to regulate other risks from biotech organisms like those from pesticides used on biotech organisms.

7.2 POTENTIAL EFFECTS OF GMB151 SOYBEAN ON T&E SPECIES AND CRITICAL HABITAT

APHIS evaluated the potential effects that a determination of nonregulated status of GMB151 Soybean may have, if any, on federally listed T&E species and species proposed for listing, as well as designated critical habitat, and habitat proposed for designation. As described in detail in this EA and in the petition (BASF 2020), BASF used recombinant DNA techniques to insert genes into soybean in its development of GMB151 Soybean. The genes inserted confer specific resistance to the nematode pest, *Heterodera glycines*, commonly referred to as the soybean cyst nematode (SCN), as well as resistance to HPPD herbicides, such as isoxaflutole. More details about the development of GMB151 Soybean are available for review in the BASF petition and the draft PPRA (BASF 2020; USDA-APHIS 2020).

Based on the information submitted by the applicant and reviewed by APHIS, GMB151 Soybean, with the exceptions of the nematode resistance and HPPD resistance, is agronomically and compositionally comparable to other biotech soybean with the same or similar traits and with conventional soybeans (BASF 2020).

APHIS has determined that the characteristics and cultivation practices required for GMB151 Soybean are indistinguishable from agronomic practices used to grow other soybean varieties. No changes in practices are expected if the Agency determines that it should no longer regulate the article. APHIS considered the potential for GMB151 Soybean to extend the range of soybean production and also the potential to extend agricultural production into new natural areas. Although GMB151 Soybean may replace certain other varieties of soybean that are cultivated currently, APHIS does not expect the introduction of GMB151 Soybean to result in new soybean acreage to be planted in areas of the United States that are not currently used. Therefore, the issues considered in this review focus on the potential of a determination of nonregulated status for GMB151 Soybean for effects on T&E species in the areas where soybeans are currently grown. APHIS obtained and reviewed the USFWS list (Appendix A) of T&E species (listed and proposed) from the USFWS Environmental Conservation Online System for all states and U.S. territories where soybeans are produced (USFWS 2021).

For its analysis on T&E plants and critical habitat, APHIS focused on: the agronomic differences between the regulated article and soybean varieties currently grown; the potential for increased weediness; and the potential for gene movement to native plants, listed species, and species proposed for listing.

For its analysis of potential effects on T&E animals, APHIS focused on the implications of exposure to the Bt derived Cry protein expressed in GMB151 Soybean, and the ability of the transgenic plants to serve as a host for a T&E species. The novel genes and traits associated with GMB151 Soybean are summarized in Table 9.

Table 9. Proteins Produced by GMB151 Soybean that are Novel in Soybean

Regulated Article	Protein	Desired Phenotypic Effects
BASF GMB151 Soybean	Cry14Ab-1 protein encoded by <i>cry14Ab-1.b</i> gene from <i>Bacillus thuringiensis</i>	Toxicity to nematodes, including the soybean pest, soybean cyst nematode (SCN) (<i>Heterodera glycines</i>)
	HPPD-4 protein encoded by <i>hppdPf-4Pa</i> gene from <i>Pseudomonas fluorescens</i>	Resistance to herbicides with HPPD-inhibitor active ingredients such as isoxaflutole

Source: (BASF 2020)

7.2.1 Threatened and Endangered Plant Species and Critical Habitat

The agronomic and morphologic characteristics data provided by BASF (BASF 2020) were used in the APHIS analysis of the weediness potential for GMB151 Soybean and evaluated for the potential to impact T&E species and critical habitat. Agronomic studies conducted by BASF evaluated the performance of GMB151 Soybean in 11 field sites in different soybean growing regions of the United States.

Based on comparative assessments, agronomic performance of GMB151 Soybean was the same as or similar to conventional comparators and reference varieties. No substantive differences were detected between GMB151 Soybean and conventional soybeans varieties when comparing hardiness, persistence, seed dormancy, germination, or susceptibility to pests and diseases. These data supported the conclusion that the weediness potential of GMB151 Soybean is unchanged with respect to conventional soybean varieties and that GMB151 Soybean lacks weediness potential and plant pest risk (BASF 2020). These results and a subsequent analysis of the findings as reported in the Draft PPRA indicated that, other than the intended effect of nematode resistance to SCN and resistance to HPPD-inhibiting herbicides, there were no differences detected between GMB151 Soybean, and non-transgenic soybean varieties (BASF 2020; USDA-APHIS 2020). The analysis also found that there were no differences between GMB151 Soybean and other biotech crop varieties including transgenic soybeans that express the same enzyme for HPPD-inhibitor herbicide resistance that were previously evaluated by APHIS and determined to have no effect on T&E species (USDA-APHIS 2018, 2013, 2014).

Cultivated soybeans and its wild progenitor, *Glycine soja*, are native to Asia. Wild species in the subgenus *Glycine* are also found in Australia, including perennial species, but wild relatives of soybeans do not grow in the United States (Sherman-Broyles et al. 2014a). Soybeans do not have any of the attributes of successful weeds, and have been cultivated (USDA-APHIS 2014) worldwide without any reports of becoming weedy, or forming non-weedy persistent feral populations (see “Weediness” Section, Chapter 3 for more details). The Global Invasive Species Database confirms that there are no *Glycine* species or closely related taxa that are listed as invasive weeds (ISSG 2019). Any risk is further limited because soybeans seeds are not frost tolerant, so they do not overwinter well (Meyer and Badaruddin 2001). The seeds are heavy so exhibit poor seed dispersal characteristics, and plants do not reproduce vegetatively. Despite

these limitations, soybeans sometimes produce volunteers that interfere with corn or other rotational crops, but are not considered to be difficult to manage (Owen and Zelaya 2005b). Because soybeans are essentially autogamous, transgene movement by outcrossing attributable to pollen dispersal is negligible (Owen and Zelaya 2005b).

APHIS evaluated the potential of GMB151 Soybean to cross with listed T&E species. From a review of the list of T&E plant species (Appendix A) in the states where soybeans are grown, APHIS determined that GMB151 Soybean is not sexually compatible with any listed T&E plant species or plants proposed for listing. None of the listed plants are in the same genus as soybeans, nor are any known to cross pollinate with any species in the genus, *Glycine*.

Based on available information, APHIS concluded that if GMB151 Soybean will have no effect on T&E plant species or on critical habitats in the United States.

6.2.2 Threatened and Endangered Animal Species and Critical Habitat

Threatened and endangered animal species that may be exposed to the components of GMB151 Soybean would include those T&E species that inhabit soybean fields and potentially feed on GMB151 Soybean. To identify potential effects on T&E animal species, APHIS analyzed the risks to them from consuming GMB151 Soybean.

Exposure of T&E species is only likely if they occur in the areas where soybeans are grown because soybean plant parts (seeds, pollen, crop debris) are not transported long distances without human intervention. Therefore, the effects analysis evaluated those animal species that frequent soybean fields, or may be present in their immediate vicinity.

Nutrition, Composition, Agronomics, and Food Safety of GMB151 Soybean

BASF performed compositional analyses on GMB151 Soybean. BASF analyzed tissues of GMB151 Soybean and compared the compositional data to those of the conventional counterpart and reference varieties. The agronomic performance of GMB151 Soybean was evaluated at 11 field sites in different soybean growing regions of United States. The results of the compositional analysis and the comparative assessment demonstrated that GMB151 Soybean grain and forage grain are comparable to the non-transgenic reference varieties (BASF 2020).

BASF evaluated the potential toxicity and allergenicity of the HPPD and Cry14Ab-1 proteins expressed in GMB151 Soybean in particular and showed that they have no significant homology to known protein toxins and allergens. They are the same as or similar to proteins present in many plant species including edible plants, indicating a history of prior exposure and a history of safe use. Also, levels of HPPD and Cry14Ab-1 proteins in seed and forage tissues of GMB151 Soybean grown under field trial conditions were extremely low—low enough to be regarded as negligible (BASF 2020). BASF presented data for this conclusion in its consultation with the FDA for its molecular, compositional, nutritional, and food and feed safety assessment for GMB151 soybean (BASF 2020). In addition, on December 4, 2017, EPA published its final rule that exempts the HPPD-4 protein from the requirement of a food and feed tolerance, based on its analyses and determinations of absence of risk (82FR57137):

<https://www.govinfo.gov/content/pkg/FR-2017-12-04/pdf/FR-2017-12-04.pdf>

On June 8, 2020, EPA concluded that Cry14Ab-1 residue in or on soybean food and feed commodities is exempt from a tolerance if it results from a PIP in soybean plants (85 FR 35008; 40 CFR 174.540). These rules are based on extensive analysis of toxicity and exposure.

APHIS reviewed all available evidence (referenced above) and concluded that except for the *cry14Ab-1.b* and *hppdPf-4Pa* gene inserts and the proteins they express, GMB151 soybean is compositionally, agronomically, and nutritionally equivalent to conventional soybean varieties (BASF 2020; USDA-APHIS 2020). Therefore, except for the possible effects of its transgenic components, GMB151 Soybean would not have any impacts on any animal species including T&E animal species. APHIS also concluded from the available toxicity data that the transgenic components of GMB151 Soybean would be unlikely to have impacts on any animal species other than the intended target species (i.e., SCN). To eliminate any possibility of impacts on T&E animal species, APHIS considered in its effects analysis, possible risks from toxic effects on specific taxa of T&E species and the likelihood of their exposure.

Exposure to GMB151 Soybean and the Transgenic Genes and Proteins It Expresses

APHIS considered potential exposure to the *cry14Ab-1.b* and the *hppdPf-4Pa* gene inserts in the GMB151 Soybean genome, and to the Cry14Ab-1 and HPPD proteins expressed in the plant tissues of GMB151 Soybeans by those genes.

The *hppdPf-4Pa* gene was isolated from the bacterium, *Pseudomonas fluorescens* strainA32. *P. fluorescens* is a Gram-negative, rod-shaped, motile, asporogenous, aerobic bacterium. It is ubiquitous in the environment, including soil, water, and food (OECD 1997). Therefore, natural exposure of it and its gene products to a wide variety of animals including T&E species can be assumed. *P. fluorescens* has many beneficial uses in agriculture, human health, and bioremediation. It is not described as allergenic, toxic, or pathogenic to healthy humans and animals and has an overall history of safe use (BASF 2020). APHIS previously evaluated a transgenic soybean cultivar engineered to incorporate the same gene from *P. fluorescens* strainA32 to express the same HPPD protein and determined that it had no effects on T&E species (USDA-APHIS 2013).

All Cry proteins and *cry* genes have been isolated from *Bacillus thuringiensis*, a bacterium that is ubiquitous in the environment (OECD 1997). Therefore, natural exposure to *B. thuringiensis*, its genes and gene products to a wide variety of animals including T&E species can be assumed. None have been described as allergenic, toxic, or pathogenic to healthy humans and animals, having an overall history of safe use (Mendelsohn et al. 2003; Siegel 2001; Betz, Hammond, and Fuchs 2000; Farmer et al. 2017; Farias et al. 2014).

BASF results showed that the *cry14Ab-1.b* and *hppdPf-4Pa* genes in GMB151 Soybean do not result in any biologically meaningful differences between it and non-transgenic soybean varieties (BASF 2020), and show no evidence of health risks to humans and animals. Safety testing of the GMB151 Soybean Cry14Ab-1 and HPPD proteins showed that they are degraded rapidly and completely in simulated gastric fluid. Testing also showed that the GMB151 Soybean Cry14Ab-1 and HPPD proteins have no similarities to known allergens, and are not toxic to mammals. EPA has established a permanent exemption (82FR57137) from the requirement for a tolerance for the HPPD-4 protein expressed in all food commodities when used as an inert ingredient. In addition, EPA concluded on June 8, 2020 that *B. thuringiensis* Cry14Ab-1 protein residue in or on soybean food and feed commodities are exempt from a tolerance, when used as a plant-incorporated protectant (PIP) in soybean (85 FR 35008; 40 CFR 174.540).

The use of biotech crops expressing Cry proteins has been shown to reduce the use of broad spectrum insecticides without significant impacts on diversity of non-target insects (Romeis, Meissle, and Bigler 2006; Romeis et al. 2008; Marvier et al. 2007; Wolfenbarger et al. 2008; Naranjo 2009). Bt toxins expressed in transgenic plants for pest management are generally regarded as safe because of their rapid degradation in the environment, and their selective mode of action that makes them highly specific for a narrow spectrum of pests (Glare and O'Callaghan 2000; Sanvido, Romeis, and Bigler 2007; Romeis, Meissle, and Bigler 2006; Romeis et al. 2008). The specificity of Cry proteins for certain invertebrates and absence of toxicity for birds and mammals results from the highly specific receptors for these proteins in the gut of target species (Arora et al. 2007).

HPPD proteins are ubiquitous in nature in nearly all aerobic organisms, including the kingdoms of bacteria, fungi, plants, and animals including mammals. HPPD amino acid sequences have been determined in bacteria such as *Streptomyces avermitilis*, in fungi such as *Aspergillus fumigatus*, in plants such as *A. thaliana*, and in animals such as the free-living nematode, *Caenorhabditis elegans*; the mouse, *Mus musculus*, and humans. In particular, HPPD proteins have been characterized in organisms present in human food, such as carrots, barley, swine, and bovines (BASF 2020).

Potential Toxicity and Exposure of Vertebrate T&E Species to GMB151 Soybean

Because soybean plant parts (stems, leave, roots, seeds, pollen, and crop residues) are not transported long distances without human intervention, vertebrate animal exposure to transgenic components of GMB151 Soybean would be limited to those that occur in and around soybean fields. The review in the Affected Environment (Chapter 3 of this EA) documents evidence that demonstrates that soybean fields are poor habitats for birds and mammals in comparison to uncultivated lands. However, a few birds and mammals visit or inhabit soybean fields at various times throughout the soybean production cycle, so could be exposed to GMB 151 Soybean if it were no longer regulated.

APHIS has concluded previously that the same *hppdPf-4Pa* gene and the HPPD protein expressed by it in soybeans has no effects on T&E species (USDA-APHIS 2013). In addition, numerous studies (e.g., (Mendelsohn et al. 2003; Siegel 2001; Betz, Hammond, and Fuchs 2000; Farmer et al. 2017; Farias et al. 2014) have demonstrated that Cry proteins in general do not have toxic effects on vertebrate animals including T&E vertebrate species. Based on the absence of any known risks from toxicity to the transgenic components of GMB151 Soybean and the absence of or limited likelihood of exposure if it were not regulated, APHIS concluded that there would be no effects to T&E vertebrate species.

Potential Toxicity and Exposure of Invertebrate T&E Species to GMB151 Soybean

Because soybean plant parts (stems, leave, roots, seeds, pollen, and crop residues) are not transported long distances without human intervention, invertebrate animal exposure to transgenic components of GMB151 Soybean would be limited to those that occur in and around soybean fields.

The environmental safety assessment reported in the petition (see Appendices 3, 7, and 8) for GMB151 Soybean (BASF 2020) included an in vitro feeding assay of potential impacts to non-target organisms (NTOs). No adverse effects were observed from NTO species tested with Cry14Ab-1 protein including adult and larval honeybees, two soil-dwelling organisms

(Collembola and earthworms), three predator organisms (two species of ladybird beetle and one green lacewing species), one aquatic organism (water flea), one mammal (mouse), and one avian species (bobwhite quail). The bioassay did detect activity of the Cry14Ab-1 on the roundworm, *Caenorhabditis elegans*. However, a two-year field assessment indicated that cultivation of GMB151 Soybean was unlikely to have negative effects on non-target free-living nematodes (BASF 2020). Based on the environmental safety assessment, the cultivation of GMB151 Soybean is unlikely to pose any risk to NTOs at expected field exposure levels. According to the USFWS list of T&E species (Appendix A), no nematodes or roundworms are threatened or endangered, nor are any currently proposed for listing.

APHIS also considered the risks to non-target organisms in its Draft PPRA (pp. 15-17) from consuming GMB151 Soybean with specific reference to agriculture. Based on its review and analysis of the data collected from assays to evaluate the activity spectrum of Cry 14Ab-1, effects on free living nematodes, and effects on other non-target organisms, the Agency concluded that exposure to and/or consumption of GMB Soybean and the PIP it expresses are unlikely to have any adverse impacts to nontarget organisms beneficial to agriculture (USDA-APHIS 2020).

Because the evaluation of NTOs did not detect any adverse effects other than in vitro activity on the roundworm, *C. elegans* (BASF 2020), and the spectrum of activity observed for Cry proteins is narrow (Pigott and Ellar 2007; Wolfenbarger et al. 2008), the remainder of the APHIS T&E animal species analysis focused on taxa within the Class, Insecta. The cross-activity of Cry proteins has been reviewed by van Frankenhuyzen (2013). Cry14 proteins tend to be mostly active on nematodes, but show some activity against coleopteran species. Based on this evidence, APHIS concluded that T&E species in other insect orders (e.g., Lepidoptera, Hymenoptera, and Diptera) would not be affected by the Cry 14Ab-1 expressed by GMB151 Soybean. Therefore, the remainder of the effects analysis focused on coleopterans.

According to the USFWS list of T&E species (Appendix A), all of the listed T&E coleopterans are aquatic or predatory species in both the larval and adult stage. Based on this information, APHIS concluded that these species either could not or would not be exposed to the Cry14Ab-1 protein expressed by GMB151 Soybean, because they don't live in the same habitat or will not feed on soybeans. Therefore, they will not be affected by it. In addition, APHIS noted that BASF conducted non-target studies on two coleopteran species (ladybird beetles) and reported results that showed no effects.

Potential to Serve as a Host Plant for T&E Species

APHIS also considered the possibility that GMB151 Soybean could serve as a host plant for T&E animal species (i.e., a listed insect or other organisms that may use the soybean plant to complete its lifecycle). A review of the T&E species list did not reveal any species that would be likely to use any members of the genus *Glycine* as a host plant (USFWS 2020). Combined, the above information indicates that GMB151 Soybean and its progeny are not expected to have any effects on T&E animal species. There is no increased risk of toxicity or allergenicity impacts directly to animal species from contact with or feeding on GMB151 Soybean. Based on this analysis, APHIS concluded that contact with GMB151 Soybean plants or plant parts by T&E species is unlikely, and if it occurred, consumption would not affect any listed T&E animal species or animal species proposed for listing.

Based on the above analysis, previous analyses of similar transgenic soybean varieties by APHIS that have received a determination of non-regulated status by the Agency, information in the peer-reviewed literature and the petition, APHIS concluded that exposure to and/or consumption of GMB151 Soybean and the HPPD and Cry proteins it expresses, there will be no effects on any threatened or endangered animal species or species proposed for listing.

7.3 SUMMARY

After reviewing the possible effects of a determination of nonregulated status of GMB151 Soybean, APHIS has not identified any stressor that would or could affect the reproduction, numbers, or distribution of a listed T&E species or those proposed for listing. As a result, a detailed exposure analysis for individual species is not necessary. APHIS also considered the potential effect of a determination of nonregulated status of GMB151 Soybean on designated critical habitat or habitat proposed for designation. Compared to other soybean varieties that are currently in use, APHIS determined that GMB151 Soybean production would not differentially affect critical habitat. Like many crops, soybean has been selected for yield rather than its ability to compete and persist in the environment. GMB151 Soybean is not expected to outcompete other plants and persist outside of direct cultivation. Soybean is not sexually compatible with, and does not serve as a host species for, any T&E species or species proposed for listing. There is no evidence that any T&E species or species proposed for listing will consume GMB151 Soybean, therefore APHIS concluded that they will not be subject to any allergic or toxic reactions.

Based on this evidence, APHIS has concluded that a determination of nonregulated status of GMB151 Soybean, and the corresponding environmental release of this soybean variety will have no effect on T&E listed species or species proposed for listing, and would not affect designated habitat or habitat proposed for designation. Because of this “no-effect” determination, consultation under Section 7(a)(2) of the ESA, or the concurrence of the USFWS or NMFS is not required.

8 CONSIDERATION OF EXECUTIVE ORDERS, STANDARDS, AND TREATIES RELATING TO ENVIRONMENTAL IMPACTS

8.1 FEDERAL LAWS AND REGULATIONS

The statutes most relevant to APHIS determinations of regulatory status are the National Environmental Policy Act of 1969 (NEPA), the Clean Water Act of 1972 (CWA), the Safe Drinking Water Act of 1974 (SDWA), the Clean Air Act of 1970 (CAA), the Endangered Species Act of 1973 (ESA), and the National Historic Preservation Act of 1966 (NHPA). Compliance with the requirements of the ESA has been addressed in Chapter 6. Compliance with the requirements of the other relevant laws, NEPA, CWA, SDWA, CAA, and NHPA, is specifically addressed in the following subsections.

8.1.1 National Environmental Policy Act (NEPA)

NEPA is designed to ensure transparency and communication on the possible environmental effects of federal actions prior to implementation of a proposed federal action. The Act and implementing regulations require federal agencies to document, in advance and in detail, the potential effects of their actions on the human environment, so as to ensure that both decision makers and the public fully understanding the possible environmental outcomes of federal actions. APHIS has prepared this draft EA in order to document the potential environmental outcomes of the alternatives considered, consistent with the requirements of NEPA (42 United States Code (U.S.C) 4321, *et seq.*) and Council on Environmental Quality implementing regulations at 40 CFR parts 1500-1508.

8.1.2 Clean Water Act, Safe Drinking Water Act, and Clean Air Act

The CWA, SDWA, and CAA authorize EPA to regulate air and water quality in the United States. This EA evaluates the potential changes in soybean crop production and byproducts associated with approving the petition for a determination of nonregulated status to GMB151 Soybean. APHIS determined that the cultivation of GMB151 Soybean would not lead to the increase in or expansion of the area in soybean production. Because GMB151 Soybean is compositionally, agronomically, and phenotypically equivalent to other non-transgenic and transgenic commercially cultivated soybean varieties (BASF 2020), the potential impacts to water and air quality from the commercial cultivation of GMB151 Soybean would be no different than that of currently cultivated soybean varieties. The herbicide resistance conferred by the genetic modification of GMB151 Soybean is not expected to result in any changes in water usage for cultivation or post-harvest processing of soybean. APHIS assumes any use of isoxaflutole will be compliant with the EPA registration and label requirements. Based on these analyses, APHIS concludes that a determination of nonregulated status for GMB151 Soybean would not lead to circumstances that resulted in non-compliance with the requirements of the CWA, CAA, and SDWA.

8.1.3 National Historic Preservation Act (NHPA) of 1966 as Amended

The National Historic Preservation Act of 1966 (NHPA; Public Law 89-665; 16 U.S.C. 470 *et seq.*) designates federal agencies that are proposing federally funded or permitted projects on historic properties (buildings, archaeological sites, etc.) to consider the impacts using the required Section 106 Review process.

The NHPA and its implementing regulations (36 CFR 800) require federal agencies to: 1) determine whether activities they propose constitute "undertakings" that have the potential to cause impacts on historic properties; and 2) if so, to evaluate the impacts of such undertakings on historic resources and consult with the Advisory Council on Historic Preservation (i.e., State Historic Preservation Office, Tribal Historic Preservation Officers), as appropriate.

A determination of nonregulated status of GMB151 Soybean would not directly or indirectly cause alteration in the character or use of historic properties protected under the NHPA. It would have no impact on districts, sites, highways, structures, or objects listed in or eligible for listing in the National Register of Historic Places, nor would it likely cause any loss or destruction of important scientific, cultural, or historical resources.

Standard agricultural practices for land preparation, planting, irrigation, and harvesting of plants would be used on these agricultural lands, including the use of EPA-registered pesticides. Adherence to the EPA label use restrictions for pesticides will mitigate impacts to the human environment, including historic and cultural resources.

In general, common agricultural activities that would be used in cultivation of GMB151 Soybean do not have the potential to introduce visual, atmospheric, or noise elements to areas in which they are used that could result in impacts on the character or use of historic properties. These cultivation practices are already being conducted throughout the soybean production regions. If GMB151 Soybean were available for cultivation, it would not change any of these agronomic practices that would result in an adverse impact under the NHPA.

8.2 EXECUTIVE ORDERS WITH DOMESTIC IMPLICATIONS

The following executive orders (EOs) require consideration of the potential impacts of the federal action to various segments of the population.

EO 12898 (US-Archives 1994): "Federal Actions to Address Environmental Justice in Minority Populations and Low-Income Populations," requires federal agencies to conduct their programs, policies, and activities that substantially affect human health or the environment in a manner so as not to exclude persons and populations from participation in or benefiting from such programs. It also enforces existing statutes to prevent minority, low-income communities, and Indian Tribes from being subjected to disproportionately high and adverse human health or environmental effects.

EO 13045 (US-Archives 1997): "Protection of Children from Environmental Health Risks and Safety Risks," acknowledges that children may suffer disproportionately from environmental health and safety risks because of their developmental stage, greater metabolic activity levels, and behavior patterns, as compared to adults. This EO (to the extent permitted by law and consistent with the agency's mission) requires each federal agency to identify, assess, and address environmental health risks and safety risks that may disproportionately affect children.

EO 13175: “Consultation and Coordination with Indian Tribal Governments”

charges Executive departments and agencies with a responsibility of engaging in consultation and collaboration with tribal governments; strengthening the government-to-government relationship between the United States and Indian Tribes; and reducing the imposition of unfunded mandates upon Indian Tribes. This EO emphasizes and pledges that federal agencies will communicate and collaborate with tribal officials when proposed federal actions have potential tribal implications.

The No Action and Preferred Alternatives were analyzed with respect to EO 12898, EO 13045 and EO 13175. Neither alternative is expected to have a disproportionate adverse impact on minorities, low-income populations, children, or tribal entities. APHIS determined that the cultivation of GMB151 Soybean would not lead to the increase in or expansion of the area in soybean production. A determination of nonregulated status of GMB151 Soybean is not likely to impact cultural resources on tribal properties. Any farming activities by farmers on tribal lands are only conducted at a Tribe’s request. Thus, the Tribes would have control over any potential conflict with cultural resources on tribal properties. The Proposed action, a determination of nonregulated status of GMB151 Soybean is not expected to impact cultural resources on tribal properties.

Available mammalian toxicity data associated with the HAHB4 protein confirmed the safety of GMB151 Soybean and its products to humans, including minorities, low-income populations, and children who might be exposed to them through agricultural production and/or processing. No additional safety precautions would need to be taken with nonregulated GMB151 Soybean.

APHIS assumes that growers will adhere to herbicide use precautions and restrictions. Pesticide labels include use precautions and restrictions intended to protect workers and their families from exposures. As discussed in Chapter 4 (under Human Health), it is expected that EPA-registered pesticides, fertilizers, and other chemicals that are currently used for soybean production would continue to be used by growers on GMB151 Soybean using application rates currently approved for other transgenic and non-transgenic soybean varieties and found by the EPA not to have adverse impacts to human health when used in accordance with label instructions. Based on these factors, a determination of nonregulated status for GMB151 Soybean is not expected to have disproportionate adverse impacts on minorities, low-income populations, or children.

EO 13751: “Safeguarding the Nation from the Impacts of Invasive Species”

Invasive species are defined as those species that are both not native to the ecosystem under consideration and that also harm the environment, economy or human health. Collectively, they constitute a major concern in the United States and elsewhere. This second EO regarding invasive species directs actions to continue coordinated federal prevention and control efforts related to invasive species. This order maintains the National Invasive Species Council (Council) and the Invasive Species Advisory Committee; adds additional members to the Council; clarifies the operations of the Council; incorporates increased considerations of human and environmental health, climate change, technological innovation, and other emerging priorities into federal efforts to address invasive species; and strengthens coordinated, cost-efficient federal action.

Soybean is not listed in the United States as a noxious weed species by the U.S. government (USDA-NRCS 2013b), nor is it listed as an invasive species by major invasive plant data bases. Cultivated soybean seed does not usually exhibit dormancy and requires specific environmental

conditions to grow as a volunteer the following year (OECD 2000). Any volunteers that may become established do not compete well with the succeeding planted crop and are easily managed using standard weed control practices. Field trials and laboratory tests indicate GMB151 Soybean has no plant pathogenic properties or weediness characteristics. The agronomic, compositional, and reproductive characteristics of GMB151 Soybean are substantially equivalent to other transgenic and non-transgenic soybean varieties. The trait for increased yield is not expected to contribute to increased weediness without changes in a combination of other characteristics associated with weediness, such as hard seed and increased lodging, among other characteristics. Non-engineered soybean, as well as other HR soybean varieties, are widely grown in the United States. Based on historical experience with these varieties and the data submitted by the applicant and reviewed by APHIS, GMB151 Soybean plants are sufficiently similar in fitness characteristics to other soybean varieties currently grown and are not expected to become weedy or invasive.

EO 13186 (US-Archives 2001): “Responsibilities of Federal Agencies to Protect Migratory Birds,” states that federal agencies taking actions that have, or are likely to have a measurable negative impact on migratory bird populations are directed to develop and implement, within two years, a Memorandum of Understanding (MOU) with the Fish and Wildlife Service that shall promote the conservation of migratory bird populations.

Migratory birds may be found in soybean fields. While soybean does not meet the nutritional requirements for many migratory birds (Krapu, Brandt, and Cox 2004), they may forage for insects and weed seeds found in and adjacent to soybean fields. As described in Chapter 4 (under Animal Communities), data submitted by the applicant has shown no difference in compositional and nutritional quality of GMB151 Soybean compared with other transgenic soybean or non-transgenic soybean varieties, apart from the presence of the SCN and isoxaflutole resistance traits. GMB151 Soybean is not expected to be allergenic, toxic, or pathogenic to wildlife. The results provided by BASF indicate that the expression of the Cry14Ab-1 protein is unlikely to be a toxin in animal diets.

Based on the Agency’s assessment of GMB151 Soybean, APHIS concluded it is unlikely that a determination of nonregulated status of GMB151 Soybean would have any negative effects on migratory bird populations.

8.3 INTERNATIONAL IMPLICATIONS

EO 12114 (US-Archives 2010), “Environmental Effects Abroad of Major Federal Actions” requires federal officials to take into consideration any potential environmental impacts outside the United States, its territories, and possessions that result from actions being taken.

APHIS has given this EO careful consideration and does not expect a significant environmental impact outside the United States if it makes a determination of nonregulated status for GMB151 Soybean. All existing national and international regulatory authorities and phytosanitary regimes that currently apply to introductions of new soybean varieties internationally apply equally to those covered by an APHIS determination of nonregulated status under 7 CFR part 340.

Any international trade of GMB151 Soybean subsequent to a determination of nonregulated status would be fully subject to national phytosanitary requirements and be in accordance with

phytosanitary standards developed under the International Plant Protection Convention (IPPC) (IPPC 2013). The purpose of the IPPC “is to secure a common and effective action to prevent the spread and introduction of pests of plants and plant products and to promote appropriate measures for their control” (IPPC 2013). The protection it affords extends to natural flora and plant products and includes both direct and indirect damage by pests, including weeds.

The IPPC establishes a standard for the reciprocal acceptance of phytosanitary certification among the nations that have signed or acceded to the Convention. There are currently 183 IPPC¹⁰ countries. In April 2004, a standard for Plant Risk Analysis of living modified organisms (LMOs) was adopted at a meeting of the governing body of the IPPC as a supplement to an existing standard: International Standard for Phytosanitary Measure No. 11 (ISPM-11, Pest Risk Analysis for Quarantine Pests). The standard acknowledges that all LMOs will not present a pest risk and that a determination needs to be made early in the Plant Risk Analysis for importation as to whether the LMO poses a potential pest risk resulting from the genetic modification. APHIS pest risk assessment procedures for transgenic organisms are consistent with the guidance developed under the IPPC. In addition, issues that may relate to commercialization and transboundary movement of particular agricultural commodities produced through biotechnology are being addressed in other international forums and through national regulations.

The *Cartagena Protocol on Biosafety* is a treaty under the United Nations Convention on Biodiversity that established a framework for the safe transboundary movement, with respect to the environment and biodiversity, of LMOs, which include those modified through biotechnology. The Protocol became effective on September 11, 2003, and currently, there are 198 parties¹¹ that have signed the Protocol. Although the United States is not a party to the Convention on Biodiversity, and thus not a party to the Cartagena Protocol on Biosafety, U.S. exporters will still need to comply with those regulations that importing countries which are Parties to the Protocol have promulgated to comply with their obligations. The first intentional transboundary movement of LMOs intended for environmental release (field trials or commercial planting) will require consent from the importing country under an advanced informed agreement provision, which includes a requirement for a risk assessment consistent with Annex III of the Protocol and the required documentation.

APHIS continues to work toward harmonization of biosafety and biotechnology consensus documents, guidelines, and regulations, including within the North American Plant Protection Organization (NAPPO), which includes Mexico, Canada, and the United States, and within the OECD. NAPPO has completed four modules for releasing transgenic plants NAPPO member countries (NAPPO 2003).

APHIS also participates in the *North American Biotechnology Initiative*, a forum for information exchange and cooperation on agricultural biotechnology issues for the United States, Mexico, and Canada. In addition, bilateral discussions on biotechnology regulatory issues are held regularly with other countries including Argentina, Brazil, Japan, China, and Korea.

¹⁰For a list of countries, go to: <https://www.ippc.int/en/countries/all/list-countries/>

¹¹For a list of signers, go to: <http://bch.cbd.int/protocol/parties/>

8.4 IMPACTS ON UNIQUE CHARACTERISTICS OF GEOGRAPHIC AREAS

A determination of nonregulated status of GMB151 Soybean is not expected to impact unique characteristics of geographic areas such as park lands, prime farmlands, wetlands, wild and scenic areas, or ecologically critical areas.

BASF has presented results of agronomic field trials for GMB151 Soybean. The results of these field trials demonstrate there are no differences in agronomic practices between GMB151 Soybean and non-transgenic hybrids needed for their cultivation. The common agricultural practices that would be carried out in the cultivation of GMB151 Soybean are not expected to deviate from current practices, including the use of EPA-registered pesticides. The product is expected to be grown on agricultural land currently suitable for production of soybean and would only replace existing varieties; it is not expected to increase the acreage of soybean production.

There are no proposed major ground disturbances; no new physical destruction or damage to property; no alterations of property, wildlife habitat, or landscapes; and no prescribed sale, lease, or transfer of ownership of any property. This action is limited to a determination of nonregulated status of GMB151 Soybean. This action would not convert land use to nonagricultural use and, therefore, would have no adverse impact on prime farmland. Standard agricultural practices for land preparation, planting, irrigation, and harvesting of plants would be used on agricultural lands planted to GMB151 Soybean, including the use of EPA-registered pesticides. Adherence by growers to EPA label requirements for all pesticides will prevent adverse effects on the human environment.

Based on these findings, including the assumption that pesticide label requirements are in place to protect unique geographic areas and that those requirements will be adhered to, a determination of nonregulated status for GMB151 Soybean will not impact unique characteristics of geographic areas such as park lands, prime farm lands, wetlands, wild and scenic areas, or ecologically critical areas.

9 List of Preparers

USDA-APHIS	
Name, Title, Project Function	Education and Experience
Elizabeth Nelson Chief, Environmental Risk Analysis Services Reviewer	Ph.D., Public Health, Capella University MBA, University of Maryland University College M.S., Health Care Administration, University of Maryland University College B.S., Biology, Bowie State University 18 years of professional experience in environmental compliance, policy, and management, including preparation of NEPA documentation
Joseph Vorgetts Environmental Protection Specialist, Biotechnology Environmental Risk Analysis Services Project Manager	Ph.D., Entomology, Clemson University M.S., Entomology, Rutgers University B.S., Environmental Science, Rutgers University 14 years of experience in environmental risk assessment and regulatory development and analysis. 25 years of experience in insect survey, suppression and management with pesticides and biological control organisms 9 years of professional experience in environmental risk assessment of transgenic organisms
Frederick David Environmental Protection Specialist, Biotechnology Environmental Risk Analysis Services Reviewer and Assistance with Chapter 3 Preparation	M.S., Environmental Management, University of Maryland University College B.S., Financial Management, University of Maryland University College 10 years of professional experience in environmental management, regulatory compliance and policy, including preparation and review of NEPA documentations
Omar Gardner Environmental Protection Specialist, Biotechnology Environmental Risk Analysis Services Assistance with Chapter 2 Preparation and Reference Library Maintenance	M.S., Environmental Sciences & Policy, Johns Hopkins University B.S., Environmental Science, CUNY Medgar Evers College 4 years of professional experience in environmental risk assessment of transgenic organisms

10 References

- 51 FR 23302. 1986. "Coordinated Framework for Regulation of Biotechnology." Office of Science and Technology Policy Executive Office of the President. Washington, DC.
- 77 FR 13258. 2012. "Biotechnology Regulatory Services; Changes Regarding the Solicitation of Public Comment for Petitions for Determinations of Nonregulated Status for Genetically Engineered Organisms." U.S. of Agriculture Department.
- 81 FR 65414. 2017. "Modernizing the Regulatory System for Biotechnology Products: Final Version of the 2017 Update to the Coordinated Framework for the Regulation of Biotechnology."
- 85 FR 32004. 2020. "BASF Corporation; Petition for a Determination of Nonregulated Status for Plant-Parasitic Nematode-Protected and Herbicide Resistant Soybean." Department of Agriculture. Riverdale Park, MD: Animal and Plant Health Inspection Service.
- Abud, S., P.I.M de Souza, G.R. Vianna, E. Leonardecz, C.T. Moreira, F.G. Faleiro, J.N. Júnior, P.M.F.O. Monteiro, E.L. Rech, and F.J.L. Aragão. 2007. 'Gene flow from transgenic to nontransgenic soybean plants in the Cerrado Region of Brazil', *Genetics and Molecular Research*, 6: 445-52.
- Ahrent, D.K., and C.E. Caviness. 1994. 'Natural cross-pollination of twelve soybean cultivars in Arkansas', *Crop Science*, 34: 376-78.
- Altieri, MA. 1999. 'The ecological role of biodiversity in agroecosystems', *Agriculture, Ecosystems and Environment*, 74: 19-31.
- Altieri, Miguel A., and Deborah K. Letourneau. 1982. 'Vegetation management and biological control in agroecosystems', *Crop Protection*, 1: 405-30.
- AOSCA. 2012a. "National Soybean Variety Review Board." Moline, IL: Association of Official Seed Certifying Agencies.
- . 2012b. "National Soybean Variety Review Board " Moline, IL: Association of Official Seed Certifying Agencies.
- . 2012c. 'Seed Certification'. <http://www.aosca.org/seed%20certification.htm>.
- . 2013. 'IP Handbook - 99.5% Non-GMO Soybean Grain Program Requirements', Accessed 2/13/2013. <http://www.identitypreserved.com/handbook/aosca-nongmosoy.htm>.
- . 2019. "Programs and Services--Association News." Association of Official Seed Certifying Agencies.
- Aref, S, and D Pike. 1998. 'Midwest farmers' perceptions of crop pest infestation', *Agronomy Journal*, 90: 819-25.
- Arora, Naresh, Neema Agrawal, Vimala Yerramilli, and Raj K Bhatnagar. 2007. 'Biology and Applications of Bacillus Thuringiensis in Integrated Pest Management.' in A Ciancio and K G Mukerji (eds.), *General Concepts in Integrated Pest Management and Disease Management* (Springer).
- ASA. 2010a. 'Soy Stats 2010, Domestic Utilization, U.S. Fats & Oils Edible Consumption 2009', American Soybean Association. <http://www.soystats.com/2010/Default-frames.htm>.
- . 2010b. 'U.S. Fats & Oils Edible Consumption 2009', American Soybean Association. <http://www.soystats.com/2010/Default-frames.htm>.
- . 2012a. 'Domestic Utilization: U.S. Biodiesel Production 1999-2011', American Soybean Association. <http://www.soystats.com/2012/Default-frames.htm>.

- . 2012b. "SoyStats 2012: U.S. Soybean Meal Use By Livestock 2011." American Soybean Association.
- Aumaitre, A, K Aulrich, A Chesson, G Flachowsky, and G. Piva. 2002. 'New feeds from genetically modified plants: substantial equivalence, nutritional equivalence, digestibility, and safety for animals and the food chain', *Livestock Production Science*, 74: 223-38.
- BASF. 2020. "Petition for a Determination of Nonregulated Status for Plant-Parasitic, Nematode-Protected and Herbicide Tolerant GMB151 Soybean."
- Beck, D.P. , and D.N. Munns. 1984. 'Phosphate Nutrition of *Rhizobium* spp', *Applied and Environmental Microbiology*, 47: 278-82.
- Beckie, H.J. 2006. 'Herbicide-resistant weeds: Management tactics and practices', *Weed Technology*, 20: 793-814.
- Benbrook, C. 2009. 'Impacts of Genetically Engineered Crops on Pesticide Use: The First Thirteen Years', The Organic Center. http://www.organic-center.org/reportfiles/13Years20091126_ExSumFrontMatter.pdf.
- Bender, R.R., J.W. Haegerle, M.L. Ruffo, and F.E. Below. 2013. 'Modern Corn Hybrids' Nutrient Uptake Patterns', *Better Crops*, 97: 7-10.
- Benedict, L, Hollier, CA., Padgett, GB., Schneider, RW. 2011. "Soybean Disease Management." *Louisiana Agriculture Magazine*. Louisiana Agricultural Extension Service.
- Berglund, D.R., and T.C. Helms. 2003a. "Soybean Production." Fargo: North Dakota Extension Service.
- Berglund, Duane R., and Ted C. Helms. 2003b. 'Soybean Production (Report: A-250 Revised)', North Dakota State University Extension Service. <http://www.ag.ndsu.edu/pubs/plantsci/rowcrops/a250w.htm>.
- Betz, FS, BG Hammond, and RL Fuchs. 2000. 'Safety and Advantages of *Bacillus thuringiensis* - Protected Plants to Control Insect Pests', *Regulatory Toxicology and Pharmacology*, 32: 156 - 73.
- Blount, A., D. Wright, R. Sprenkel, T. Hewitt, and R. Myer. 2009. 'Forage Soybeans for Grazing, Hay and Silage', Florida Cooperative Extension Service, Institute of Food and Agricultural Sciences, University of Florida. <http://edis.ifas.ufl.edu/pdffiles/AG/AG18400.pdf>.
- Boerboom, C.M. 1999. 'Nonchemical options for delaying weed resistance to herbicides in Midwest cropping system', *Weed Technology*, 13: 636-42.
- Bradford, K. 2006. 'Methods to maintain genetic purity of seed stocks', *Agricultural Biotechnology in California Series*, 8189: 1-5.
- Bradley, CA; Wise KA. 2018. " ID-249: A Comprehensive Guide to Soybean Management in Kentucky "
- Brady, S. J. 2007. "Effects of Cropland Conservation Practices on Fish and Wildlife Habitat." *The Wildlife Society Bulletin*. 9-23..
- Brookes, G., and P. Barfoot. 2010. *GM Crops: Global Socio-Economic and Environmental Impacts 1996-2008* (PG Economics Ltd, UK: Dorchester).
- Brown, J.R. 2003. 'Ancient horizontal gene transfer', *Nature Reviews/Genetics*, 4: 121-32.
- Carpenter, Janet, Allan Felsot, Timothy Goode, Michael Hammig, David Onstad, and Sujatha Sankula. 2002. "Comparative Environmental Impacts of Biotechnology-derived and Traditional Soybean, Corn, and Cotton Crops." Council for Agricultural Science and Technology.

- Cassman, K.G., A.S. Whitney, and K.R. Stockinger. 1980. 'Root Growth and Dry Matter Distribution of Soybean as Affected by Phosphorus Stress, Nodulation, and Nitrogen Source', *Crop Science*, 20: 239-44.
- CAST. 2009. "Sustainability of U.S. Soybean Production: Conventional, Transgenic, and Organic Production Systems." Council for Agricultural Science and Technology.
- . 2012. "Herbicide-resistant Weeds Threaten Soil Conservation Gains: Finding a Balance for Soil and Farm Sustainability." Council for Agricultural Science and Technology.
- Caviness, C. 1966. 'Estimates of natural cross-pollination in Jackson soybeans in Arkansas', *Crop Science*, 6: 211-12.
- CBD. 2020. "Sustaining Life on Earth." Montreal, Quebec Canada: Secretariat of the Convention on Biological Diversity.
- Chung, Gyuhwa, and Ram J. Singh. 2008. 'Broadening the Genetic Base of Soybean: A Multidisciplinary Approach', *Critical Reviews in Plant Sciences*, 27: 295-341.
- Clevenger, B. 2010. 'Tips for Evaluating Agronomic Inputs', The Ohio State University, . <http://ohioagmanager.osu.edu/uncategorized/tips-for-evaluating-agronomic-inputs>.
- CLI. 2015. 'Integrated Weed Management', CropLife-International (CLI). <https://croplife.org/plant-biotechnology/stewardship-2/resistance-management/integrated-weed-management/>.
- Clive, J. 2011. "Global Status of Commercialized Biotech/GM Crops: 2011." International Service for the Acquisition of Agri-Biotech Applications.
- Conley, S, and E Christmas. 2005. "Utilizing Inoculates in a Corn-Soybean Rotation (SPS-100-W)." Purdue University Cooperative Extension Service.
- Corn and Soybean Digest. 2012. "Soil Test Results May Not Reveal Accurate Field Nutrient Availability." Penton Media, Inc.
- . 2013. 'Cover Your Cover Crops | A Guide to Which Programs are Most Likely to Partially Reimburse You for Cover Cropping', Penton Media, Inc. <http://cornandsoybeandigest.com/conservation/cover-your-cover-crops-guide-which-programs-are-most-likely-partially-reimburse-you-cov>.
- Cox, M.S., P.D. Gerard, M.C. Wardlaw, and M.J. Abshire. 2003. 'Variability of selected soil properties and their relationships with soybean yield', *Soil Science Society of America Journal*, 67: 1296–302.
- Curtis, T.P., W.T. Sloan, and J.W. Scannell. 2002. 'Estimating prokaryotic diversity and its limits', *Proceedings of the National Academy of Science*, 99: 10494-99.
- De Bruin, Jason, and Palle Pedersen. 2009. 'Growth, Yield, and Yield Component Changes Among Old and New Soybean Cultivars', *Agronomy Journal*, 101: 124-30.
- Dill, G. M., CA CaJacob, and SR Padgett. 2008. 'Glyphosate-resistant crops: Adoption, use and future considerations', *Pest Management Science*, 64: 326-31.
- Doran, J., M. Sarrantonio, and M. Liebig. 1996. 'Soil health and sustainability', *Advances in Agronomy*, 56: 1-54.
- Dubelman, Samuel, Bonnie R Ayden, Brenda M Bader, Christopher R Brown, Changjian Jiang, and Demetra Vlachos. 2005. 'Cry1Ab Protein Does Not Persist in Soil After 3 Years of Sustained Bt Corn Use', *Environmental Entomology*, 34: 915 - 21.
- Duffy, M. 2011. "Estimated Cost of Crop Production in Iowa-2011." Iowa State University
- Duke, S. 2005. 'Taking stock of herbicide-resistant crops ten years after introduction', *Pest Management Science*, 61: 211-18.

- Duke, S.O., and S.B. Powles. 2009. 'Glyphosate-resistant crops and weeds: Now and in the future', *AgBioForum*, 12: 346-57.
- Durgan, B.R. , and J.L. Gunsolus. 2003. "Developing Weed Management Strategies that Address Weed Species Shifts and Herbicide Resistant Weeds. Department of Agronomy and Plant Genetics University of Minnesota."
- Ebelhar, S. A., E. C. Varsa, T. D. Wyciskalla, and C. D. Hart. 2004a. "Effects of Annual Versus Biennial Phosphorus and Potassium Applications in Corn-Soybean Rotation." *Illinois Fertilizer Conference January 26 - 28, 2004*.
- Ebelhar, S.A., E.C. Varsa, T.D. Wyciskalla, and C.D. Hart. 2004b. "Effects of Annual Versus Biennial Phosphorus and Potassium Applications in Corn–Soybean Rotation." *Illinois Fertilizer Conference*. University of Illinois at Urbana-Champaign.
- Egli, D.B. 2008. 'Comparison of Corn and Soybean Yields in the United States: Historical Trends and Future Prospects', *Agronomy Journal*, 100: S79-S88.
- Elbehri, A. 2007. "The Changing Face of the U.S. Grain System Differentiation and Identity Preservation Trends." U.S. Department of Agriculture–Economic Research Service.
- Ellstrand, N. C., H. C. Prentice, and J.K. Hancock. 1999. 'Gene flow and introgression from domesticated plants into their wild relatives', *Annual Review of Ecology and Systematics*, 30: 539-63.
- Elmore, R. W. 1984. "Soybean Inoculation -- When Is It Necessary?" University of Nebraska-Lincoln Extension.
- ETIPCC. 2016. "National Strategy for Modernizing the Regulatory System for Biotechnology Products." https://www.epa.gov/sites/production/files/2016-12/documents/biotech_national_strategy_final.pdf
- FAO. 1997. "Pesticide Residues in Food - 2,4-D and its Salts and Esters." Food and Agriculture Organization of the United Nations.
- . 2009. *Codex Alimentarius, Foods Derived from Modern Biotechnology, 2nd Edition* (World Health Organization, Food and Agriculture Organization of the United Nations: Rome).
- Farias, DF, MP Viana, G Ramos de Oliveira, MA Beneventi, BM Soares, C Pessoa, IP Pessoa, LP Silva, IM Vasconcelos, MF Maria Fátima Grossi de Sá, and AFU Carvalho. 2014. 'Evaluation of Cytotoxic and Antimicrobial Effects of Two Bt Cry Proteins on a GMO Safety Perspective I, Vol 2014 (2014).Publisher .', *BioMed Research International*, 2014: 14.
- Farmer, DR, TC Edrington, CR Kessinich, C Wang, and JS Petrick. 2017. 'Improving insect control protein activity for GM crops: A case study demonstrating that increased target insect potency can be achieved without impacting mammalian safety', *Regulatory Toxicology and Pharmacology*, 89: 155-64.
- Farnham, D. 2001. 'Corn Planting Guide', Iowa State University University Extension. <http://www.extension.iastate.edu/publications/pm1885.pdf>.
- Faust, M. 2002. 'New feeds from genetically modified plants: The US approach to safety for animals and the food chain', *Livestock Production Science*, 74: 239-54.
- Fawcett, R., and S. Caruana. 2001. "Better Soils Better Yield: A Guidebook to Improving Soil Organic Matter and Infiltration With Continuous No-Till." Conservation Technology Information Center.

- Fawcett, Richard , and Dan Towery. 2002. 'Conservation Tillage and Plant Biotechnology: How New Technologies Can Improve the Environment By Reducing the Need to Plow', Conservation Technology Information Center.
<http://www.whybiotech.com/resources/tps/ConservationTillageandPlantBiotechnology.pdf>.
- Feng, Yuanjiao, Lin Ling, Huizhi Fan, Yinghu Liu, Fengxiao Tan, Yinghua Shu, and Jianwu Wang. 2011. 'Effects of temperature, water content and pH on degradation of Cry1Ab protein released from Bt corn straw in soil', *Soil Biology and Biochemistry*, 43: 1600 - 06.
- Fernandez-Cornejo, J. , and W. McBride. 2000. "Genetically Engineered Crops for Pest Management in U.S. Agriculture: Farm-Level Effects." Washington, DC: U.S. Department of Agriculture–Economic Research Service.
- Fernandez-Cornejo, J., C. Hendricks, and A.K. Mishra. 2005. 'Technology adoption and off-farm household income: The case of herbicide-tolerant soybeans', *Journal of Agricultural and Applied Economics*, 37: 549-63.
- Fernandez-Cornejo, J., and M. Caswell. 2006. "The First Decade of Genetically Engineered Crops in the United States. Economic Information Bulletin Number 11." U.S. Department of Agriculture–Economic Research Service.
- Fernandez-Cornejo, J., and W. McBride. 2002. "Adoption of Bioengineered Crops." Washington: U.S. Department of Agriculture–Economic Research Service.
- Fernandez, F.G., and C. White. 2012. 'No-Till and Strip-Till Corn Production with Broadcast and Subsurface-Band Phosphorus and Potassium', *Agronomy Journal*, 104: 996-1005.
- Flachowsky, G, A Chesson, and K Aulrich. 2005. 'Animal nutrition with feeds from genetically modified plants', *Archives of Animal Nutrition*, 59: 1-40.
- FOCUS. 2008. "Pesticides in Air: Considerations for Exposure Assessment." European Commission Forum for Coordination of Pesticide Fate Models and Their Use Working Group on Pesticides in Air.
- Frank, K.D. 2000. 'Fertility Principles: Calcium and Magnesium.' in RB Ferguson and KM De Groot (eds.), *Nutrient Management for Agronomic Crops in Nebraska* (The University of Nebraska Institute of Agriculture and Natural Resources.).
- Franzen, D. W. 2019. 'Soybean Soil Fertility (Report: SF-1164)', North Dakota State University Extension Service. <https://www.ag.ndsu.edu/publications/crops/soybean-soil-fertility/sf1164.pdf>.
- Franzen, D.W. 1999. "Soybean Soil Fertility." North Dakota State University Extension Service.
- Frisvold, G.B., T.M. Hurley, and P.D. Mitchell. 2009. 'Adoption of best management practices to control weed resistance by corn, cotton, and soybean Growers', *AgBioForum*, 12: 370-81.
- Gage, D.J. 2004. 'Infection and invasion of roots by symbiotic, nitrogen-fixing rhizobia during nodulation of temperate legumes', *Microbiology and Molecular Biology Reviews*, 68: 280-300.
- Garbeva, P. , J.D. van Elsas, and J.A. van Veen. 2008. 'Rhizosphere microbial community and its response to plant species and soil history', *Plant Soil*, 302: 19-32.
- Garbeva, P., J. A. van Veen, and J. D. van Elsas. 2004. 'Microbial diversity in soil: Selection of microbial populations by plant and soil type and implications for disease suppressiveness', *Annual Review of Phytopathology*, 42: 243-70.

- Garrison, A.J., A.D. Miller, M.R. Ryan, S.H. Roxburgh, and S. Katriona. 2014. 'Stacked Crop Rotations Exploit Weed-Weed Competition for Sustainable Weed Management', *Weed Science*, 62: 166-76.
- Gianessi, LP, and NP Reigner. 2007. 'The value of herbicides in U.S. crop production', *Weed Technology*: 559-66.
- Givens, W. A., David R. Shaw, Greg R. Kruger, William G. Johnson, Stephen C. Weller, Bryan G. Young, Robert G. Wilson, Micheal D. K. Owen, and David Jordan. 2009. 'Survey of tillage trends following the adoption of glyphosate-resistant crops', *Weed Technology*, 23: 150-55.
- Glare, Travis, and Maureen O'Callaghan. 2000. *Bacillus thuringiensis: Biology, Ecology, and Safety* (Wiley: New York).
- Gonzalez, P., and P. Burch. 2019. "Forages for Summer Grazing." North Carolina Cooperative Extension Service. Raleigh, NC: North Carolina State University.
- Green, J.D. , and J.R. Martin. 1996. "Dealing with Perennial Broadleaf Weeds in Conservation Tillage Systems." University of Kentucky.
- Gunsolus, J.L. 2002. 'Herbicide Resistant Weeds', University of Minnesota Extension. <http://www.extension.umn.edu/distribution/cropsystems/DC6077.html>.
- Hammond, B. G., and J. M. Jez. 2011. 'Impact of food processing on the safety assessment for proteins introduced into biotechnology-derived soybean and corn crops', *Food and chemical toxicology : an international journal published for the British Industrial Biological Research Association*, 49: 711-21.
- Harlan, J. R. 1975. 'Our vanishing genetic resources', *Science*, 188: 618-21.
- Harris, G. 2011. 'Fertilization and Liming.' in J Whitaker (ed.), *2011 Georgia Soybean Production Guide* (University of Georgia).
- Hart, C. . 2006. "Feeding the Ethanol Boom: Where Will the Corn Come From?" *Iowa Ag Review*. Iowa State University.
- Hartman, Glen L., Ellen D. West, and Theresa K. Herman. 2011. 'Crops that feed the World 2. Soybean—worldwide production, use, and constraints caused by pathogens and pests', *Food Security*, 3: 5-17.
- Head, Graham, James B. Surber, Jon A. Watson, John W. Martin, and Jian J. Duan. 2002. 'No Detection of Cry1Ac Protein in Soil After Multiple Years of Transgenic Bt Cotton (Bollgard) Use', *Environmental Entomology*, 31: 30-36.
- Heap, I M. 2021. 'International Survey of Herbicide Resistant Weeds'. <http://www.weedscience.org/Home.aspx>.
- Heatherly, L. 2012. "USB's Kitchen Sink Project – Part II." Mississippi Soybean Promotion Board.
- Heatherly, L., A. Dorrance, R. Hoefl, D. Onstad, J. Orf, P. Porter, S. Spurlock, and B. Young. 2009. "Sustainability of U.S. Soybean Production: Conventional, Transgenic, and Organic Production Systems." Council for Agricultural Science and Technology.
- Heckenmueller, M., D. Narita, and G. Klepper. 2014. 'Global Availability of Phosphorus and Its Implications for Global Food Supply: An Economic Overview', *Kiel Working Papers*, 1897.

- Management." The North Carolina Cooperative Extension Service, College of Agriculture and Life Sciences, North Carolina State University.
- Higgins, R. 1997. "Soybean Insects-Soybean Production Handbook." Kansas State University. Manhattan: Kansas State University.
- Higley, L. G., and D. J. Boethel. 1994. "Handbook of Soybean Insect Pests." Entomological Society of America.
- Hoefst, RG, ED Nafziger, RR Johnson, and SR Aldrich. 2000. *Modern Corn and Soybean Production (1st Ed)* (MCSP Publications: Champaign).
- Hoekema, A, PR Hirsch, PJJ Hooykaas, and RA. Schilperoort. 1983. 'A binary plant vector strategy based on separation of vir- and T-region of the *Agrobacterium tumefaciens* Ti-plasmid. ', *Nature*, 303: 179-80.
- Hoorman, J., R. Islam, and A. Sundermeier. 2009. "Sustainable Crop Rotations with Cover Crops." Ohio State University.
- Hoorman, J., R. Islam, A. Sundermeier, and R. Reeder. 2009. "Using Cover Crops to Convert to No-till." Ohio State University Extension.
- Hubbs, T. 2018. "Outlook for U.S. Corn and Soybean Exports." In *Successful Farming*. Meredith Corporation.
- Hurley, T.M., P.D. Mitchell, and G.B. Frisvold. 2009. 'Effects of weed-resistance concerns and resistance-management practices on the value of Roundup Ready® crops', *AgBioForum*, 12: 291-302.
- Hymowitz, T. 1970. 'On the domestication of the soybean', *Economic Botany*, 24: 408-21.
- Hymowitz, T., and C. A. Newell. 1981. 'Taxonomy of the Genus *Glycine*, Domestication and Uses of Soybeans', *Economic Botany*, 35: 272-88.
- Illinois Crop Improvement Association. 2013. "Seed Directory 2013 Spring Planting, Illinois Certified Seed." Illinois Crop Improvement Association, Inc.
- . 2019. "Seed Directory 2019 Spring Planting." Illinois Certified Seed. Illinois Crop Improvement Association, Inc.
https://www.ilcrop.com/images/Publications/2019_Spring_Seed_Dir.pdf."
- IPM. 2015. 'IPM-Centers: Crop Profiles and Timelines'.
<http://www.ipmcenters.org/cropprofiles/>.
- IPM, Southern. 2019. "National Integrated Pest Management Database," Raleigh, NC. NC: Southern IPM Center, North Carolina State University.
<https://ipmdata.ipmcenters.org/>."
- IPNI. 1998. "Influence of Potassium on Nitrogen Fixation." International Plant Nutrition Institute.
- IPPC. 2013. 'Protecting the World's Plant Resources from Pests', International Plant Protection Convention. <https://www.ippc.int/IPPC/En/default.jsp>.
- Irwin, S.H., and D.L. Good. 2009. 'Market Instability in a New Era of Corn, Soybean, and Wheat Prices', *Choices*, 24: 6-11.
- Israel, D.W. 1987. 'Investigation of the Role of Phosphorus in Symbiotic Dinitrogen Fixation', *Plant Physiology*, 87: 835-41.
- ISSG. 2019. "Invasive Species Specialist Group " *Global Invasive Species Database*. . Species Survival Commission (SSC) of the International Union for Conservation of Nature (IUCN).
- Jardine, D. 1997. "Soybean Diseases-Soybean Production Handbook." Kansas State University.
- Johnson, S, J. Dunphy, and M. Poore. 2007. 'Soybeans as Forage for Grazing, Haying or Silage',

- North Carolina Cooperative Extension Service, North Carolina State University, .
<http://www.ces.ncsu.edu/disaster/drought/Soybeans10-9-07.pdf>.
- Kandel, H, Endres, G (Ed.) 2019. "Disease Management and Identification." North Dakota State University Extension.
- Kaneko, T, Y. Nakamura, S. Sato, E. Asamizu, T. Kato, S. Sasamoto, A. Watanabe, K. Idesawa, A. Ishikawa, K. Kawashima, T. Kimura, Y. Kishida, C. Kiyokawa, M. Kohara, M. Matsumoto, A. Matsuno, Y. Mochizuki, S. Nakayama, N. Nakazaki, S. Shimpo, M. Sugimoto, C. Takeuchi, M. Yamada, and S. Tabata. 2000. 'Complete genome structure of the nitrogen-fixing symbiotic bacterium *Mesorhizobium loti*', *DNA Research*, 7: 331-38.
- Kaneko, T, Y. Nakamura, S. Sato, K. Minamisawa, T. Uchiumi, S. Sasamoto, A. Watanabe, K. Idesawa, M. Iriguchi, K. Kawashima, M. Kohara, M. Matsumoto, S. Shimpo, H. Tsuruoka, T. Wada, M. Yamada, and S. Tabata. 2002. 'Complete genomic sequence of nitrogen-fixing symbiotic bacterium *Bradyrhizobium japonicum* USDA110', *DNA Research*, 9: 189-97.
- Keese, P. 2008. 'Risks from GMOs due to Horizontal Gene Transfer', *Environmental Biosafety Research*, 7: 123-49.
- Kok, H., D. Fjell, and G. Kilgore. 1997. "Seedbed Preparation and Planting Practices-Soybean Production Handbook." Kansas State University.
- Koonin, E.V , K.S. Makarova, and L. Aravind. 2001. 'Horizontal gene transfer in prokaryotes: Quantification and classification', *Annual Review of Microbiology*, 55: 709-42.
- Kowalchuk, G.A., M. Bruinsma, and J.A. van Veen. 2003. 'Assessing responses of soil microorganisms to GM plants', *Trends in Ecology and Evolution*, 18: 403-10.
- Krapu, G. L., D.A. Brandt, and R.R. Cox. 2004. 'Less waste corn, more land in soybeans, and the switch to genetically modified crops: Trends with important implications for wildlife management', *Wildlife Society Bulletin*, 32: 127-36.
- Krausz, R.F. , B.G. Young, G. Kapusta, and J.L. Matthews. 2001. 'Influence of weed competition and herbicides on glyphosate-resistant soybean (*Glycine max*)', *Weed Technology*, 15: 530-34.
- Langemeier, L. . 1997. "Profit Prospects-Soybean Production Handbook." Kansas State University.
- Lee, Dewey, Julia Gaskin, Harry Schomberg, Gary Hawkins, Glen Harris, and Barbara Bellows. No Date. 'Success with Cover Crops', University of Georgia Cooperative Extension.
<http://www.caes.uga.edu/commodities/fieldcrops/gagrains/documents/SuccesswithCovercropswithnotes.pdf>.
- Locke, M. A., R. M. Zablotowicz, and K. N. Reddy. 2008. 'Integrating soil conservation practices and glyphosate-resistant crops: Impacts on soil', *Pest Management Science*, 64: 457-69.
- Lorenz, Gus, Donald R. Johnson, Glenn Stuebaker, Charles Allen, and Seth Young III. 2006. 'Soybean Insect Management', University of Arkansas Division of Agriculture Cooperative Extension Service.
<http://www.aragriculture.org/insects/soybean/default.htm>.
- Lovett, S., P. Price, and J. Lovett. 2003. "Managing Riparian Lands in the Cotton Industry." Cotton Research and Development Corporation.
- Mallarino, A., R. Oltmans, J. Prater, C. Villavicencio, and L. Thompson. 2011. "Nutrient uptake

- by corn and soybean, removal, and recycling with crop residue." *Integrated Crop Management Conference*. Iowa State University.
- Mallory-Smith, C., and M. Zapiola. 2008. 'Gene flow from glyphosate-resistant crops', *Pest Management Science*, 64: 428-40.
- Marra, M.C., N.E. Piggott, and G.A. Carlson. 2004. "The Net Benefits, Including Convenience, Of Roundup Ready® Soybeans: Results from a National Survey." NSF Center for Integrated Pest Management.
- Marvier, Michelle, Chanel McCreedy, James Regetz, and Peter Kareiva. 2007. 'A Meta-Analysis of Effects of Bt Cotton and Maize on Nontarget Invertebrates', *Science*, 316: 1475-77.
- May, M.J., and R.G. Wilson. 2006. 'Weeds and weed control.' in A.P Draycott (ed.), *In Sugar Beet* (Blackwell Publishing: Oxford).
- McBride, W., and C. Greene. 2009. 'The profitability of organic soybean production.', *Renewable Agriculture and Food Systems*, 23: 276-84.
- McMahon, K. 2011. "Commodity Prices Right to Double-Crop Soybeans in Wheat Stubble." Farm Industry Newsletter.
- MDA. 2012. "Cover Crops." Minnesota Department of Agriculture.
- Mendelsohn, M., J. Kough, Zigfridas Vaituzis, and K. Matthews. 2003. 'Are Bt crops safe?', *Nature Biotechnology*, 21: 1003-09.
- Mengel, K., M-R. Haghparast, and K. Kock. 1974. 'The Effect of Potassium on the Fixation of Molecular Nitrogen by Root Nodules of *Vicia faba*', *Plant Physiology*, 54: 535-38.
- Meyer, DW, and M Badaruddin. 2001. 'Frost Tolerance of Ten Seedling Legume Species at Four Growth Stages', *Crop Science*, 41(6): 1338-42.
- Mississippi Crop Improvement Association. 2008. "General Seed Certification Standards." Mississippi Crop Improvement Association.
- . 2015. "General Seed Certification Standards "
- Missouri University of Science and Technology. No Date. '*Bradyrhizobium japonicum*', Missouri University of Science and Technology Accessed November 7.
<http://web.mst.edu/~djwesten/Bj.html>.
- Monsanto. 2010. "Planting Soybeans after Soybeans." *Agronomic Spotlight*, 2. Monsanto.
- Montgomery, David R. 2007. 'Soil erosion and agricultural sustainability', *Proceedings of the National Academy of Sciences*, 104: 13268–72.
- Mullen, M.D., D.W. Israel, and A.G. Wollum. 1988. 'Effects of *Bradyrhizobium japonicum* and Soybean (*Glycine max* (L.) Merr.). Phosphorus Nutrition on Nodulation and Dinitrogen Fixation', *Applied and Environmental Microbiology*, 54.
- NAPPO. 2003. "NAPPO Regional Standards for Phytosanitary Measures (RSPM) " North American Plant Protection Organization.
- Naranjo, Steven E. 2009. 'Impacts of Bt crops on non-target invertebrates and insecticide use patterns ', *CAB Reviews: Perspectives in Agriculture, Veterinary Science, Nutrition, and Natural Resources*, 4.
- NCAT. 2012. "Guide for Organic Crop Producers." National Center for Appropriate Technology.
- NCSRP. No Date. 'Soybean Cyst Management Guide--5th Edition', North Central Soybean Research Program. <http://www.ncipmc.org/pmsp/>.
- Nelson, B. 2011. 'Soybean Diseases in North Dakota', North Dakota State University.
http://www.ndsu.edu/pubweb/~bernelso/soybean_diseases/.
- Nelson, B.D., and C.A. Bradley. 2003. 'Soybean Cyst Nematode (SCN)', North Dakota State

- University. <http://www.ndsu.edu/pubweb/~bernelso/soydiseseases/cyst.shtml>.
- NGWA. 2017. "Ground Water and Agriculture." National Groundwater Association.
- Nichols, C.I., and M.A. Altieri. 2012. 'Plant biodiversity enhances bees and other insect pollinators in agroecosystems. A review', *Agronomic Sustainable Development*, 33: 257-74.
- NRC. 2004. *Safety of Genetically Engineered Foods: Approaches to Assessing Unintended Health Effects* (National Resource Council - The National Academies Press: Washington, D.C.).
- . 2010. 'The Impact of Genetically Engineered Crops on Farm Sustainability in the United States.' in National Research Council (ed.) (National Academies Press: Washington).
- NSRL. No Date. 'Soybean Production Basics', National Soybean Research Laboratory. <http://www.nsrl.illinois.edu/general/soyprod.html>.
- NWF. 2012. "Roadmap to Increased Cover Crop Adoption." National Wildlife Federation.
- OECD-FAO. 2008. "Agricultural Outlook 2008-2017." Organisation for Economic Co-operation and Development – Food and Agriculture Organization of the United Nations.
- . 2020. "OECD-FAO Agricultural Outlook 2020-2029 OECD-FAO Agricultural Outlook 2020, OECD Publishing. ." Organization for Co-Operative Economic Development and Food Agriculture Organization of the UN.
- OECD. 1997. "Series on Harmonization of Regulatory Oversight in Biotechnology No. 6. Consensus document on information used in the assessment of environmental applications involving *Pseudomonas*." Organisation for Economic Co-Operation and Development
- . 2000. "Consensus Document on the Biology of *Glycine max* (L.) Merr. (Soybean)." Organisation for Economic Co-operation and Development
- OMAFRA. 2011. "Agronomy Guide for Field Crops (Publication 118) - Soybean: Planting and Crop Development." Ontario Ministry of Agriculture, Food and Rural Affairs.
- Oplinger, E., and E Amberson. 1986. "Soybean Seed Quality and Certified Seed." University of Wisconsin Extension
- OSTP. 2001. "Case Studies of Environmental Regulation for Biotechnology: Herbicide-Tolerant Soybean." Washington, DC: The White House Office of Science and Technology Policy.
- Owen, M., and IA Zelaya. 2005a. 'Herbicide-resistant crops and weed resistance to herbicides', *Pest Management Science*, 61: 301-11.
- Owen, M. D. K. 2011a. 'Weed resistance development and management in herbicide-tolerant crops: Experiences from the USA', *Journal of Consumer Protection and Food Safety*, 6: 85-89.
- Owen, M. D. K. . 2011b. 'Weed resistance development and management in herbicide-tolerant crops: Experiences from the USA ', *Journal of Consumer Protection and Food Safety*, 6: 85-89.
- Owen, M. D. K., B.G. Young, D.R. Shaw, R.G. Wilson, D.L. Jordan, P.M. Dixon, and S.C. Weller. 2011. 'Benchmark study on glyphosate-resistant cropping systems in the United States. Part 2: Perspectives', *Pest Management Science*, 67: 747-57.
- Owen, M.D.K. 2008. 'Weed species shifts in glyphosate-resistant crops', *Pest Management Science*, 64: 377-87.
- Owen, MDK, and IA Zelaya. 2005b. 'Herbicide-resistant Crops and Weed Resistance to Herbicides', *Pest Management Science*, 61: 301 - 11.
- Palmer, W. E., P. T. Bromley, and J. R. Anderson, Jr. No Date. "Pesticides and Wildlife –

- Soybeans " North Carolina Cooperative Extension Service.
- Palmer, WE, Bromley, PT, Anderson Jr. , JR., 2012. 'Pesticides and Wildlife - Soybeans', North Carolina Extension Integrated Pest Management Program.
http://ipm.ncsu.edu/wildlife/soybeans_wildlife.html.
- Pedersen, P. 2007. 'Soybean Production: Seed Inoculation', Iowa State University Extension Service. http://extension.agron.iastate.edu/soybean/production_seedinoc.html.
- . 2008. "Managing Soybean for High Yield. Iowa State University, Department of Agronomy." Iowa State University, Department of Agronomy.
- Pedersen, P., J. Lauer, C. Grau, and J. Gaska. 2001. "Raising Non-rotation Soybean." *Wisconsin Crop Management Conference*. University of Wisconsin-Madison: Department of Soil Science.
- Peterson. 2016. "Weed Management." Kansas State University Agricultural Experiment Station and Cooperative Extension Service
- Peterson, D. 1997. "Weed Management-Soybean Production Handbook." Kansas State University.
- Pigott, Craig R, and David J Ellar. 2007. 'Role of Receptors in *Bacillus thuringiensis* Crystal Toxin Activity', *Microbiology and Molecular Biology Reviews*, 71: 255-81.
- Powles, S. B. 2008a. 'Evolution in action: Glyphosate-resistant weeds threaten world crops', *Outlooks on Pest Management*, 12: 256-59.
- Powles, S.B. 2008b. 'Evolved glyphosate-resistant weeds around the world: Lessons to be learnt', *Pest Management Science*, 64: 360-65.
- PPI. 2003. "Phosphorus Nutrition of Wheat—Optimize Production." Phosphate and Potash Institute.
- Program, North Central Soybean Research. 2014. "Soybean cyst nematode management guide. ." *Plant Health Initiative*.
- Ray, J., T. Kilen, C. Abel, and R Paris. 2003. 'Soybean natural cross-pollination rates under field conditions', *Environmental Biosafety Research*, 2: 133-38.
- Robertson, G.P., and S.M. Swinton. 2005. 'Reconciling agricultural productivity and environmental integrity: A grand challenge for agriculture', *Frontiers in Agriculture and the Environment*, 3: 38-46.
- Robinson, A. P., and S. P. Conley. 2007. 'Soybean Production Systems: Plant Populations and Seeding Rates for Soybeans', Purdue University Extension.
<http://www.ces.purdue.edu/extmedia/AY/AY-217-W.pdf>.
- Romeis, Jorg, Detlef Bartsch, Franz Bigler, Marco P. Candolfi, Marco M. C. Gielkens, Susan E. Hartley, Richard L. Hellmich, Joseph E. Huesing, Paul C. Jepson, Raymond Layton, Hector Quemada, Alan Raybould, Robyn I. Rose, Joachim Schiemann, Mark K. Sears, Anthony M. Shelton, Jeremy Sweet, Zigfridas Vaituzis, and Jeffrey D. Wolt. 2008. 'Assessment of risk of insect-resistant transgenic crops to nontarget arthropods', *Nat Biotech*, 26: 203-08.
- Romeis, Jorg, Michael Meissle, and Franz Bigler. 2006. 'Transgenic crops expressing *Bacillus thuringiensis* toxins and biological control', *Nature Biotechnology*, 24: 63-71.
- Ronald, P, and B Fouche. 2006. "Genetic Engineering and Organic Production Systems." *Agricultural Biotechnology in California Series*, 1-5. Oakland, CA: University of California, Division of Agriculture and Natural Resources.
- Ruhl, G. 2007. 'Crop Diseases in Corn, Soybean, and Wheat', Purdue University, Department of

- Botany and Plant Pathology.
<http://www.btny.purdue.edu/Extension/Pathology/CropDiseases/Soybean/Soybean.html>.
- Sa, T-M. , and D.W. Israel. 1991. 'Energy Status and Functioning of Phosphorus-Deficient Soybean Nodules', *Plant Physiology*, 97: 928-35.
- Sammons, R.D., D.C. Heering, N. Dinicola, H. Glick, and G.A. Elmore. 2007. 'Sustainability and stewardship of glyphosate and glyphosate-resistant crops', *Weed Technology*, 21: 347-54.
- Sankula, S. 2006. "Quantification of the Impacts on US Agriculture of Biotechnology-Derived Crops Planted In 2005." Washington: National Center for Food and Agricultural Policy.
- Sanvido, Olivier, Jörg Romeis, and Franz Bigler. 2007. 'Ecological Impacts of Genetically Modified Crops: Ten Years of Field Research and Commercial Cultivation', *Advances in Biochemical Engineering/Biotechnology*, 107: 235 - 78.
- SARE. 2012. 'Breaking Costly Pest Cycles with Cover Crops', Sustainable Agriculture Research and Education. <http://www.sare.org/Newsroom/Press-Releases/Archives/Breaking-Costly-Pest-Cycles-with-Cover-Crops>.
- Scherr, Sara J, and Jeffrey A McNeely. 2008. 'Biodiversity conservation and agricultural sustainability: towards a new paradigm of 'ecoagriculture' landscapes', *Phil. Trans. R. Soc. B* 363: 477-94.
- Sedivy, E. J., F. Wu, and Y. Hanzawa. 2017. 'Soybean domestication: the origin, genetic architecture and molecular bases', *New Phytol*, 214: 539-53.
- Sellers, B.A., J.A. Ferrell, and G.E. MacDonald. 2011. "Herbicide Resistant Weeds." In.: University of Florida, Institute of Food and Agricultural Sciences Extension.
- Senseman, S.A. 2007. *Herbicide Handbook, Ninth Edition* (Weed Science Society of America).
- Sharpe, T. 2010. 'Cropland Management ' in M. D. Jones and J. S. Braden (eds.), *Tarheel Wildlife: A Guide for Managing Wildlife on Private Lands in North Carolina* (North Carolina Wildlife Resources Commission: Raleigh).
- Shaw, D. R., M. D. Owen, P. M. Dixon, S. C. Weller, B. G. Young, R. G. Wilson, and D. L. Jordan. 2011. 'Benchmark study on glyphosate-resistant cropping systems in the United States. Part 1: Introduction to 2006-2008', *Pest Manag Sci*, 67: 741-6.
- Sherman-Broyles, S, A Bombarely, AF Powell, JL Doyle, SN Egan, JE Coate, and JJ Doyle. 2014a. 'The Wild Side of a Major Crop: Soybean's Perennial Cousins from Down Under', *American Journal Of Botany*, 101: 1651-65.
- Sherman-Broyles, S., A. Bombarely, A. F. Powell, J. L. Doyle, A. N. Egan, J. E. Coate, and J. J. Doyle. 2014b. 'The wild side of a major crop: soybean's perennial cousins from Down Under', *Am J Bot*, 101: 1651-65.
- Shoup, D; Duncan, S; Ciampitti, IA. 2016. " Soybean Production Handbook." Kansas State University Agricultural Experiment Station and Cooperative Extension Service.
- Siegel, JP. 2001. 'The mammalian safety of *Bacillus thuringiensis*-based insecticides.', *Journal of Invertebrate Pathology*, 77, 13–21.
- Silva, George. 2011. 'Nutrient removal rates in grain crops', Michigan State University Extension. http://msue.anr.msu.edu/news/nutrient_removal_rates_in_grain_crops.
- Snow, A.A., D.A. Andow, P. Gepts, E.M. Hallerman, A. Power, J.M. Tiede, and L.L. Wolfenbarger. 2005. 'Genetically Engineered Organisms and the Environment: Current Status and Recommendations', *Ecological Applications*, 15: 377-404.
- Snyder, C.S. 2000. "Raise Soybean Yields and Profit Potential with Phosphorus and Potassium

- Fertilization." Potash & Phosphate Institute.
- South Dakota Crop Improvement Association. 2011. "South Dakota Seed Certification Standards." South Dakota Crop Improvement Association.
- Specht, James E., Angela Bastidas, Fernando Salvagiotti, Tri Setiyono, Adam Liska, Achim Dobermann, Daniel T. Walters, and Kenneth G. Cassman. 2006. "Soybean Yield Potential and Management Practices Required to Achieve It." Department of Agronomy and Horticulture, University of Nebraska, Lincoln.
- SSCA. No Date-a. "General Seed Certification Standards." Southern Seed Certification Association, Inc.
- . No Date-b. "Standards and Regulations for Certified Seed Production." Southern Seed Certification association, Inc.
- SSDW. No Date. 'Soybean Disease Atlas 2nd Edition', Southern Soybean Disease Workers, . <http://cipm.ncsu.edu/ent/ssdw/soyatlas.htm>.
- Sundstrom, FJ, J Williams, A Van Deynze, and KJ Bradford. 2002. 'Identity preservation of agricultural commodities', *Agricultural Biotechnology in California Series*, 8077: 1-15.
- Tharayil-Santhakumar, N. . 2003. "Mechanisms of Herbicide Resistance in Weeds." University of Massachusetts, Plant & Soil Sciences.
- Towery, Dan, and Steve Werblow. 2010. "Facilitating Conservation Farming Practices and Enhancing Environmental Sustainability with Agricultural Biotechnology." Conservation Technology Information Center.
- Trostle, R. 2008a. "Fluctuating Food Commodity Prices: A Complex Issue With No Easy Answers." U.S. Department of Agriculture–Economic Research Service.
- Trostle, R. . 2008b. "Global Agricultural Supply and Demand: Factors Contributing to the Recent Increase in Food Commodity Prices." Washington: U.S. Department of Agriculture–Economic Research Service.
- Tsvetkova, G.E. , and G.I. Georgiev. 2003. 'Effect of Phosphorus Nutrition on the Nodulation, Nitrogen Fixation and Nutrient - Use Efficiency of *Bradyrhizobium japonicum* – Soybean (*Glycine max* L. Merr.) Symbiosis', *Bulgarin Journal of Plant Physiology*, Special Issue 2003: 331-35.
- Tu, M., C. Hurd, and JM Randall. 2001. *Weed Control Methods Handbook: Tools and Techniques for Use in Natural Areas* (UC Davis, CA.).
- Tylka, G. L., and M. P. Mullaney. 2018. 'Soybean cyst nematode-resistant soybean varieties for Iowa. ', *Extension and Outreach Publications*. 100.: 26.
- University of Arkansas. 2006. "Soil and Water Management, Soybeans – Crop Irrigation." Fayetteville, Arkansas: University of Arkansas Division of Agriculture Cooperative Extension Service.
- University of Arkansas Cooperative Extension Service. 2021. "Arkansas Plant Disease Control Products Guide - MP154."
- University of Georgia Soybean Team. 2019. "Soyban Production Guide." College of Agriculture and Environmental Sciences. Athens, GA: Univeristy of Georgia.
- US-Archives. 1994. "Executive Order 12898 of February 11, 1994.
- Federal Actions To Address Environmental Justice in Minority Populations and Low-Income Populations." "EO 13045 - Protection of Children from Environmental Health Risks and Safety Risks."
- . 2001. 'Executive Order 13186, Responsibilities of Federal Agencies to Protect

- Migratory Birds', Accessed 04/04/2018. <https://www.fws.gov/birds/policies-and-regulations/executive-orders/e0-13186.php>.
- . 2010. 'Executive Order 12114- Environmental effects abroad of major Federal actions', Accessed 04/04/2018. <https://www.archives.gov/federal-register/codification/executive-order/12114.html>.
- US-EPA. 2000. "Profile of the Agricultural Crop Production Industry." Washington: U.S. Environmental Protection Agency-Office of Compliance Sector Notebook Profile.
- . 2005. "Reregistration Eligibility Decision for 2,4-D." Washington: U.S. Environmental Protection Agency.
- . 2009. "Risks of 2,4-D Use to the Federally Threatened California Red-legged Frog (*Rana aurora draytonii*) and Alameda Whipsnake (*Masticophis lateralis euryxanthus*) - Pesticide Effects Determination." Washington: U.S. Environmental Protection Agency.
- . 2010a. 'Pesticide Registration Manual: Chapter 11 - Tolerance Petitions', U.S. Environmental Protection Agency. <http://www.epa.gov/pesticides/bluebook/chapter11.html>.
- . 2010b. "A Set of Scientific Issues Being Considered by the Environmental Protection Agency Regarding: Field Volatilization of Conventional Pesticides." U.S. Environmental Protection Agency, Scientific Advisory Panel.
- . 2011a. 'Agricultural Burning', U.S. Environmental Protection Agency. <http://www.epa.gov/agriculture/tburn.html#Air>.
- . 2011b. 'Pesticide Tolerance', U.S. Environmental Protection Agency. <http://www.epa.gov/pesticides/regulating/tolerances.htm>.
- . 2011c. 'Sole Source Aquifer Protection Program', U.S. Environmental Protection Agency. <http://water.epa.gov/infrastructure/drinkingwater/sourcewater/protection/solesourceaquifer.cfm>.
- . 2012. "Overview of Impaired Waters and Total Maximum Daily Loads Program." U.S. Environmental Protection Agency.
- . 2014. "Biopesticide Oversight and Registration at the U.S. Environmental Protection Agency." U.S. Environmental Protection Agency. Washington, DC: American Chemical Society.
- . 2017a. 'Agriculture: Programs, Best Management Practices and Topics of Interest', U.S. Environmental Protection Agency. <https://www.epa.gov/agriculture/agriculture-about-epas-national-agriculture-center>.
- . 2017b. "PR Notice 2017-2, Guidance for Herbicide-Resistance Management, Labeling, Education, Training and Stewardship." U.S. Environmental Protection Agency.
- . 2017c. 'Worker Protection Standard: How to Comply Manual'.———. 2018. "Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA) and Federal Facilities."
- . 2020. "Federal Food, Drug, and Cosmetic Act (FFDCA) Safety Determination for Cry14Ab-1 Protein." U.S. Environmental Protection Agency, 5. Washington, DC: Office of Pesticide Programs.
- US-FDA. 1998. 'Guidance for Industry: Use of Antibiotic Resistance Marker Genes in Transgenic Plants', U.S. Food and Drug Administration, Accessed November 10. <http://www.fda.gov/Food/GuidanceComplianceRegulatoryInformation/GuidanceDocuments/Biotechnology/ucm096135.htm>.

- USB. 2011a. "A Farmer's Guide to Soy's Top Customers." United Soybean Board.
- . 2011b. "A Farmer's Guide to Soy's Top Customers. News from the Checkoff " United Soybean Board.
- . 2011c. 'Poultry Sector Remains Top Meal Market Despite Challenges', United Soybean Board. <http://unitedsoybean.org/topics/animal-ag>.
- . 2011d. 'U.S. Soy Growers Poised for Growth in Global Animal-Feed Industry', United Soybean Board. <http://unitedsoybean.org/topics/animal-ag>.
- . 2012. "Soybean Market Scan. A report for United Soybean Board." Agralytica Consulting.
- . 2018. "Food and Feed. News from the Checkoff." United Soybean Board.
- USDA-AMS. 2008. 'National Organic Program', U.S. Department of Agriculture-Agricultural Marketing Service. <http://www.ams.usda.gov/AMSV1.0/nop>
- . 2011. 'Organic Production and Handling Standards', U.S. Department of Agriculture-Agricultural Marketing Service. <http://www.ams.usda.gov/AMSV1.0/getfile?dDocName=STELDEV3004445&acct=nopgeninfo>.
- . 2016. 'Pesticide Data Program', Accessed 06/01/2018. <https://www.ams.usda.gov/datasets/pdp>.
- . 2020. "National Organic Program." U.S. Department of Agriculture. Washington, DC: Agricultural Marketing Service.
- USDA-APHIS. 2011. 'Plant Pest Risk Assessment for Monsanto MON 87701 Soybeans', U.S. Department of Agriculture–Animal and Plant Health Inspection Service. http://www.aphis.usda.gov/biotechnology/not_reg.html.
- . 2013. "Bayer Petition 09-328-01 for Determination of Non-regulated Status of Double Herbicide-tolerant Soybean (Glycine Max) Event FG72." U.S. Department of Agriculture. Washington, DC: Animal and Plant Health Inspection Service.
- . 2014. "Final Environtal Assessment for Syngenta and Bayer Petition 12-215-01p for Determination of Non-regulated Status of Herbicide Tolerant Event SYHT0H2 Soybean (Glycine max)." U.S. Department of Agriculture, 205. Washington, DC: Animal and Plant Helth Inspection Service.
- . 2018. "Bayer CropScience LP Petition 17-138-01p for Determination of Nonregulated Status for Glyphosate Resistant and HPPD Inhibitor Resistant GHB811 Cotton." U.S. Department of Agriculture. Washington, DC: Animal and Plant Health Inspection Service.
- . 2020. "Draft Plant Pest Risk Assessment for BASF Petition (19-317-01p)." Washington, DC: USDA-APHIS.
- USDA-ERS. 1997. 'Cropping Management.' in M. Anderson and R Magleby (eds.), *Agricultural Resources and Environmental Indicators 1996-1997* (U.S. Department of Agriculture–Economic Research Service: Washington).
- . 2006. "Soybean Backgrounder." U.S. Department of Agriculture–Economic Research Service.
- . 2010a. "Table 7-U.S. Certified Organic Beans by State, 2008." U.S. Department of Agriculture–Economic Research Service.
- . 2010b. "USDA Soybean Baseline, 2010-19." U.S. Department of Agriculture–Economic Research Service.
- . 2011a. 'Adoption of Genetically Engineered Crops in the U.S.: Soybeans Varieties', U.S.

- Department of Agriculture–Economic Research Service.
<http://www.ers.usda.gov/Data/BiotechCrops/ExtentofAdoptionTable3.htm>.
- . 2011b. "ARMS Farm Financial and Crop Production Practices: Tailored Reports - Crop Production Practices for Soybeans." U.S. Department of Agriculture–Economic Research Service.
 - . 2011c. "Major Uses of Land in the United States, 2007, Economic Information Bulletin Number 89." U.S. Department of Agriculture–Economic Research Service.
 - . 2011d. "Oil Crops Outlook." Washington, DC: U.S. Department of Agriculture–Economic Research Service.
 - . 2012a. "Genetically Engineered Varieties of Corn, Upland Cotton, Soybeans by State and for the United States, 2000-2012." U.S. Department of Agriculture–Economic Research Service.
 - . 2012b. "Oil Crops Yearbook, Table 3: Soybeans: Supply, Disappearance, and Price, U.S., 1980/81-2011/12." U.S. Department of Agriculture–Economic Research Service.
 - . 2016. "Corn and Soybean Production Costs and Export Competitiveness in Argentina, Brazil, and the United States."
 - . 2017. "U.S. Agricultural Trade in 2016: Major Commodities and Trends."
 - . 2020a. "Adoption of Genetically Engineered Crops in the U.S." U.S. Department of Agriculture. Washington, DC: Economic Research Service.
 - . 2020b. "Commodity Costs and Returns--Soybeans: 1997-2019." U.S. Department of Agriculture. Washington, DC: Economic Research Service.
- USDA-FAS. 2013. "Oilseeds: World Market and Trade." Washington: U.S. Department of Agriculture–Foreign Agricultural Service.
- . 2019a. "Oilseeds: World Market and Trade. ." U.S. Department of Agriculture. Washington, DC: Foreign Agricultural Service.
 - . 2019b. "World Agricultural Production." U.S. Department of Agriculture. Washington, DC: Foreign Agricultural Service.
- USDA-FS. 2009. 'Invasive Species', U.S. Department of Agriculture–Forest Service, Pacific Northwest Research Station. <http://www.fs.fed.us/pnw/invasives/index.shtml>.
- USDA-NASS. 2010. '2008 Farm and Ranch Irrigation Survey -Table 27 Crops Harvested from Irrigated Farms: 2008 and 2003', U.S. Department of Agriculture–National Agricultural Statistics Service.
http://www.agcensus.usda.gov/Publications/2007/Online_Highlights/Farm_and_Ranch_Irrigation_Survey/index.asp.
- . 2011. "U.S. & All States Data - Crops Planted, Harvested, Yield, Production, Price (MYA), Value of Production 1991-2011." U.S. Department of Agriculture–National Agricultural Statistics Service.
 - . 2012a. "2011 Certified Organic Production Survey." U.S. Department of Agriculture–National Agriculture Statistics Service.
 - . 2012b. "Farms, Land in Farms, and Livestock Operations 2011 Summary." U.S. Department of Agriculture–National Agriculture Statistics Service.
 - . 2012c. "Indiana Crop & Weather Report." U.S. Department of Agriculture–National Agriculture Statistics Service.
 - . 2012d. "Quick Stats – Soybean Production (National Totals)." U.S. Department of Agriculture–National Agriculture Statistics Service.
 - . 2012e. "U.S. Soybean Acres." U.S. Department of Agriculture–National Agriculture

- Statistics Service.
- . 2016a. "Soybean Production Acres by County." U.S. Department of Agriculture–National Agriculture Statistics Service.
 - . 2017. "Soybean Yield by Year." U.S. Department of Agriculture–National Agriculture Statistics Service.
 - . 2017 "Soybean Production Acreage by Year." U.S. Department of Agriculture–National Agriculture Statistics Service.
 - . 2018a. "Agricultural Chemical Use--Soybeans." U.S. Department of Agriculture–National Agriculture Statistics Service.
 - . 2018b. "Agricultural Chemical Use--Soybeans." *NASS Highlights*.
 - . 2018c. "Corn Planted Acreage Down 1 Percent from 2017, Soybean Acreage Down 1 Percent, All Wheat Acreage Up 4 Percent, All Cotton Acreage Up 7 Percent." U.S. Department of Agriculture–National Agriculture Statistics Service.
 - . 2018d. "National Agricultural Statistics Service June Agricultural Survey." U.S. Department of Agriculture. Washington, DC: National Agricultural Statistic Service.
 - . 2018e. 'National Agricultural Statistics Service, Quick Stats'.
 - . 2018f. "Soybeans & Oil Crops—Overview." U.S. Department of Agriculture–National Agriculture Statistics Service.
 - . 2019. "Crop Progress." U.S. Department of Agriculture. Washington, DC: National Agriculture Statistics Service.
 - . 2020. "U.S. Soybean Acres." U.S. Department of Agriculture. Washington, DC: National Agricultural Statistics Service.
- USDA-NIFA. 2017. 'Sustainable Agriculture Program', United States Department of Agriculture, National Institute of Food and Agriculture. <https://nifa.usda.gov/program/sustainable-agriculture-program>.
- USDA-NRCS. 1996. "Effects of Residue Management and No-Till on Soil Quality." U.S. Department of Agriculture–Natural Resources Conservation Service.
- . 1999a. 'Conservation Tillage Systems and Wildlife', U.S. Department of Agriculture–Natural Resources Conservation Service. http://www.nrcs.usda.gov/Internet/FSE_DOCUMENTS/nrcs143_022212.pdf.
 - . 1999b. 'Soil Taxonomy: A Basic System of Soil Classification for Making and Interpreting Soil Surveys', U.S. Department of Agriculture–Natural Resources Conservation Service. <http://soils.usda.gov/technical/classification/taxonomy/>.
 - . 2001. 'Soil Quality - Introduction', U.S. Department of Agriculture–Natural Resources Conservation Service. <http://soils.usda.gov/sqi/publications/publications.html#agy>.
 - . 2004. "Soil Biology and Land Management." U.S. Department of Agriculture–Natural Resources Conservation Service.
 - . 2005. 'Conservation Practices that Save: Crop Residue Management', U.S. Department of Agriculture–Natural Resources Conservation Service. http://www.nrcs.usda.gov/wps/portal/nrcs/detailfull/?ss=16&navtype=BROWSEBYSUBJECT&cid=nrcs143_023637&navid=1700000000000000&pnavid=null&position=Not%20Yet%20Determined.Html&ttype=detailfull&pname=Conservation%20Practices%20that%20Save:%20Crop%20Residue%20Management%20%20NRCS.
 - . 2006a. "Conservation Resource Brief Soil Erosion." U.S. Department of Agriculture–Natural Resources Conservation Service.
 - . 2006b. "Conservation Resource Brief: Soil Quality." U.S. Department of Agriculture–

- Natural Resources Conservation Service.
- . 2010a. 'Conservation Practice Standard, Conservation Crop Rotation (Ac.) Code 328', U.S. Department of Agriculture–Natural Resources Conservation Service. <http://efotg.sc.egov.usda.gov/references/public/MN/328mn.pdf>.
- . 2010b. 'Keys to Soil Taxonomy', U.S. Department of Agriculture–Natural Resources Conservation Service. http://soils.usda.gov/technical/classification/tax_keys/.
- . 2011a. "In Wet and Dry Years Cover Crops Are a Valuable Practice." Department of Agriculture–Natural Resources Conservation Service.
- . 2011b. "Managing Nonpoint Source Pollution from Agriculture." U.S. Environmental Protection Agency.
- . 2013a. "Energy Consumption Awareness Tool: Tillage." U.S. Department of Agriculture–Natural Resources Conservation Service.
- . 2013b. "Federal Noxious Weeds." U.S. Department of Agriculture–Natural Resources Conservation Service.
- . 2016. "Wildlife Plan (Grassland CRP CP-87/CP-88). Iowa Job Sheet." U.S. Department of Agriculture. Washington, DC: Natural Resources Conservation Service.
- USDA-OCE. 2012. "USDA Agricultural Projections to 2021." U.S. Department of Agriculture–Office of the Chief Economist.
- . 2018. 'USDA Agricultural Projections to 2027', *Office of the Chief Economist, World Agricultural Outlook Board, USDA Interagency Agricultural Projections Committee*, 1: 117.
- USDA. 2011. 'Pest Management Strategic Plans', National Information System of Regional Integrated Pest Management Centers. <http://www.ipmcenters.org/pmsp/index.cfm>.
- . 2020. "USDA Agricultural Projections to 2029." U.S. Department of Agriculture. Washington, DC: Office of the Chief Economist, World Agricultural Outlook Board.
- USDA NASS. 2017. "Organic Survey." U.S. Department of Agriculture. Washington, DC: National Agricultural Statistics Service.
- . 2020a. "Corn and Soybean Area Planted, Left to be Planted, and Harvested – United States 2019 and 2020." United States Department of Agriculture. Washington, DC: National Agricultural Statistics Service (NASS), Agricultural Statistics Board, United States Department of Agriculture.
- . 2020b. "U.S. Soybean Yield." U.S. Department of Agriculture. Washington, DC: National Agricultural Statistics Service.
- van Frankenhuyzen, K. Insecticidal activity of *Bacillus thuringiensis* crystal proteins. *Journal of Invertebrate Pathology* 2009, **101**:1 - 16.
- Van Acker, R.C., and R.C. Swanton. 1993. 'The critical period of weed control in soybean (*Glycine max* (L.) Merr.)', *Weed Science*, 41: 194-200.
- Vencill, W.K., R.L. Nichols, T.M. Webster, J.K. Soteres, C. Mallory-Smith, N.R. Burgos, W.G. Johnson, and M.R. McClelland. 2012. 'Herbicide Resistance: Toward an Understanding of Resistance Development and the Impact of Herbicide-Resistant Crops', *Weed Science 2012 Special Issue*: 2-30.
- Virginia Crop Improvement Association. 2013. "Soybean Certification Standards." Virginia Crop Improvement Association.
- . No Date. "Virginia Handbook of Seed Certification Standards." Virginia Crop Improvement Association.
- Vitosh, M.L., J.W. Johnson, and D.B. Mengel. 2007. "Tri-State Fertilizer Recommendations for

- Corn, Soybeans, Wheat and Alfalfa." Purdue University Cooperative Extension.
- Weirich, JW, DR Shaw, MDK Owen, PM Dixon, SC Weller, BG Young, RG Wilson, and DL Jordan. 2011. 'Benchmark study on glyphosate-resistant cropping systems in the United States. Part 5: Effects of glyphosate-based weed management programs on farm-level profitability', *Pest Management Science*, 67: 781-84.
- Whitaker, J. 2017. "Georgia Soybean Production Guide " University of Georgia College of Agriculture and Environmental Science.
- Whitworth, J and Schwarting, H. 2016. "Soybean Production Handbook " Kansas State University Agricultural Experiment Station and Cooperative Extension Service.
- Whitworth, R. J., J.P. Michaud, and Holly N. Davis. 2011. 'Soybean Insect Management 2011', Kansas State University Agricultural Experiment Station and Cooperative Extension. <http://www.ksre.ksu.edu/library/entml2/mf743.pdf>.
- Whitworth, R. Jeff, J P Michaud, and Holly N Davis. 2012. "Soybean Insect Management (Report MF743)." Kansas State University.
- Wilson, E. 1988. *Biodiversity* (National Academy Press: Washington).
- Wilson, R. S., N. Hooker, M. Tucker, J. LeJeune, and D. Doohan. 2009. 'Targeting the farmer decision making process: A pathway to increased adoption of integrated weed management', *Crop Protection*, 28: 756-64.
- Wolfenbarger, L. LaReesa, Steven E. Naranjo, Jonathan G. Lundgren, Royce J. Bitzer, and Lidia S. Watrud. 2008. 'Bt Crop Effects on Functional Guilds of Non-Target Arthropods: A Meta-Analysis', *PLoS ONE*, 3: e2118.
- Wood, D.W., J. C. Setubal, R. Kaul, D. E. Monks, J. P. Kitajima, V. K. Okura, Y. Zhou, L. Chen, G. E. Wood, N. F. Almeida Jr., Y. Chen L. Woo, I. T. Paulsen, J. A. Eisen, P. D. Karp, D. Bovee Sr., P. Chapman, J. Clendenning, G. Deatherage, W. Gillet, C. Grant, T. Kuttyavin, R. Levy, M Li, E. McClelland, A. Palmieri, C. Raymond, G. Rouse, C. Saenphimmachak, Z. Wu, P. Romero, D. Gordon, S. Zhang, H. Yoo, Y. Tao, P. Biddle, M. Jung, W. Krespan, M. Perry, B. Gordon-Kamm, L. Liao, S. Kim, C. Hendrick, Z. Zhao, M. Dolan, F. Chumley, S. V. Tingey, J. Tomb, M. P. Gordon, M. V. Olson, and E. W. Nester. 2001. 'The genome of the natural genetic engineer *Agrobacterium tumefaciens* C58', *Science*, 294: 2317-23.
- WSSA. 2011. 'Resistance', Weed Science Society of America. <http://www.wssa.net/Weeds/Resistance/index.htm>.
- . 2013. 'Resistance and Tolerance Definitions', Weed Science Society of America, <http://wssa.net/wssa/weed/resistance/herbicide-resistance-and-herbicide-tolerance-definitions/>.
- York, A. C., J.B. Beam, and A. S. Culpepper. 2005. 'Control of volunteer glyphosate-resistant soybean in cotton', *Journal of Cotton Science*, 9: 102-09.
- Yoshimura, Y. 2011. 'Wind tunnel and field assessment of pollen dispersal in soybean [*Glycine max* (L.) Merr.]', *Journal of Plant Science*, 124: 109-14.
- Yoshimura, Y, K Matsuo, and K Yasuda. 2006. 'Gene flow from GM glyphosate-tolerant to conventional soybeans under field conditions in Japan', *Environmental Biosafety Research*, 5: 169-73.
- Zapiola, M.L., C. K. Campbell, M. D. Butler, and C. A. Mallory-Smith. 2008. 'Escape and establishment of transgenic glyphosateresistant creeping bentgrass *Agrostis stolonifera* in Oregon, USA: A 4-year study', *Journal of Applied Ecology*, 45: 486–94.

APPENDIX A

Threatened or Endangered Species



U.S. Fish & Wildlife Service

ECOS

[ECOS](#) / [Species Reports](#) / [Species Search](#)

U.S. Listed Threatened, Endangered, and Proposed Species as of 11/15/20

U.S. Listed Threatened, Endangered, and Proposed Species as of 11/15/20

1653 Records

Parameters:

Listing Statuses: Endangered, Threatened, Not Listed in the US

Scientific Name	Common Name	Family	Federal Listing Status
<u>Abronia macrocarpa</u>	Large-fruited sand-verbena	Nyctaginaceae	Endangered
<u>Abutilon eremitopetalum</u>	No common name	Malvaceae	Endangered
<u>Abutilon menziesii</u>	Ko`oloa`ula	Malvaceae	Endangered
<u>Abutilon sandwicense</u>	No common name	Malvaceae	Endangered
<u>Acaena exigua</u>	Liliwai	Rosaceae	Endangered
<u>Acanthomintha ilicifolia</u>	San Diego thornmint	Lamiaceae	Threatened
<u>Acanthomintha obovata ssp. duttonii</u>	San Mateo thornmint	Lamiaceae	Endangered
<u>Accipiter striatus venator</u>	Puerto Rican sharp-shinned hawk	Accipitridae	Endangered
<u>Achatinella spp.</u>	Oahu tree snails	Achatinellidae	Endangered
<u>Achyranthes mutica</u>	No common name	Amaranthaceae	Endangered

<u><i>Achyranthes splendens</i> var. <i>rotundata</i></u>	Round-leaved chaff-flower	Amaranthaceae	Endangered
<u><i>Acipenser brevirostrum</i></u>	Shortnose sturgeon	Acipenseridae	Endangered
<u><i>Acipenser medirostris</i></u>	green sturgeon	Acipenseridae	Threatened
<u><i>Acipenser oxyrinchus</i></u> <u>(=<i>oxyrhynchus</i>) <i>desotoi</i></u>	Atlantic sturgeon (Gulf subspecies)	Acipenseridae	Threatened
<u><i>Acipenser oxyrinchus oxyrinchus</i></u>	Atlantic sturgeon	Acipenseridae	Endangered
<u><i>Acipenser oxyrinchus oxyrinchus</i></u>	Atlantic sturgeon	Acipenseridae	Threatened

Scientific Name	Common Name	Family	Federal Listing Status
<u><i>Acipenser transmontanus</i></u>	White sturgeon	Acipenseridae	Endangered 1
<u><i>Acmispon dendroideus</i> var. <i>traskiae</i> (= <i>Lotus d. ssp. traskiae</i>)</u>	San Clemente Island lotus (=broom)	Fabaceae	Threatened 1
<u><i>Aconitum noveboracense</i></u>	Northern wild monkshood	Ranunculaceae	Threatened 1
<u><i>Acrocephalus familiaris kingi</i></u>	Nihoa millerbird (old world warbler)	Muscicapidae	Endangered 1
<u><i>Acrocephalus luscini</i></u>	Nightingale reed warbler (old world warbler)	Sylviidae	Endangered 1
<u><i>Acropora cervicornis</i></u>	Staghorn coral	Acroporidae	Threatened 1
<u><i>Acropora palmata</i></u>	Elkhorn coral	Acroporidae	Threatened 1
<u><i>Adelocosa anops</i></u>	Kauai cave wolf or pe'e pe'e maka 'ole spider	Lycosidae	Endangered 1
<u><i>Adenophorus periens</i></u>	Pendant kihi fern	Polypodiaceae	Endangered 1
<u><i>Adiantum vivesii</i></u>	No common name	Adiantaceae	Endangered 1
<u><i>Aerodramus vanikorensis bartschi</i></u>	Mariana gray swiftlet	Apodidae	Endangered 1
<u><i>Aeschynomene virginica</i></u>	Sensitive joint-vetch	Fabaceae	Threatened 1
<u><i>Agalinis acuta</i></u>	Sandplain gerardia	Scrophulariaceae	Endangered 1
<u><i>Agave eggersiana</i></u>	No common name	Agavaceae	Endangered 1
<u><i>Agelaius xanthomus</i></u>	Yellow-shouldered blackbird	Icteridae	Endangered 1
<u><i>Akialoa stejnegeri</i></u>	Kauai akialoa (honeycreeper)	Drepanidinae	Endangered 1
<u><i>Alasmidonta atropurpurea</i></u>	Cumberland elktoe	Unionidae	Endangered 1
<u><i>Alasmidonta heterodon</i></u>	Dwarf wedgemussel	Unionidae	Endangered 1
<u><i>Alasmidonta raveneliana</i></u>	Appalachian elktoe	Unionidae	Endangered 1
<u><i>Alectryon macrococcus</i></u>	Mahoe	Sapindaceae	Endangered 1
<u><i>Allium munzii</i></u>	Munz's onion	Liliaceae	Endangered 1
<u><i>Alopecurus aequalis</i> var. <i>sonomensis</i></u>	Sonoma alopecurus	Poaceae	Endangered 1
<u><i>Amaranthus brownii</i></u>	No common name	Amaranthaceae	Endangered 1
<u><i>Amaranthus pumilus</i></u>	Seabeach amaranth	Amaranthaceae	Threatened 1

Scientific Name	Common Name	Family	Federal Listing Status
<u><i>Amazona vittata</i></u>	Puerto Rican parrot	Psittacidae	Endangered ¹
<u><i>Amblema neislerii</i></u>	Fat threeridge (mussel)	Unionidae	Endangered ¹
<u><i>Amblyopsis rosae</i></u>	Ozark cavefish	Amblyopsidae	Threatened ¹
<u><i>Ambrosia cheiranthifolia</i></u>	South Texas ambrosia	Asteraceae	Endangered ¹
<u><i>Ambrosia pumila</i></u>	San Diego ambrosia	Asteraceae	Endangered ¹
<u><i>Ambrysus amargosus</i></u>	Ash Meadows naucorid	Naucoridae	Threatened ¹
<u><i>Ambystoma bishopi</i></u>	Reticulated flatwoods salamander	Ambystomatidae	Endangered ¹
<u><i>Ambystoma californiense</i></u>	California tiger Salamander	Ambystomatidae	Endangered ¹
<u><i>Ambystoma californiense</i></u>	California tiger Salamander	Ambystomatidae	Threatened ¹
<u><i>Ambystoma cingulatum</i></u>	Frosted Flatwoods salamander	Ambystomatidae	Threatened ¹
<u><i>Ambystoma macrodactylum croceum</i></u>	Santa Cruz long-toed salamander	Ambystomatidae	Endangered ¹
<u><i>Ambystoma tigrinum stebbinsi</i></u>	Sonora tiger Salamander	Ambystomatidae	Endangered ¹
<u><i>Ameiva polops</i></u>	St. Croix ground lizard	Teiidae	Endangered ¹
<u><i>Ammodramus maritimus mirabilis</i></u>	Cape Sable seaside sparrow	Emberizidae	Endangered ¹
<u><i>Ammodramus savannarum floridanus</i></u>	Florida grasshopper sparrow	Emberizidae	Endangered ¹
<u><i>Amorpha crenulata</i></u>	Crenulate lead-plant	Fabaceae	Endangered ¹
<u><i>Amphianthus pusillus</i></u>	Little amphianthus	Scrophulariaceae	Threatened ¹
<u><i>Amphispiza belli clementeae</i></u>	San Clemente sage sparrow	Emberizidae	Threatened ¹
<u><i>Amsinckia grandiflora</i></u>	Large-flowered fiddleneck	Boraginaceae	Endangered ¹
<u><i>Amsonia kearneyana</i></u>	Kearney's blue-star	Apocynaceae	Endangered ¹
<u><i>Anaea troglodyta floridalis</i></u>	Florida leafwing Butterfly	Nymphalidae	Endangered ¹
<u><i>Anas laysanensis</i></u>	Laysan duck	Anatidae	Endangered ¹
<u><i>Anas wyvilliana</i></u>	Hawaiian (=koloa) Duck	Anatidae	Endangered ¹
<u><i>Anaxyrus californicus</i></u>	Arroyo (=arroyo southwestern) toad	Bufonidae	Endangered ¹
<u><i>Anaxyrus canorus</i></u>	Yosemite toad	Bufonidae	Threatened ¹

Scientific Name	Common Name	Family	Federal Listing Status
<u><i>Anguispira picta</i></u>	Painted snake coiled forest snail	Stylommatophora	Threatened
<u><i>Anolis roosevelti</i></u>	Culebra Island giant anole	Iguanidae	Endangered
<u><i>Anoxypristis cuspidata</i></u>	Narrow sawfish	Pristidae	Endangered
<u><i>Antilocapra americana sonoriensis</i></u>	Sonoran pronghorn	Antilocapridae	Endangered
<u><i>Antrobia culveri</i></u>	Tumbling Creek cavesnail	Hydrobiidae	Endangered
<u><i>Antrolana lira</i></u>	Madison Cave isopod	Cirolanidae	Threatened
<u><i>Aphelocoma coerulescens</i></u>	Florida scrub-jay	Corvidae	Threatened
<u><i>Apios priceana</i></u>	Prices potato-bean	Fabaceae	Threatened
<u><i>Aplodontia rufa nigra</i></u>	Point Arena mountain beaver	Aplodontidae	Endangered
<u><i>Apodemia mormo langei</i></u>	Lange's metalmark butterfly	Lycaenidae	Endangered
<u><i>Arabis georgiana</i></u>	Georgia rockcress	Brassicaceae	Threatened
<u><i>Arabis hoffmannii</i></u>	Hoffmann's rock-cress	Brassicaceae	Endangered
<u><i>Arabis macdonaldiana</i></u>	McDonald's rock-cress	Brassicaceae	Endangered
<u><i>Arabis perstellata</i></u>	Braun's rock-cress	Brassicaceae	Endangered
<u><i>Arabis serotina</i></u>	Shale barren rock cress	Brassicaceae	Endangered
<u><i>Arctocephalus townsendi</i></u>	Guadalupe fur seal	Phocidae	Threatened
<u><i>Arctomecon humilis</i></u>	Dwarf Bear-poppy	Papaveraceae	Endangered
<u><i>Arctostaphylos confertiflora</i></u>	Santa Rosa Island manzanita	Ericaceae	Endangered
<u><i>Arctostaphylos franciscana</i></u>	Franciscan manzanita	Ericaceae	Endangered
<u><i>Arctostaphylos glandulosa ssp. crassifolia</i></u>	Del Mar manzanita	Ericaceae	Endangered
<u><i>Arctostaphylos hookeri var. ravenii</i></u>	Presidio Manzanita	Ericaceae	Endangered
<u><i>Arctostaphylos morroensis</i></u>	Morro manzanita	Ericaceae	Threatened
<u><i>Arctostaphylos myrtifolia</i></u>	Ione manzanita	Ericaceae	Threatened
<u><i>Arctostaphylos pallida</i></u>	Pallid manzanita	Ericaceae	Threatened
<u><i>Arenaria cumberlandensis</i></u>	Cumberland sandwort	Caryophyllaceae	Endangered
<u><i>Arenaria paludicola</i></u>	Marsh Sandwort	Caryophyllaceae	Endangered

Scientific Name	Common Name	Family	Federal Listing Status
<u><i>Arenaria ursina</i></u>	Bear Valley sandwort	Caryophyllaceae	Threatened
<u><i>Argemone pleiacantha ssp. pinnatisecta</i></u>	Sacramento prickly poppy	Papaveraceae	Endangered
<u><i>Argyroxiphium kauense</i></u>	Mauna Loa (=Ka'u) silversword	Asteraceae	Endangered
<u><i>Argyroxiphium sandwicense ssp. macrocephalum</i></u>	`Ahinahina	Asteraceae	Threatened
<u><i>Argyroxiphium sandwicense ssp. sandwicense</i></u>	`Ahinahina	Asteraceae	Endangered
<u><i>Argythamnia blodgettii</i></u>	Blodgett's silverbush	Euphorbiaceae	Threatened
<u><i>Aristida chaseae</i></u>	No common name	Poaceae	Endangered
<u><i>Aristida portoricensis</i></u>	Pelos del diablo	Poaceae	Endangered
<u><i>Arkansia wheeleri</i></u>	Ouachita rock pocketbook	Unionidae	Endangered
<u><i>Asclepias meadii</i></u>	Mead's milkweed	Asclepiadaceae	Threatened
<u><i>Asclepias welshii</i></u>	Welsh's milkweed	Asclepiadaceae	Threatened
<u><i>Asimina tetramera</i></u>	Four-petal pawpaw	Annonaceae	Endangered
<u><i>Asplenium dielirectum</i></u>	Asplenium-leaved diellia	Aspleniaceae	Endangered
<u><i>Asplenium dielfalcatum</i></u>	No common name	Aspleniaceae	Endangered
<u><i>Asplenium diellaciniatum</i></u>	No common name	Aspleniaceae	Endangered
<u><i>Asplenium dielmannii</i></u>	No common name	Aspleniaceae	Endangered
<u><i>Asplenium dielpallidum</i></u>	No common name	Aspleniaceae	Endangered
<u><i>Asplenium peruvianum var. insulare</i></u>	No common name	Aspleniaceae	Endangered
<u><i>Asplenium scolopendrium var. americanum</i></u>	American hart's-tongue fern	Aspleniaceae	Threatened
<u><i>Asplenium unisorum</i></u>	No common name	Aspleniaceae	Endangered
<u><i>Assiminea pecos</i></u>	Pecos assiminea snail	Assimineidae	Endangered
<u><i>Astelia waialealae</i></u>	Pa`iniu	Liliaceae	Endangered
<u><i>Astragalus albens</i></u>	Cushenbury milk-vetch	Fabaceae	Endangered
<u><i>Astragalus ampullarioides</i></u>	Shivwits milk-vetch	Fabaceae	Endangered
<u><i>Astragalus applegatei</i></u>	Applegate's milk-vetch	Fabaceae	Endangered

Scientific Name	Common Name	Family	Federal Listing Status
<u><i>Astragalus bibullatus</i></u>	Guthrie's (=Pyne's) ground-plum	Fabaceae	Endangered 1
<u><i>Astragalus brauntonii</i></u>	Braunton's milk-vetch	Fabaceae	Endangered 1
<u><i>Astragalus clarianus</i></u>	Clara Hunt's milk-vetch	Fabaceae	Endangered 1
<u><i>Astragalus cremnophylax</i></u> <u>var. <i>cremnophylax</i></u>	Sentry milk-vetch	Fabaceae	Endangered 1
<u><i>Astragalus holmgreniorum</i></u>	Holmgren milk-vetch	Fabaceae	Endangered 1
<u><i>Astragalus humillimus</i></u>	Mancos milk-vetch	Fabaceae	Endangered 1
<u><i>Astragalus jaegerianus</i></u>	Lane Mountain milk-vetch	Fabaceae	Endangered 1
<u><i>Astragalus lentiginosus</i></u> var. <u><i>coachellae</i></u>	Coachella Valley milk-vetch	Fabaceae	Endangered 1
<u><i>Astragalus lentiginosus</i></u> var. <u><i>piscinensis</i></u>	Fish Slough milk-vetch	Fabaceae	Threatened 1
<u><i>Astragalus magdalenae</i></u> var. <u><i>peirsonii</i></u>	Peirson's milk-vetch	Fabaceae	Threatened 1
<u><i>Astragalus montii</i></u>	Heliotrope milk-vetch	Fabaceae	Threatened 1
<u><i>Astragalus osterhoutii</i></u>	Osterhout milkvetch	Fabaceae	Endangered 1
<u><i>Astragalus phoenix</i></u>	Ash meadows milk-vetch	Fabaceae	Threatened 1
<u><i>Astragalus pycnostachyus</i></u> var. <u><i>lanosissimus</i></u>	Ventura Marsh Milk-vetch	Fabaceae	Endangered 1
<u><i>Astragalus robbinsii</i></u> var. <u><i>jesupi</i></u>	Jesup's milk-vetch	Fabaceae	Endangered 1
<u><i>Astragalus schmolliae</i></u>	Chapin Mesa milkvetch	Fabaceae	Proposed Threatened 1
<u><i>Astragalus tener</i></u> var. <u><i>titi</i></u>	Coastal dunes milk-vetch	Fabaceae	Endangered 1
<u><i>Astragalus tricarinatus</i></u>	Triple-ribbed milk-vetch	Fabaceae	Endangered 1
<u><i>Astrophytum asterias</i></u>	Star cactus	Cactaceae	Endangered 1
<u><i>Athearnia anthonyi</i></u>	Anthony's riversnail	Pleuroceridae	Endangered 1
<u><i>Atlantea tulita</i></u>	Puerto Rico harlequin butterfly	Nymphalidae	Proposed Threatened 1
<u><i>Atriplex coronata</i></u> var. <u><i>notator</i></u>	San Jacinto Valley crownscale	Chenopodiaceae	Endangered 1
<u><i>Auerodendron pauciflorum</i></u>	No common name	Rhamnaceae	Endangered 1

Scientific Name	Common Name	Family	Federal Listing Status
<u><i>Ayenia limitaris</i></u>	Texas ayenia	Sterculiaceae	Endangered
<u><i>Baccharis vanessae</i></u>	Encinitas baccharis	Asteraceae	Threatened
<u><i>Balaena mysticetus</i></u>	Bowhead whale	Balaenidae	Endangered
<u><i>Balaenoptera borealis</i></u>	Sei whale	Balaenopteridae	Endangered
<u><i>Balaenoptera musculus</i></u>	Blue whale	Balaenopteridae	Endangered
<u><i>Balaenoptera physalus</i></u>	Finback whale	Balaenopteridae	Endangered
<u><i>Banara vanderbiltii</i></u>	Palo de ramon	Flacourtiaceae	Endangered
<u><i>Baptisia arachnifera</i></u>	Hairy rattleweed	Fabaceae	Endangered
<u><i>Batrachoseps aridus</i></u>	Desert slender salamander	Plethodontidae	Endangered
<u><i>Batrisodes texanus</i></u>	Coffin Cave mold beetle	Pselaphidae	Endangered
<u><i>Batrisodes venyivi</i></u>	Helotes mold beetle	Pselaphidae	Endangered
<u><i>Berberis nevinii</i></u>	Nevin's barberry	Berberidaceae	Endangered
<u><i>Berberis pinnata ssp. insularis</i></u>	Island Barberry	Berberidaceae	Endangered
<u><i>Betula uber</i></u>	Virginia round-leaf birch	Betulaceae	Threatened
<u><i>Bidens amplexans</i></u>	Ko`oko`olau	Asteraceae	Endangered
<u><i>Bidens campylotheca ssp. pentamera</i></u>	Ko`oko`olau	Asteraceae	Endangered
<u><i>Bidens campylotheca ssp. waihoiensis</i></u>	Ko`oko`olau	Asteraceae	Endangered
<u><i>Bidens conjuncta</i></u>	Ko`oko`olau	Asteraceae	Endangered
<u><i>Bidens hillebrandiana ssp. hillebrandiana</i></u>	kookoolau	Asteraceae	Endangered
<u><i>Bidens micrantha ssp. ctenophylla</i></u>	Ko`oko`olau	Asteraceae	Endangered
<u><i>Bidens micrantha ssp. kalealaha</i></u>	Ko`oko`olau	Asteraceae	Endangered
<u><i>Bidens wiebkei</i></u>	Ko`oko`olau	Asteraceae	Endangered
<u><i>Bison bison athabasca</i></u>	Wood Bison	Bovidae	Threatened
<u><i>Blennosperma bakeri</i></u>	Sonoma sunshine	Asteraceae	Endangered
<u><i>Boloria acrocynema</i></u>	Uncompahgre fritillary butterfly	Nymphalidae	Endangered

Scientific Name	Common Name	Family	Federal Listing Status
<u><i>Boltonia decurrens</i></u>	Decurrent false aster	Asteraceae	Threatened
<u><i>Bombus affinis</i></u>	Rusty patched bumble bee	Apidae	Endangered
<u><i>Bombus franklini</i></u>	Franklin's bumblebee	Apidae	Proposed Endangered
<u><i>Bonamia grandiflora</i></u>	Florida bonamia	Convolvulaceae	Threatened
<u><i>Bonamia menziesii</i></u>	No common name	Convolvulaceae	Endangered
<u><i>Brachylagus idahoensis</i></u>	Columbia Basin Pygmy Rabbit	Leporidae	Endangered
<u><i>Brachyramphus marmoratus</i></u>	Marbled murrelet	Alcidae	Threatened
<u><i>Branchinecta conservatio</i></u>	Conservancy fairy shrimp	Branchinectidae	Endangered
<u><i>Branchinecta longiantenna</i></u>	Longhorn fairy shrimp	Branchinectidae	Endangered
<u><i>Branchinecta lynchi</i></u>	Vernal pool fairy shrimp	Branchinectidae	Threatened
<u><i>Branchinecta sandiegonensis</i></u>	San Diego fairy shrimp	Branchinectidae	Endangered
<u><i>Branta</i> (=Nesochen) <i>sandvicensis</i></u>	Hawaiian goose	Anatidae	Threatened
<u><i>Brickellia mosieri</i></u>	Florida brickell-bush	Asteraceae	Endangered
<u><i>Brighamia insignis</i></u>	Olulu	Campanulaceae	Endangered
<u><i>Brighamia rockii</i></u>	Pua `ala	Campanulaceae	Endangered
<u><i>Brodiaea filifolia</i></u>	Thread-leaved brodiaea	Liliaceae	Threatened
<u><i>Brodiaea pallida</i></u>	Chinese Camp brodiaea	Liliaceae	Threatened
<u><i>Brychius hungerfordi</i></u>	Hungerford's crawling water Beetle	Halipilidae	Endangered
<u><i>Bufo hemiophrys baxteri</i></u>	Wyoming Toad	Bufonidae	Endangered
<u><i>Bufo houstonensis</i></u>	Houston toad	Bufonidae	Endangered
<u><i>Bulbophyllum guamense</i></u>	Cebello halumtano	Orchidaceae	Threatened
<u><i>Buteo platypterus brunneescens</i></u>	Puerto Rican broad-winged hawk	Accipitridae	Endangered
<u><i>Buxus vahlii</i></u>	Vahl's boxwood	Buxaceae	Endangered
<u><i>Calamagrostis expansa</i></u>	Maui reedgrass	Poaceae	Endangered
<u><i>Calamagrostis hillebrandii</i></u>	Hillebrand's reedgrass	Poaceae	Endangered
<u><i>Calidris canutus rufa</i></u>	Red knot	Scolopacidae	Threatened

Scientific Name	Common Name	Family	Federal Listing Status
<u><i>Callicarpa ampla</i></u>	Capa rosa	Verbenaceae	Endangered
<u><i>Callirhoe scabriuscula</i></u>	Texas poppy-mallow	Malvaceae	Endangered
<u><i>Callophrys mossii bayensis</i></u>	San Bruno elfin butterfly	Lycaenidae	Endangered
<u><i>Calochortus tiburonensis</i></u>	Tiburon mariposa lily	Liliaceae	Threatened
<u><i>Calyptranthes thomasiana</i></u>	No common name	Myrtaceae	Endangered
<u><i>Calyptridium pulchellum</i></u>	Mariposa pussypaws	Portulacaceae	Threatened
<u><i>Calyptronomia rivalis</i></u>	Palma de manaca	Arecaceae	Threatened
<u><i>Calystegia stebbinsii</i></u>	Stebbins' morning-glory	Convolvulaceae	Endangered
<u><i>Cambarus aculabrum</i></u>	Benton County cave crayfish	Cambaridae	Endangered
<u><i>Cambarus callainus</i></u>	Big Sandy crayfish	Cambaridae	Threatened
<u><i>Cambarus cracens</i></u>	Slenderclaw crayfish	Cambaridae	Proposed Threatened
<u><i>Cambarus veteranus</i></u>	Guyandotte River crayfish	Cambaridae	Endangered
<u><i>Cambarus zophonastes</i></u>	Hell Creek Cave crayfish	Cambaridae	Endangered
<u><i>Camissonia benitensis</i></u>	San Benito evening-primrose	Onagraceae	Threatened
<u><i>Campanula robinsiae</i></u>	Brooksville bellflower	Campanulaceae	Endangered
<u><i>Campeloma decampi</i></u>	Slender campeloma	Viviparidae	Endangered
<u><i>Campephilus principalis</i></u>	Ivory-billed woodpecker	Picidae	Endangered
<u><i>Canavalia molokaiensis</i></u>	`Awikiwiki	Fabaceae	Endangered
<u><i>Canavalia napaliensis</i></u>	`Awikiwiki	Fabaceae	Endangered
<u><i>Canavalia pubescens</i></u>	`Awikiwiki	Fabaceae	Endangered
<u><i>Canis lupus baileyi</i></u>	Mexican wolf	Canidae	Endangered
<u><i>Canis rufus</i></u>	Red wolf	Canidae	Endangered
<u><i>Caprimulgus noctitherus</i></u>	Puerto Rican nightjar	Caprimulgidae	Endangered
<u><i>Cardamine micranthera</i></u>	Small-anthered bittercress	Brassicaceae	Endangered
<u><i>Caretta caretta</i></u>	Loggerhead sea turtle	Cheloniidae	Endangered
<u><i>Caretta caretta</i></u>	Loggerhead sea turtle	Cheloniidae	Threatened
<u><i>Carex albida</i></u>	White sedge	Cyperaceae	Endangered
<u><i>Carex lutea</i></u>	Golden sedge	Cyperaceae	Endangered
<u><i>Carex specuicola</i></u>	Navajo sedge	Cyperaceae	Threatened

Scientific Name	Common Name	Family	Federal Listing Status
<u><i>Castilleja affinis ssp. neglecta</i></u>	Tiburon paintbrush	Scrophulariaceae	Endangered ①
<u><i>Castilleja campestris ssp. succulenta</i></u>	Fleshy owl's-clover	Scrophulariaceae	Threatened ①
<u><i>Castilleja cinerea</i></u>	Ash-grey paintbrush	Scrophulariaceae	Threatened ①
<u><i>Castilleja grisea</i></u>	San Clemente Island Paintbrush	Scrophulariaceae	Threatened ①
<u><i>Castilleja levisecta</i></u>	golden paintbrush	Orobanchaceae	Threatened ①
<u><i>Castilleja mollis</i></u>	Soft-leaved paintbrush	Scrophulariaceae	Endangered ①
<u><i>Catesbaea melanocarpa</i></u>	No common name	Rubiaceae	Endangered ①
<u><i>Catostomus discobolus yarrowi</i></u>	Zuni bluehead Sucker	Catostomidae	Endangered ①
<u><i>Catostomus santaanae</i></u>	Santa Ana sucker	Catostomidae	Threatened ①
<u><i>Catostomus warnerensis</i></u>	Warner sucker	Catostomidae	Threatened ①
<u><i>Caulanthus californicus</i></u>	California jewelflower	Brassicaceae	Endangered ①
<u><i>Ceanothus ferrisiae</i></u>	Coyote ceanothus	Rhamnaceae	Endangered ①
<u><i>Ceanothus ophiochilus</i></u>	Vail Lake ceanothus	Rhamnaceae	Threatened ①
<u><i>Ceanothus roderickii</i></u>	Pine Hill ceanothus	Rhamnaceae	Endangered ①
<u><i>Cenchrus agrimonioides</i></u>	Kamanomano	Poaceae	Endangered ①
<u><i>Centaurium namophilum</i></u>	Spring-loving centaury	Gentianaceae	Threatened ①
<u><i>Centrocercus minimus</i></u>	Gunnison sage-grouse	Phasianidae	Threatened ①
<u><i>Cercocarpus traskiae</i></u>	Catalina Island mountain-mahogany	Rosaceae	Endangered ①
<u><i>Cereus eriophorus var. fragrans</i></u>	Fragrant prickly-apple	Cactaceae	Endangered ①
<u><i>Chamaecrista glandulosa var. mirabilis</i></u>	No common name	Fabaceae	Endangered ①
<u><i>Chamaecrista lineata keyensis</i></u>	Big Pine partridge pea	Fabaceae	Endangered ①
<u><i>Chamaesyce deltoidea pinetorum</i></u>	Pineland sandmat	Euphorbiaceae	Threatened ①
<u><i>Chamaesyce deltoidea serpyllum</i></u>	Wedge spurge	Euphorbiaceae	Endangered ①

Scientific Name	Common Name	Family	Federal Listing Status
<u><i>Chamaesyce deltoidea</i> ssp. <i>deltoidea</i></u>	Deltoid spurge	Euphorbiaceae	Endangered 1
<u><i>Chamaesyce garberi</i></u>	Garber's spurge	Euphorbiaceae	Threatened 1
<u><i>Chamaesyce hooveri</i></u>	Hoover's spurge	Euphorbiaceae	Threatened 1
<u><i>Charadrius melodus</i></u>	Piping Plover	Charadriidae	Endangered 1
<u><i>Charadrius melodus</i></u>	Piping Plover	Charadriidae	Threatened 1
<u><i>Charadrius nivosus nivosus</i></u>	Western snowy plover	Charadriidae	Threatened 1
<u><i>Charpentiera densiflora</i></u>	Papala	Amaranthaceae	Endangered 1
<u><i>Chasiempis ibidis</i></u>	Oahu elepaio	Monarchidae	Endangered 1
<u><i>Chasmistes brevirostris</i></u>	Shortnose Sucker	Catostomidae	Endangered 1
<u><i>Chasmistes cujus</i></u>	Cui-ui	Catostomidae	Endangered 1
<u><i>Chasmistes liorus</i></u>	June sucker	Catostomidae	Endangered 1
<u><i>Chelonia mydas</i></u>	Green sea turtle	Cheloniidae	Endangered 1
<u><i>Chelonia mydas</i></u>	Green sea turtle	Cheloniidae	Threatened 1
<u><i>Chilabothrus granti</i></u>	Virgin Islands tree boa	Boidae	Endangered 1
<u><i>Chionanthus pygmaeus</i></u>	Pygmy fringe-tree	Oleaceae	Endangered 1
<u><i>Chlorogalum purpureum</i></u>	Purple amole	Agavaceae	Threatened 1
<u><i>Chorizanthe howellii</i></u>	Howell's spineflower	Polygonaceae	Endangered 1
<u><i>Chorizanthe orcuttiana</i></u>	Orcutt's spineflower	Polygonaceae	Endangered 1
<u><i>Chorizanthe pungens</i> var. <i>hartwegiana</i></u>	Ben Lomond spineflower	Polygonaceae	Endangered 1
<u><i>Chorizanthe pungens</i> var. <i>pungens</i></u>	Monterey spineflower	Polygonaceae	Threatened 1
<u><i>Chorizanthe robusta</i> var. <i>hartwegii</i></u>	Scotts Valley spineflower	Polygonaceae	Endangered 1
<u><i>Chorizanthe robusta</i> var. <i>robusta</i></u>	Robust spineflower	Polygonaceae	Endangered 1
<u><i>Chorizanthe valida</i></u>	Sonoma spineflower	Polygonaceae	Endangered 1
<u><i>Chromolaena frustrata</i></u>	Cape Sable Thoroughwort	Asteraceae	Endangered 1
<u><i>Chrosomus saylora</i></u>	Laurel dace	Cyprinidae	Endangered 1
<u><i>Chrysopsis floridana</i></u>	Florida golden aster	Asteraceae	Endangered 1

Scientific Name	Common Name	Family	Federal Listing Status
<u><i>Cicindela dorsalis dorsalis</i></u>	Northeastern beach tiger beetle	Cicindelidae	Threatened
<u><i>Cicindela nevadica lincolniiana</i></u>	Salt Creek Tiger beetle	Cicindelidae	Endangered
<u><i>Cicindela ohlone</i></u>	Ohlone tiger beetle	Cicindelidae	Endangered
<u><i>Cicindela puritana</i></u>	Puritan tiger beetle	Cicindelidae	Threatened
<u><i>Cicindelidia floridana</i></u>	Miami tiger beetle	Carabidae	Endangered
<u><i>Cicurina baronia</i></u>	Robber Baron Cave Meshweaver	Dictynidae	Endangered
<u><i>Cicurina madla</i></u>	Madla Cave Meshweaver	Dictynidae	Endangered
<u><i>Cicurina venii</i></u>	Braken Bat Cave Meshweaver	Dictynidae	Endangered
<u><i>Cicurina vespera</i></u>	Government Canyon Bat Cave Meshweaver	Dictynidae	Endangered
<u><i>Cirsium fontinale</i> var. <i>fontinale</i></u>	Fountain thistle	Asteraceae	Endangered
<u><i>Cirsium fontinale</i> var. <i>obispoense</i></u>	Chorro Creek bog thistle	Asteraceae	Endangered
<u><i>Cirsium hydrophilum</i> var. <i>hydrophilum</i></u>	Suisun thistle	Asteraceae	Endangered
<u><i>Cirsium loncholepis</i></u>	La Graciosa thistle	Asteraceae	Endangered
<u><i>Cirsium pitcheri</i></u>	Pitcher's thistle	Asteraceae	Threatened
<u><i>Cirsium vinaceum</i></u>	Sacramento Mountains thistle	Asteraceae	Threatened
<u><i>Cladonia perforata</i></u>	Florida perforate cladonia	Cladoniaceae	Endangered
<u><i>Clarkia franciscana</i></u>	Presidio clarkia	Onagraceae	Endangered
<u><i>Clarkia imbricata</i></u>	Vine Hill clarkia	Onagraceae	Endangered
<u><i>Clarkia speciosa</i> ssp. <i>immaculata</i></u>	Pismo clarkia	Onagraceae	Endangered
<u><i>Clarkia springvillensis</i></u>	Springville clarkia	Onagraceae	Threatened
<u><i>Clematis morefieldii</i></u>	Morefields leather flower	Ranunculaceae	Endangered
<u><i>Clematis socialis</i></u>	Alabama leather flower	Ranunculaceae	Endangered
<u><i>Clemmys muhlenbergii</i></u>	bog turtle	Emydidae	Threatened

Scientific Name	Common Name	Family	Federal Listing Status
<u><i>Clermontia drepanomorpha</i></u>	`Oha wai	Campanulaceae	Endangered
<u><i>Clermontia lindseyana</i></u>	`Oha wai	Campanulaceae	Endangered
<u><i>Clermontia oblongifolia</i> ssp. <i>brevipes</i></u>	`Oha wai	Campanulaceae	Endangered
<u><i>Clermontia oblongifolia</i> ssp. <i>mauiensis</i></u>	`Oha wai	Campanulaceae	Endangered
<u><i>Clermontia peleana</i></u>	`Oha wai	Campanulaceae	Endangered
<u><i>Clermontia pyrularia</i></u>	`Oha wai	Campanulaceae	Endangered
<u><i>Clermontia samuelii</i></u>	`Oha wai	Campanulaceae	Endangered
<u><i>Clitoria fragrans</i></u>	Pigeon wings	Fabaceae	Threatened
<u><i>Coccyzus americanus</i></u>	Yellow-billed Cuckoo	Cuculidae	Threatened
<u><i>Colinus virginianus ridgwayi</i></u>	Masked bobwhite (quail)	Phasianidae	Endangered
<u><i>Colubrina oppositifolia</i></u>	Kauila	Rhamnaceae	Endangered
<u><i>Columba inornata wetmorei</i></u>	Puerto Rican plain Pigeon	Columbidae	Endangered
<u><i>Conradina brevifolia</i></u>	Short-leaved rosemary	Lamiaceae	Endangered
<u><i>Conradina etonia</i></u>	Etonia rosemary	Lamiaceae	Endangered
<u><i>Conradina glabra</i></u>	Apalachicola rosemary	Lamiaceae	Endangered
<u><i>Conradina verticillata</i></u>	Cumberland rosemary	Lamiaceae	Threatened
<u><i>Consolea corallicola</i></u>	Florida semaphore Cactus	Cactaceae	Endangered
<u><i>Cordia bellonis</i></u>	No common name	Boraginaceae	Endangered
<u><i>Cordylanthus maritimus</i> ssp. <i>maritimus</i></u>	Salt marsh bird's-beak	Scrophulariaceae	Endangered
<u><i>Cordylanthus mollis</i> ssp. <i>mollis</i></u>	Soft bird's-beak	Scrophulariaceae	Endangered
<u><i>Cordylanthus palmatus</i></u>	Palmate-bracted bird's beak	Scrophulariaceae	Endangered
<u><i>Cordylanthus tenuis</i> ssp. <i>capillaris</i></u>	Pennell's bird's-beak	Scrophulariaceae	Endangered
<u><i>Cornutia obovata</i></u>	Palo de nigua	Verbenaceae	Endangered
<u><i>Corvus hawaiiensis</i></u>	Hawaiian (= 'alala) Crow	Corvidae	Endangered
<u><i>Corvus kubaryi</i></u>	Mariana (=aga) Crow	Corvidae	Endangered
<u><i>Corvus leucognaphalus</i></u>	White-necked crow	Corvidae	Endangered

Scientific Name	Common Name	Family	Federal Listing Status
<u><i>Corynorhinus (=Plecotus) townsendii ingens</i></u>	Ozark big-eared bat	Vespertilionidae	Endangered
<u><i>Corynorhinus (=Plecotus) townsendii virginianus</i></u>	Virginia big-eared bat	Vespertilionidae	Endangered
<u><i>Coryphantha minima</i></u>	Nellie cory cactus	Cactaceae	Endangered
<u><i>Coryphantha ramillosa</i></u>	Bunched cory cactus	Cactaceae	Threatened
<u><i>Coryphantha robbinsorum</i></u>	Cochise pincushion cactus	Cactaceae	Threatened
<u><i>Coryphantha scheeri</i> var. <i>robustispina</i></u>	Pima pineapple cactus	Cactaceae	Endangered
<u><i>Coryphantha sneedii</i> var. <i>leei</i></u>	Lee pincushion cactus	Cactaceae	Threatened
<u><i>Coryphantha sneedii</i> var. <i>sneedii</i></u>	Sneed pincushion cactus	Cactaceae	Endangered
<u><i>Cottus paulus (=pygmaeus)</i></u>	Pygmy Sculpin	Cottidae	Threatened
<u><i>Cottus specus</i></u>	Grotto Sculpin	Cottidae	Endangered
<u><i>Cranichis ricartii</i></u>	No common name	Orchidaceae	Endangered
<u><i>Crenichthys baileyi baileyi</i></u>	White River springfish	Cyprinodontidae	Endangered
<u><i>Crenichthys baileyi grandis</i></u>	Hiko White River springfish	Cyprinodontidae	Endangered
<u><i>Crenichthys nevadae</i></u>	Railroad Valley springfish	Cyprinodontidae	Threatened
<u><i>Crescentia portoricensis</i></u>	Higuero de sierra	Bignoniaceae	Endangered
<u><i>Crocodylus acutus</i></u>	American crocodile	Crocodylidae	Threatened
<u><i>Crotalaria avonensis</i></u>	Avon Park harebells	Fabaceae	Endangered
<u><i>Crotalus willardi obscurus</i></u>	New Mexican ridge-nosed rattlesnake	Crotalidae	Threatened
<u><i>Cryptantha crassipes</i></u>	Terlingua Creek cat's-eye	Boraginaceae	Endangered
<u><i>Cryptobranchus alleganiensis alleganiensis</i></u>	Eastern Hellbender Missouri DPS	Cryptobranchidae	Proposed Endangered
<u><i>Cryptobranchus alleganiensis bishopi</i></u>	Ozark Hellbender	Cryptobranchidae	Endangered
<u><i>Crystallaria cincotta</i></u>	diamond Darter	Percidae	Endangered
<u><i>Ctenitis squamigera</i></u>	Pauoa	Aspleniaceae	Endangered
<u><i>Cucurbita okeechobeensis</i> ssp. <i>okeechobeensis</i></u>	Okeechobee gourd	Cucurbitaceae	Endangered

Scientific Name	Common Name	Family	Federal Listing Status
<u><i>Cumberlandia monodonta</i></u>	Spectaclecase (mussel)	Margaritiferidae	Endangered
<u><i>Cupressus abramsiana</i></u>	Santa Cruz cypress	Cupressaceae	Threatened
<u><i>Cupressus goveniana ssp. goveniana</i></u>	Gowen cypress	Cupressaceae	Threatened
<u><i>Cyanea acuminata</i></u>	Haha	Campanulaceae	Endangered
<u><i>Cyanea asarifolia</i></u>	Haha	Campanulaceae	Endangered
<u><i>Cyanea asplenifolia</i></u>	Haha	Campanulaceae	Endangered
<u><i>Cyanea calycina</i></u>	Haha	Campanulaceae	Endangered
<u><i>Cyanea copelandii ssp. copelandii</i></u>	Haha	Campanulaceae	Endangered
<u><i>Cyanea copelandii ssp. haleakalaensis</i></u>	Haha	Campanulaceae	Endangered
<u><i>Cyanea crispa</i></u>	haha	Campanulaceae	Endangered
<u><i>Cyanea dolichopoda</i></u>	Haha	Campanulaceae	Endangered
<u><i>Cyanea dunbariae</i></u>	haha	Campanulaceae	Endangered
<u><i>Cyanea duvalliorum</i></u>	haha	Campanulaceae	Endangered
<u><i>Cyanea eleeleensis</i></u>	Haha	Campanulaceae	Endangered
<u><i>Cyanea gibsonii</i></u>	haha	Campanulaceae	Endangered
<u><i>Cyanea glabra</i></u>	Haha	Campanulaceae	Endangered
<u><i>Cyanea grimesiana ssp. grimesiana</i></u>	Haha	Campanulaceae	Endangered
<u><i>Cyanea grimesiana ssp. obatae</i></u>	Haha	Campanulaceae	Endangered
<u><i>Cyanea hamatiflora ssp. carlsonii</i></u>	Haha	Campanulaceae	Endangered
<u><i>Cyanea hamatiflora ssp. hamatiflora</i></u>	Haha	Campanulaceae	Endangered
<u><i>Cyanea horrida</i></u>	haha nui	Campanulaceae	Endangered
<u><i>Cyanea humboldtiana</i></u>	Haha	Campanulaceae	Endangered
<u><i>Cyanea kauaulaensis</i></u>	No common name	Campanulaceae	Endangered
<u><i>Cyanea kolekoleensis</i></u>	Haha	Campanulaceae	Endangered

Scientific Name	Common Name	Family	Federal Listing Status
<u><i>Cyanea koolauensis</i></u>	Haha	Campanulaceae	Endangered
<u><i>Cyanea kuhihewa</i></u>	Haha	Campanulaceae	Endangered
<u><i>Cyanea kunthiana</i></u>	Haha	Campanulaceae	Endangered
<u><i>Cyanea lanceolata</i></u>	Haha	Campanulaceae	Endangered
<u><i>Cyanea lobata</i></u>	Haha	Campanulaceae	Endangered
<u><i>Cyanea longiflora</i></u>	Haha	Campanulaceae	Endangered
<u><i>Cyanea magnicalyx</i></u>	haha	Campanulaceae	Endangered
<u><i>Cyanea mannii</i></u>	Haha	Campanulaceae	Endangered
<u><i>Cyanea maritae</i></u>	haha	Campanulaceae	Endangered
<u><i>Cyanea marksii</i></u>	Haha	Campanulaceae	Endangered
<u><i>Cyanea mauiensis</i></u>	haha	Campanulaceae	Endangered
<u><i>Cyanea mceldowneyi</i></u>	Haha	Campanulaceae	Endangered
<u><i>Cyanea munroi</i></u>	haha	Campanulaceae	Endangered
<u><i>Cyanea obtusa</i></u>	Haha	Campanulaceae	Endangered
<u><i>Cyanea pinnatifida</i></u>	Haha	Campanulaceae	Endangered
<u><i>Cyanea platyphylla</i></u>	`aku`aku	Campanulaceae	Endangered
<u><i>Cyanea procera</i></u>	Haha	Campanulaceae	Endangered
<u><i>Cyanea profuga</i></u>	Haha	Campanulaceae	Endangered
<u><i>Cyanea purpurellifolia</i></u>	Haha	Campanulaceae	Endangered
<u><i>Cyanea recta</i></u>	Haha	Campanulaceae	Threatened
<u><i>Cyanea remyi</i></u>	Haha	Campanulaceae	Endangered
<u><i>Cyanea rivularis</i></u>	Haha	Campanulaceae	Endangered
<u><i>Cyanea shipmanii</i></u>	Haha	Campanulaceae	Endangered
<u><i>Cyanea solanacea</i></u>	Popolo	Campanulaceae	Endangered
<u><i>Cyanea st.-johnii</i></u>	Haha	Campanulaceae	Endangered
<u><i>Cyanea stictophylla</i></u>	Haha	Campanulaceae	Endangered
<u><i>Cyanea superba</i></u>	Haha	Campanulaceae	Endangered
<u><i>Cyanea tritomantha</i></u>	`aku	Campanulaceae	Endangered
<u><i>Cyanea truncata</i></u>	Haha	Campanulaceae	Endangered

Scientific Name	Common Name	Family	Federal Listing Status
<u><i>Cyanea undulata</i></u>	Haha	Campanulaceae	Endangered
<u><i>Cyathea dryopteroides</i></u>	Elfin tree fern	Cyatheaceae	Endangered
<u><i>Cycas micronesica</i></u>	Fadang	Cycadaceae	Threatened
<u><i>Cycladenia humilis</i> var. <i>jonesii</i></u>	Jones Cycladenia	Apocynaceae	Threatened
<u><i>Cyclargus</i> (= <i>Hemiargus</i>) <i>thomasi bethunebakeri</i></u>	Miami Blue Butterfly	Lycaenidae	Endangered
<u><i>Cyclosorus boydiae</i></u>	Kupukupu makalii	Thelypteridaceae	Endangered
<u><i>Cyclura stejnegeri</i></u>	Mona ground Iguana	Iguanidae	Threatened
<u><i>Cynomys parvidens</i></u>	Utah prairie dog	Sciuridae	Threatened
<u><i>Cyperus fauriei</i></u>	No common name	Cyperaceae	Endangered
<u><i>Cyperus neokunthianus</i></u>	No common name	Cyperaceae	Endangered
<u><i>Cyperus pennatiformis</i></u>	No common name	Cyperaceae	Endangered
<u><i>Cyperus trachysanthos</i></u>	Pu`uka`a	Cyperaceae	Endangered
<u><i>Cyprinella caerulea</i></u>	Blue shiner	Cyprinidae	Threatened
<u><i>Cyprinella formosa</i></u>	Beautiful shiner	Cyprinidae	Threatened
<u><i>Cyprinodon bovinus</i></u>	Leon Springs pupfish	Cyprinodontidae	Endangered
<u><i>Cyprinodon diabolis</i></u>	Devils Hole pupfish	Cyprinodontidae	Endangered
<u><i>Cyprinodon elegans</i></u>	Comanche Springs pupfish	Cyprinodontidae	Endangered
<u><i>Cyprinodon macularius</i></u>	Desert pupfish	Cyprinodontidae	Endangered
<u><i>Cyprinodon nevadensis mionectes</i></u>	Ash Meadows Amargosa pupfish	Cyprinodontidae	Endangered
<u><i>Cyprinodon nevadensis pectoralis</i></u>	Warm Springs pupfish	Cyprinodontidae	Endangered
<u><i>Cyprinodon radiosus</i></u>	Owens pupfish	Cyprinodontidae	Endangered
<u><i>Cyprogenia stegaria</i></u>	Fanshell	Unionidae	Endangered
<u><i>Cyrtandra crenata</i></u>	Ha`iwale	Gesneriaceae	Endangered
<u><i>Cyrtandra cyaneoides</i></u>	Mapele	Gesneriaceae	Endangered
<u><i>Cyrtandra dentata</i></u>	Ha`iwale	Gesneriaceae	Endangered
<u><i>Cyrtandra ferripilosa</i></u>	haiwale	Gesneriaceae	Endangered
<u><i>Cyrtandra filipes</i></u>	Ha`iwale	Gesneriaceae	Endangered

Scientific Name	Common Name	Family	Federal Listing Status
<u><i>Cyrtandra giffardii</i></u>	Ha`iwale	Gesneriaceae	Endangered
<u><i>Cyrtandra gracilis</i></u>	Haiwale	Gesneriaceae	Endangered
<u><i>Cyrtandra hematos</i></u>	Ha`iwale	Gesneriaceae	Endangered
<u><i>Cyrtandra kaulantha</i></u>	Ha`iwale	Gesneriaceae	Endangered
<u><i>Cyrtandra limahuliensis</i></u>	Ha`iwale	Gesneriaceae	Threatened
<u><i>Cyrtandra munroi</i></u>	Ha`iwale	Gesneriaceae	Endangered
<u><i>Cyrtandra nanawaleensis</i></u>	haiwale	Gesneriaceae	Endangered
<u><i>Cyrtandra oenobarba</i></u>	Ha`iwale	Gesneriaceae	Endangered
<u><i>Cyrtandra oxybapha</i></u>	Ha`iwale	Gesneriaceae	Endangered
<u><i>Cyrtandra paliku</i></u>	Haiwale	Gesneriaceae	Endangered
<u><i>Cyrtandra polyantha</i></u>	Ha`iwale	Gesneriaceae	Endangered
<u><i>Cyrtandra sessilis</i></u>	Ha`iwale	Gesneriaceae	Endangered
<u><i>Cyrtandra subumbellata</i></u>	Ha`iwale	Gesneriaceae	Endangered
<u><i>Cyrtandra tintinnabula</i></u>	Ha`iwale	Gesneriaceae	Endangered
<u><i>Cyrtandra viridiflora</i></u>	Ha`iwale	Gesneriaceae	Endangered
<u><i>Cyrtandra wagneri</i></u>	haiwale	Gesneriaceae	Endangered
<u><i>Cyrtandra waiolani</i></u>	Haiwale	Gesneriaceae	Endangered
<u><i>Dalea carthagenensis</i></u> <u><i>floridana</i></u>	Florida prairie-clover	Fabaceae	Endangered
<u><i>Dalea foliosa</i></u>	Leafy prairie-clover	Fabaceae	Endangered
<u><i>Daphnopsis helleriana</i></u>	No common name	Thymelaeaceae	Endangered
<u><i>Deeringothamnus pulchellus</i></u>	Beautiful pawpaw	Annonaceae	Endangered
<u><i>Deeringothamnus rugelii</i></u>	Rugel's pawpaw	Annonaceae	Endangered
<u><i>Deinandra (=Hemizonia)</i></u> <u><i>conjugens</i></u>	Otay tarplant	Asteraceae	Threatened
<u><i>Deinandra increscens ssp.</i></u> <u><i>villosa</i></u>	Gaviota Tarplant	Asteraceae	Endangered
<u><i>Delissea rhytidosperma</i></u>	No common name	Campanulaceae	Endangered
<u><i>Delissea subcordata</i></u>	Oha	Campanulaceae	Endangered
<u><i>Delissea undulata</i></u>	No common name	Campanulaceae	Endangered

Scientific Name	Common Name	Family	Federal Listing Status
<u><i>Delphinapterus leucas</i></u>	beluga whale	Monodontidae	Endangered
<u><i>Delphinium bakeri</i></u>	Baker's larkspur	Ranunculaceae	Endangered
<u><i>Delphinium luteum</i></u>	Yellow larkspur	Ranunculaceae	Endangered
<u><i>Delphinium variegatum ssp. kinkiense</i></u>	San Clemente Island larkspur	Ranunculaceae	Endangered
<u><i>Deltistes luxatus</i></u>	Lost River sucker	Catostomidae	Endangered
<u><i>Dendrobium guamense</i></u>	No common name	Orchidaceae	Threatened
<u><i>Dendrogyra cylindrus</i></u>	Pillar Coral	Meandrinidae	Threatened
<u><i>Dendroica chrysoparia</i></u>	Golden-cheeked warbler (=wood)	Parulidae	Endangered
<u><i>Deparia kaalaana</i></u>	No common name	Woodsiaceae	Endangered
<u><i>Dermochelys coriacea</i></u>	Leatherback sea turtle	Dermochelyidae	Endangered
<u><i>Desmocerus californicus dimorphus</i></u>	Valley elderberry longhorn beetle	Cesambycidae	Threatened
<u><i>Dicerandra christmanii</i></u>	Garrett's mint	Lamiaceae	Endangered
<u><i>Dicerandra cornutissima</i></u>	Longspurred mint	Lamiaceae	Endangered
<u><i>Dicerandra frutescens</i></u>	Scrub mint	Lamiaceae	Endangered
<u><i>Dicerandra immaculata</i></u>	Lakela's mint	Lamiaceae	Endangered
<u><i>Digitaria pauciflora</i></u>	Florida pineland crabgrass	Poaceae	Threatened
<u><i>Dinacoma caseyi</i></u>	Casey's June Beetle	Scarabaeidae	Endangered
<u><i>Dionda diaboli</i></u>	Devils River minnow	Cyprinidae	Threatened
<u><i>Diplacus vandenbergensis</i></u>	Vandenberg monkeyflower	Phrymaceae	Endangered
<u><i>Diplazium molokaiense</i></u>	No common name	Woosiaceae	Endangered
<u><i>Dipodomys heermanni morroensis</i></u>	Morro Bay kangaroo rat	Heteromyidae	Endangered
<u><i>Dipodomys ingens</i></u>	Giant kangaroo rat	Heteromyidae	Endangered
<u><i>Dipodomys merriami parvus</i></u>	San Bernardino Merriam's kangaroo rat	Heteromyidae	Endangered
<u><i>Dipodomys nitratoides exilis</i></u>	Fresno kangaroo rat	Heteromyidae	Endangered
<u><i>Dipodomys nitratoides nitratoides</i></u>	Tipton kangaroo rat	Heteromyidae	Endangered

Scientific Name	Common Name	Family	Federal Listing Status
<u><i>Dipodomys stephensi</i> (incl. <i>D. cascus</i>)</u>	Stephens' kangaroo rat	Heteromyidae	Endangered
<u><i>Discus macclintocki</i></u>	Iowa Pleistocene snail	Discidae	Endangered
<u><i>Dodecahema leptoceras</i></u>	Slender-horned spineflower	Polygonaceae	Endangered
<u><i>Doryopteris angelica</i></u>	No common name	Pteridaceae	Endangered
<u><i>Doryopteris takeuchii</i></u>	No common name	Pteridaceae	Endangered
<u><i>Drepanis coccinea</i></u>	Tiwi	Fringillidae	Threatened
<u><i>Dromus dromas</i></u>	Dromedary pearlymussel	Unionidae	Endangered
<u><i>Drosophila aglaia</i></u>	Hawaiian picture-wing fly	Drosophilidae	Endangered
<u><i>Drosophila differens</i></u>	Hawaiian picture-wing fly	Drosophilidae	Endangered
<u><i>Drosophila digressa</i></u>	Hawaiian picture-wing fly	Drosophilidae	Endangered
<u><i>Drosophila hemipeza</i></u>	Hawaiian picture-wing fly	Drosophilidae	Endangered
<u><i>Drosophila heteroneura</i></u>	Hawaiian picture-wing fly	Drosophilidae	Endangered
<u><i>Drosophila montgomeryi</i></u>	Hawaiian picture-wing fly	Drosophilidae	Endangered
<u><i>Drosophila mulli</i></u>	Hawaiian picture-wing fly	Drosophilidae	Threatened
<u><i>Drosophila musaphilia</i></u>	Hawaiian picture-wing fly	Drosophilidae	Endangered
<u><i>Drosophila neoclavisetae</i></u>	Hawaiian picture-wing fly	Drosophilidae	Endangered
<u><i>Drosophila obatai</i></u>	Hawaiian picture-wing fly	Drosophilidae	Endangered
<u><i>Drosophila ochrobasis</i></u>	Hawaiian picture-wing fly	Drosophilidae	Endangered
<u><i>Drosophila sharpi</i></u>	Hawaiian picture-wing fly	Drosophilidae	Endangered
<u><i>Drosophila substenoptera</i></u>	Hawaiian picture-wing fly	Drosophilidae	Endangered
<u><i>Drosophila tarphytrichia</i></u>	Hawaiian picture-wing fly	Drosophilidae	Endangered
<u><i>Drymarchon corais couperi</i></u>	Eastern indigo snake	Colubridae	Threatened
<u><i>Dryopteris crinalis</i> var. <i>podosorus</i></u>	Palapalai aumakua	Dryopteridaceae	Endangered
<u><i>Dryopteris glabra</i> var. <i>pusilla</i></u>	Hohiu	Dryopteridaceae	Endangered
<u><i>Dubautia herbstobatae</i></u>	Na`ena`e	Asteraceae	Endangered
<u><i>Dubautia imbricata</i> ssp. <i>imbricata</i></u>	Na`ena`e	Asteraceae	Endangered
<u><i>Dubautia kalalauensis</i></u>	Naenae	Asteraceae	Endangered

Scientific Name	Common Name	Family	Federal Listing Status
<u><i>Dubautia kenwoodii</i></u>	Naenae	Asteraceae	Endangered
<u><i>Dubautia latifolia</i></u>	Koholapehu	Asteraceae	Endangered
<u><i>Dubautia pauciflora</i></u>	Na`ena`e	Asteraceae	Endangered
<u><i>Dubautia plantaginea ssp. humilis</i></u>	Na`ena`e	Asteraceae	Endangered
<u><i>Dubautia plantaginea ssp. magnifolia</i></u>	Na`ena`e	Asteraceae	Endangered
<u><i>Dubautia waialealae</i></u>	Na`ena`e	Asteraceae	Endangered
<u><i>Dudleya abramsii ssp. parva</i></u>	Conejo dudleya	Crassulaceae	Threatened
<u><i>Dudleya cymosa ssp. marcescens</i></u>	Marcescent dudleya	Crassulaceae	Threatened
<u><i>Dudleya cymosa ssp. ovatifolia</i></u>	Santa Monica Mountains dudleyea	Crassulaceae	Threatened
<u><i>Dudleya nesiotica</i></u>	Santa Cruz Island dudleya	Crassulaceae	Threatened
<u><i>Dudleya setchellii</i></u>	Santa Clara Valley dudleya	Crassulaceae	Endangered
<u><i>Dudleya stolonifera</i></u>	Laguna Beach liveforever	Crassulaceae	Threatened
<u><i>Dudleya traskiae</i></u>	Santa Barbara Island liveforever	Crassulaceae	Endangered
<u><i>Dudleya verityi</i></u>	Verity's dudleya	Crassulaceae	Threatened
<u><i>Echinacea laevigata</i></u>	Smooth coneflower	Asteraceae	Endangered
<u><i>Echinocactus horizonthalonius var. nicholii</i></u>	Nichol's Turk's head cactus	Cactaceae	Endangered
<u><i>Echinocereus chisoensis var. chisoensis</i></u>	Chisos Mountain hedgehog Cactus	Cactaceae	Threatened
<u><i>Echinocereus fendleri var. kuenzleri</i></u>	Kuenzler hedgehog cactus	Cactaceae	Threatened
<u><i>Echinocereus reichenbachii var. albertii</i></u>	Black lace cactus	Cactaceae	Endangered
<u><i>Echinocereus triglochidiatus var. arizonicus</i></u>	Arizona hedgehog cactus	Cactaceae	Endangered
<u><i>Echinocereus viridiflorus var. davisii</i></u>	Davis' green pitaya	Cactaceae	Endangered
<u><i>Echinomastus erectocentrus var. acunensis</i></u>	Acuna Cactus	Cactaceae	Endangered

Scientific Name	Common Name	Family	Federal Listing Status
<u><i>Echinomastus mariposensis</i></u>	Lloyd's Mariposa cactus	Cactaceae	Threatened
<u><i>Elaphoglossum serpens</i></u>	No common name	Lomariopsidaceae	Endangered
<u><i>Elaphrus viridis</i></u>	Delta green ground beetle	Carabidae	Threatened
<u><i>Elassoma alabamae</i></u>	Spring pygmy sunfish	Elassomatidae	Threatened
<u><i>Eleutherodactylus cooki</i></u>	Guajon	Leptodactylidae	Threatened
<u><i>Eleutherodactylus jasperi</i></u>	Golden coqui	Leptodactylidae	Threatened
<u><i>Eleutherodactylus juanariveroi</i></u>	Llanero Coqui	Leptodactylidae	Endangered
<u><i>Elimia crenatella</i></u>	Lacy elimia (snail)	Pleuroceridae	Threatened
<u><i>Elliptio chipolaensis</i></u>	Chipola slabshell	Unionidae	Threatened
<u><i>Elliptio lanceolata</i></u>	Yellow lance	Unionidae	Threatened
<u><i>Elliptio spinosa</i></u>	Altamaha Spiny mussel	Unionidae	Endangered
<u><i>Elliptio steinstansana</i></u>	Tar River spiny mussel	Unionidae	Endangered
<u><i>Elliptoideus sloatianus</i></u>	Purple bankclimber (mussel)	Unionidae	Threatened
<u><i>Emballonura semicaudata rotensis</i></u>	Pacific sheath-tailed Bat	Emballonuridae	Endangered
<u><i>Emballonura semicaudata semicaudata</i></u>	Pacific sheath-tailed Bat	Emballonuridae	Endangered
<u><i>Emoia slevini</i></u>	Slevin's skink	Scincidae	Endangered
<u><i>Empetrichthys latos</i></u>	Pahrump poolfish	Cyprinodontidae	Endangered
<u><i>Empidonax traillii extimus</i></u>	Southwestern willow flycatcher	Tyrannidae	Endangered
<u><i>Enceliopsis nudicaulis var. corrugata</i></u>	Ash Meadows sunray	Asteraceae	Threatened
<u><i>Enhydra lutris kenyon</i></u>	Northern Sea Otter	Mustelidae	Threatened
<u><i>Enhydra lutris nereis</i></u>	Southern sea otter	Mustelidae	Threatened
<u><i>Epicrates inornatus</i></u>	Puerto Rican boa	Boidae	Endangered
<u><i>Epicrates monensis monensis</i></u>	Mona boa	Boidae	Threatened
<u><i>Epinephelus striatus</i></u>	Nassau grouper	Serranidae	Threatened
<u><i>Epioblasma brevidens</i></u>	Cumberlandian combshell	Unionidae	Endangered
<u><i>Epioblasma capsaeformis</i></u>	Oyster mussel	Unionidae	Endangered

Scientific Name	Common Name	Family	Federal Listing Status
<u><i>Epioblasma florentina curtisii</i></u>	Curtis pearlymussel	Unionidae	Endangered
<u><i>Epioblasma florentina florentina</i></u>	Yellow blossom (pearlymussel)	Unionidae	Endangered
<u><i>Epioblasma florentina walkeri</i> (=E. walkeri)</u>	Tan riffleshell	Unionidae	Endangered
<u><i>Epioblasma metastriata</i></u>	Upland combshell	Unionidae	Endangered
<u><i>Epioblasma obliquata obliquata</i></u>	Purple Cat's paw (=Purple Cat's paw pearlymussel)	Unionidae	Endangered
<u><i>Epioblasma obliquata perobliqua</i></u>	White catspaw (pearlymussel)	Unionidae	Endangered
<u><i>Epioblasma othcaloogensis</i></u>	Southern acornshell	Unionidae	Endangered
<u><i>Epioblasma penita</i></u>	Southern combshell	Unionidae	Endangered
<u><i>Epioblasma torulosa gubernaculum</i></u>	Green blossom (pearlymussel)	Unionidae	Endangered
<u><i>Epioblasma torulosa rangiana</i></u>	Northern riffleshell	Unionidae	Endangered
<u><i>Epioblasma torulosa torulosa</i></u>	Tubercled blossom (pearlymussel)	Unionidae	Endangered
<u><i>Epioblasma triquetra</i></u>	Snuffbox mussel	Unionidae	Endangered
<u><i>Epioblasma turgidula</i></u>	Turgid blossom (pearlymussel)	Unionidae	Endangered
<u><i>Eragrostis fosbergii</i></u>	Fosberg's love grass	Poaceae	Endangered
<u><i>Eremalche kernensis</i></u>	Kern mallow	Malvaceae	Endangered
<u><i>Eremichthys acros</i></u>	Desert dace	Cyprinidae	Threatened
<u><i>Eremophila alpestris strigata</i></u>	Streaked Horned lark	Alaudidae	Threatened
<u><i>Eretmochelys imbricata</i></u>	Hawksbill sea turtle	Cheloniidae	Endangered
<u><i>Eriastrum densifolium ssp. sanctorum</i></u>	Santa Ana River woolly-star	Polemoniaceae	Endangered
<u><i>Erigeron decumbens</i></u>	Willamette daisy	Asteraceae	Endangered
<u><i>Erigeron parishii</i></u>	Parish's daisy	Asteraceae	Threatened
<u><i>Erigeron rhizomatus</i></u>	Zuni fleabane	Asteraceae	Threatened
<u><i>Erignathus barbatus nauticus</i></u>	bearded Seal	Phocidae	Threatened
<u><i>Erimonax monachus</i></u>	Spotfin Chub	Cyprinidae	Threatened

Scientific Name	Common Name	Family	Federal Listing Status
<u><i>Erimystax cahni</i></u>	Slender chub	Cyprinidae	Threatened 1
<u><i>Erinna newcombi</i></u>	Newcomb's snail	Lymnaeidae	Threatened 1
<u><i>Eriodictyon altissimum</i></u>	Indian Knob mountainbalm	Namaceae	Endangered 1
<u><i>Eriodictyon capitatum</i></u>	Lompoc yerba santa	Hydrophyllaceae	Endangered 1
<u><i>Eriogonum apricum</i> (incl. var. <i>prostratum</i>)</u>	Ione (incl. Irish Hill) buckwheat	Polygonaceae	Endangered 1
<u><i>Eriogonum codium</i></u>	Umtanum Desert buckwheat	Polygonaceae	Threatened 1
<u><i>Eriogonum gypsophilum</i></u>	Gypsum wild-buckwheat	Polygonaceae	Threatened 1
<u><i>Eriogonum kennedyi</i> var. <i>austromontanum</i></u>	Southern mountain wild-buckwheat	Polygonaceae	Threatened 1
<u><i>Eriogonum longifolium</i> var. <i>gnaphalifolium</i></u>	Scrub buckwheat	Polygonaceae	Threatened 1
<u><i>Eriogonum ovalifolium</i> var. <i>vineum</i></u>	Cushenbury buckwheat	Polygonaceae	Endangered 1
<u><i>Eriogonum ovalifolium</i> var. <i>williamsiae</i></u>	Steamboat buckwheat	Polygonaceae	Endangered 1
<u><i>Eriogonum pelinophilum</i></u>	Clay-Loving wild buckwheat	Polygonaceae	Endangered 1
<u><i>Eriophyllum latilobum</i></u>	San Mateo woolly sunflower	Asteraceae	Endangered 1
<u><i>Eryngium aristulatum</i> var. <i>parishii</i></u>	San Diego button-celery	Apiaceae	Endangered 1
<u><i>Eryngium constancei</i></u>	Loch Lomond coyote thistle	Apiaceae	Endangered 1
<u><i>Eryngium cuneifolium</i></u>	Snakeroot	Apiaceae	Endangered 1
<u><i>Erysimum capitatum</i> var. <i>angustatum</i></u>	Contra Costa wallflower	Brassicaceae	Endangered 1
<u><i>Erysimum menziesii</i></u>	Menzies' wallflower	Brassicaceae	Endangered 1
<u><i>Erysimum teretifolium</i></u>	Ben Lomond wallflower	Brassicaceae	Endangered 1
<u><i>Erythronium propullans</i></u>	Minnesota dwarf trout lily	Liliaceae	Endangered 1
<u><i>Etheostoma akatulo</i></u>	bluemask darter	Percidae	Endangered 1
<u><i>Etheostoma boschungii</i></u>	Slackwater darter	Percidae	Threatened 1
<u><i>Etheostoma chermocki</i></u>	Vermilion darter	Percidae	Endangered 1
<u><i>Etheostoma chienense</i></u>	Relict darter	Percidae	Endangered 1

1

1

Scientific Name	Common Name	Family	Federal Listing Status
<u><i>Etheostoma etowahae</i></u>	Etowah darter	Percidae	Endangered
<u><i>Etheostoma fonticola</i></u>	Fountain darter	Percidae	Endangered
<u><i>Etheostoma moorei</i></u>	Yellowcheek Darter	Percidae	Endangered
<u><i>Etheostoma nianguae</i></u>	Niangua darter	Percidae	Threatened
<u><i>Etheostoma nuchale</i></u>	Watercress darter	Percidae	Endangered
<u><i>Etheostoma okaloosae</i></u>	Okaloosa darter	Percidae	Threatened
<u><i>Etheostoma osburni</i></u>	Candy darter	Percidae	Endangered
<u><i>Etheostoma percnurum</i></u>	Duskytail darter	Percidae	Endangered
<u><i>Etheostoma phytophilum</i></u>	Rush Darter	Percidae	Endangered
<u><i>Etheostoma rubrum</i></u>	Bayou darter	Percidae	Threatened
<u><i>Etheostoma scotti</i></u>	Cherokee darter	Percidae	Threatened
<u><i>Etheostoma sellare</i></u>	Maryland darter	Percidae	Endangered
<u><i>Etheostoma spilotum</i></u>	Kentucky arrow darter	Percidae	Threatened
<u><i>Etheostoma susanae</i></u>	Cumberland darter	Percidae	Endangered
<u><i>Etheostoma trisella</i></u>	Trispot darter	Percidae	Threatened
<u><i>Etheostoma wapiti</i></u>	Boulder darter	Percidae	Endangered
<u><i>Eua zebrina</i></u>	Snail [no common name]	Partulidae	Endangered
<u><i>Eubalaena glacialis</i></u>	North Atlantic Right Whale	Balaenidae	Endangered
<u><i>Eubalaena japonica</i></u>	North Pacific Right Whale	Balaenidae	Endangered
<u><i>Euchloe ausonides insulanus</i></u>	Island marble Butterfly	Pieridae	Endangered
<u><i>Eucyclogobius newberryi</i></u>	Tidewater goby	Gobiidae	Endangered
<u><i>Eugenia bryanii</i></u>	No common name	Myrtaceae	Endangered
<u><i>Eugenia haematocarpa</i></u>	Uvillo	Myrtaceae	Endangered
<u><i>Eugenia koolauensis</i></u>	Nioi	Myrtaceae	Endangered
<u><i>Eugenia woodburyana</i></u>	No common name	Myrtaceae	Endangered
<u><i>Eumeces egregius lividus</i></u>	Bluetail mole skink	Scincidae	Threatened
<u><i>Eumetopias jubatus</i></u>	Steller sea lion	Otariidae	Endangered
<u><i>Eumops floridanus</i></u>	Florida bonneted bat	Molossidae	Endangered
<u><i>Euphilotes battoides allyni</i></u>	El Segundo blue butterfly	Lycaenidae	Endangered

Scientific Name	Common Name	Family	Federal Listing Status
<u><i>Euphilotes enoptes smithi</i></u>	Smith's blue butterfly	Lycaenidae	Endangered
<u><i>Euphorbia celastroides</i> var. <i>kaenana</i></u>	`Akoko	Euphorbiaceae	Endangered
<u><i>Euphorbia deppeana</i></u>	`Akoko	Euphorbiaceae	Endangered
<u><i>Euphorbia eleanoriae</i></u>	`Akoko	Euphorbiaceae	Endangered
<u><i>Euphorbia haeleeleana</i></u>	`Akoko	Euphorbiaceae	Endangered
<u><i>Euphorbia halemanui</i></u>	"Akoko	Euphorbiaceae	Endangered
<u><i>Euphorbia herbstii</i></u>	`Akoko	Euphorbiaceae	Endangered
<u><i>Euphorbia kuwaleana</i></u>	`Akoko	Euphorbiaceae	Endangered
<u><i>Euphorbia remyi</i> var. <i>kauaiensis</i></u>	`Akoko	Euphorbiaceae	Endangered
<u><i>Euphorbia remyi</i> var. <i>remyi</i></u>	`Akoko	Euphorbiaceae	Endangered
<u><i>Euphorbia rockii</i></u>	`Akoko	Euphorbiaceae	Endangered
<u><i>Euphorbia skottsbergii</i> var. <i>skottsbergii</i></u>	Ewa Plains `akoko	Euphorbiaceae	Endangered
<u><i>Euphorbia telephioides</i></u>	Telephus spurge	Euphorbiaceae	Threatened
<u><i>Euphydryas editha bayensis</i></u>	Bay checkerspot butterfly	Nymphalidae	Threatened
<u><i>Euphydryas editha quino</i></u> (= <i>E. e. wrighti</i>)	Quino checkerspot butterfly	Nymphalidae	Endangered
<u><i>Euphydryas editha taylori</i></u>	Taylor's (=whulge) Checkerspot	Nymphalidae	Endangered
<u><i>Euproserpinus euterpe</i></u>	Kern primrose sphinx moth	Sphingidae	Threatened
<u><i>Eurycea chisholmensis</i></u>	Salado Salamander	Plethodontidae	Threatened
<u><i>Eurycea nana</i></u>	San Marcos salamander	Plethodontidae	Threatened
<u><i>Eurycea naufragia</i></u>	Georgetown Salamander	Plethodontidae	Threatened
<u><i>Eurycea sosorum</i></u>	Barton Springs salamander	Plethodontidae	Endangered
<u><i>Eurycea tonkawae</i></u>	Jollyville Plateau Salamander	Plethodontidae	Threatened
<u><i>Eurycea waterlooensis</i></u>	Austin blind Salamander	Plethodontidae	Endangered
<u><i>Eutrema penlandii</i></u>	Penland alpine fen mustard	Brassicaceae	Threatened
<u><i>Exocarpos luteolus</i></u>	Heau	Santalaceae	Endangered
<u><i>Exocarpos menziesii</i></u>	Heau	Santalaceae	Endangered

Scientific Name	Common Name	Family	Federal Listing Status
<u><i>Falco femoralis septentrionalis</i></u>	Northern Aplomado Falcon	Falconidae	Endangered 1
<u><i>Faxonius peruncus</i></u>	Big Creek Crayfish	Cambaridae	Proposed Threatened 1
<u><i>Faxonius quadruncus</i></u>	St. Francis River Crayfish	Cambaridae	Proposed Threatened 1
<u><i>Festuca hawaiiensis</i></u>	No common name	Poaceae	Endangered 1
<u><i>Festuca ligulata</i></u>	Guadalupe fescue	Poaceae	Endangered 1
<u><i>Festuca molokaiensis</i></u>	No common name	Poaceae	Endangered 1
<u><i>Flueggea neowawraea</i></u>	Mehamehame	Phyllanthaceae	Endangered 1
<u><i>Fremontodendron californicum ssp. decumbens</i></u>	Pine Hill flannelbush	Sterculiaceae	Endangered 1
<u><i>Fremontodendron mexicanum</i></u>	Mexican flannelbush	Sterculiaceae	Endangered 1
<u><i>Fritillaria gentneri</i></u>	Gentner's Fritillary	Liliaceae	Endangered 1
<u><i>Fulica americana alai</i></u>	Hawaiian coot	Rallidae	Endangered 1
<u><i>Fundulus julisia</i></u>	Barrens topminnow	Cyprinodontidae	Endangered 1
<u><i>Fusconaia burkei</i></u>	Tapered pigtoe	Unionidae	Threatened 1
<u><i>Fusconaia cor</i></u>	Shiny pigtoe	Unionidae	Endangered 1
<u><i>Fusconaia cuneolus</i></u>	Finerayed pigtoe	Unionidae	Endangered 1
<u><i>Fusconaia escambia</i></u>	Narrow pigtoe	Unionidae	Threatened 1
<u><i>Fusconaia masoni</i></u>	Atlantic pigtoe	Unionidae	Proposed Threatened 1
<u><i>Fusconaia rotulata</i></u>	Round Ebonyshell	Unionidae	Endangered 1
<u><i>Fusconaia subrotunda</i></u>	Longsolid	Unionidae	Proposed Threatened 1
<u><i>Galactia smallii</i></u>	Small's milkpea	Fabaceae	Endangered 1
<u><i>Galium buxifolium</i></u>	Island bedstraw	Rubiaceae	Endangered 1
<u><i>Galium californicum ssp. sierrae</i></u>	El Dorado bedstraw	Rubiaceae	Endangered 1
<u><i>Gallicolumba stairi</i></u>	Friendly Ground-Dove	Columbidae	Endangered 1
<u><i>Gallinula chloropus guami</i></u>	Mariana common moorhen	Rallidae	Endangered 1
<u><i>Gallinula galeata sandvicensis</i></u>	Hawaiian common gallinule	Rallidae	Endangered 1
<u><i>Gambelia silus</i></u>	Blunt-nosed leopard lizard	Crotaphytidae	Endangered 1
<u><i>Gambusia gaigei</i></u>	Big Bend gambusia	Poeciliidae	Endangered

Scientific Name	Common Name	Family	Federal Listing Status
<u><i>Gambusia georgei</i></u>	San Marcos gambusia	Poeciliidae	Endangered
<u><i>Gambusia heterochir</i></u>	Clear Creek gambusia	Poeciliidae	Endangered
<u><i>Gambusia nobilis</i></u>	Pecos gambusia	Poeciliidae	Endangered
<u><i>Gammarus acherondytes</i></u>	Illinois cave amphipod	Gammaridae	Endangered
<u><i>Gammarus desperatus</i></u>	Noel's Amphipod	Gammaridae	Endangered
<u><i>Gammarus hyalleloides</i></u>	Diminutive Amphipod	Gammaridae	Endangered
<u><i>Gammarus pecos</i></u>	Pecos amphipod	Gammaridae	Endangered
<u><i>Gardenia brighamii</i></u>	Hawaiian gardenia (=Na`u)	Rubiaceae	Endangered
<u><i>Gardenia mannii</i></u>	Nanu	Rubiaceae	Endangered
<u><i>Gardenia remyi</i></u>	Nanu	Rubiaceae	Endangered
<u><i>Gasterosteus aculeatus williamsoni</i></u>	Unarmored threespine stickleback	Gasterosteidae	Endangered
<u><i>Geocarpon minimum</i></u>	No common name	Caryophyllaceae	Threatened
<u><i>Geranium arboreum</i></u>	Nohoanu	Geraniaceae	Endangered
<u><i>Geranium hanaense</i></u>	Nohoanu	Geraniaceae	Endangered
<u><i>Geranium hillebrandii</i></u>	Nohoanu	Geraniaceae	Endangered
<u><i>Geranium kauaiense</i></u>	Nohoanu	Geraniaceae	Endangered
<u><i>Geranium multiflorum</i></u>	Nohoanu	Geraniaceae	Endangered
<u><i>Gesneria pauciflora</i></u>	No common name	Gesneriaceae	Threatened
<u><i>Geum radiatum</i></u>	Spreading avens	Rosaceae	Endangered
<u><i>Gila bicolor ssp.</i></u>	Hutton tui chub	Cyprinidae	Threatened
<u><i>Gila bicolor ssp. mohavensis</i></u>	Mohave tui chub	Cyprinidae	Endangered
<u><i>Gila bicolor ssp. snyderi</i></u>	Owens Tui Chub	Cyprinidae	Endangered
<u><i>Gila cypha</i></u>	Humpback chub	Cyprinidae	Endangered
<u><i>Gila ditaenia</i></u>	Sonora chub	Cyprinidae	Threatened
<u><i>Gila elegans</i></u>	Bonytail	Cyprinidae	Endangered
<u><i>Gila intermedia</i></u>	Gila chub	Cyprinidae	Endangered
<u><i>Gila nigrescens</i></u>	Chihuahua chub	Cyprinidae	Threatened
<u><i>Gila purpurea</i></u>	Yaqui chub	Cyprinidae	Endangered

Scientific Name	Common Name	Family	Federal Listing Status
<u><i>Gila robusta jordani</i></u>	Pahranagat roundtail chub	Cyprinidae	Endangered
<u><i>Gila seminuda (=robusta)</i></u>	Virgin River Chub	Cyprinidae	Endangered
<u><i>Gilia tenuiflora ssp. arenaria</i></u>	Monterey gilia	Polemoniaceae	Endangered
<u><i>Gilia tenuiflora ssp. hoffmannii</i></u>	Hoffmann's slender- flowered gilia	Polemoniaceae	Endangered
<u><i>Glaucomys sabrinus coloratus</i></u>	Carolina northern flying squirrel	Sciuridae	Endangered
<u><i>Glaucopsyche lygdamus palosverdesensis</i></u>	Palos Verdes blue butterfly	Lycaenidae	Endangered
<u><i>Goetzea elegans</i></u>	Beautiful goetzea	Solanaceae	Endangered
<u><i>Gonocalyx concolor</i></u>	No common name	Ericaceae	Endangered
<u><i>Gopherus agassizii</i></u>	Desert tortoise	Testudinidae	Threatened
<u><i>Gopherus polyphemus</i></u>	Gopher tortoise	Testudinidae	Threatened
<u><i>Gouania hillebrandii</i></u>	No common name	Rhamnaceae	Endangered
<u><i>Gouania meyenii</i></u>	No common name	Rhamnaceae	Endangered
<u><i>Gouania vitifolia</i></u>	No common name	Rhamnaceae	Endangered
<u><i>Graptemys flavimaculata</i></u>	Yellow-blotched map turtle	Emydidae	Threatened
<u><i>Graptemys oculifera</i></u>	Ringed map turtle	Emydidae	Threatened
<u><i>Graptopetalum bartramii</i></u>	Bartram stonecrop	Crassulaceae	Proposed Threatened
<u><i>Grindelia fraxinipratensis</i></u>	Ash Meadows gumplant	Asteraceae	Threatened
<u><i>Grus americana</i></u>	Whooping crane	Gruidae	Endangered
<u><i>Grus canadensis pulla</i></u>	Mississippi sandhill crane	Gruidae	Endangered
<u><i>Gymnoderma lineare</i></u>	Rock gnome lichen	Cladoniaceae	Endangered
<u><i>Gymnogyps californianus</i></u>	California condor	Cathartidae	Endangered
<u><i>Gymnomyza samoensis</i></u>	Mao (= maomao) (honeyeater)	Meliphagidae	Endangered
<u><i>Hackelia venusta</i></u>	Showy stickseed	Boraginaceae	Endangered
<u><i>Halcyon cinnamomina cinnamomina</i></u>	Guam Micronesian kingfisher	Alcedinidae	Endangered
<u><i>Haliotis cracherodii</i></u>	Black Abalone	Haliotidae	Endangered
<u><i>Haliotis sorenseni</i></u>	White Abalone	Haliotidae	Endangered

Scientific Name	Common Name	Family	Federal Listing Status
<u><i>Halophila johnsonii</i></u>	Johnson's seagrass	Hydrocharitaceae	Threatened
<u><i>Hamiota australis</i></u>	Southern sandshell	Unionidae	Threatened
<u><i>Haplostachys haplostachya</i></u>	Honohono	Lamiaceae	Endangered
<u><i>Harperocallis flava</i></u>	Harper's beauty	Liliaceae	Endangered
<u><i>Harrisia</i> (=Cereus) <i>aboriginum</i> (=gracilis)</u>	Aboriginal Prickly-apple	Cactaceae	Endangered
<u><i>Harrisia portoricensis</i></u>	Higo Chumbo	Cactaceae	Threatened
<u><i>Hedeoma todsenii</i></u>	Todsen's pennyroyal	Lamiaceae	Endangered
<u><i>Hedyotis megalantha</i></u>	Paudedo	Rubiaceae	Endangered
<u><i>Hedyotis purpurea</i> var. <i>montana</i></u>	Roan Mountain bluet	Rubiaceae	Endangered
<u><i>Helenium virginicum</i></u>	Virginia sneezeweed	Asteraceae	Threatened
<u><i>Helianthemum greenei</i></u>	Island rush-rose	Cistaceae	Threatened
<u><i>Helianthus paradoxus</i></u>	Pecos (=puzzle, =paradox) sunflower	Asteraceae	Threatened
<u><i>Helianthus schweinitzii</i></u>	Schweinitz's sunflower	Asteraceae	Endangered
<u><i>Helianthus verticillatus</i></u>	Whorled Sunflower	Asteraceae	Endangered
<u><i>Helminthoglypta walkeriana</i></u>	Morro shoulderband (=Banded dune) snail	Helminthoglyptida	Endangered
<u><i>Helonias bullata</i></u>	Swamp pink	Liliaceae	Threatened
<u><i>Hemignathus affinis</i></u>	Maui nukupuu	Fringillidae	Endangered
<u><i>Hemignathus hanapepe</i></u>	Kauai nukupuu	Fringillidae	Endangered
<u><i>Hemignathus wilsoni</i></u>	akiapolaau	Drepanidinae	Endangered
<u><i>Hemistena lata</i></u>	Cracking pearlymussel	Unionidae	Endangered
<u><i>Heraclides aristodemus</i> <i>ponceanus</i></u>	Schaus swallowtail butterfly	Papilionidae	Endangered
<u><i>Heritiera longipetiolata</i></u>	Ufa-halomtano	Sterculiaceae	Endangered
<u><i>Herpailurus</i> (=Felis) <i>yagouaroundi cacomitli</i></u>	Gulf Coast jaguarundi	Felidae	Endangered
<u><i>Herpailurus</i> (=Felis) <i>yagouaroundi tolteca</i></u>	Sinaloan Jaguarundi	Felidae	Endangered

Scientific Name	Common Name	Family	Federal Listing Status
<u><i>Hesperia dacotae</i></u>	Dakota Skipper	Hesperiidae	Threatened
<u><i>Hesperia leonardus montana</i></u>	Pawnee montane skipper	Hesperiidae	Threatened
<u><i>Hesperolinon congestum</i></u>	Marin dwarf-flax	Linaceae	Threatened
<u><i>Hesperomannia arborescens</i></u>	No common name	Asteraceae	Endangered
<u><i>Hesperomannia arbuscula</i></u>	No common name	Asteraceae	Endangered
<u><i>Hesperomannia lydgatei</i></u>	No common name	Asteraceae	Endangered
<u><i>Heterelmis comalensis</i></u>	Comal Springs riffle beetle	Elmidae	Endangered
<u><i>Hexastylis naniflora</i></u>	Dwarf-flowered heartleaf	Aristolochiaceae	Threatened
<u><i>Hibiscadelphus distans</i></u>	Kauai hau kuahiwi	Malvaceae	Endangered
<u><i>Hibiscadelphus giffardianus</i></u>	Hau kuahiwi	Malvaceae	Endangered
<u><i>Hibiscadelphus hualalaiensis</i></u>	Hau kuahiwi	Malvaceae	Endangered
<u><i>Hibiscadelphus woodii</i></u>	Hau kuahiwi	Malvaceae	Endangered
<u><i>Hibiscus arnottianus</i> ssp. <i>immaculatus</i></u>	Koki`o ke`oke`o	Malvaceae	Endangered
<u><i>Hibiscus brackenridgei</i></u>	(=Native yellow hibiscus) ma`o hau hele	Malvaceae	Endangered
<u><i>Hibiscus clayi</i></u>	Clay's hibiscus	Malvaceae	Endangered
<u><i>Hibiscus dasycalyx</i></u>	Neches River rose-mallow	Malvaceae	Threatened
<u><i>Hibiscus waimeae</i> ssp. <i>hannerae</i></u>	Koki`o ke`oke`o	Malvaceae	Endangered
<u><i>Himantopus mexicanus knudseni</i></u>	Hawaiian stilt	Recurvirostridae	Endangered
<u><i>Hoffmannseggia tenella</i></u>	Slender rush-pea	Fabaceae	Endangered
<u><i>Holocarpha macradenia</i></u>	Santa Cruz tarplant	Asteraceae	Threatened
<u><i>Howellia aquatilis</i></u>	Water howellia	Campanulaceae	Threatened
<u><i>Hudsonia montana</i></u>	Mountain golden heather	Cistaceae	Threatened
<u><i>Huperzia mannii</i></u>	Wawae`iole	Lycopodiaceae	Endangered
<u><i>Huperzia nutans</i></u>	Wawae`iole	Lycopodiaceae	Endangered
<u><i>Huperzia stemmermanniae</i></u>	No common name	Lycopodiaceae	Endangered
<u><i>Hybognathus amarus</i></u>	Rio Grande Silvery Minnow	Cyprinidae	Endangered

Scientific Name	Common Name	Family	Federal Listing Status
<u><i>Hylaeus anthracinus</i></u>	Anthricinan yellow-faced bee	Hylaeidae	Endangered
<u><i>Hylaeus assimulans</i></u>	Assimulans yellow-faced bee	Hylaeidae	Endangered
<u><i>Hylaeus facilis</i></u>	Easy yellow-faced bee	Hylaeidae	Endangered
<u><i>Hylaeus hiliaris</i></u>	Hilaris yellow-faced bee	Hylaeidae	Endangered
<u><i>Hylaeus kuakea</i></u>	Hawaiian yellow-faced bee	Colletidae	Endangered
<u><i>Hylaeus longiceps</i></u>	Hawaiian yellow-faced bee	Hylaeidae	Endangered
<u><i>Hylaeus mana</i></u>	Hawaiian yellow-faced bee	Colletidae	Endangered
<u><i>Hymenoxys herbacea</i></u>	Lakeside daisy	Asteraceae	Threatened
<u><i>Hymenoxys texana</i></u>	Texas prairie dawn-flower	Asteraceae	Endangered
<u><i>Hypericum cumulicola</i></u>	Highlands scrub hypericum	Hypericaceae	Endangered
<u><i>Hypolepis hawaiiensis</i> var. <i>mauiensis</i></u>	olua	Dennstaedtiaceae	Endangered
<u><i>Hypolimnys octocula marianensis</i></u>	Mariana eight-spot butterfly	Nymphalidae	Endangered
<u><i>Hypomesus transpacificus</i></u>	Delta smelt	Osmeridae	Threatened
<u><i>Icaricia (Plebejus) shasta charlestonensis</i></u>	Mount Charleston blue butterfly	Lycaenidae	Endangered
<u><i>Icaricia icarioides fenderi</i></u>	Fender's blue butterfly	Lycaenidae	Endangered
<u><i>Icaricia icarioides missionensis</i></u>	Mission blue butterfly	Lycaenidae	Endangered
<u><i>Ictalurus pricei</i></u>	Yaqui catfish	Ictaluridae	Threatened
<u><i>Ilex cookii</i></u>	Cook's holly	Aquifoliaceae	Endangered
<u><i>Ilex sintenisii</i></u>	No common name	Aquifoliaceae	Endangered
<u><i>Iliamna corei</i></u>	Peter's Mountain mallow	Malvaceae	Endangered
<u><i>Ipomopsis polyantha</i></u>	Pagosa skyrocket	Polemoniaceae	Endangered
<u><i>Ipomopsis sancti-spiritus</i></u>	Holy Ghost ipomopsis	Polemoniaceae	Endangered
<u><i>Iris lacustris</i></u>	Dwarf lake iris	Iridaceae	Threatened
<u><i>Ischaemum byrone</i></u>	Hilo ischaemum	Poaceae	Endangered
<u><i>Ischnura luta</i></u>	Rota blue damselfly	coenagrionidae	Endangered
<u><i>Isodendron hosakae</i></u>	Aupaka	Violaceae	Endangered

Scientific Name	Common Name	Family	Federal Listing Status
<u><i>Isodendron laurifolium</i></u>	Aupaka	Violaceae	Endangered
<u><i>Isodendron longifolium</i></u>	Aupaka	Violaceae	Threatened
<u><i>Isodendron pyrifolium</i></u>	Kula wahine noho	Violaceae	Endangered
<u><i>Isoetes louisianensis</i></u>	Louisiana quillwort	Isoetaceae	Endangered
<u><i>Isoetes melanospora</i></u>	Black spored quillwort	Isoetaceae	Endangered
<u><i>Isoetes tegetiformans</i></u>	Mat-forming quillwort	Isoetaceae	Endangered
<u><i>Isotria medeoloides</i></u>	Small whorled pogonia	Orchidaceae	Threatened
<u><i>Ivesia kingii</i> var. <i>eremica</i></u>	Ash Meadows ivesia	Rosaceae	Threatened
<u><i>Ivesia webberi</i></u>	Webber's ivesia	Rosaceae	Threatened
<u><i>Jacquemontia reclinata</i></u>	Beach jacquemontia	Convolvulaceae	Endangered
<u><i>Joinvillea ascendens</i></u> <u><i>ascendens</i></u>	`Ohe	Joinvilleaceae	Endangered
<u><i>Juglans jamaicensis</i></u>	West Indian Walnut (=Nogal)	Juglandaceae	Endangered
<u><i>Justicia cooleyi</i></u>	Cooley's water-willow	Acanthaceae	Endangered
<u><i>Juturnia kosteri</i></u>	Koster's springsnail	Hydrobiidae	Endangered
<u><i>Kadua cookiana</i></u>	'Awiwi	Rubiaceae	Endangered
<u><i>Kadua cordata</i> <i>remyi</i></u>	kopa	Rubiaceae	Endangered
<u><i>Kadua coriacea</i></u>	Kio`ele	Rubiaceae	Endangered
<u><i>Kadua degeneri</i></u>	No common name	Rubiaceae	Endangered
<u><i>Kadua fluviatilis</i></u>	Kamapua`a	Rubiaceae	Endangered
<u><i>Kadua haupuensis</i></u>	No common name	Rubiaceae	Endangered
<u><i>Kadua laxiflora</i></u>	pilo	Rubiaceae	Endangered
<u><i>Kadua parvula</i></u>	No common name	Rubiaceae	Endangered
<u><i>Kadua st.-johnii</i></u>	No common name	Rubiaceae	Endangered
<u><i>Kanaloa kahoolawensis</i></u>	Kohe malama malama o kanaloa	Fabaceae	Endangered
<u><i>Keysseria</i> (= <i>Lagenifera</i>) <i>erici</i></u>	No common name	Asteraceae	Endangered
<u><i>Keysseria</i> (= <i>Lagenifera</i>) <i>helenae</i></u>	No common name	Asteraceae	Endangered

Scientific Name	Common Name	Family	Federal Listing Status
<u><i>Kinosternon sonoriense longifemorale</i></u>	Sonoyta mud turtle	Kinosternidae	Endangered
<u><i>Kokia cookei</i></u>	Cooke's koki`o	Malvaceae	Endangered
<u><i>Kokia drynarioides</i></u>	Koki`o	Malvaceae	Endangered
<u><i>Kokia kauaiensis</i></u>	Koki`o	Malvaceae	Endangered
<u><i>Korthalsella degeneri</i></u>	Hulumoa	Santalaceae	Endangered
<u><i>Labordia cyrtandrae</i></u>	Kamakahala	Loganiaceae	Endangered
<u><i>Labordia helleri</i></u>	Kamakahala	Loganiaceae	Endangered
<u><i>Labordia lorenciana</i></u>	No common name	Loganiaceae	Endangered
<u><i>Labordia lydgatei</i></u>	Kamakahala	Loganiaceae	Endangered
<u><i>Labordia pumila</i></u>	Kamakahala	Loganiaceae	Endangered
<u><i>Labordia tinifolia</i> var. <i>lanaiensis</i></u>	Kamakahala	Loganiaceae	Endangered
<u><i>Labordia tinifolia</i> var. <i>wahiawaensis</i></u>	Kamakahala	Loganiaceae	Endangered
<u><i>Labordia triflora</i></u>	Kamakahala	Loganiaceae	Endangered
<u><i>Lampsilis abrupta</i></u>	Pink mucket (pearlymussel)	Unionidae	Endangered
<u><i>Lampsilis altilis</i></u>	Finelined pocketbook	Unionidae	Threatened
<u><i>Lampsilis higginsii</i></u>	Higgins eye (pearlymussel)	Unionidae	Endangered
<u><i>Lampsilis perovalis</i></u>	Orangenacre mucket	Unionidae	Threatened
<u><i>Lampsilis powellii</i></u>	Arkansas fatmucket	Unionidae	Threatened
<u><i>Lampsilis rafinesqueana</i></u>	Neosho Mucket	Unionidae	Endangered
<u><i>Lampsilis streckeri</i></u>	Speckled pocketbook	Unionidae	Endangered
<u><i>Lampsilis subangulata</i></u>	Shinyrayed pocketbook	Unionidae	Endangered
<u><i>Lampsilis virescens</i></u>	Alabama lampmussel	Unionidae	Endangered
<u><i>Lanius ludovicianus mearnsi</i></u>	San Clemente loggerhead shrike	Laniidae	Endangered
<u><i>Lanx sp.</i></u>	Banbury Springs limpet	Lymnaeidae	Endangered
<u><i>Lasiurus cinereus semotus</i></u>	Hawaiian hoary bat	Vespertilionidae	Endangered
<u><i>Lasmigona decorata</i></u>	Carolina heelsplitter	Unionidae	Endangered

Scientific Name	Common Name	Family	Federal Listing Status
<u><i>Lasthenia burkei</i></u>	Burke's goldfields	Asteraceae	Endangered
<u><i>Lasthenia conjugens</i></u>	Contra Costa goldfields	Asteraceae	Endangered
<u><i>Laterallus jamaicensis ssp. jamaicensis</i></u>	Eastern Black rail	Rallidae	Threatened
<u><i>Layia carnosa</i></u>	Beach layia	Asteraceae	Endangered
<u><i>Leavenworthia crassa</i></u>	Fleshy-fruit glade cress	Brassicaceae	Endangered
<u><i>Leavenworthia exigua laciniata</i></u>	Kentucky glade cress	Brassicaceae	Threatened
<u><i>Leavenworthia texana</i></u>	Texas golden Glade cress	Brassicaceae	Endangered
<u><i>Lednia tumana</i></u>	Meltwater lednian stonefly	Nemouridae	Threatened
<u><i>Lemiox rimosus</i></u>	Birdwing pearly mussel	Unionidae	Endangered
<u><i>Leopardus (=Felis) pardalis</i></u>	Ocelot	Felidae	Endangered
<u><i>Leopardus (=Felis) wiedii</i></u>	Margay	Felidae	Endangered
<u><i>Lepanthes eltoroensis</i></u>	No common name	Orchidaceae	Endangered
<u><i>Lepidium arbuscula</i></u>	Anaunau	Brassicaceae	Endangered
<u><i>Lepidium barnebyanum</i></u>	Barneby ridge-cress	Brassicaceae	Endangered
<u><i>Lepidium orbiculare</i></u>	No common name	Brassicaceae	Endangered
<u><i>Lepidium papilliferum</i></u>	Slickspot peppergrass	Brassicaceae	Threatened
<u><i>Lepidochelys kempii</i></u>	Kemp's ridley sea turtle	Cheloniidae	Endangered
<u><i>Lepidochelys olivacea</i></u>	Olive ridley sea turtle	Cheloniidae	Threatened
<u><i>Lepidomeda albivallis</i></u>	White River spinedace	Cyprinidae	Endangered
<u><i>Lepidomeda mollispinis pratensis</i></u>	Big Spring spinedace	Cyprinidae	Threatened
<u><i>Lepidomeda vittata</i></u>	Little Colorado spinedace	Cyprinidae	Threatened
<u><i>Lepidurus packardii</i></u>	Vernal pool tadpole shrimp	Caenestheriidae	Endangered
<u><i>Leptocereus grantianus</i></u>	No common name	Cactaceae	Endangered
<u><i>Leptodea leptodon</i></u>	Scaleshell mussel	Unionidae	Endangered
<u><i>Leptonycteris nivalis</i></u>	Mexican long-nosed bat	Phyllostomidae	Endangered
<u><i>Leptoxis ampla</i></u>	Round rocksnail	Pleuroceridae	Threatened

Scientific Name	Common Name	Family	Federal Listing Status
<u><i>Leptoxis foremani</i></u>	Interrupted (=Georgia) Rocksnailed	Pleuroceridae	Endangered 1
<u><i>Leptoxis plicata</i></u>	Plicate rocksnailed	Pleuroceridae	Endangered 1
<u><i>Leptoxis taeniata</i></u>	Painted rocksnailed	Pleuroceridae	Threatened 1
<u><i>Lepyrium showalteri</i></u>	Flat pebblesnailed	Hydrobiidae	Endangered 1
<u><i>Lespedeza leptostachya</i></u>	Prairie bush-clover	Fabaceae	Threatened 1
<u><i>Lesquerella congesta</i></u>	Dudley Bluffs bladderpod	Brassicaceae	Threatened 1
<u><i>Lesquerella kingii ssp. bernardina</i></u>	San Bernardino Mountains bladderpod	Brassicaceae	Endangered 1
<u><i>Lesquerella lyrata</i></u>	Lyrate bladderpod	Brassicaceae	Threatened 1
<u><i>Lesquerella pallida</i></u>	White bladderpod	Brassicaceae	Endangered 1
<u><i>Lesquerella perforata</i></u>	Spring Creek bladderpod	Brassicaceae	Endangered 1
<u><i>Lesquerella thamnophila</i></u>	Zapata bladderpod	Brassicaceae	Endangered 1
<u><i>Lesquerella tumulosa</i></u>	Kodachrome bladderpod	Brassicaceae	Endangered 1
<u><i>Lessingia germanorum</i></u> (= <i>L. g. var. germanorum</i>)	San Francisco lessingia	Asteraceae	Endangered 1
<u><i>Liatris helleri</i></u>	Heller's blazingstar	Asteraceae	Threatened 1
<u><i>Liatris ohlingerae</i></u>	Scrub blazingstar	Asteraceae	Endangered 1
<u><i>Lilaeopsis schaffneriana var. recurva</i></u>	Huachuca water-umbel	Apiaceae	Endangered 1
<u><i>Lilium occidentale</i></u>	Western lily	Liliaceae	Endangered 1
<u><i>Lilium pardalinum ssp. pitkinense</i></u>	Pitkin Marsh lily	Liliaceae	Endangered 1
<u><i>Limnanthes floccosa ssp. californica</i></u>	Butte County meadowfoam	Limnanthaceae	Endangered 1
<u><i>Limnanthes pumila ssp. grandiflora</i></u>	Large-flowered woolly meadowfoam	Limnanthaceae	Endangered 1
<u><i>Limnanthes vinculans</i></u>	Sebastopol meadowfoam	Limnanthaceae	Endangered 1
<u><i>Lindera melissifolia</i></u>	Pondberry	Lauraceae	Endangered 1
<u><i>Linum arenicola</i></u>	Sand flax	Linaceae	Endangered 1
<u><i>Linum carteri carteri</i></u>	Carter's small-flowered flax	Linaceae	Endangered 1

Scientific Name	Common Name	Family	Federal Listing Status
<u><i>Lioplax cyclostomaformis</i></u>	Cylindrical lioplax (snail)	Viviparidae	Endangered
<u><i>Lipochaeta fauriei</i></u>	Nehe	Asteraceae	Endangered
<u><i>Lipochaeta lobata</i> var. <i>leptophylla</i></u>	Nehe	Asteraceae	Endangered
<u><i>Lipochaeta micrantha</i></u>	Nehe	Asteraceae	Endangered
<u><i>Lipochaeta venosa</i></u>	No common name	Asteraceae	Endangered
<u><i>Lipochaeta waimeaensis</i></u>	Nehe	Asteraceae	Endangered
<u><i>Lirceus usdagalun</i></u>	Lee County cave isopod	Asellidae	Endangered
<u><i>Lithophragma maximum</i></u>	San Clemente Island woodland-star	Saxifragaceae	Endangered
<u><i>Lobelia koolauensis</i></u>	No common name	Campanulaceae	Endangered
<u><i>Lobelia monostachya</i></u>	No common name	Campanulaceae	Endangered
<u><i>Lobelia niihauensis</i></u>	No common name	Campanulaceae	Endangered
<u><i>Lobelia oahuensis</i></u>	No common name	Campanulaceae	Endangered
<u><i>Lomatium bradshawii</i></u>	Bradshaw's desert-parsley	Apiaceae	Endangered
<u><i>Lomatium cookii</i></u>	Cook's lomatium	Apiaceae	Endangered
<u><i>Loxioides bailleui</i></u>	Palila (honeycreeper)	Drepanidinae	Endangered
<u><i>Loxops caeruleirostris</i></u>	Akekee	Fringillidae	Endangered
<u><i>Loxops coccineus</i></u>	Hawaii akepa	Drepanidinae	Endangered
<u><i>Loxops ochraceus</i></u>	Maui akepa	Drepanidinae	Endangered
<u><i>Lupinus aridorum</i></u>	Scrub lupine	Fabaceae	Endangered
<u><i>Lupinus nipomensis</i></u>	Nipomo Mesa lupine	Fabaceae	Endangered
<u><i>Lupinus sulphureus</i> ssp. <i>kincaidii</i></u>	Kincaid's Lupine	Fabaceae	Threatened
<u><i>Lupinus tidestromii</i></u>	Clover lupine	Fabaceae	Endangered
<u><i>Lycaeides argyrognomon</i> <i>lotis</i></u>	Lotis blue butterfly	Lycaenidae	Endangered
<u><i>Lycaeides melissa samuelis</i></u>	Karner blue butterfly	Lycaenidae	Endangered
<u><i>Lycaena hermes</i></u>	Hermes copper butterfly	Lycaenidae	Proposed Threatened
<u><i>Lynx canadensis</i></u>	Canada Lynx	Felidae	Threatened
<u><i>Lyonia truncata</i> var. <i>proctorii</i></u>	No common name	Ericaceae	Endangered

Scientific Name	Common Name	Family	Federal Listing Status
<u><i>Lysimachia asperulaefolia</i></u>	Rough-leaved loosestrife	Primulaceae	Endangered 1
<u><i>Lysimachia daphnoides</i></u>	lehua makanoe	Primulaceae	Endangered 1
<u><i>Lysimachia filifolia</i></u>	No common name	Primulaceae	Endangered
<u><i>Lysimachia iniki</i></u>	No common name	Primulaceae	Endangered
<u><i>Lysimachia lydgatei</i></u>	No common name	Primulaceae	Endangered
<u><i>Lysimachia maxima</i></u>	No common name	Primulaceae	Endangered
<u><i>Lysimachia pendens</i></u>	No common name	Primulaceae	Endangered
<u><i>Lysimachia scopulensis</i></u>	No common name	Primulaceae	Endangered
<u><i>Lysimachia venosa</i></u>	No common name	Primulaceae	Endangered 1
<u><i>Macbridea alba</i></u>	White birds-in-a-nest	Lamiaceae	Threatened 1
<u><i>Maesa walkeri</i></u>	No common name	Myrsinaceae	Threatened 1
<u><i>Malacothamnus clementinus</i></u>	San Clemente Island bush-mallow	Malvaceae	Endangered 1
<u><i>Malacothamnus fasciculatus</i></u> <u>var. nesioticus</u>	Santa Cruz Island bush-mallow	Malvaceae	Endangered 1
<u><i>Malacothrix indecora</i></u>	Santa Cruz Island malacothrix	Asteraceae	Endangered 1
<u><i>Malacothrix squalida</i></u>	Island malacothrix	Asteraceae	Endangered 1
<u><i>Manduca blackburni</i></u>	Blackburn's sphinx moth	Sphingidae	Endangered 1
<u><i>Manihot walkerae</i></u>	Walker's manioc	Euphorbiaceae	Endangered 1
<u><i>Margaritifera hembeli</i></u>	Louisiana pearlshell	Unionidae	Threatened 1
<u><i>Margaritifera marrianae</i></u>	Alabama pearlshell	Margaritiferidae	Endangered 1
<u><i>Marshallia mohrii</i></u>	Mohr's Barbara's buttons	Asteraceae	Threatened 1
<u><i>Marsilea villosa</i></u>	Ihi`ihi	Marsileaceae	Endangered 1
<u><i>Martes caurina</i></u>	Pacific Marten, Coastal Distinct Population Segment	Mustelidae	Threatened 1
<u><i>Masticophis lateralis euryxanthus</i></u>	Alameda whipsnake (=striped racer)	Colubridae	Threatened 1
<u><i>Meda fulgida</i></u>	Spikedace	Cyprinidae	Endangered 1
<u><i>Medionidus acutissimus</i></u>	Alabama moccasinshell	Unionidae	Threatened 1
<u><i>Medionidus parvulus</i></u>	Coosa moccasinshell	Unionidae	Endangered

Scientific Name	Common Name	Family	Federal Listing Status
<u><i>Medionidus penicillatus</i></u>	Gulf moccasinshell	Unionidae	Endangered 1
<u><i>Medionidus simpsonianus</i></u>	Ochlockonee moccasinshell	Unionidae	Endangered 1
<u><i>Medionidus walkeri</i></u>	Suwannee moccasinshell	Unionidae	Threatened 1
<u><i>Megalagrion leptodemas</i></u>	Crimson Hawaiian damselfly	Coenagrionidae	Endangered 1
<u><i>Megalagrion nesiotes</i></u>	Flying earwig Hawaiian damselfly	Coenagrionidae	Endangered 1
<u><i>Megalagrion nigrohamatum</i></u> <u><i>nigrolineatum</i></u>	Blackline Hawaiian damselfly	Coenagrionidae	Endangered 1
<u><i>Megalagrion oceanicum</i></u>	Oceanic Hawaiian damselfly	Coenagrionidae	Endangered 1
<u><i>Megalagrion pacificum</i></u>	Pacific Hawaiian damselfly	Coenagrionidae	Endangered 1
<u><i>Megalagrion xanthomelas</i></u>	Orangeblack Hawaiian damselfly	Coenagrionidae	Endangered 1
<u><i>Megapodius laperouse</i></u>	Micronesian megapode	Megapodiidae	Endangered 1
<u><i>Megaptera novaeangliae</i></u>	Humpback whale	Balaenopteridae	Threatened 1
<u><i>Melamprosops phaeosoma</i></u>	Po'ouli (honeycreeper)	Drepanidinae	Endangered 1
<u><i>Melanthera kamolensis</i></u>	nehe	Asteraceae	Endangered 1
<u><i>Melanthera tenuifolia</i></u>	Nehe	Asteraceae	Endangered 1
<u><i>Melicope adscendens</i></u>	Alani	Rutaceae	Endangered 1
<u><i>Melicope balloui</i></u>	Alani	Rutaceae	Endangered 1
<u><i>Melicope christophersenii</i></u>	Alani	Rutaceae	Endangered 1
<u><i>Melicope degeneri</i></u>	Alani	Rutaceae	Endangered 1
<u><i>Melicope haupuensis</i></u>	Alani	Rutaceae	Endangered 1
<u><i>Melicope hiiakae</i></u>	Alani	Rutaceae	Endangered 1
<u><i>Melicope knudsenii</i></u>	Alani	Rutaceae	Endangered 1
<u><i>Melicope lydgatei</i></u>	Alani	Rutaceae	Endangered 1
<u><i>Melicope makahae</i></u>	Alani	Rutaceae	Endangered 1
<u><i>Melicope mucronulata</i></u>	Alani	Rutaceae	Endangered 1
<u><i>Melicope munroi</i></u>	Alani	Rutaceae	Endangered 1
<u><i>Melicope ovalis</i></u>	Alani	Rutaceae	Endangered 1
<u><i>Melicope pallida</i></u>	Alani	Rutaceae	Endangered 1

Scientific Name	Common Name	Family	Federal Listing Status
<u>Melicope paniculata</u>	Alani	Rutaceae	Endangered
<u>Melicope puberula</u>	Alani	Rutaceae	Endangered
<u>Melicope quadrangularis</u>	Alani	Rutaceae	Endangered
<u>Melicope reflexa</u>	Alani	Rutaceae	Endangered
<u>Melicope saint-johnii</u>	Alani	Rutaceae	Endangered
<u>Melicope zahlbruckneri</u>	Alani	Rutaceae	Endangered
<u>Menidia extensa</u>	Waccamaw silverside	Atherinidae	Threatened
<u>Mentzelia leucophylla</u>	Ash Meadows blazingstar	Loasaceae	Threatened
<u>Mesodon clarki nantahala</u>	noonday snail	Polygyridae	Threatened
<u>Mezoneuron kavaiense</u>	Uhi uhi	Fabaceae	Endangered
<u>Microhexura montivaga</u>	Spruce-fir moss spider	Dipluridae	Endangered
<u>Microlepidia strigosa var. mauiensis</u>	No common name	Dennstaedtiaceae	Endangered
<u>Microtus californicus scirpensis</u>	Amargosa vole	Cricetidae	Endangered
<u>Microtus pennsylvanicus dukecampbelli</u>	Florida salt marsh vole	Cricetidae	Endangered
<u>Mimulus michiganensis</u>	Michigan monkey-flower	Scrophulariaceae	Endangered
<u>Mirabilis macfarlanei</u>	MacFarlane's four-o'clock	Nyctaginaceae	Threatened
<u>Mitracarpus maxwelliae</u>	No common name	Rubiaceae	Endangered
<u>Mitracarpus polycladus</u>	No common name	Rubiaceae	Endangered
<u>Moapa coriacea</u>	Moapa dace	Cyprinidae	Endangered
<u>Moho braccatus</u>	Kauai `o`o (honeyeater)	Meliphagidae	Endangered
<u>Monachus schauinslandi</u>	Hawaiian monk seal	Phocidae	Endangered
<u>Monardella viminea</u>	Willowy monardella	Lamiaceae	Endangered
<u>Monolopia (=Lembertia) congdonii</u>	San Joaquin wooly-threads	Asteraceae	Endangered
<u>Mucuna sloanei var. persericea</u>	sea bean	Fabaceae	Endangered
<u>Mustela nigripes</u>	Black-footed ferret	Mustelidae	Endangered
<u>Myadestes lanaiensis rutha</u>	Molokai thrush	Muscicapidae	Endangered

Scientific Name	Common Name	Family	Federal Listing Status
<u><i>Myadestes myadestinus</i></u>	Large Kauai (=kamao) Thrush	Muscicapidae	Endangered 1
<u><i>Myadestes palmeri</i></u>	Small Kauai (=puaiohi) Thrush	Muscicapidae	Endangered 1
<u><i>Mycetophyllia ferox</i></u>	Rough Cactus Coral	Mussidae	Threatened 1
<u><i>Mycteria americana</i></u>	Wood stork	Ciconiidae	Threatened 1
<u><i>Myotis grisescens</i></u>	Gray bat	Vespertilionidae	Endangered 1
<u><i>Myotis septentrionalis</i></u>	Northern Long-Eared Bat	Vespertilionidae	Threatened 1
<u><i>Myotis sodalis</i></u>	Indiana bat	Vespertilionidae	Endangered 1
<u><i>Myrcia paganii</i></u>	No common name	Myrtaceae	Endangered 1
<u><i>Myrsine fosbergii</i></u>	Kolea	Myrsinaceae	Endangered 1
<u><i>Myrsine juddii</i></u>	Kolea	Primulaceae	Endangered 1
<u><i>Myrsine knudsenii</i></u>	Kolea	Primulaceae	Endangered 1
<u><i>Myrsine linearifolia</i></u>	Kolea	Primulaceae	Threatened 1
<u><i>Myrsine mezii</i></u>	Kolea	Primulaceae	Endangered 1
<u><i>Myrsine vaccinioides</i></u>	Kolea	Primulaceae	Endangered 1
<u><i>Navarretia fossalis</i></u>	Spreading navarretia	Polemoniaceae	Threatened 1
<u><i>Navarretia leucocephala ssp. pauciflora</i></u> (=N. pauciflora)	Few-flowered navarretia	Polemoniaceae	Endangered 1
<u><i>Navarretia leucocephala ssp. plieantha</i></u>	Many-flowered navarretia	Polemoniaceae	Endangered 1
<u><i>Necturus alabamensis</i></u>	Black warrior (=Sipsey Fork) Waterdog	Proteidae	Endangered 1
<u><i>Necturus lewisi</i></u>	Neuse River waterdog	Proteidae	Proposed Threatened 1
<u><i>Neoleptoneta microps</i></u>	Government Canyon Bat Cave Spider	Leptonetidae	Endangered 1
<u><i>Neoleptoneta myopica</i></u>	Tooth Cave Spider	Leptonetidae	Endangered 1
<u><i>Neonympha mitchellii francisci</i></u>	Saint Francis' satyr butterfly	Nymphalidae	Endangered 1
<u><i>Neonympha mitchellii mitchellii</i></u>	Mitchell's satyr Butterfly	Nymphalidae	Endangered 1
<u><i>Neoseps reynoldsi</i></u>	Sand skink	Scincidae	Threatened 1

Scientific Name	Common Name	Family	Federal Listing Status
<u><i>Neostapfia colusana</i></u>	Colusa grass	Poaceae	Threatened
<u><i>Neotoma floridana smalli</i></u>	Key Largo woodrat	Cricetidae	Endangered
<u><i>Neotoma fuscipes riparia</i></u>	Riparian woodrat (=San Joaquin Valley)	Muridae	Endangered
<u><i>Neraudia angulata</i></u>	No common name	Urticaceae	Endangered
<u><i>Neraudia ovata</i></u>	No common name	Urticaceae	Endangered
<u><i>Neraudia sericea</i></u>	No common name	Urticaceae	Endangered
<u><i>Nerodia clarkii taeniata</i></u>	Atlantic salt marsh snake	Colubridae	Threatened
<u><i>Nerodia erythrogaster neglecta</i></u>	Copperbelly water snake	Colubridae	Threatened
<u><i>Nervilia jacksoniae</i></u>	No common name	Orchidaceae	Threatened
<u><i>Nesogenes rotensis</i></u>	No common name	Verbenaceae	Endangered
<u><i>Newcombia cumingi</i></u>	Newcomb's Tree snail	Achatinellidae	Endangered
<u><i>Nicrophorus americanus</i></u>	American burying beetle	Silphidae	Threatened
<u><i>Nitrophila mohavensis</i></u>	Amargosa niterwort	Chenopodiaceae	Endangered
<u><i>Nolina brittoniana</i></u>	Britton's beargrass	Agavaceae	Endangered
<u><i>Nothocestrum breviflorum</i></u>	`Aiea	Solanaceae	Endangered
<u><i>Nothocestrum latifolium</i></u>	`Aiea	Solanaceae	Endangered
<u><i>Nothocestrum peltatum</i></u>	`Aiea	Solanaceae	Endangered
<u><i>Nototrichium humile</i></u>	Kulu`i	Amaranthaceae	Endangered
<u><i>Notropis albizonatus</i></u>	Palezone shiner	Cyprinidae	Endangered
<u><i>Notropis buccula</i></u>	Smalleye Shiner	Cyprinidae	Endangered
<u><i>Notropis cahabae</i></u>	Cahaba shiner	Cyprinidae	Endangered
<u><i>Notropis girardi</i></u>	Arkansas River shiner	Cyprinidae	Threatened
<u><i>Notropis mekistocholas</i></u>	Cape Fear shiner	Cyprinidae	Endangered
<u><i>Notropis oxyrhynchus</i></u>	Sharpnose Shiner	Cyprinidae	Endangered
<u><i>Notropis simus pecosensis</i></u>	Pecos bluntnose shiner	Cyprinidae	Threatened
<u><i>Notropis topeka (=tristis)</i></u>	Topeka shiner	Cyprinidae	Endangered
<u><i>Noturus baileyi</i></u>	Smoky madtom	Ictaluridae	Endangered

Scientific Name	Common Name	Family	Federal Listing Status
<u><i>Noturus crypticus</i></u>	Chucky Madtom	Ictaluridae	Endangered 1
<u><i>Noturus flavipinnis</i></u>	Yellowfin madtom	Ictaluridae	Threatened 1
<u><i>Noturus furiosus</i></u>	Carolina madtom	Ictaluridae	Proposed Endangered 1
<u><i>Noturus placidus</i></u>	Neosho madtom	Ictaluridae	Threatened 1
<u><i>Noturus stanauli</i></u>	Pygmy madtom	Ictaluridae	Endangered 1
<u><i>Noturus trautmani</i></u>	Scioto madtom	Ictaluridae	Endangered 1
<u><i>Numenius borealis</i></u>	Eskimo curlew	Scolopacidae	Endangered 1
<u><i>Oarisma poweshiek</i></u>	Poweshiek skipperling	Hesperiidae	Endangered 1
<u><i>Obovaria retusa</i></u>	Ring pink (mussel)	Unionidae	Endangered 1
<u><i>Obovaria subrotunda</i></u>	Round hickorynut	Unionidae	Proposed Threatened 1
<u><i>Oceanodroma castro</i></u>	Band-rumped storm-petrel	Hydrobatidae	Endangered 1
<u><i>Ochrosia haleakalae</i></u>	Holei	Apocynaceae	Endangered 1
<u><i>Ochrosia kilaueaensis</i></u>	Holei	Apocynaceae	Endangered 1
<u><i>Odocoileus virginianus clavium</i></u>	Key deer	Cervidae	Endangered 1
<u><i>Odocoileus virginianus leucurus</i></u>	Columbian white-tailed deer	Cervidae	Threatened 1
<u><i>Oenothera deltoides ssp. howellii</i></u>	Antioch Dunes evening-primrose	Onagraceae	Endangered 1
<u><i>Oncorhynchus (=Salmo) kisutch</i></u>	Coho salmon	Salmonidae	Endangered 1
<u><i>Oncorhynchus (=Salmo) kisutch</i></u>	Coho salmon	Salmonidae	Threatened 1
<u><i>Oncorhynchus (=Salmo) mykiss</i></u>	Steelhead	Salmonidae	Endangered 1
<u><i>Oncorhynchus (=Salmo) mykiss</i></u>	Steelhead	Salmonidae	Threatened 1
<u><i>Oncorhynchus (=Salmo) nerka</i></u>	Sockeye salmon	Salmonidae	Endangered 1
<u><i>Oncorhynchus (=Salmo) nerka</i></u>	Sockeye salmon	Salmonidae	Threatened 1
<u><i>Oncorhynchus (=Salmo) tshawytscha</i></u>	Chinook salmon	Salmonidae	Endangered 1

Scientific Name	Common Name	Family	Federal Listing Status
<u><i>Oncorhynchus</i> (=Salmo) tshawytscha</u>	Chinook salmon	Salmonidae	Threatened ①
<u><i>Oncorhynchus aguabonita whitei</i></u>	Little Kern golden trout	Salmonidae	Threatened ①
<u><i>Oncorhynchus apache</i></u>	Apache trout	Salmonidae	Threatened ①
<u><i>Oncorhynchus clarkii henshawi</i></u>	Lahontan cutthroat trout	Salmonidae	Threatened ①
<u><i>Oncorhynchus clarkii seleniris</i></u>	Paiute cutthroat trout	Salmonidae	Threatened ①
<u><i>Oncorhynchus clarkii stomias</i></u>	Greenback Cutthroat trout	Salmonidae	Threatened ①
<u><i>Oncorhynchus gilae</i></u>	Gila trout	Salmonidae	Threatened ①
<u><i>Oncorhynchus keta</i></u>	Chum salmon	Salmonidae	Threatened ①
<u><i>Opuntia treleasei</i></u>	Bakersfield cactus	Cactaceae	Endangered ①
<u><i>Orbicella annularis</i></u>	Lobed Star Coral	Faviidae	Threatened ①
<u><i>Orbicella faveolata</i></u>	Mountainous Star Coral	Faviidae	Threatened ①
<u><i>Orbicella franksi</i></u>	Boulder star coral	Merulinidae	Threatened ①
<u><i>Orcinus orca</i></u>	Killer whale	Delphinidae	Endangered ①
<u><i>Orconectes shoupi</i></u>	Nashville crayfish	Cambaridae	Endangered ①
<u><i>Orcuttia californica</i></u>	California Orcutt grass	Poaceae	Endangered ①
<u><i>Orcuttia inaequalis</i></u>	San Joaquin Orcutt grass	Poaceae	Threatened ①
<u><i>Orcuttia pilosa</i></u>	Hairy Orcutt grass	Poaceae	Endangered ①
<u><i>Orcuttia tenuis</i></u>	Slender Orcutt grass	Poaceae	Threatened ①
<u><i>Orcuttia viscida</i></u>	Sacramento Orcutt grass	Poaceae	Endangered ①
<u><i>Oreomystis bairdi</i></u>	Akikiki	Fringillidae	Endangered ①
<u><i>Oreomystis mana</i></u>	Hawaii creeper	Drepanidinae	Endangered ①
<u><i>Orthalicus reses</i> (not incl. nesodryas)</u>	Stock Island tree snail	Bulimulidae	Threatened ①
<u><i>Oryzomys palustris natator</i></u>	Silver rice rat	Muridae	Endangered ①
<u><i>Osmoxylon mariannense</i></u>	No common name	Araliaceae	Endangered ①
<u><i>Ostodes strigatus</i></u>	Snail [no common name]	Potariidae	Endangered ①
<u><i>Ottoschulzia rhodoxylon</i></u>	Palo de rosa	Icacinaceae	Endangered

Scientific Name	Common Name	Family	Federal Listing Status
<u><i>Ovis canadensis nelsoni</i></u>	Peninsular bighorn sheep	Bovidae	Endangered
<u><i>Ovis canadensis sierrae</i></u>	Sierra Nevada bighorn sheep	Bovidae	Endangered
<u><i>Oxyloma haydeni kanabensis</i></u>	Kanab ambersnail	Succineidae	Endangered
<u><i>Oxypolis canbyi</i></u>	Canby's dropwort	Apiaceae	Endangered
<u><i>Oxytheca parishii</i> var. <i>goodmaniana</i></u>	Cushenbury oxytheca	Polygonaceae	Endangered
<u><i>Oxytropis campestris</i> var. <i>chartacea</i></u>	Fassett's locoweed	Fabaceae	Threatened
<u><i>Pacifastacus fortis</i></u>	Shasta crayfish	Cambaridae	Endangered
<u><i>Packera franciscana</i></u>	San Francisco Peaks ragwort	Asteraceae	Threatened
<u><i>Palaemonetes cummingsi</i></u>	Squirrel Chimney Cave shrimp	Palaemonidae	Threatened
<u><i>Palaemonias alabamiae</i></u>	Alabama cave shrimp	Atyidae	Endangered
<u><i>Palaemonias ganteri</i></u>	Kentucky cave shrimp	Atyidae	Endangered
<u><i>Palmeria dolei</i></u>	crested honeycreeper (Akohekohe)	Drepanidinae	Endangered
<u><i>Panicum fauriei</i> var. <i>carteri</i></u>	Carter's panicgrass	Poaceae	Endangered
<u><i>Panicum niihauense</i></u>	Lau `ehu	Poaceae	Endangered
<u><i>Panthera onca</i></u>	Jaguar	Felidae	Endangered
<u><i>Paronychia chartacea</i></u>	Papery whitlow-wort	Caryophyllaceae	Threatened
<u><i>Paroreomyza flammea</i></u>	Molokai creeper	Drepanidinae	Endangered
<u><i>Paroreomyza maculata</i></u>	Oahu creeper	Drepanidinae	Endangered
<u><i>Partula gibba</i></u>	Humped tree snail	Partulidae	Endangered
<u><i>Partula langfordi</i></u>	Langford's tree snail	Partulidae	Endangered
<u><i>Partula radiolata</i></u>	Guam tree snail	Partulidae	Endangered
<u><i>Partulina semicarinata</i></u>	Lanai tree snail	Achatinellidae	Endangered
<u><i>Partulina variabilis</i></u>	Lanai tree snail	Achatinellidae	Endangered
<u><i>Parvisedum leiocarpum</i></u>	Lake County stonecrop	Crassulaceae	Endangered
<u><i>Pectis imberbis</i></u>	Beardless chinch weed	Asteraceae	Proposed Endangered
<u><i>Pedicularis furbishiae</i></u>	Furbish lousewort	Scrophulariaceae	Endangered

Scientific Name	Common Name	Family	Federal Listing Status
<u><i>Pediocactus</i></u> (= <i>Echinocactus</i> ,= <i>Utahia</i>) <u><i>sileri</i></u>	Siler pincushion cactus	Cactaceae	Threatened
<u><i>Pediocactus bradyi</i></u>	Brady pincushion cactus	Cactaceae	Endangered
<u><i>Pediocactus despainii</i></u>	San Rafael cactus	Cactaceae	Endangered
<u><i>Pediocactus knowltonii</i></u>	Knowlton's cactus	Cactaceae	Endangered
<u><i>Pediocactus peeblesianus</i></u> <u><i>fickeiseniae</i></u>	Fickeisen plains cactus	Cactaceae	Endangered
<u><i>Pediocactus peeblesianus</i></u> <u>var. <i>peeblesianus</i></u>	Peebles Navajo cactus	Cactaceae	Endangered
<u><i>Pediocactus winkleri</i></u>	Winkler cactus	Cactaceae	Threatened
<u><i>Pegias fabula</i></u>	Littlewing pearlymussel	Unionidae	Endangered
<u><i>Pekania pennanti</i></u>	Fisher	Mustelidae	Endangered
<u><i>Peltophryne lemur</i></u>	Puerto Rican crested toad	Bufonidae	Threatened
<u><i>Penstemon debilis</i></u>	Parachute beardtongue	Plantaginaceae	Threatened
<u><i>Penstemon haydenii</i></u>	Blowout penstemon	Scrophulariaceae	Endangered
<u><i>Penstemon penlandii</i></u>	Penland beardtongue	Scrophulariaceae	Endangered
<u><i>Pentachaeta bellidiflora</i></u>	White-rayed pentachaeta	Asteraceae	Endangered
<u><i>Pentachaeta lyonii</i></u>	Lyon's pentachaeta	Asteraceae	Endangered
<u><i>Peperomia subpetiolata</i></u>	`Ala `ala wai nui	Piperaceae	Endangered
<u><i>Peperomia wheeleri</i></u>	Wheeler's peperomia	Piperaceae	Endangered
<u><i>Percina antesella</i></u>	Amber darter	Percidae	Endangered
<u><i>Percina aurolineata</i></u>	Goldline darter	Percidae	Threatened
<u><i>Percina aurora</i></u>	Pearl darter	Percidae	Threatened
<u><i>Percina jenkinsi</i></u>	Conasauga logperch	Percidae	Endangered
<u><i>Percina pantherina</i></u>	Leopard darter	Percidae	Threatened
<u><i>Percina rex</i></u>	Roanoke logperch	Percidae	Endangered
<u><i>Percina tanasi</i></u>	Snail darter	Percidae	Threatened
<u><i>Perognathus longimembris</i></u> <u><i>pacificus</i></u>	Pacific pocket mouse	Heteromyidae	Endangered
<u><i>Peromyscus gossypinus</i></u> <u><i>allapaticola</i></u>	Key Largo cotton mouse	Muridae	Endangered

Scientific Name	Common Name	Family	Federal Listing Status
<u><i>Peromyscus polionotus allopshys</i></u>	Choctawhatchee beach mouse	Muridae	Endangered
<u><i>Peromyscus polionotus ammobates</i></u>	Alabama beach mouse	Muridae	Endangered
<u><i>Peromyscus polionotus niveiventris</i></u>	Southeastern beach mouse	Muridae	Threatened
<u><i>Peromyscus polionotus peninsularis</i></u>	St. Andrew beach mouse	Muridae	Endangered
<u><i>Peromyscus polionotus phasma</i></u>	Anastasia Island beach mouse	Muridae	Endangered
<u><i>Peromyscus polionotus trissyllepsis</i></u>	Perdido Key beach mouse	Muridae	Endangered
<u><i>Peucedanum sandwicense</i></u>	Makou	Apiaceae	Threatened
<u><i>Phacelia argillacea</i></u>	Clay phacelia	Hydrophyllaceae	Endangered
<u><i>Phacelia formosula</i></u>	North Park phacelia	Hydrophyllaceae	Endangered
<u><i>Phacelia insularis ssp. insularis</i></u>	Island phacelia	Hydrophyllaceae	Endangered
<u><i>Phacelia submutica</i></u>	DeBeque phacelia	Hydrophyllaceae	Threatened
<u><i>Phaeognathus hubrichti</i></u>	Red Hills salamander	Plethodontidae	Threatened
<u><i>Phlox hirsuta</i></u>	Yreka phlox	Polemoniaceae	Endangered
<u><i>Phlox nivalis ssp. texensis</i></u>	Texas trailing phlox	Polemoniaceae	Endangered
<u><i>Phoca (=Pusa) hispida botnica</i></u>	Ringed Seal	Phocidae	Threatened
<u><i>Phoca (=Pusa) hispida hispida</i></u>	Ringed Seal	Phocidae	Threatened
<u><i>Phoca (=Pusa) hispida ladogensis</i></u>	Ringed seal	Phocidae	Endangered
<u><i>Phoca (=Pusa) hispida ochotensis</i></u>	Ringed Seal	Phocidae	Threatened
<u><i>Phoca largha</i></u>	Spotted Seal	Phocidae	Threatened
<u><i>Phoebastria (=Diomedea) albatrus</i></u>	Short-tailed albatross	Diomedidae	Endangered
<u><i>Phoxinus cumberlandensis</i></u>	Blackside dace	Cyprinidae	Threatened
<u><i>Phyllanthus saffordii</i></u>	No common name	Phyllanthaceae	Endangered
<u><i>Phyllostegia bracteata</i></u>	No common name	Lamiaceae	Endangered

Scientific Name	Common Name	Family	Federal Listing Status
<i>Phyllostegia brevidens</i>	No common name	Lamiaceae	Endangered 1
<i>Phyllostegia floribunda</i>	No common name	Lamiaceae	Endangered 1
<i>Phyllostegia glabra</i> var. <i>lanaiensis</i>	No common name	Lamiaceae	Endangered 1
<i>Phyllostegia haliakalae</i>	No common name	Lamiaceae	Endangered 1
<i>Phyllostegia helleri</i>	No common name	Lamiaceae	Endangered 1
<i>Phyllostegia hirsuta</i>	No common name	Lamiaceae	Endangered 1
<i>Phyllostegia hispida</i>	No common name	Lamiaceae	Endangered 1
<i>Phyllostegia kaalaensis</i>	No common name	Lamiaceae	Endangered 1
<i>Phyllostegia knudsenii</i>	No common name	Lamiaceae	Endangered 1
<i>Phyllostegia mannii</i>	No common name	Lamiaceae	Endangered 1
<i>Phyllostegia mollis</i>	No common name	Lamiaceae	Endangered 1
<i>Phyllostegia parviflora</i>	No common name	Lamiaceae	Endangered 1
<i>Phyllostegia pilosa</i>	No common name	Lamiaceae	Endangered 1
<i>Phyllostegia racemosa</i>	Kiponapona	Lamiaceae	Endangered 1
<i>Phyllostegia renovans</i>	No common name	Lamiaceae	Endangered 1
<i>Phyllostegia stachyoides</i>	No common name	Lamiaceae	Endangered 1
<i>Phyllostegia velutina</i>	No common name	Lamiaceae	Endangered 1
<i>Phyllostegia waimeae</i>	No common name	Lamiaceae	Endangered 1
<i>Phyllostegia warshaueri</i>	No common name	Lamiaceae	Endangered 1
<i>Phyllostegia wawrana</i>	No common name	Lamiaceae	Endangered 1
<i>Physa natricina</i>	Snake River physa snail	Physidae	Endangered 1
<i>Physaria douglasii</i> ssp. <i>tuplashensis</i>	White Bluffs bladderpod	Brassicaceae	Threatened 1
<i>Physaria filiformis</i>	Missouri bladderpod	Brassicaceae	Threatened 1
<i>Physaria globosa</i>	Short's bladderpod	Brassicaceae	Endangered 1
<i>Physaria obcordata</i>	Dudley Bluffs twinpod	Brassicaceae	Threatened 1
<i>Physeter catodon</i> (= <i>macrocephalus</i>)	Sperm whale	Physeteridae	Endangered 1
<i>Picoides borealis</i>	Red-cockaded woodpecker	Picidae	Endangered 1

Scientific Name	Common Name	Family	Federal Listing Status
<u><i>Pilosocereus robinii</i></u>	Key tree cactus	Cactaceae	Endangered
<u><i>Pinguicula ionantha</i></u>	Godfrey's butterwort	Lentibulariaceae	Threatened
<u><i>Piperia yadonii</i></u>	Yadon's piperia	Orchidaceae	Endangered
<u><i>Pipilo crissalis eremophilus</i></u>	Inyo California towhee	Emberizidae	Threatened
<u><i>Pittosporum halophilum</i></u>	Hoawa	Pittosporaceae	Endangered
<u><i>Pittosporum hawaiiense</i></u>	Hoawa	Pittosporaceae	Endangered
<u><i>Pittosporum napaliense</i></u>	Ho`awa	Pittosporaceae	Endangered
<u><i>Pituophis melanoleucus lodingi</i></u>	Black pine snake	Colubridae	Threatened
<u><i>Pituophis ruthveni</i></u>	Louisiana pinesnake	Colubridae	Threatened
<u><i>Pityopsis ruthii</i></u>	Ruth's golden aster	Asteraceae	Endangered
<u><i>Plagiobothrys hirtus</i></u>	rough popcornflower	Boraginaceae	Endangered
<u><i>Plagiobothrys strictus</i></u>	Calistoga allocarya	Boraginaceae	Endangered
<u><i>Plagopterus argentissimus</i></u>	Woundfin	Cyprinidae	Endangered
<u><i>Plantago hawaiiensis</i></u>	Kuahiwi laukahi	Plantaginaceae	Endangered
<u><i>Plantago princeps</i></u>	Kuahiwi laukahi	Plantaginaceae	Endangered
<u><i>Platanthera holochila</i></u>	No common name	Orchidaceae	Endangered
<u><i>Platanthera integrilabia</i></u>	White fringeless orchid	Orchidaceae	Threatened
<u><i>Platanthera leucophaea</i></u>	Eastern prairie fringed orchid	Orchidaceae	Threatened
<u><i>Platanthera praeclara</i></u>	Western prairie fringed Orchid	Orchidaceae	Threatened
<u><i>Platydesma cornuta</i> var. <i>cornuta</i></u>	No common name	Rutaceae	Endangered
<u><i>Platydesma cornuta</i> var. <i>decurrens</i></u>	No common name	Rutaceae	Endangered
<u><i>Platydesma remyi</i></u>	No common name	Rutaceae	Endangered
<u><i>Platydesma rostrata</i></u>	Pilo kea lau li`i	Rutaceae	Endangered
<u><i>Pleodendron macranthum</i></u>	Chupacallos	Canellaceae	Endangered
<u><i>Pleomele fernaldii</i></u>	Hala pepe	Agavaceae	Endangered
<u><i>Pleomele forbesii</i></u>	Hala pepe	Agavaceae	Endangered

Scientific Name	Common Name	Family	Federal Listing Status
<u><i>Pleomele hawaiiensis</i></u>	Hala pepe	Asparagaceae	Endangered
<u><i>Plethobasus cicatricosus</i></u>	White wartyback (pearlymussel)	Unionidae	Endangered
<u><i>Plethobasus cooperianus</i></u>	Orangefoot pimpleback (pearlymussel)	Unionidae	Endangered
<u><i>Plethobasus cyphyus</i></u>	Sheepnose Mussel	Unionidae	Endangered
<u><i>Plethodon neomexicanus</i></u>	Jemez Mountains salamander	Plethodontidae	Endangered
<u><i>Plethodon nettingi</i></u>	Cheat Mountain salamander	Plethodontidae	Threatened
<u><i>Plethodon shenandoah</i></u>	Shenandoah salamander	Plethodontidae	Endangered
<u><i>Pleurobema athearni</i></u>	Canoe Creek Clubshell	Unionidae	Proposed Endangered
<u><i>Pleurobema clava</i></u>	Clubshell	Unionidae	Endangered
<u><i>Pleurobema collina</i></u>	James spinymussel	Unionidae	Endangered
<u><i>Pleurobema curtum</i></u>	Black clubshell	Unionidae	Endangered
<u><i>Pleurobema decisum</i></u>	Southern clubshell	Unionidae	Endangered
<u><i>Pleurobema furvum</i></u>	Dark pigtoe	Unionidae	Endangered
<u><i>Pleurobema georgianum</i></u>	Southern pigtoe	Unionidae	Endangered
<u><i>Pleurobema gibberum</i></u>	Cumberland pigtoe	Unionidae	Endangered
<u><i>Pleurobema hanleyianum</i></u>	Georgia pigtoe	Unionidae	Endangered
<u><i>Pleurobema marshalli</i></u>	Flat pigtoe	Unionidae	Endangered
<u><i>Pleurobema perovatum</i></u>	Ovate clubshell	Unionidae	Endangered
<u><i>Pleurobema plenum</i></u>	Rough pigtoe	Unionidae	Endangered
<u><i>Pleurobema pyriforme</i></u>	Oval pigtoe	Unionidae	Endangered
<u><i>Pleurobema strodeanum</i></u>	Fuzzy pigtoe	Unionidae	Threatened
<u><i>Pleurobema taitianum</i></u>	Heavy pigtoe	Unionidae	Endangered
<u><i>Pleurocera foremani</i></u>	Rough hornsnail	Pleuroceridae	Endangered
<u><i>Pleuonaia dolabelloides</i></u>	Slabside Pearlymussel	Unionidae	Endangered
<u><i>Poa atropurpurea</i></u>	San Bernardino bluegrass	Poaceae	Endangered
<u><i>Poa mannii</i></u>	Mann's bluegrass	Poaceae	Endangered
<u><i>Poa napensis</i></u>	Napa bluegrass	Poaceae	Endangered

Scientific Name	Common Name	Family	Federal Listing Status
<u><i>Poa sandvicensis</i></u>	Hawaiian bluegrass	Poaceae	Endangered
<u><i>Poa siphonoglossa</i></u>	No common name	Poaceae	Endangered
<u><i>Poeciliopsis occidentalis</i></u>	Gila topminnow (incl. Yaqui)	Poeciliidae	Endangered
<u><i>Pogogyne abramsii</i></u>	San Diego mesa-mint	Lamiaceae	Endangered
<u><i>Pogogyne nudiuscula</i></u>	Otay mesa-mint	Lamiaceae	Endangered
<u><i>Polioptila californica californica</i></u>	Coastal California gnatcatcher	Muscicapidae	Threatened
<u><i>Polyborus plancus audubonii</i></u>	Audubon's crested caracara	Falconidae	Threatened
<u><i>Polygala lewtonii</i></u>	Lewton's polygala	Polygalaceae	Endangered
<u><i>Polygala smallii</i></u>	Tiny polygala	Polygalaceae	Endangered
<u><i>Polygonella basiramia</i></u>	Wireweed	Polygonaceae	Endangered
<u><i>Polygonella myriophylla</i></u>	Sandlace	Polygonaceae	Endangered
<u><i>Polygonum hickmanii</i></u>	Scotts Valley Polygonum	Polygonaceae	Endangered
<u><i>Polygyriscus virginianus</i></u>	Virginia fringed mountain snail	Helicodiscidae	Endangered
<u><i>Polyphylla barbata</i></u>	Mount Hermon June beetle	Scarabaeidae	Endangered
<u><i>Polyscias bisattenuata</i></u>	No common name	Araliaceae	Endangered
<u><i>Polyscias flynnii</i></u>	No common name	Araliaceae	Endangered
<u><i>Polyscias gymnocarpa</i></u>	`Ohe`ohe	Araliaceae	Endangered
<u><i>Polyscias lydgatei</i></u>	No common name	Araliaceae	Endangered
<u><i>Polyscias racemosa</i></u>	No common name	Araliaceae	Endangered
<u><i>Polystichum aleuticum</i></u>	Aleutian shield fern	Dryopteridaceae	Endangered
<u><i>Polystichum calderonense</i></u>	No common name	Dryopteridaceae	Endangered
<u><i>Polysticta stelleri</i></u>	Steller's Eider	Anatidae	Threatened
<u><i>Popenaias popeii</i></u>	Texas Hornshell	Unionidae	Endangered
<u><i>Portulaca sclerocarpa</i></u>	Po`e	Portulacaceae	Endangered
<u><i>Portulaca villosa</i></u>	Ihi	Portulacaceae	Endangered
<u><i>Potamilus capax</i></u>	Fat pocketbook	Unionidae	Endangered
<u><i>Potamilus inflatus</i></u>	Inflated heelsplitter	Unionidae	Threatened

Scientific Name	Common Name	Family	Federal Listing Status
<u>Potamogeton clystocarpus</u>	Little Aguja (=Creek) Pondweed	Potamogetonaceae	Endangered 1
<u>Potentilla hickmanii</u>	Hickman's potentilla	Rosaceae	Endangered 1
<u>Primula maguirei</u>	Maguire primrose	Primulaceae	Threatened 1
<u>Pristis clavata</u>	Dwarf sawfish	Pristidae	Endangered 1
<u>Pristis pectinata</u>	Smalltooth sawfish	Pristidae	Endangered 1
<u>Pristis pristis</u>	Large-tooth Sawfish	Pristidae	Endangered 1
<u>Pritchardia aylmer-robinsonii</u>	Wahane	Arecaceae	Endangered 1
<u>Pritchardia bakeri</u>	Baker's Loulu	Arecaceae	Endangered 1
<u>Pritchardia hardyi</u>	Lo`ulu	Asteraceae	Endangered 1
<u>Pritchardia kaalae</u>	Lo`ulu	Arecaceae	Endangered 1
<u>Pritchardia lanigera</u>	Lo`ulu	Arecaceae	Endangered 1
<u>Pritchardia maideniana</u>	Lo`ulu	Arecaceae	Endangered 1
<u>Pritchardia munroi</u>	Lo`ulu	Arecaceae	Endangered 1
<u>Pritchardia napaliensis</u>	Lo`ulu	Arecaceae	Endangered 1
<u>Pritchardia remota</u>	Lo`ulu	Arecaceae	Endangered 1
<u>Pritchardia schattaueri</u>	Lo`ulu	Arecaceae	Endangered 1
<u>Pritchardia viscosa</u>	Lo`ulu	Arecaceae	Endangered 1
<u>Procambarus econfinae</u>	Panama City crayfish	Cambaridae	Proposed Threatened 1
<u>Procaris hawaiana</u>	Anchialine pool Shrimp	Procarididae	Endangered 1
<u>Prunus geniculata</u>	Scrub plum	Rosaceae	Endangered 1
<u>Pseudemys alabamensis</u>	Alabama red-bellied turtle	Emydidae	Endangered 1
<u>Pseudemys rubriventris bangsi</u>	Plymouth Redbelly Turtle	Emydidae	Endangered 1
<u>Pseudobahia bahiifolia</u>	Hartweg's golden sunburst	Asteraceae	Endangered 1
<u>Pseudobahia peirsonii</u>	San Joaquin adobe sunburst	Asteraceae	Threatened 1
<u>Pseudocopaeodes eunus obscurus</u>	Carson wandering skipper	Hesperiidae	Endangered 1
<u>Pseudognaphalium sandwicensium var. molokaiense</u>	Ena`ena	Asteraceae	Endangered 1

Scientific Name	Common Name	Family	Federal Listing Status
<u><i>Pseudonestor xanthophrys</i></u>	Maui parrotbill (Kiwikiu)	Drepanidinae	Endangered
<u><i>Pseudorca crassidens</i></u>	false killer whale	Delphinidae	Endangered
<u><i>Pseudotryonia adamantina</i></u>	Diamond Tryonia	Hydrobiidae	Endangered
<u><i>Psittirostra psittacea</i></u>	`O`u (honeycreeper)	Drepanidinae	Endangered
<u><i>Psychotria grandiflora</i></u>	Kopiko	Rubiaceae	Endangered
<u><i>Psychotria hexandra ssp. oahuensis</i></u>	Kopiko	Rubiaceae	Endangered
<u><i>Psychotria hobdyi</i></u>	Kopiko	Rubiaceae	Endangered
<u><i>Psychotria malaspinae</i></u>	Aplokating-palaoan	Rubiaceae	Endangered
<u><i>Pteralyxia kauaiensis</i></u>	Kaulu	Apocynaceae	Endangered
<u><i>Pteralyxia macrocarpa</i></u>	Kaulu	Apocynaceae	Endangered
<u><i>Pteris lidgatei</i></u>	No common name	Adiantaceae	Endangered
<u><i>Pterodroma cahow</i></u>	Bermuda petrel	Procellariidae	Endangered
<u><i>Pterodroma hasitata</i></u>	Black-capped petrel	Procellariidae	Proposed Threatened
<u><i>Pterodroma sandwichensis</i></u>	Hawaiian petrel	Procellariidae	Endangered
<u><i>Pteropus mariannus mariannus</i></u>	Mariana fruit Bat (=Mariana flying fox)	Pteropodidae	Threatened
<u><i>Pteropus tokudae</i></u>	Little Mariana fruit Bat	Pteropidae	Endangered
<u><i>Ptilimnium nodosum</i></u>	Harperella	Apiaceae	Endangered
<u><i>Ptychobranhus greenii</i></u>	Triangular Kidneyshell	Unionidae	Endangered
<u><i>Ptychobranhus jonesi</i></u>	Southern kidneyshell	Unionidae	Endangered
<u><i>Ptychobranhus subtentus</i></u>	Fluted kidneyshell	Unionidae	Endangered
<u><i>Ptychocheilus lucius</i></u>	Colorado pikeminnow (=squawfish)	Cyprinidae	Endangered
<u><i>Puffinus auricularis newelli</i></u>	Newell's Townsend's shearwater	Procellariidae	Threatened
<u><i>Puma (=Felis) concolor coryi</i></u>	Florida panther	Felidae	Endangered
<u><i>Purshia (=Cowania) subintegra</i></u>	Arizona Cliffrose	Rosaceae	Endangered
<u><i>Pyrgulopsis (=Marstonia) pachyta</i></u>	Armored snail	Hydrobiidae	Endangered

Scientific Name	Common Name	Family	Federal Listing Status
<u><i>Pyrgulopsis bernardina</i></u>	San Bernardino springsnail	Hydrobiidae	Threatened ①
<u><i>Pyrgulopsis bruneauensis</i></u>	Bruneau Hot springsnail	Hydrobiidae	Endangered ①
<u><i>Pyrgulopsis chupaderae</i></u>	Chupadera springsnail	Hydrobiidae	Endangered ①
<u><i>Pyrgulopsis neomexicana</i></u>	Socorro springsnail	Hydrobiidae	Endangered ①
<u><i>Pyrgulopsis ogmorhappe</i></u>	Royal marstonia (snail)	Hydrobiidae	Endangered ①
<u><i>Pyrgulopsis roswellensis</i></u>	Roswell springsnail	Hydrobiidae	Endangered ①
<u><i>Pyrgulopsis texana</i></u>	Phantom Springsnail	Hydrobiidae	Endangered ①
<u><i>Pyrgulopsis trivialis</i></u>	Three Forks Springsnail	Hydrobiidae	Endangered ①
<u><i>Pyrgus ruralis lagunae</i></u>	Laguna Mountains skipper	Hesperiidae	Endangered ①
<u><i>Quadrula cylindrica cylindrica</i></u>	Rabbitsfoot	Unionidae	Threatened ①
<u><i>Quadrula cylindrica strigillata</i></u>	Rough rabbitsfoot	Unionidae	Endangered ① ①
<u><i>Quadrula fragosa</i></u>	Winged Mapleleaf	Unionidae	Endangered ①
<u><i>Quadrula intermedia</i></u>	Cumberland monkeyface (pearlymussel)	Unionidae	Endangered ①
<u><i>Quadrula sparsa</i></u>	Appalachian monkeyface (pearlymussel)	Unionidae	Endangered ①
<u><i>Quadrula stapes</i></u>	Stirrupshell	Unionidae	Endangered ①
<u><i>Quercus hinckleyi</i></u>	Hinckley oak	Fagaceae	Threatened ①
<u><i>Rallus longirostris levipes</i></u>	Light-footed clapper rail	Rallidae	Endangered ①
<u><i>Rallus longirostris obsoletus</i></u>	California clapper rail	Rallidae	Endangered ①
<u><i>Rallus obsoletus</i></u> <u>[=<i>longirostris</i>] <i>yumanensis</i></u>	Yuma Ridgways (clapper) rail	Rallidae	Endangered ①
<u><i>Rallus owstoni</i></u>	Guam rail	Rallidae	Endangered ①
<u><i>Rana chiricahuensis</i></u>	Chiricahua leopard frog	Ranidae	Threatened ①
<u><i>Rana draytonii</i></u>	California red-legged frog	Ranidae	Threatened ①
<u><i>Rana muscosa</i></u>	Mountain yellow-legged frog	Ranidae	Endangered ①
<u><i>Rana pretiosa</i></u>	Oregon spotted frog	Ranidae	Threatened ①
<u><i>Rana sevosa</i></u>	dusky gopher frog	Ranidae	Endangered ①
<u><i>Rana sierrae</i></u>	Sierra Nevada Yellow-legged Frog	Ranidae	Endangered

Scientific Name	Common Name	Family	Federal Listing Status
<u><i>Rangifer tarandus caribou</i></u>	Woodland Caribou	Cervidae	Endangered
<u><i>Ranunculus aestivalis</i></u> (= <i>acriformis</i>)	Autumn Buttercup	Ranunculaceae	Endangered
<u><i>Ranunculus hawaiiensis</i></u>	Makou	Ranunculaceae	Endangered
<u><i>Ranunculus mauiensis</i></u>	Makou	Ranunculaceae	Endangered
<u><i>Reithrodontomys raviventris</i></u>	Salt marsh harvest mouse	Cricetidae	Endangered
<u><i>Remya kauaiensis</i></u>	No common name	Asteraceae	Endangered
<u><i>Remya mauiensis</i></u>	Maui remya	Asteraceae	Endangered
<u><i>Remya montgomeryi</i></u>	No common name	Asteraceae	Endangered
<u><i>Rhadine exilis</i></u>	[no common name] Beetle	Carabidae	Endangered
<u><i>Rhadine infernalis</i></u>	[no common name] Beetle	Carabidae	Endangered
<u><i>Rhadine persephone</i></u>	Tooth Cave ground beetle	Carabidae	Endangered
<u><i>Rhaphiomidas terminatus abdominalis</i></u>	Delhi Sands flower-loving fly	Mydidae	Endangered
<u><i>Rhinichthys osculus lethoporus</i></u>	Independence Valley speckled dace	Cyprinidae	Endangered
<u><i>Rhinichthys osculus nevadensis</i></u>	Ash Meadows speckled dace	Cyprinidae	Endangered
<u><i>Rhinichthys osculus oligoporus</i></u>	Clover Valley speckled dace	Cyprinidae	Endangered
<u><i>Rhinichthys osculus thermalis</i></u>	Kendall Warm Springs dace	Cyprinidae	Endangered
<u><i>Rhodiola integrifolia ssp. leedyi</i></u>	Leedy's roseroot	Crassulaceae	Threatened
<u><i>Rhododendron chapmanii</i></u>	Chapman rhododendron	Ericaceae	Endangered
<u><i>Rhus michauxii</i></u>	Michaux's sumac	Anacardiaceae	Endangered
<u><i>Rhynchopsitta pachyrhyncha</i></u>	Thick-billed parrot	Psittacidae	Endangered
<u><i>Rhynchospora knieskernii</i></u>	Knieskern's Beaked-rush	Cyperaceae	Threatened
<u><i>Ribes echinellum</i></u>	Miccosukee gooseberry	Saxifragaceae	Threatened
<u><i>Rorippa gambellii</i></u>	Gambel's watercress	Brassicaceae	Endangered
<u><i>Rostrhamus sociabilis plumbeus</i></u>	Everglade snail kite	Accipitridae	Endangered

Scientific Name	Common Name	Family	Federal Listing Status
<u><i>Sagittaria fasciculata</i></u>	Bunched arrowhead	Alismataceae	Endangered
<u><i>Sagittaria secundifolia</i></u>	Kral's water-plantain	Alismataceae	Threatened
<u><i>Salmo salar</i></u>	Atlantic salmon	Salmonidae	Endangered
<u><i>Salvelinus confluentus</i></u>	Bull Trout	Salmonidae	Threatened
<u><i>Samoana fragilis</i></u>	Fragile tree snail	Partulidae	Endangered
<u><i>Sanicula mariversa</i></u>	No common name	Apiaceae	Endangered
<u><i>Sanicula purpurea</i></u>	No common name	Apiaceae	Endangered
<u><i>Sanicula sandwicensis</i></u>	No common name	Apiaceae	Endangered
<u><i>Santalum haleakalae</i> var. <i>lanaiense</i></u>	Lanai sandalwood (= `iliahi)	Santalaceae	Endangered
<u><i>Santalum involutum</i></u>	No common name	Santalaceae	Endangered
<u><i>Sarracenia oreophila</i></u>	Green pitcher-plant	Sarraceniaceae	Endangered
<u><i>Sarracenia rubra</i> ssp. <i>alabamensis</i></u>	Alabama canebrake pitcher-plant	Sarraceniaceae	Endangered
<u><i>Sarracenia rubra</i> ssp. <i>jonesii</i></u>	Mountain sweet pitcher-plant	Sarraceniaceae	Endangered
<u><i>Scaevola coriacea</i></u>	Dwarf naupaka	Goodeniaceae	Endangered
<u><i>Scaphirhynchus albus</i></u>	Pallid sturgeon	Acipenseridae	Endangered
<u><i>Scaphirhynchus suttkusi</i></u>	Alabama sturgeon	Acipenseridae	Endangered
<u><i>Schenkia sebaeoides</i></u>	Awiwi	Gentianaceae	Endangered
<u><i>Schiedea adamantis</i></u>	Diamond Head schiedea	Caryophyllaceae	Endangered
<u><i>Schiedea apokremnos</i></u>	Ma`oli`oli	Caryophyllaceae	Endangered
<u><i>Schiedea attenuata</i></u>	No common name	Caryophyllaceae	Endangered
<u><i>Schiedea diffusa</i> ssp. <i>macraei</i></u>	No common name	Caryophyllaceae	Endangered
<u><i>Schiedea diffusa</i> subsp. <i>diffusa</i></u>	No common name	Caryophyllaceae	Endangered
<u><i>Schiedea haleakalensis</i></u>	No common name	Caryophyllaceae	Endangered
<u><i>Schiedea hawaiiensis</i></u>	Ma`oli`oli	Caryophyllaceae	Endangered
<u><i>Schiedea helleri</i></u>	No common name	Caryophyllaceae	Endangered
<u><i>Schiedea hookeri</i></u>	No common name	Caryophyllaceae	Endangered

Scientific Name	Common Name	Family	Federal Listing Status
<u><i>Schiedea jacobii</i></u>	No common name	Caryophyllaceae	Endangered
<u><i>Schiedea kaalae</i></u>	No common name	Caryophyllaceae	Endangered
<u><i>Schiedea kauaiensis</i></u>	No common name	Caryophyllaceae	Endangered
<u><i>Schiedea kealiae</i></u>	Ma`oli`oli	Caryophyllaceae	Endangered
<u><i>Schiedea laui</i></u>	No common name	Caryophyllaceae	Endangered
<u><i>Schiedea lychnoides</i></u>	Kuawawaenohu	Caryophyllaceae	Endangered
<u><i>Schiedea lydgatei</i></u>	No common name	Caryophyllaceae	Endangered
<u><i>Schiedea membranacea</i></u>	No common name	Caryophyllaceae	Endangered
<u><i>Schiedea nuttallii</i></u>	No common name	Caryophyllaceae	Endangered
<u><i>Schiedea obovata</i></u>	No common name	Caryophyllaceae	Endangered
<u><i>Schiedea pubescens</i></u>	Ma`oli`oli	Caryophyllaceae	Endangered
<u><i>Schiedea salicaria</i></u>	No common name	Caryophyllaceae	Endangered
<u><i>Schiedea sarmentosa</i></u>	No common name	Caryophyllaceae	Endangered
<u><i>Schiedea spergulina</i> var. <i>leiopoda</i></u>	No common name	Caryophyllaceae	Endangered
<u><i>Schiedea spergulina</i> var. <i>spergulina</i></u>	No common name	Caryophyllaceae	Threatened
<u><i>Schiedea stellarioides</i></u>	Laulihilihi	Caryophyllaceae	Endangered
<u><i>Schiedea trinervis</i></u>	No common name	Caryophyllaceae	Endangered
<u><i>Schiedea verticillata</i></u>	No common name	Caryophyllaceae	Endangered
<u><i>Schiedea viscosa</i></u>	No common name	Caryophyllaceae	Endangered
<u><i>Schoenocrambe argillacea</i></u>	Clay reed-mustard	Brassicaceae	Threatened
<u><i>Schoenocrambe barnebyi</i></u>	Barneby reed-mustard	Brassicaceae	Endangered
<u><i>Schoenocrambe suffrutescens</i></u>	Shrubby reed-mustard	Brassicaceae	Endangered
<u><i>Schoepfia arenaria</i></u>	No common name	Olacaceae	Threatened
<u><i>Schwalbea americana</i></u>	American chaffseed	Scrophulariaceae	Endangered
<u><i>Scirpus ancistrochaetus</i></u>	Northeastern bulrush	Cyperaceae	Endangered
<u><i>Sclerocactus brevihamatus</i> ssp. <i>tobuschii</i></u>	Tobusch fishhook cactus	Cactaceae	Threatened
<u><i>Sclerocactus brevispinus</i></u>	Pariette cactus	Cacatacea	Threatened

Scientific Name	Common Name	Family	Federal Listing Status
<u><i>Sclerocactus glaucus</i></u>	Colorado hookless Cactus	Cactaceae	Threatened
<u><i>Sclerocactus mesae-verdae</i></u>	Mesa Verde cactus	Cactaceae	Threatened
<u><i>Sclerocactus wetlandicus</i></u>	Uinta Basin hookless cactus	Cactaceae	Threatened
<u><i>Sclerocactus wrightiae</i></u>	Wright fishhook cactus	Cactaceae	Endangered
<u><i>Scutellaria floridana</i></u>	Florida skullcap	Lamiaceae	Threatened
<u><i>Scutellaria montana</i></u>	Large-flowered skullcap	Lamiaceae	Threatened
<u><i>Sebastes paucispinis</i></u>	Bocaccio	Sebastidae	Endangered
<u><i>Sebastes pinniger</i></u>	canary rockfish	Sebastidae	Threatened
<u><i>Sebastes ruberrimus</i></u>	yelloweye rockfish	Sebastidae	Threatened
<u><i>Senecio layneae</i></u>	Layne's butterweed	Asteraceae	Threatened
<u><i>Serianthes nelsonii</i></u>	Hayun Iagu (=Guam), Tronkon guafi (Rota))	Fabaceae	Endangered
<u><i>Sesbania tomentosa</i></u>	Ohai	Fabaceae	Endangered
<u><i>Setophaga angelae</i></u>	Elfin-woods warbler	Parulidae	Threatened
<u><i>Sibara filifolia</i></u>	Santa Cruz Island rockcress	Brassicaceae	Endangered
<u><i>Sicyos albus</i></u>	`Anunu	Cucurbitaceae	Endangered
<u><i>Sicyos lanceoloideus</i></u>	No common name	Cucurbitaceae	Endangered
<u><i>Sicyos macrophyllus</i></u>	`Anunu	Cucurbitaceae	Endangered
<u><i>Sidalcea keckii</i></u>	Keck's Checker-mallow	Malvaceae	Endangered
<u><i>Sidalcea nelsoniana</i></u>	Nelson's checker-mallow	Malvaceae	Threatened
<u><i>Sidalcea oregana ssp. valida</i></u>	Kenwood Marsh checker-mallow	Malvaceae	Endangered
<u><i>Sidalcea oregana var. calva</i></u>	Wenatchee Mountains checkermallow	Malvaceae	Endangered
<u><i>Sidalcea pedata</i></u>	Pedate checker-mallow	Malvaceae	Endangered
<u><i>Sideroxylon reclinatum ssp. austrofloridense</i></u>	Everglades bully	Sapotaceae	Threatened
<u><i>Silene alexandri</i></u>	No common name	Caryophyllaceae	Endangered
<u><i>Silene hawaiiensis</i></u>	No common name	Caryophyllaceae	Threatened
<u><i>Silene lanceolata</i></u>	No common name	Caryophyllaceae	Endangered

Scientific Name	Common Name	Family	Federal Listing Status
<u><i>Silene perlmanii</i></u>	No common name	Caryophyllaceae	Endangered
<u><i>Silene polypetala</i></u>	Fringed campion	Caryophyllaceae	Endangered
<u><i>Silene spaldingii</i></u>	Spalding's Catchfly	Caryophyllaceae	Threatened
<u><i>Sistrurus catenatus</i></u>	Eastern Massasauga (=rattlesnake)	Viperidae	Threatened
<u><i>Sisyrinchium dichotomum</i></u>	White irisette	Iridaceae	Endangered
<u><i>Solanum conocarpum</i></u>	Marron bacora	Solanaceae	Proposed Endangered
<u><i>Solanum dryophilum</i></u>	Erubia	Solanaceae	Endangered
<u><i>Solanum guamense</i></u>	Berenghenas halomtano	Solanaceae	Endangered
<u><i>Solanum incompletum</i></u>	Popolo ku mai	Solanaceae	Endangered
<u><i>Solanum nelsonii</i></u>	Popolo	Solanaceae	Endangered
<u><i>Solanum sandwicense</i></u>	`Aiakeakua, popolo	Solanaceae	Endangered
<u><i>Solidago houghtonii</i></u>	Houghton's goldenrod	Asteraceae	Threatened
<u><i>Solidago shortii</i></u>	Short's goldenrod	Asteraceae	Endangered
<u><i>Solidago spithamea</i></u>	Blue Ridge goldenrod	Asteraceae	Threatened
<u><i>Somateria fischeri</i></u>	Spectacled eider	Anatidae	Threatened
<u><i>Somatochlora hineana</i></u>	Hine's emerald dragonfly	Corduliidae	Endangered
<u><i>Sorex ornatus relictus</i></u>	Buena Vista Lake ornate Shrew	Soricidae	Endangered
<u><i>Spelaeorchestia koloana</i></u>	Kauai cave amphipod	Talitridae	Endangered
<u><i>Speoplatyrhinus poulsoni</i></u>	Alabama cavefish	Amblyopsidae	Endangered
<u><i>Spermolepis hawaiiensis</i></u>	No common name	Apiaceae	Endangered
<u><i>Speyeria callippe callippe</i></u>	Callippe silverspot butterfly	Nymphalidae	Endangered
<u><i>Speyeria zerene behrensi</i></u>	Behren's silverspot butterfly	Nymphalidae	Endangered
<u><i>Speyeria zerene hippolyta</i></u>	Oregon silverspot butterfly	Nymphalidae	Threatened
<u><i>Speyeria zerene myrtleae</i></u>	Myrtle's silverspot butterfly	Nymphalidae	Endangered
<u><i>Sphaeralcea gierischii</i></u>	Gierisch mallow	malvaceae	Endangered
<u><i>Sphyrna lewini</i></u>	Scalloped Hammerhead Shark	Sphyrnidae	Endangered
<u><i>Spigelia gentianoides</i></u>	Gentian pinkroot	Loganiaceae	Endangered

Scientific Name	Common Name	Family	Federal Listing Status
<u><i>Spiraea virginiana</i></u>	Virginia spiraea	Rosaceae	Threatened
<u><i>Spiranthes delitescens</i></u>	Canelo Hills ladies-tresses	Orchidaceae	Endangered
<u><i>Spiranthes diluvialis</i></u>	Ute ladies'-tresses	Orchidaceae	Threatened
<u><i>Spiranthes parksii</i></u>	Navasota ladies-tresses	Orchidaceae	Endangered
<u><i>Stahlia monosperma</i></u>	Cobana negra	Fabaceae	Threatened
<u><i>Stenogyne angustifolia</i> var. <i>angustifolia</i></u>	No common name	Lamiaceae	Endangered
<u><i>Stenogyne bifida</i></u>	No common name	Lamiaceae	Endangered
<u><i>Stenogyne campanulata</i></u>	No common name	Lamiaceae	Endangered
<u><i>Stenogyne cranwelliae</i></u>	No common name	Lamiaceae	Endangered
<u><i>Stenogyne kaalae</i> ssp. <i>sherffii</i></u>	No common name	Lamiaceae	Endangered
<u><i>Stenogyne kanehoana</i></u>	No common name	Lamiaceae	Endangered
<u><i>Stenogyne kauaulaensis</i></u>	No common name	Lamiaceae	Endangered
<u><i>Stenogyne kealiae</i></u>	No common name	Lamiaceae	Endangered
<u><i>Stephanomeria malheurensis</i></u>	Malheur wire-lettuce	Asteraceae	Endangered
<u><i>Sterna antillarum</i></u>	Least tern	Laridae	Endangered
<u><i>Sterna antillarum browni</i></u>	California least tern	Laridae	Endangered
<u><i>Sterna dougallii dougallii</i></u>	Roseate tern	Laridae	Endangered
<u><i>Sterna dougallii dougallii</i></u>	Roseate tern	Laridae	Threatened
<u><i>Sternotherus depressus</i></u>	Flattened musk turtle	Kinosternidae	Threatened
<u><i>Streptanthus albidus</i> ssp. <i>albidus</i></u>	Metcalf Canyon jewelflower	Brassicaceae	Endangered
<u><i>Streptanthus niger</i></u>	Tiburon jewelflower	Brassicaceae	Endangered
<u><i>Streptocephalus woottoni</i></u>	Riverside fairy shrimp	Branchinectidae	Endangered
<u><i>Strix occidentalis caurina</i></u>	Northern spotted owl	Strigidae	Threatened
<u><i>Strix occidentalis lucida</i></u>	Mexican spotted owl	Strigidae	Threatened
<u><i>Strymon acis bartrami</i></u>	Bartram's hairstreak Butterfly	Lycaenidae	Endangered
<u><i>Stygobromus</i> (= <i>Stygonectes</i>) <i>pecki</i></u>	Peck's cave amphipod	Crangonyctidae	Endangered

Scientific Name	Common Name	Family	Federal Listing Status
<u><i>Stygobromus hayi</i></u>	Hay's Spring amphipod	Crangonyctidae	Endangered
<u><i>Stygoparnus comalensis</i></u>	Comal Springs dryopid beetle	Dryopidae	Endangered
<u><i>Styrax portoricensis</i></u>	Palo de jazmin	Styracaceae	Endangered
<u><i>Styrax texanus</i></u>	Texas snowbells	Styracaceae	Endangered
<u><i>Suaeda californica</i></u>	California seablite	Chenopodiaceae	Endangered
<u><i>Succinea chittenangoensis</i></u>	Chittenango ovate amber snail	Succineidae	Threatened
<u><i>Swallenia alexandrae</i></u>	Eureka Dune grass	Poaceae	Threatened
<u><i>Sylvilagus bachmani riparius</i></u>	Riparian brush rabbit	Leporidae	Endangered
<u><i>Sylvilagus palustris hefneri</i></u>	Lower Keys marsh rabbit	Leporidae	Endangered
<u><i>Syncaris pacifica</i></u>	California freshwater shrimp	Palaemonidae	Endangered
<u><i>Tabernaemontana rotensis</i></u>	No common name	Apocynaceae	Threatened
<u><i>Tamiasciurus hudsonicus grahamensis</i></u>	Mount Graham red squirrel	Sciuridae	Endangered
<u><i>Taraxacum californicum</i></u>	California taraxacum	Asteraceae	Endangered
<u><i>Tartarocreagris texana</i></u>	Tooth Cave pseudoscorpion	Neobisiidae	Endangered
<u><i>Taylorconcha serpenticola</i></u>	Bliss Rapids snail	Hydrobiidae	Threatened
<u><i>Tectaria estremerana</i></u>	No common name	Dryopteridaceae	Endangered
<u><i>Telespyza cantans</i></u>	Laysan finch (honeycreeper)	Drepanidinae	Endangered
<u><i>Telespyza ultima</i></u>	Nihoa finch (honeycreeper)	Drepanidinae	Endangered
<u><i>Ternstroemia luquillensis</i></u>	Palo colorado	Theaceae	Endangered
<u><i>Ternstroemia subsessilis</i></u>	No common name	Theaceae	Endangered
<u><i>Tetramolopium arenarium</i></u>	No common name	Asteraceae	Endangered
<u><i>Tetramolopium capillare</i></u>	Pamakani	Asteraceae	Endangered
<u><i>Tetramolopium filiforme</i></u>	No common name	Asteraceae	Endangered
<u><i>Tetramolopium lepidotum ssp. lepidotum</i></u>	No common name	Asteraceae	Endangered
<u><i>Tetramolopium remyi</i></u>	No common name	Asteraceae	Endangered
<u><i>Tetramolopium rockii</i></u>	No common name	Asteraceae	Threatened

Scientific Name	Common Name	Family	Federal Listing Status
<u><i>Texamaurops reddelli</i></u>	Kretschmarr Cave mold beetle	Pselaphidae	Endangered
<u><i>Texella cokendolpheri</i></u>	Cokendolpher Cave Harvestman	Phalangodidae	Endangered
<u><i>Texella reddelli</i></u>	Bee Creek Cave harvestman	Phalangodidae	Endangered
<u><i>Texella reyesi</i></u>	Bone Cave harvestman	Phalangodidae	Endangered
<u><i>Thaleichthys pacificus</i></u>	Eulachon	Osmeridae	Threatened
<u><i>Thalictrum cooleyi</i></u>	Cooley's meadowrue	Ranunculaceae	Endangered
<u><i>Thamnophis eques megalops</i></u>	Northern Mexican gartersnake	Colubridae	Threatened
<u><i>Thamnophis gigas</i></u>	Giant garter snake	Colubridae	Threatened
<u><i>Thamnophis rufipunctatus</i></u>	Narrow-headed gartersnake	Colubridae	Threatened
<u><i>Thamnophis sirtalis tetrataenia</i></u>	San Francisco garter snake	Colubridae	Endangered
<u><i>Thelypodium howellii ssp. spectabilis</i></u>	Howell's spectacular thelypody	Brassicaceae	Threatened
<u><i>Thelypodium stenopetalum</i></u>	Slender-petaled mustard	Brassicaceae	Endangered
<u><i>Thelypteris inabonensis</i></u>	No common name	Thelypteridaceae	Endangered
<u><i>Thelypteris pilosa var. alabamensis</i></u>	Alabama streak-sorus fern	Thelypteridaceae	Threatened
<u><i>Thelypteris verecunda</i></u>	No common name	Thelypteridaceae	Endangered
<u><i>Thelypteris yaucoensis</i></u>	No common name	Thelypteridaceae	Endangered
<u><i>Thermosphaeroma thermophilus</i></u>	Socorro isopod	Sphaeromatidae	Endangered
<u><i>Thlaspi californicum</i></u>	Kneeland Prairie penny-cress	Brassicaceae	Endangered
<u><i>Thomomys mazama glacialis</i></u>	Roy Prairie pocket gopher	Geomyidae	Threatened
<u><i>Thomomys mazama pugetensis</i></u>	Olympia pocket gopher	Geomyidae	Threatened
<u><i>Thomomys mazama tumuli</i></u>	Tenino pocket gopher	Geomyidae	Threatened
<u><i>Thomomys mazama yelmensis</i></u>	Yelm pocket gopher	Geomyidae	Threatened
<u><i>Thymophylla tephroleuca</i></u>	Ashy dogweed	Asteraceae	Endangered

Scientific Name	Common Name	Family	Federal Listing Status
<u><i>Thysanocarpus conchuliferus</i></u>	Santa Cruz Island fringe-pod	Brassicaceae	Endangered
<u><i>Tiaroga cobitis</i></u>	Loach minnow	Cyprinidae	Endangered
<u><i>Tinospora homosepala</i></u>	No common name	Menispermaceae	Endangered
<u><i>Torreya taxifolia</i></u>	Florida torreya	Taxaceae	Endangered
<u><i>Townsendia aprica</i></u>	Last Chance townsendia	Asteraceae	Threatened
<u><i>Toxolasma cylindrellus</i></u>	Pale lilliput (pearly mussel)	Unionidae	Endangered
<u><i>Trematolobelia singularis</i></u>	No common name	Campanulaceae	Endangered
<u><i>Trichechus manatus</i></u>	West Indian Manatee	Trichechidae	Threatened
<u><i>Trichilia triacantha</i></u>	Bariaco	Meliaceae	Endangered
<u><i>Trichomanes punctatum ssp. floridanum</i></u>	Florida bristle fern	Hymenophyllaceae	Endangered
<u><i>Trifolium amoenum</i></u>	Showy Indian clover	Fabaceae	Endangered
<u><i>Trifolium stoloniferum</i></u>	Running buffalo clover	Fabaceae	Endangered
<u><i>Trifolium trichocalyx</i></u>	Monterey clover	Fabaceae	Endangered
<u><i>Trillium persistens</i></u>	Persistent trillium	Liliaceae	Endangered
<u><i>Trillium reliquum</i></u>	Relict trillium	Liliaceae	Endangered
<u><i>Trimerotropis infantilis</i></u>	Zayante band-winged grasshopper	Acrididae	Endangered
<u><i>Triodopsis platysayoides</i></u>	Flat-spined three-toothed Snail	Polygyridae	Threatened
<u><i>Tryonia alamosae</i></u>	Alamosa springsnail	Hydrobiidae	Endangered
<u><i>Tryonia cheatumi</i></u>	Phantom Tryonia	Hydrobiidae	Endangered
<u><i>Tryonia circumstriata</i></u> (= <i>stocktonensis</i>)	Gonzales tryonia	Hydrobiidae	Endangered
<u><i>Tuberolabium guamense</i></u>	No common name	Orchidaceae	Threatened
<u><i>Tuctoria greenei</i></u>	Greene's tuctoria	Poaceae	Endangered
<u><i>Tuctoria mucronata</i></u>	Solano grass	Poaceae	Endangered
<u><i>Tulotoma magnifica</i></u>	Tulotoma snail	Viviparidae	Threatened
<u><i>Tympanuchus cupido attwateri</i></u>	Attwater's greater prairie-chicken	Phasianidae	Endangered
<u><i>Typhlomolge rathbuni</i></u>	Texas blind salamander	Plethodontidae	Endangered

Scientific Name	Common Name	Family	Federal Listing Status
<u><i>Uma inornata</i></u>	Coachella Valley fringe-toed lizard	Phrynosomatidae	Threatened
<u><i>Urera kaalae</i></u>	Opuhe	Urticaceae	Endangered
<u><i>Urocitellus brunneus</i></u>	Northern Idaho Ground Squirrel	Sciuridae	Threatened
<u><i>Urocyon littoralis catalinae</i></u>	Santa Catalina Island Fox	Canidae	Threatened
<u><i>Ursus arctos horribilis</i></u>	Grizzly bear	Ursidae	Threatened
<u><i>Ursus maritimus</i></u>	Polar bear	Ursidae	Threatened
<u><i>Vagrans egistina</i></u>	Mariana wandering butterfly	Nymphalidae	Endangered
<u><i>Varronia rupicola</i></u>	No common name	Boraginaceae	Threatened
<u><i>Verbena californica</i></u>	Red Hills vervain	Verbenaceae	Threatened
<u><i>Verbesina dissita</i></u>	Big-leaved crownbeard	Asteraceae	Threatened
<u><i>Vermivora bachmanii</i></u>	Bachman's warbler (=wood)	Emberizidae	Endangered
<u><i>Vernonia proctorii</i></u>	No common name	Asteraceae	Endangered
<u><i>Vetericaris chaceorum</i></u>	Anchialine pool shrimp	Procaridae	Endangered
<u><i>Vicia menziesii</i></u>	Hawaiian vetch	Fabaceae	Endangered
<u><i>Vigna o-wahuensis</i></u>	No common name	Fabaceae	Endangered
<u><i>Villosa choctawensis</i></u>	Choctaw bean	Unionidae	Endangered
<u><i>Villosa fabalis</i></u>	Rayed Bean	Unionidae	Endangered
<u><i>Villosa perpurpurea</i></u>	Purple bean	Unionidae	Endangered
<u><i>Villosa trabalis</i></u>	Cumberland bean (pearlymussel)	Unionidae	Endangered
<u><i>Viola chamissoniana ssp. chamissoniana</i></u>	Pamakani	Violaceae	Endangered
<u><i>Viola helenae</i></u>	No common name	Violaceae	Endangered
<u><i>Viola kauaiensis var. wahiawaensis</i></u>	Nani wai`ale`ale	Violaceae	Endangered
<u><i>Viola lanaiensis</i></u>	No common name	Violaceae	Endangered
<u><i>Viola oahuensis</i></u>	No common name	Violaceae	Endangered
<u><i>Vireo bellii pusillus</i></u>	Least Bell's vireo	Vireonidae	Endangered
<u><i>Vulpes macrotis mutica</i></u>	San Joaquin kit fox	Canidae	Endangered

Scientific Name	Common Name	Family		Federal Listing Status
<u><i>Vulpes vulpes necator</i></u>	Sierra Nevada red fox	Canidae	⊕	Proposed Endangered
<u><i>Warea amplexifolia</i></u>	Wide-leaf warea	Brassicaceae	⊕	Endangered
<u><i>Warea carteri</i></u>	Carter's mustard	Brassicaceae	⊕	Endangered
<u><i>Wikstroemia skottsbergiana</i></u>	No common name	Thymelaeaceae	⊕	Endangered
<u><i>Wikstroemia villosa</i></u>	No common name	Thymelaeaceae	⊕	Endangered
<u><i>Wilkesia hobdyi</i></u>	Dwarf iliau	Asteraceae	⊕	Endangered
<u><i>Xylosma crenatum</i></u>	No common name	Salicaceae	⊕	Endangered
<u><i>Xyrauchen texanus</i></u>	Razorback sucker	Catostomidae	⊕	Endangered
<u><i>Xyris tennesseensis</i></u>	Tennessee yellow-eyed grass	Xyridaceae	⊕	Endangered
<u><i>Yermo xanthocephalus</i></u>	Desert yellowhead	Asteraceae	⊕	Threatened
<u><i>Zanthoxylum dipetalum</i> var. <i>tomentosum</i></u>	A`e	Rutaceae	⊕	Endangered
<u><i>Zanthoxylum hawaiiense</i></u>	A`e	Rutaceae	⊕	Endangered
<u><i>Zanthoxylum oahuense</i></u>	A`e	Rutaceae	⊕	Endangered
<u><i>Zanthoxylum thomsonianum</i></u>	St. Thomas prickly-ash	Rutaceae	⊕	Endangered
<u><i>Zapada glacier</i></u>	Western glacier stonefly	Nemouridae	⊕	Threatened
<u><i>Zapus hudsonius luteus</i></u>	New Mexico meadow jumping mouse	Zapodidae	⊕	Endangered
<u><i>Zapus hudsonius preblei</i></u>	Preble's meadow jumping mouse	Zapodidae	⊕	Threatened
<u><i>Zizania texana</i></u>	Texas wild-rice	Poaceae		Endangered
<u><i>Ziziphus celata</i></u>	Florida ziziphus	Rhamnaceae	⊕	Endangered
<u><i>Zosterops conspicillatus</i></u>	Bridled white-eye	Zosteropidae	⊕	Endangered
<u><i>Zosterops rotensis</i></u>	Rota bridled White-eye	Zosteropidae	⊕	Endangered

Showing 1 to 1,653 of 1,653 entries

