



Summary of Studies Supporting USDA Product Licensure

Establishment Name	Zoetis Inc.
USDA Vet Biologics Establishment Number	190
Product Code	19K5.R1
True Name	Porcine Circovirus Vaccine, Type 1 - Type 2 Chimera, Killed Virus
Tradename(s) / Distributor or Subsidiary (if different from manufacturer)	Fostera Gold PCV - No distributor specified Fostera Gold PCV - Zoetis (Thailand) Limited Fostera Gold PCV - Zoetis Australia Pty Ltd Fostera Gold PCV - Zoetis Colombia S.A.S. Fostera Gold PCV - Zoetis Industria Productos Veterinarios Ltda. Fostera Gold PCV - Zoetis Japan Inc. Fostera Gold PCV - Zoetis Mexico Not Listed - Zoetis Korea
Date of Compilation Summary	September 07, 2022

Disclaimer: Do not use the following studies to compare one product to another. Slight differences in study design and execution can render the comparisons meaningless.

Study Type	Efficacy																
Pertaining to	Circovirus, porcine, type 2 (PCV2)																
Study Purpose	Efficacy against Porcine Circovirus type 2b																
Product Administration	Two doses administered intramuscularly 3 weeks apart.																
Study Animals	PCV2 serologically negative, commercial pigs, 3 days of age at first administration; 24 vaccinates and 25 controls.																
Challenge Description	Porcine Circovirus 2 (PCV2b), 3 weeks after second vaccination.																
Interval observed after challenge	<p>PCV2 viremia (virus in serum) and PCV2 shedding in feces were evaluated twice weekly post challenge for 3 weeks.</p> <p>Lymphoid tissues were evaluated 3 weeks post challenge for the presence of PCV2 and characteristic changes attributable to PCV2 infection (lymphoid depletion).</p>																
Results	<table border="1"> <thead> <tr> <th>Variable</th> <th>Vaccinates Ever Positive*</th> <th>Controls Ever Positive*</th> </tr> </thead> <tbody> <tr> <td>Viremia (PCV2 detection in serum)</td> <td>3/24 (13%)</td> <td>19/25 (76%)</td> </tr> <tr> <td>Fecal Shedding (PCV2 detection in feces)</td> <td>0/24 (0%)</td> <td>20/25 (80%)</td> </tr> <tr> <td>Lymphoid Depletion (PCV2 lesions)</td> <td>1/24 (4%)</td> <td>8/25 (32%)</td> </tr> <tr> <td>Lymphoid Colonization (PCV2 in lymphoid tissues)</td> <td>1/24 (4%)</td> <td>11/25 (44%)</td> </tr> </tbody> </table> <p>*A pig was considered positive for PCV2 detection in serum or feces if it had a positive result at any sampling point.</p> <p>Raw data to follow:</p>		Variable	Vaccinates Ever Positive*	Controls Ever Positive*	Viremia (PCV2 detection in serum)	3/24 (13%)	19/25 (76%)	Fecal Shedding (PCV2 detection in feces)	0/24 (0%)	20/25 (80%)	Lymphoid Depletion (PCV2 lesions)	1/24 (4%)	8/25 (32%)	Lymphoid Colonization (PCV2 in lymphoid tissues)	1/24 (4%)	11/25 (44%)
Variable	Vaccinates Ever Positive*	Controls Ever Positive*															
Viremia (PCV2 detection in serum)	3/24 (13%)	19/25 (76%)															
Fecal Shedding (PCV2 detection in feces)	0/24 (0%)	20/25 (80%)															
Lymphoid Depletion (PCV2 lesions)	1/24 (4%)	8/25 (32%)															
Lymphoid Colonization (PCV2 in lymphoid tissues)	1/24 (4%)	11/25 (44%)															

**Results
(continued)**

Control ID	viremia	fecal shedding	lymphoid colonization	lymphoid depletion	Vaccinate ID	viremia	fecal shedding	lymphoid colonization	lymphoid depletion
	13	+	+	-		-	1	-	-
16	+	+	+	-	10	-	-	-	-
2	+	+	-	-	11	-	-	-	-
29	-	-	-	-	12	-	-	-	-
30	-	-	-	-	14	-	-	-	-
31	+	+	-	-	15	-	-	-	-
33	-	+	-	-	23	-	-	-	-
34	+	+	+	+	26	-	-	-	-
35	+	+	+	+	27	-	-	-	-
38	+	+	+	+	28	-	-	-	-
4	+	+	-	-	36	-	-	-	-
48	+	-	-	-	43	-	-	-	-
5	+	+	-	+	50	-	-	-	-
52	+	+	+	-	51	-	-	-	-
6	-	-	-	-	55	-	-	-	-
60	-	+	+	+	65	+	-	+	+
62	+	+	+	-	66	+	-	-	-
68	+	+	+	+	67	+	-	-	-
69	+	+	+	-	7	-	-	-	-
70	+	-	-	-	72	-	-	-	-
76	+	+	-	-	74	-	-	-	-
78	+	+	+	+	79	-	-	-	-
8	-	+	-	-	86	-	-	-	-
83	+	+	+	+	9	-	-	-	-
88	+	+	-	-					

Viremia, observed over time:

Control	Day 42-46	Day 46-50	Day 49-53	Day 54-58	Day 56-60	Day 60-64
13	-	-	+	-	+	-
16	-	+	+	+	+	+
2	-	+	+	-	+	-
29	-	-	-	-	-	-
30	-	-	-	-	-	-
31	-	-	+	-	-	-
33	-	-	-	-	-	-
34	-	+	+	+	+	+
35	-	-	+	+	-	-
38	-	-	+	+	+	-
4	-	+	+	+	-	-
48	-	-	+	-	-	-
5	-	+	+	+	+	-
52	-	-	+	+	-	-
6	-	-	-	-	-	-
60	-	-	-	-	-	-
62	-	+	+	+	+	-
68	-	+	+	+	+	-
69	-	+	+	+	+	-
70	-	-	-	+	-	-
76	-	-	+	+	+	-
78	-	-	+	+	+	+
8	-	-	-	-	-	-
83	-	-	+	+	+	-
88	-	+	+	+	+	-

Vaccinate	Day 42-46	Day 46-50	Day 49-53	Day 54-58	Day 56-60	Day 60-64
1	-	-	-	-	-	-
10	-	-	-	-	-	-
11	-	-	-	-	-	-
12	-	-	-	-	-	-
14	-	-	-	-	-	-
15	-	-	-	-	-	-
23	-	-	-	-	-	-
26	-	-	-	-	-	-
27	-	-	-	-	-	-
28	-	-	-	-	-	-
36	-	-	-	-	-	-
43	-	-	-	-	-	-
50	-	-	-	-	-	-

51	-	-	-	-	-	-
55	-	-	-	-	-	-
65	-	-	-	+	+	-
66	-	-	+	-	+	-
67	-	+	-	+	-	-
7	-	-	-	-	-	-
72	-	-	-	-	-	-
74	-	-	-	-	-	-
79	-	-	-	-	-	-
86	-	-	-	-	-	-
9	-	-	-	-	-	-

Fecal shedding, observed over time:

Control	Day 42-46	Day 46-50	Day 49-53	Day 54-58	Day 56-60	Day 60-64
13	-	-	+	+	-	-
16	-	+	+	+	+	+
2	-	-	+	+	-	-
29	-	-	-	-	-	-
30	-	-	-	-	-	-
31	-	-	+	+	-	-
33	-	-	+	-	-	+
34	-	+	+	+	+	+
35	-	-	+	+	+	-
38	-	-	+	+	+	-
4	-	+	+	+	-	+
48	-	-	-	-	-	-
5	-	-	-	+	+	-
52	-	-	+	+	+	+
6	-	-	-	-	-	-
60	-	-	-	+	-	-
62	-	-	+	+	+	+
68	-	-	-	+	+	+
69	-	-	+	+	+	+
70	-	-	-	-	-	-
76	-	-	+	+	+	+
78	-	+	+	+	+	+
8	-	-	-	+	-	+
83	-	-	+	+	+	+
88	-	-	+	+	-	+

All vaccinates were negative at all sampling points

**USDA Approval
Date**

April 07, 2017

Study Type	Efficacy
Pertaining to	Circovirus, porcine, type 2 (PCV2)
Study Purpose	Pivotal efficacy against Porcine Circovirus type 2b
Product Administration	One dose, administered intramuscularly
Study Animals	PCV2 serologically negative, commercial pigs, 21 days of age at administration; 26 vaccinates and 14 controls.
Challenge Description	Porcine Circovirus 2 (PCV2b), 3 weeks after vaccination
Interval observed after challenge	PCV2 viremia (virus in serum) and PCV2 fecal shedding were evaluated twice weekly post challenge for 3 weeks. Lymphoid tissues were evaluated 3 weeks post challenge for the presence of PCV2 and characteristic changes attributable to PCV2 infection (lymphoid depletion).

Results	Results:		
	Variable	Vaccinates Ever Positive*	Controls Ever Positive*
	Viremia (PCV2 detection in serum)	3/26 (12%)	10/14 (71%)
	Fecal Shedding (PCV2 detection in feces)	7/26 (27%)	12/14 (86%)
	Lymphoid Depletion (PCV2 lesions)	6/26 (23%)	11/14 (79%)
	Lymphoid Colonization (PCV2 in lymphoid tissues)	8/26 (31%)	12/14 (86%)
	*A pig was considered positive for PCV2 detection in serum or feces if it had a positive result at any sampling point.		

Results (Continued)	Control ID	viremia	fecal shedding	lymphoid colonization	lymphoid depletion	Vaccinate ID	viremia	fecal shedding	lymphoid colonization	lymphoid depletion
		122	+	+	+	+	121	-	-	-
	146	+	+	-	-	123	-	-	-	-
	153	+	+	+	+	142	-	-	-	+
	167	+	+	+	+	147	-	-	+	-
	188	+	+	+	+	152	-	-	-	-
	193	-	-	-	-	155	-	+	-	-
	202	-	+	+	+	156	-	-	-	-
	214	+	+	+	+	158	-	-	-	-
	25	+	+	+	+	166	-	-	+	+
	358	-	-	+	+	170	+	+	+	+
	368	+	+	+	-	183	-	+	-	-
	378	+	+	+	+	187	+	+	+	+
	48	+	+	+	+	192	-	-	-	-
	78	-	+	+	+	201	-	-	-	-
						205	-	-	-	-
						21	-	+	-	-
						213	+	-	-	-
						357	-	-	+	+
						369	-	-	-	-
						372	-	+	-	-
						377	-	-	-	-
						41	-	-	+	-
						46	-	+	+	-
						77	-	-	+	+
						84	-	-	-	-
						89	-	-	-	-
	Raw data, by study day, shown below.									
USDA Approval Date	09/15/2017									

Viremia, observed over time:

		Study Day					
	Animal	24	28	31	35	38	42
Control	25	-	+	+	+	+	+
Control	48	-	+	+	+	+	+
Control	78	-	-	-	-	-	-
Control	122	-	-	+	+	+	-
Control	146	-	-	+	+	+	-
Control	153	-	-	+	+	+	+
Control	167	-	-	+	+	+	+
Control	188	-	-	-	+	+	+
Control	193	-	-	-	-	-	-
Control	202	-	-	-	-	-	-
Control	214	-	-	+	+	+	+
Control	358	-	-	-	-	-	-
Control	368	-	-	+	-	-	-
Control	378	-	-	-	+	+	-
Vaccinate	21	-	-	-	-	-	-
Vaccinate	41	-	-	-	-	-	-
Vaccinate	46	-	-	-	-	-	-
Vaccinate	77	-	-	-	-	-	-
Vaccinate	84	-	-	-	-	-	-
Vaccinate	89	-	-	-	-	-	-
Vaccinate	121	-	-	-	-	-	-
Vaccinate	123	-	-	-	-	-	-
Vaccinate	142	-	-	-	-	-	-
Vaccinate	147	-	-	-	-	-	-
Vaccinate	152	-	-	-	-	-	-
Vaccinate	155	-	-	-	-	-	-
Vaccinate	156	-	-	-	-	-	-
Vaccinate	158	-	-	-	-	-	-
Vaccinate	166	-	-	-	-	-	-

		Study Day					
	Animal	24	28	31	35	38	42
Vaccinate	170	-	-	+	-	-	-
Vaccinate	183	-	-	-	-	-	-
Vaccinate	187	-	+	+	-	-	-
Vaccinate	192	-	-	-	-	-	-
Vaccinate	201	-	-	-	-	-	-
Vaccinate	205	-	-	-	-	-	-
Vaccinate	213	-	-	+	-	-	-
Vaccinate	357	-	-	-	-	-	-
Vaccinate	369	-	-	-	-	-	-
Vaccinate	372	-	-	-	-	-	-
Vaccinate	377	-	-	-	-	-	-

Fecal shedding, observed over time:

		Study Day					
	Animal	24	28	31	35	38	42
Control	25	-	+	+	+	+	+
Control	48	-	+	+	+	+	+
Control	78	-	-	+	+	+	+
Control	122	-	-	+	-	+	-
Control	146	-	-	-	-	+	+
Control	153	-	-	-	+	+	+
Control	167	-	-	+	+	+	+
Control	188	-	-	-	+	+	-
Control	193	-	-	-	-	-	-
Control	202	-	-	-	+	-	-
Control	214	-	-	+	+	+	+
Control	358	-	-	-	-	-	-
Control	368	-	-	+	+	-	-
Control	378	-	-	-	+	+	+

		Study Day					
	Animal	24	28	31	35	38	42
Vaccinate	21	-	-	-	-	+	-
Vaccinate	41	-	-	-	-	-	-
Vaccinate	46	+	-	-	-	-	-
Vaccinate	77	-	-	-	-	-	-
Vaccinate	84	-	-	-	-	-	-
Vaccinate	89	-	-	-	-	-	-
Vaccinate	121	-	-	-	-	-	-
Vaccinate	123	-	-	-	-	-	-
Vaccinate	142	-	-	-	-	-	-
Vaccinate	147	-	-	-	-	-	-
Vaccinate	152	-	-	-	-	-	-
Vaccinate	155	-	-	-	-	-	+
Vaccinate	156	-	-	-	-	-	-
Vaccinate	158	-	-	-	-	-	-
Vaccinate	166	-	-	-	-	-	-
Vaccinate	170	-	-	+	-	-	+
Vaccinate	183	-	-	-	-	-	+
Vaccinate	187	-	-	+	-	-	-
Vaccinate	192	-	-	-	-	-	-
Vaccinate	201	-	-	-	-	-	-
Vaccinate	205	-	-	-	-	-	-
Vaccinate	213	-	-	-	-	-	-
Vaccinate	357	-	-	-	-	-	-
Vaccinate	369	-	-	-	-	-	-
Vaccinate	372	-	-	-	+	-	-
Vaccinate	377	-	-	-	-	-	-

Study Type	Efficacy															
Pertaining to	Circovirus, porcine, type 2 (PCV2)															
Study Purpose	Pivotal efficacy against Porcine Circovirus type 2a															
Product Administration	Two doses, administered intramuscularly, 3 weeks apart															
Study Animals	PCV2 serologically negative, commercial pigs, 3 days of age at first administration; 30 vaccinates and 29 controls.															
Challenge Description	Porcine Circovirus 2 (PCV2a), 3 weeks after second vaccination															
Interval observed after challenge	PCV2 viremia (virus in serum) and PCV2 shedding in feces were evaluated twice weekly post challenge for 3 weeks. Lymphoid tissues were evaluated 3 weeks post challenge for the presence of PCV2 and characteristic changes attributable to PCV2 infection (lymphoid depletion and histiocytic replacement).															
Results	<p>Results:</p> <table border="1"> <thead> <tr> <th>Variable</th> <th>Vaccinates Ever Positive*</th> <th>Controls Ever Positive*</th> </tr> </thead> <tbody> <tr> <td>Viremia (PCV2 detection in serum)</td> <td>1/30 (3%)</td> <td>24/29 (83%)</td> </tr> <tr> <td>Fecal Shedding (PCV2 detection in feces)</td> <td>5/30 (17%)</td> <td>25/29 (86%)</td> </tr> <tr> <td>Lymphoid Depletion (PCV2 lesions)</td> <td>1/30 (3%)</td> <td>12/29 (41%)</td> </tr> <tr> <td>Lymphoid Colonization (PCV2 in lymphoid tissues)</td> <td>2/30 (7%)</td> <td>14/29 (48%)</td> </tr> </tbody> </table> <p>*A pig was considered positive for PCV2 detection in serum or feces if it had a positive result at any sampling point.</p>	Variable	Vaccinates Ever Positive*	Controls Ever Positive*	Viremia (PCV2 detection in serum)	1/30 (3%)	24/29 (83%)	Fecal Shedding (PCV2 detection in feces)	5/30 (17%)	25/29 (86%)	Lymphoid Depletion (PCV2 lesions)	1/30 (3%)	12/29 (41%)	Lymphoid Colonization (PCV2 in lymphoid tissues)	2/30 (7%)	14/29 (48%)
Variable	Vaccinates Ever Positive*	Controls Ever Positive*														
Viremia (PCV2 detection in serum)	1/30 (3%)	24/29 (83%)														
Fecal Shedding (PCV2 detection in feces)	5/30 (17%)	25/29 (86%)														
Lymphoid Depletion (PCV2 lesions)	1/30 (3%)	12/29 (41%)														
Lymphoid Colonization (PCV2 in lymphoid tissues)	2/30 (7%)	14/29 (48%)														

Results (Continued)	Control ID					Vaccinate ID				
		viremia	fecal shedding	lymphoid colonization	lymphoid depletion		viremia	fecal shedding	lymphoid colonization	lymphoid depletion
	2	+	+	+	+	1	-	+	-	-
	21	+	+	+	+	10	-	-	-	-
	23	+	+	+	+	11	-	-	-	-
	26	+	+	-	-	12	-	+	+	-
	27	+	+	-	+	16	-	-	-	-
	28	+	+	+	+	17	-	-	-	-
	3	+	+	+	+	18	-	-	-	-
	31	+	+	-	-	19	-	+	-	-
	32	-	-	-	-	20	-	-	-	-
	35	+	+	+	+	25	-	-	-	-
	36	+	+	-	+	29	-	+	-	-
	39	+	-	-	-	30	-	-	-	-
	4	+	+	+	+	37	-	-	-	-
	45	+	+	-	-	38	-	-	-	-
	5	+	+	-	-	40	-	-	-	-
	52	-	-	-	-	47	-	-	-	-
	53	-	-	-	-	49	-	+	-	-
	57	+	+	+	-	50	-	-	-	-
	58	+	+	+	+	55	-	-	-	-
	59	+	+	+	-	56	-	-	-	-
	60	+	+	+	-	6	-	-	-	-
	63	+	+	+	-	61	-	-	-	-
	66	+	+	-	-	64	-	-	-	-
	68	-	+	-	-	65	-	-	+	-
	69	+	+	-	-	67	-	-	-	-
	71	+	+	+	+	70	-	-	-	-
	73	+	+	-	-	72	-	-	-	-
	80	-	+	-	-	8	-	-	-	+
	9	+	+	+	+	83	+	-	-	-
						84	-	-	-	-
	Raw Data, by sampling day, presented below									
USDA Approval Date	August 25, 2017									

Viremia, observed over time:

	Animal	Study Day						
		43-47	47-51	51-55	55-58	58-62	61-65	64-68
Control	2	-	-	+	+	+	+	+
Control	3	-	-	+	+	+	+	+
Control	4	-	-	+	+	+	+	+
Control	5	-	-	+	-	-	-	+
Control	9	-	-	+	+	+	+	+
Control	21	-	-	+	+	+	+	+
Control	23	-	-	-	+	+	-	-
Control	26	-	-	+	+	-	-	-
Control	27	-	+	+	+	-	-	-
Control	28	-	-	+	+	-	-	-
Control	31	-	-	+	+	-	-	-
Control	32	-	-	-	-	-	-	-
Control	35	-	-	+	+	-	-	-
Control	36	-	-	+	+	-	-	-
Control	39	-	-	-	+	-	-	-
Control	45	-	-	-	+	-	+	-
Control	52	-	-	-	-	-	-	-
Control	53	-	-	-	-	-	-	-
Control	57	-	-	+	+	+	+	+
Control	58	-	-	+	+	-	-	-
Control	59	-	-	+	+	+	+	+
Control	60	-	-	+	+	-	+	-
Control	63	-	-	+	+	+	-	+
Control	66	-	-	+	+	+	+	+
Control	68	-	-	-	-	-	-	-
Control	69	-	-	-	-	-	+	+
Control	71	-	-	+	+	+	+	+
Control	73	-	-	+	+	-	-	-
Control	80	-	-	-	-	-	-	-

		Study Day						
	Animal	43-47	47-51	51-55	55-58	58-62	61-65	64-68
Vaccinate	1	-	-	-	-	-	-	-
Vaccinate	6	-	-	-	-	-	-	-
Vaccinate	8	-	-	-	-	-	-	-
Vaccinate	10	-	-	-	-	-	-	-
Vaccinate	11	-	-	-	-	-	-	-
Vaccinate	12	-	-	-	-	-	-	-
Vaccinate	16	-	-	-	-	-	-	-
Vaccinate	17	-	-	-	-	-	-	-
Vaccinate	18	-	-	-	-	-	-	-
Vaccinate	19	-	-	-	-	-	-	-
Vaccinate	20	-	-	-	-	-	-	-
Vaccinate	25	-	-	-	-	-	-	-
Vaccinate	29	-	-	-	-	-	-	-
Vaccinate	30	-	-	-	-	-	-	-
Vaccinate	37	-	-	-	-	-	-	-
Vaccinate	38	-	-	-	-	-	-	-
Vaccinate	40	-	-	-	-	-	-	-
Vaccinate	47	-	-	-	-	-	-	-
Vaccinate	49	-	-	-	-	-	-	-
Vaccinate	50	-	-	-	-	-	-	-
Vaccinate	55	-	-	-	-	-	-	-
Vaccinate	56	-	-	-	-	-	-	-
Vaccinate	61	-	-	-	-	-	-	-
Vaccinate	64	-	-	-	-	-	-	-
Vaccinate	65	-	-	-	-	-	-	-
Vaccinate	67	-	-	-	-	-	-	-
Vaccinate	70	-	-	-	-	-	-	-
Vaccinate	72	-	-	-	-	-	-	-
Vaccinate	83	-	-	-	+	-	-	-
Vaccinate	84	-	-	-	-	-	-	-

Fecal Shedding, observed over time:

	Animal	Study Day						
		43-47	47-51	51-55	55-58	58-62	61-65	64-68
Control	2	-	-	+	+	+	+	+
Control	3	-	-	+	+	+	+	+
Control	4	-	-	+	+	+	+	+
Control	5	-	-	+	+	+	+	+
Control	9	-	-	+	+	+	+	+
Control	21	-	-	+	+	+	+	+
Control	23	-	-	+	+	+	+	-
Control	26	-	-	+	+	+	+	+
Control	27	-	+	+	+	+	-	+
Control	28	-	+	+	+	+	+	+
Control	31	-	-	+	+	+	-	+
Control	32	-	-	-	-	-	-	-
Control	35	-	-	+	+	+	-	-
Control	36	-	-	+	+	-	+	+
Control	39	-	-	-	-	-	-	-
Control	45	-	-	-	+	+	+	-
Control	52	-	-	-	-	-	-	-
Control	53	-	-	-	-	-	-	-
Control	57	-	-	+	+	+	+	+
Control	58	-	-	+	+	-	-	-
Control	59	-	-	+	+	+	+	+
Control	60	-	-	-	+	+	-	+
Control	63	-	-	+	+	+	+	+
Control	66	-	-	+	+	+	+	+
Control	68	-	-	-	-	+	+	+
Control	69	-	-	+	+	+	+	+
Control	71	-	-	+	+	+	+	+
Control	73	-	-	+	-	+	-	-
Control	80	-	+	+	-	+	-	-

		Study Day						
	Animal	43-47	47-51	51-55	55-58	58-62	61-65	64-68
Vaccinate	1	-	-	+	-	-	-	-
Vaccinate	6	-	-	-	-	-	-	-
Vaccinate	8	-	-	-	-	-	-	-
Vaccinate	10	-	-	-	-	-	-	-
Vaccinate	11	-	-	-	-	-	-	-
Vaccinate	12	-	-	-	-	+	-	-
Vaccinate	16	-	-	-	-	-	-	-
Vaccinate	17	-	-	-	-	-	-	-
Vaccinate	18	-	-	-	-	-	-	-
Vaccinate	19	-	-	+	-	-	-	-
Vaccinate	20	-	-	-	-	-	-	-
Vaccinate	25	-	-	-	-	-	-	-
Vaccinate	29	-	+	-	-	-	-	-
Vaccinate	30	-	-	-	-	-	-	-
Vaccinate	37	-	-	-	-	-	-	-
Vaccinate	38	-	-	-	-	-	-	-
Vaccinate	40	-	-	-	-	-	-	-
Vaccinate	47	-	-	-	-	-	-	-
Vaccinate	49	-	-	+	+	-	-	-
Vaccinate	50	-	-	-	-	-	-	-
Vaccinate	55	-	-	-	-	-	-	-
Vaccinate	56	-	-	-	-	-	-	-
Vaccinate	61	-	-	-	-	-	-	-
Vaccinate	64	-	-	-	-	-	-	-
Vaccinate	65	-	-	-	-	-	-	-
Vaccinate	67	-	-	-	-	-	-	-
Vaccinate	70	-	-	-	-	-	-	-
Vaccinate	72	-	-	-	-	-	-	-
Vaccinate	83	-	-	-	-	-	-	-
Vaccinate	84	-	-	-	-	-	-	-

Study Type	Efficacy																
Pertaining to	Circovirus, porcine, type 2 (PCV2)																
Study Purpose	Efficacy against Porcine Circovirus type 2a.																
Product Administration	One dose, administered intramuscularly.																
Study Animals	PCV2 serologically negative, commercial pigs, 3 weeks of age at administration; 28 vaccinates and 29 controls.																
Challenge Description	Porcine Circovirus 2 (PCV2a), 3 weeks after vaccination.																
Interval observed after challenge	<p>PCV2 viremia (virus in serum) and PCV2 fecal shedding were evaluated twice weekly post challenge for 3 weeks.</p> <p>Lymphoid tissues were evaluated 3 weeks post challenge for the presence of PCV2 and characteristic changes attributable to PCV2 infection (lymphoid depletion).</p>																
Results	<table border="1"> <thead> <tr> <th>Variable</th> <th>Vaccinates Ever Positive*</th> <th>Controls Ever Positive*</th> </tr> </thead> <tbody> <tr> <td>Viremia (PCV2 detection in serum)</td> <td>1/28 (4%)</td> <td>22/29 (76%)</td> </tr> <tr> <td>Fecal Shedding (PCV2 detection in feces)</td> <td>7/28 (25%)</td> <td>22/29 (76%)</td> </tr> <tr> <td>Lymphoid Depletion (PCV2 lesions)</td> <td>0/28 (0%)</td> <td>9/29 (31%)</td> </tr> <tr> <td>Lymphoid Colonization (PCV2 in lymphoid tissues)</td> <td>0/28 (0%)</td> <td>13/29 (45%)</td> </tr> </tbody> </table> <p>*A pig was considered positive for PCV2 detection in serum or feces if it had a positive result at any sampling point.</p> <p>Raw data to follow:</p>		Variable	Vaccinates Ever Positive*	Controls Ever Positive*	Viremia (PCV2 detection in serum)	1/28 (4%)	22/29 (76%)	Fecal Shedding (PCV2 detection in feces)	7/28 (25%)	22/29 (76%)	Lymphoid Depletion (PCV2 lesions)	0/28 (0%)	9/29 (31%)	Lymphoid Colonization (PCV2 in lymphoid tissues)	0/28 (0%)	13/29 (45%)
Variable	Vaccinates Ever Positive*	Controls Ever Positive*															
Viremia (PCV2 detection in serum)	1/28 (4%)	22/29 (76%)															
Fecal Shedding (PCV2 detection in feces)	7/28 (25%)	22/29 (76%)															
Lymphoid Depletion (PCV2 lesions)	0/28 (0%)	9/29 (31%)															
Lymphoid Colonization (PCV2 in lymphoid tissues)	0/28 (0%)	13/29 (45%)															

Results (continued)	Control ID					Vaccinate ID				
		viremia	fecal shedding	lymphoid colonization	lymphoid depletion		viremia	fecal shedding	lymphoid colonization	lymphoid depletion
	2457	+	+	+	+	2463	-	-	-	-
	2464	+	+	+	-	4851	-	+	-	-
	4850	+	+	-	-	4852	-	-	-	-
	5638	-	-	-	-	5644	-	-	-	-
	5645	+	-	-	-	5649	-	-	-	-
	5647	+	+	-	-	5654	-	-	-	-
	5650	+	+	+	+	5679	-	+	-	-
	5652	+	+	+	+	5686	-	-	-	-
	5653	+	+	+	-	5699	-	-	-	-
	5658	+	+	-	-	5705	-	-	-	-
	5684	+	+	-	+	5717	-	-	-	-
	5700	+	+	+	-	5719	-	+	-	-
	5707	-	-	-	-	5720	-	-	-	-
	5710	+	+	+	+	5725	-	+	-	-
	5713	-	+	-	-	5760	-	-	-	-
	5716	+	+	-	-	5761	-	-	-	-
	5723	+	+	+	+	5766	-	+	-	-
	5726	-	-	-	-	5767	-	-	-	-
	5762	+	-	-	-	6519	-	-	-	-
	6503	-	-	-	-	6524	-	-	-	-
	6515	-	-	-	-	6527	-	-	-	-
	6526	-	+	-	-	6534	-	+	-	-
	6542	+	+	+	-	6540	-	-	-	-
	6552	+	+	-	-	6545	+	-	-	-
	6564	+	+	+	+	6560	-	-	-	-
	6565	+	+	+	+	6568	-	-	-	-
	6572	+	+	+	-	6581	-	-	-	-
	6589	+	+	-	-	6582	-	+	-	-
	6599	+	+	+	+					
USDA Approval Date	September 26, 2016									

Study Type	Efficacy															
Pertaining to	Circovirus, porcine, type 2 (PCV2)															
Study Purpose	Pivotal efficacy against Porcine Circovirus type 2															
Product Administration	Two doses, administered intramuscularly, 3 weeks apart.															
Study Animals	PCV2 serologically negative, commercial pigs, 3 weeks of age at first administration; 29 vaccinates and 27 controls.															
Challenge Description	Porcine Circovirus 2 (PCV2a), 3 weeks after second vaccination.															
Interval observed after challenge	PCV2 viremia (virus in serum) and PCV2 fecal shedding were evaluated weekly post challenge for 3 weeks. Lymphoid tissues were evaluated 3 weeks post challenge for the presence of PCV2 and characteristic changes attributable to PCV2 infection (lymphoid depletion and histiocytic replacement).															
Results	<table border="1"> <thead> <tr> <th>Variable</th> <th>Vaccinates Ever Positive*</th> <th>Controls Ever Positive*</th> </tr> </thead> <tbody> <tr> <td>Viremia (PCV2 detection in serum)</td> <td>0/29 (0%)</td> <td>25/27 (93%)</td> </tr> <tr> <td>Fecal Shedding (PCV2 detection in feces)</td> <td>7/29 (24%)</td> <td>25/27 (93%)</td> </tr> <tr> <td>Lymphoid Depletion (PCV2 lesions)</td> <td>0/29 (0%)</td> <td>4/27 (15%)</td> </tr> <tr> <td>Lymphoid Colonization (PCV2 in lymphoid tissues)</td> <td>0/29 (0%)</td> <td>3/27 (11%)</td> </tr> </tbody> </table> <p>*A pig was considered positive for PCV2 detection in serum or feces if it had a positive result at any sampling point.</p> <p>Raw data to follow:</p>	Variable	Vaccinates Ever Positive*	Controls Ever Positive*	Viremia (PCV2 detection in serum)	0/29 (0%)	25/27 (93%)	Fecal Shedding (PCV2 detection in feces)	7/29 (24%)	25/27 (93%)	Lymphoid Depletion (PCV2 lesions)	0/29 (0%)	4/27 (15%)	Lymphoid Colonization (PCV2 in lymphoid tissues)	0/29 (0%)	3/27 (11%)
Variable	Vaccinates Ever Positive*	Controls Ever Positive*														
Viremia (PCV2 detection in serum)	0/29 (0%)	25/27 (93%)														
Fecal Shedding (PCV2 detection in feces)	7/29 (24%)	25/27 (93%)														
Lymphoid Depletion (PCV2 lesions)	0/29 (0%)	4/27 (15%)														
Lymphoid Colonization (PCV2 in lymphoid tissues)	0/29 (0%)	3/27 (11%)														
Results																

(continued)	Control ID					Vaccinate ID				
		viremia	fecal shedding	lymphoid colonization	lymphoid depletion		viremia	fecal shedding	lymphoid colonization	lymphoid depletion
	1102	+	+	-	-	1101	-	+	-	-
	1103	+	+	-	-	1104	-	+	-	-
	1107	+	+	-	-	1105	-	-	-	-
	1108	+	+	-	-	1106	-	-	-	-
	1111	+	+	+	-	1112	-	-	-	-
	1114	+	+	-	-	1115	-	-	-	-
	1117	+	+	-	+	1118	-	-	-	-
	1121	+	+	-	-	1123	-	+	-	-
	1125	-	-	-	-	1124	-	-	-	-
	1126	+	+	-	+	1127	-	+	-	-
	1131	+	+	+	+	1132	-	-	-	-
	1135	+	+	-	-	1133	-	+	-	-
	1138	+	+	-	-	1134	-	-	-	-
	1145	+	+	+	+	1143	-	-	-	-
	1146	+	+	-	-	1147	-	-	-	-
	1150	+	+	-	-	1148	-	-	-	-
	1153	-	+	-	-	1149	-	+	-	-
	1154	+	+	-	-	1151	-	-	-	-
	1159	+	+	-	-	1155	-	-	-	-
	1169	+	+	-	-	1156	-	-	-	-
	1170	+	-	-	-	1158	-	-	-	-
	1176	+	+	-	-	1167	-	-	-	-
	1178	+	+	-	-	1171	-	-	-	-
	1181	+	+	-	-	1172	-	+	-	-
	1188	+	+	-	-	1177	-	-	-	-
	1195	+	+	-	-	1180	-	-	-	-
	1199	+	+	-	-	1191	-	-	-	-
						1194	-	-	-	-
						1198	-	-	-	-
USDA Approval Date	September 13, 2013									

Study Type	Efficacy												
Pertaining to	Circovirus, porcine, type 2 (PCV2)												
Study Purpose	Duration of immunity against Porcine Circovirus 2.												
Product Administration	One dose, administered intramuscularly.												
Study Animals	PCV2 serologically negative, commercial pigs, 3 weeks of age at administration; 28 vaccinates and 26 controls.												
Challenge Description	Porcine Circovirus 2 (PCV2b), 23 weeks after vaccination.												
Interval observed after challenge	PCV2 viremia (virus in serum) was evaluated twice weekly, and fecal shedding was evaluated weekly, for 30 days after challenge. Lymphoid tissues were evaluated 30 days post challenge for the presence of PCV2 and characteristic changes attributable to PCV2 infection (lymphoid depletion and histiocytic replacement).												
Results	<table border="1"> <thead> <tr> <th>Variable</th> <th>Vaccinates Ever Positive*</th> <th>Controls Ever Positive*</th> </tr> </thead> <tbody> <tr> <td>Viremia (PCV2 detection in serum)</td> <td>0/28 (0%)</td> <td>15/26 (58%)</td> </tr> <tr> <td>Fecal Shedding (PCV2 detection in feces)</td> <td>5/28 (18%)</td> <td>16/26 (62%)</td> </tr> <tr> <td>Lymphoid Colonization (PCV2 in lymphoid tissues)</td> <td>1/28 (4%)</td> <td>9/26 (35%)</td> </tr> </tbody> </table> <p>*A pig was considered positive for PCV2 detection in serum or feces if it had a positive result at any sampling point.</p> <p>Raw data to follow:</p>	Variable	Vaccinates Ever Positive*	Controls Ever Positive*	Viremia (PCV2 detection in serum)	0/28 (0%)	15/26 (58%)	Fecal Shedding (PCV2 detection in feces)	5/28 (18%)	16/26 (62%)	Lymphoid Colonization (PCV2 in lymphoid tissues)	1/28 (4%)	9/26 (35%)
Variable	Vaccinates Ever Positive*	Controls Ever Positive*											
Viremia (PCV2 detection in serum)	0/28 (0%)	15/26 (58%)											
Fecal Shedding (PCV2 detection in feces)	5/28 (18%)	16/26 (62%)											
Lymphoid Colonization (PCV2 in lymphoid tissues)	1/28 (4%)	9/26 (35%)											

Results (continued)	Control ID				Vaccinate ID			
		viremia	fecal shedding	lymphoid colonization		viremia	fecal shedding	lymphoid colonization
	502	+	+	+	501	-	-	-
	505	+	+	-	503	-	-	-
	506	-	-	-	504	-	-	-
	507	-	+	-	508	-	-	-
	509	+	+	+	510	-	-	-
	512	-	-	-	511	-	+	-
	513	-	-	-	514	-	-	-
	515	+	+	+	516	-	-	-
	517	-	-	-	518	-	-	-
	522	+	+	-	519	-	+	-
	524	+	+	+	521	-	-	-
	526	+	+	-	523	-	+	+
	527	-	-	-	525	-	-	-
	528	+	+	+	529	-	-	-
	531	-	+	-	530	-	-	-
	533	-	+	-	532	-	-	-
	538	+	-	-	536	-	-	-
	545	+	+	-	539	-	-	-
	546	-	-	-	540	-	-	-
	547	+	+	+	542	-	-	-
	548	+	-	+	543	-	+	-
	550	-	-	-	544	-	-	-
	552	+	+	-	549	-	+	-
	558	-	-	-	551	-	-	-
	559	+	+	+	553	-	-	-
	560	+	+	+	554	-	-	-
					555	-	-	-
					556	-	-	-
USDA Approval Date	June 6, 2014							

Study Type	Efficacy															
Pertaining to	Circovirus, porcine, type 2 (PCV2)															
Study Purpose	Duration of Immunity against Porcine Circovirus Type 2b															
Product Administration	One dose, administered intramuscularly.															
Study Animals	PCV2 serologically negative, commercial pigs, 3 weeks of age at administration; 25 vaccinates and 25 controls.															
Challenge Description	Porcine Circovirus Type 2b, 23 weeks after vaccination															
Interval observed after challenge	PCV2 viremia and PCV2 Fecal Shedding were evaluated twice weekly post challenge for 3 weeks. Lymphoid tissues were evaluated 3 weeks post challenge for the presence of PCV2 in lymphoid tissues and characteristic lesions attributable to PCV2 disease (lymphoid depletion and histiocytic replacement).															
Results	<table border="1"> <thead> <tr> <th>Variable</th> <th>Vaccinates Ever Positive*</th> <th>Controls Ever Positive*</th> </tr> </thead> <tbody> <tr> <td>Viremia (PCV2 detection in serum)</td> <td>4/25 (16%)</td> <td>13/25 (52%)</td> </tr> <tr> <td>Fecal Shedding (PCV2 detection in feces)</td> <td>3/25 (12%)</td> <td>20/25 (80%)</td> </tr> <tr> <td>Lymphoid Depletion (PCV2 lesions)</td> <td>3/25 (12%)</td> <td>11/25 (44%)</td> </tr> <tr> <td>Lymphoid Colonization (PCV2 in lymphoid tissues)</td> <td>0/25 (0%)</td> <td>6/25 (24%)</td> </tr> </tbody> </table> <p>*A pig was considered positive for PCV2 detection in serum or feces if it had a positive result at any sampling point.</p> <p>Raw data to follow:</p>	Variable	Vaccinates Ever Positive*	Controls Ever Positive*	Viremia (PCV2 detection in serum)	4/25 (16%)	13/25 (52%)	Fecal Shedding (PCV2 detection in feces)	3/25 (12%)	20/25 (80%)	Lymphoid Depletion (PCV2 lesions)	3/25 (12%)	11/25 (44%)	Lymphoid Colonization (PCV2 in lymphoid tissues)	0/25 (0%)	6/25 (24%)
Variable	Vaccinates Ever Positive*	Controls Ever Positive*														
Viremia (PCV2 detection in serum)	4/25 (16%)	13/25 (52%)														
Fecal Shedding (PCV2 detection in feces)	3/25 (12%)	20/25 (80%)														
Lymphoid Depletion (PCV2 lesions)	3/25 (12%)	11/25 (44%)														
Lymphoid Colonization (PCV2 in lymphoid tissues)	0/25 (0%)	6/25 (24%)														
USDA Approval Date	04/12/17															

Control ID					Vaccinate ID				
	viremia	fecal shedding	lymphoid colonization	lymphoid depletion		viremia	fecal shedding	lymphoid colonization	lymphoid depletion
622	+	+	-	-	624	+	-	-	-
617	+	+	-	-	618	-	-	-	-
623	-	+	-	-	620	-	-	-	-
664	+	+	+	+	625	-	-	-	+
668	+	+	-	+	665	-	+	-	-
986	-	+	-	-	670	-	-	-	-
610	-	+	-	+	675	-	-	-	-
611	-	-	-	-	995	-	+	-	-
612	-	-	-	-	606	+	-	-	-
615	+	+	-	+	607	-	-	-	-
616	-	+	-	+	613	-	-	-	+
990	-	-	-	+	614	-	-	-	-
994	-	-	-	-	991	-	-	-	-
673	+	+	+	+	992	+	-	-	-
674	+	+	+	+	667	-	-	-	-
741	-	-	-	-	669	-	-	-	-
979	-	+	-	-	672	-	-	-	-
981	+	+	-	-	977	-	-	-	-
982	+	+	-	-	978	-	-	-	-
985	-	+	-	-	983	-	-	-	+
744	-	+	-	-	742	-	-	-	-
746	+	+	+	+	743	-	-	-	-
748	+	+	+	+	745	-	-	-	-
750	+	+	+	+	747	-	+	-	-
976	+	+	-	-	749	+	-	-	-

Study Type	Efficacy
Pertaining to	Circovirus, porcine, type 2 (PCV2)
Study Purpose	23-week Duration of immunity against Porcine Circovirus 2.
Product Administration	One dose, administered intramuscularly at 3 weeks of age or two doses, administered intramuscularly 21 days apart at 3 days of age
Study Animals	Pigs
Challenge Description	
Interval observed after challenge	
Results	Study data are not available.

Study Type	Safety																																	
Pertaining to	ALL																																	
Study Purpose	To demonstrate safety under field conditions.																																	
Product Administration	Two doses, administered intramuscularly 2 weeks apart.																																	
Study Animals	Commercial pigs at three different geographical locations, 3 weeks of age at first administration; 602 vaccinates and 148 controls were enrolled.																																	
Challenge Description	N/A																																	
Interval observed after challenge	Animals were observed daily as a group and individually post vaccination(s) until three weeks after the final vaccination.																																	
Results	<p>All 3 sites were combined for this summary.</p> <p>Number of Animals with Clinical Signs</p> <table border="1"> <thead> <tr> <th></th> <th>Abnormal γ</th> <th>Cough</th> <th>Depression</th> <th>Diarrhea</th> <th>Thin</th> <th>Other **</th> <th>Death ***</th> </tr> </thead> <tbody> <tr> <td>Control</td> <td>14 (10%)</td> <td>0 (0%)</td> <td>2 (1%)</td> <td>4 (3%)</td> <td>7 (5%)</td> <td>6 (4%)</td> <td>0 (0%)</td> </tr> <tr> <td>Vaccinate</td> <td>48 (8%)</td> <td>0 (0%)</td> <td>12 (2%)</td> <td>17 (3%)</td> <td>24 (4%)</td> <td>14 (2%)</td> <td>8 (1%)</td> </tr> </tbody> </table> <p>^γ If animals were marked “abnormal;” specific clinical observations were marked or described as “other.” **Rash, lame, greasy pig, unable to stand were classified as “other.” ***Death affirmed by licensee to have cause other than vaccination. Percent was rounded to the nearest whole number.</p> <p>Number of Animals with Injection Site Reactions</p> <table border="1"> <thead> <tr> <th></th> <th>1st Vaccination</th> <th>2nd Vaccination</th> </tr> </thead> <tbody> <tr> <td>Control</td> <td>0 (0%)</td> <td>0 (0%)</td> </tr> <tr> <td>Vaccinate</td> <td>2 (1%)</td> <td>10 (2%)</td> </tr> </tbody> </table> <p>Percent was rounded to the nearest whole number. Reactions resolved within 3 days.</p>		Abnormal γ	Cough	Depression	Diarrhea	Thin	Other **	Death ***	Control	14 (10%)	0 (0%)	2 (1%)	4 (3%)	7 (5%)	6 (4%)	0 (0%)	Vaccinate	48 (8%)	0 (0%)	12 (2%)	17 (3%)	24 (4%)	14 (2%)	8 (1%)		1st Vaccination	2nd Vaccination	Control	0 (0%)	0 (0%)	Vaccinate	2 (1%)	10 (2%)
	Abnormal γ	Cough	Depression	Diarrhea	Thin	Other **	Death ***																											
Control	14 (10%)	0 (0%)	2 (1%)	4 (3%)	7 (5%)	6 (4%)	0 (0%)																											
Vaccinate	48 (8%)	0 (0%)	12 (2%)	17 (3%)	24 (4%)	14 (2%)	8 (1%)																											
	1st Vaccination	2nd Vaccination																																
Control	0 (0%)	0 (0%)																																
Vaccinate	2 (1%)	10 (2%)																																
USDA Approval Date	September 27, 2013																																	

Study Type	Safety																														
Pertaining to	ALL																														
Study Purpose	To demonstrate safety under field conditions.																														
Product Administration	One dose administered intramuscularly.																														
Study Animals	Commercial pigs at three different geographical locations, 3 weeks of age at administration; 158 controls and 632 vaccinates were enrolled.																														
Challenge Description	N/A																														
Interval observed after challenge	Animals were observed daily as a group and individually post vaccination(s) until three weeks after the final vaccination.																														
Results	<p>All 3 sites were combined for this summary.</p> <p>Number of Animals with Clinical Signs</p> <table border="1"> <thead> <tr> <th></th> <th>Abnormal^γ</th> <th>Anorexia</th> <th>Depression</th> <th>Diarrhea</th> <th>Dyspnea</th> <th>Other^{**}</th> <th>Death^{***}</th> </tr> </thead> <tbody> <tr> <td>Control</td> <td>20 (13%)</td> <td>8 (5%)</td> <td>3 (2%)</td> <td>6 (4%)</td> <td>2 (1%)</td> <td>9 (6%)</td> <td>0 (0%)</td> </tr> <tr> <td>Vaccinate</td> <td>108 (17%)</td> <td>29 (5%)</td> <td>13 (2%)</td> <td>17 (3%)</td> <td>3 (1%)</td> <td>81 (13%)</td> <td>2 (0%)</td> </tr> </tbody> </table> <p>^γ If animals were marked "abnormal;" specific clinical observations were marked or described as "other." ^{**}Cough, thin, lameness, lethargy, rough coat, jowl abscess were classified as "other." ^{***}Death affirmed by licensee to have cause other than vaccination. Percent was rounded to the nearest whole number.</p> <p>Number of Animals with Injection Site Reactions</p> <table border="1"> <thead> <tr> <th></th> <th>Animals Ever with Reaction</th> </tr> </thead> <tbody> <tr> <td>Control</td> <td>0 (0%)</td> </tr> <tr> <td>Vaccinate</td> <td>7 (1%)</td> </tr> </tbody> </table> <p>Percent was rounded to the nearest whole number. Reactions resolved within 3 days.</p>		Abnormal ^γ	Anorexia	Depression	Diarrhea	Dyspnea	Other ^{**}	Death ^{***}	Control	20 (13%)	8 (5%)	3 (2%)	6 (4%)	2 (1%)	9 (6%)	0 (0%)	Vaccinate	108 (17%)	29 (5%)	13 (2%)	17 (3%)	3 (1%)	81 (13%)	2 (0%)		Animals Ever with Reaction	Control	0 (0%)	Vaccinate	7 (1%)
	Abnormal ^γ	Anorexia	Depression	Diarrhea	Dyspnea	Other ^{**}	Death ^{***}																								
Control	20 (13%)	8 (5%)	3 (2%)	6 (4%)	2 (1%)	9 (6%)	0 (0%)																								
Vaccinate	108 (17%)	29 (5%)	13 (2%)	17 (3%)	3 (1%)	81 (13%)	2 (0%)																								
	Animals Ever with Reaction																														
Control	0 (0%)																														
Vaccinate	7 (1%)																														
USDA Approval Date	June 14, 2013																														

Study Type	Safety																											
Pertaining to	ALL																											
Study Purpose	To demonstrate safety under field conditions.																											
Product Administration	One dose administered intramuscularly at 3 weeks of age or two doses administered intramuscularly 3 weeks apart with first dose at 3 days of age.																											
Study Animals	Commercial pigs at three different geographical locations, a total of 600 vaccinates (200 receiving one-dose and 400 receiving two-doses) and 150 non-treated controls (50 per site) were enrolled.																											
Challenge Description	N/A																											
Interval observed after challenge	Animals were observed daily as a group and individually post vaccination(s) until three weeks after the final vaccination.																											
Results	<p>Site A – One dose administration Sites B & C –Two dose administration</p> <p>Injection Site Reactions Ever Present (# of Animals (% of Animals))</p> <table border="1"> <thead> <tr> <th rowspan="2"></th> <th colspan="2">Site A</th> <th colspan="2">Site B ⁺</th> <th colspan="2">Site C</th> </tr> <tr> <th>1st</th> <th>2nd</th> <th>1st</th> <th>2nd</th> <th>1st</th> <th>2nd</th> </tr> </thead> <tbody> <tr> <td>Non-Treated Control</td> <td>0</td> <td>n/a</td> <td>2 (4%)</td> <td>8 (17%)</td> <td>0</td> <td>0</td> </tr> <tr> <td>Vaccinate</td> <td>0</td> <td>n/a</td> <td>11 (6%)</td> <td>38 (20%)</td> <td>0</td> <td>2 (1%)</td> </tr> </tbody> </table> <p>Injection site reactions ranged from 0.5 to 5.0 cm. Reactions resolved within 7 days of first vaccination and within 14 days of second vaccination. Percent was rounded to the nearest whole number. n/a = Not Applicable ⁺ All site B animals were dosed with an autogenous vaccine in the same targeted vaccination site prior to the 2nd dose. Some animals (non-treated controls and vaccinates) had injection site reactions at the site of vaccination prior to the 2nd dose. Therefore it is not clear if the injection site reactions were attributable to the test article at site B.</p>		Site A		Site B ⁺		Site C		1 st	2 nd	1 st	2 nd	1 st	2 nd	Non-Treated Control	0	n/a	2 (4%)	8 (17%)	0	0	Vaccinate	0	n/a	11 (6%)	38 (20%)	0	2 (1%)
	Site A		Site B ⁺		Site C																							
	1 st	2 nd	1 st	2 nd	1 st	2 nd																						
Non-Treated Control	0	n/a	2 (4%)	8 (17%)	0	0																						
Vaccinate	0	n/a	11 (6%)	38 (20%)	0	2 (1%)																						

	Observations Ever Present (# of Animals (% of Animals))					
	Site A		Site B		Site C	
	Control	Vaccinate	Control	Vaccinate	Control	Vaccinate
None	38 (76)	166 (83)	40 (80)	144 (72)	37 (74)	146 (73)
Cough	2 (4)	4 (2)	1 (2)	4 (2)	.	.
Incoordination	3 (6)	6 (3)
Increased respiratory rate	8 (16)	16 (8)	.	.	.	2 (1)
Joint swelling	5 (10)	4 (2)	.	.	1 (2)	1 (1)
Lameness	5 (10)	16 (8)	1 (2)	3 (2)	1 (2)	3 (2)
Lethargy	3 (6)	7 (4)	1 (2)	6 (3)	1 (2)	8 (4)
Not eating	2 (4)	1 (1)
Swollen limb	2 (4)	1 (1)
Unthrifty	2 (4)	1 (1)	5 (10)	38 (19)	1 (2)	12 (6)
Central nervous system disorder	.	2 (1)
Depression	.	1 (1)
Other ear disorder	.	2 (1)
Abdominal cavity hernia	.	.	1 (2)	.	.	2 (1)
Abrasion	.	.	1 (2)	.	.	1 (1)
Diarrhea	.	.	3 (6)	17 (9)	6 (12)	22 (11)
Scar	.	.	1 (2)	6 (3)	.	.
Sore Eye	.	.	1 (2)	.	.	.
Claw / hoof / nail disorder	.	.	.	1 (1)	.	.
Dermatitis	1 (2)	1 (1)
Head tilt - neurological disorder	2 (4)	.
Infectious disease	1 (2)	3 (2)
Scour	7 (14)	25 (13)
Fracture	1 (1)
Paraplegia	1 (1)
Weakness	1 (1)
Found Dead*	1 (2)	2 (1)	.	.	1 (2)	11 (6)
Death by Euthanasia*	.	.	2 (4)	9 (5)	.	2 (1)
Uncoded Sign	1 (1)
All observations were VeDDRA coded (Veterinary Dictionary for Drug Related Affairs). Percent was rounded to the nearest whole number. * All deaths were affirmed by Investigators to NOT be attributable to the product.						
USDA Approval Date	November 1, 2017					

Study Type	Safety																																																						
Pertaining to	ALL																																																						
Study Purpose	To demonstrate safety under field conditions.																																																						
Product Administration	One dose administered intramuscularly.																																																						
Study Animals	Commercial sows and gilts at two different geographical locations, in each of three gestational stages (23-34 days, 61-75 days and 79-89 days); 60 controls and 299 vaccinates were enrolled.																																																						
Challenge Description	Not applicable																																																						
Interval observed after vaccination	No challenge. Animals were observed approximately four hours post-vaccination, one day post-vaccination and daily from 2 to 14 days post-vaccination for any health observation, and for any veterinary interventions from 15 days post-vaccination until 7 days post-farrowing. Injection sites were observed on day 0 and day 1. Reproductive performance through farrowing was evaluated. Piglets were observed for 7 days post-farrowing.																																																						
Results	<table border="1"> <thead> <tr> <th></th> <th>Controls (%)</th> <th>Vaccinates (%)</th> </tr> </thead> <tbody> <tr> <td>Total Number</td> <td>60 (100)</td> <td>299 (100)</td> </tr> <tr> <td>Normal</td> <td>47 (78.3)</td> <td>268 (89.6)</td> </tr> <tr> <td>Cough</td> <td>1 (1.7)</td> <td>0</td> </tr> <tr> <td>Decreased Appetite</td> <td>2 (3.3)</td> <td>13 (4.3)</td> </tr> <tr> <td>Depression</td> <td>6 (10)</td> <td>17 (5.7)</td> </tr> <tr> <td>Joint stiffness***</td> <td>2 (3.3)</td> <td>4 (1.3)</td> </tr> <tr> <td>Joint Swelling***</td> <td>2 (3.3)</td> <td>3 (1)</td> </tr> <tr> <td>Lameness***</td> <td>1 (1.7)</td> <td>10 (3.3)</td> </tr> <tr> <td>Metritis</td> <td>3 (5)</td> <td>6 (2)</td> </tr> <tr> <td>Vaginal Hemorrhage</td> <td>0</td> <td>1 (0.3)</td> </tr> <tr> <td>Abortion</td> <td>1 (1.7)</td> <td>2 (0.7)</td> </tr> <tr> <td>Rectal Prolapse</td> <td>1 (1.7)</td> <td>2 (0.7)</td> </tr> <tr> <td>Dystocia</td> <td>1 (1.7)</td> <td>4 (1.3)</td> </tr> <tr> <td>Found dead**</td> <td>0</td> <td>3 (1)</td> </tr> <tr> <td>Vaginal Prolapse</td> <td>0</td> <td>2 (0.7)</td> </tr> <tr> <td>Mammary Gland Edema</td> <td>1 (1.7)</td> <td>0</td> </tr> <tr> <td>Mastitis NOS</td> <td>1 (1.7)</td> <td>0</td> </tr> </tbody> </table> <p>Adverse Health Events* * Pigs observed as abnormal may have exhibited more than one clinical sign. ** The Investigator affirmed that deaths were unrelated to product use. *** Nine of these events were observed at one site following housing changes on day -1.</p>		Controls (%)	Vaccinates (%)	Total Number	60 (100)	299 (100)	Normal	47 (78.3)	268 (89.6)	Cough	1 (1.7)	0	Decreased Appetite	2 (3.3)	13 (4.3)	Depression	6 (10)	17 (5.7)	Joint stiffness***	2 (3.3)	4 (1.3)	Joint Swelling***	2 (3.3)	3 (1)	Lameness***	1 (1.7)	10 (3.3)	Metritis	3 (5)	6 (2)	Vaginal Hemorrhage	0	1 (0.3)	Abortion	1 (1.7)	2 (0.7)	Rectal Prolapse	1 (1.7)	2 (0.7)	Dystocia	1 (1.7)	4 (1.3)	Found dead**	0	3 (1)	Vaginal Prolapse	0	2 (0.7)	Mammary Gland Edema	1 (1.7)	0	Mastitis NOS	1 (1.7)	0
	Controls (%)	Vaccinates (%)																																																					
Total Number	60 (100)	299 (100)																																																					
Normal	47 (78.3)	268 (89.6)																																																					
Cough	1 (1.7)	0																																																					
Decreased Appetite	2 (3.3)	13 (4.3)																																																					
Depression	6 (10)	17 (5.7)																																																					
Joint stiffness***	2 (3.3)	4 (1.3)																																																					
Joint Swelling***	2 (3.3)	3 (1)																																																					
Lameness***	1 (1.7)	10 (3.3)																																																					
Metritis	3 (5)	6 (2)																																																					
Vaginal Hemorrhage	0	1 (0.3)																																																					
Abortion	1 (1.7)	2 (0.7)																																																					
Rectal Prolapse	1 (1.7)	2 (0.7)																																																					
Dystocia	1 (1.7)	4 (1.3)																																																					
Found dead**	0	3 (1)																																																					
Vaginal Prolapse	0	2 (0.7)																																																					
Mammary Gland Edema	1 (1.7)	0																																																					
Mastitis NOS	1 (1.7)	0																																																					

Injection Site Reactions

Injection sites were observed on Day 0 and 1. Since there were no injection site reactions observed on Day 1, further observations were not necessary.

	Score 0 (normal)	Score 1 (≥ 0.5 cm and ≤ 1.5 cm)	Score 2 (≥ 1.6 cm and ≤ 3 cm)	Score 3 (> 3 cm)
Controls	60	0	0	0
Vaccinates	299	0	0	0

Reproductive Outcomes -Sow Outcome

	Controls (%)	Vaccinates (%)
Total Number	60 (100)	299 (100)
Farrowed	57 (95)	286 (95.7)
Abortion*+	1 (1.7)	2 (0.7)
Not Pregnant*	2 (3.3)	8 (2.7)
Culled/ Dead+	0	3 (1)

* Events occurred at one site

+Same as reported on the Adverse Events table

Farrowing Outcomes

	Controls (%)	Vaccinates (%)
Total Number of Piglets	876	4557
Born Normal	698 (80)	3740 (82)
Low viability	38 (4)	191 (4)
Stillborn	123 (14)	510 (11)
Mummies	17 (2)	116 (3)

Litter Health Observations

Five number summary of percent of litter that is Normal (not mummified, stillborn, or low viability)

Group	Min	Q1	Median	Q3	Max
Vaccinates	26.3	75.0	84.4	93.3	100.0
Controls	0.0	71.4	83.3	88.2	100.0

USDA Approval Date

February 10, 2022