



Summary of Studies Supporting USDA Product Licensure

Establishment Name	Biomune Company
USDA Vet Biologics Establishment Number	368
Product Code	1A91.R0
True Name	Bursal Disease-Marek's Disease-Newcastle Disease Vaccine, Serotypes 2 & 3, Live Virus, Live Marek's Disease Vector
Tradename(s) / Distributor or Subsidiary (if different from manufacturer)	Ultifend IBD ND + SB1 - no distributor specified
Date of Compilation Summary	March 26, 2018

Disclaimer: Do not use the following studies to compare one product to another. Slight differences in study design and execution can render the comparisons meaningless.

Study Type	Efficacy
Pertaining to	Infectious Bursal Disease Virus (IBDV)
Study Purpose	To demonstrate effectiveness against IBDV
Product Administration	One dose administered by the <i>in ovo</i> route
Study Animals	<ol style="list-style-type: none"> 1. 30 SPF chicken embryos per treatment group vaccinated at 18 days of incubation with product (<i>in ovo</i> vaccinate) 2. 30 SPF chickens per treatment group vaccinated at at 18 days of incubation with placebo-matched vaccine (positive control) 3. 30 SPF chickens per treatment group vaccinated at at 18 days of incubation with placebo-matched vaccine (negative control)
Challenge Description	IBDV USDA Standard strain at five weeks of age for all except negative control group
Interval observed after challenge	Daily observation for 4 days post challenge; Tissues examined at 4 days post challenge
Results	<p>A chicken was considered affected by the challenge (positive) if grossly observable lesions caused by the IBDV USDA Standard challenge were present.</p> <p>3/30 (10%) vaccinates, 30/30 (100%) positive controls and 0/30 (0%) negative controls were affected by the challenge.</p> <p>Raw data are shown on the attached page.</p>
USDA Approval Date	March 9, 2016

Vaccinate ID	Infectious Bursal Disease Lesions ¹	Positive Control ID	Infectious Bursal Disease Lesions ¹	Negative Control ID	Infectious Bursal Disease Lesions ¹
1	NA	31	P,A,Y	61	NA
2	NA	32	P,Y	62	NA
3	NA	33	E	63	NA
4	NA	34	P,A,Y	64	NA
5	NA	35	P,A,Y	65	NA
6	NA	36	A,E	66	NA
7	NA	37	P,E	67	NA
8	NA	38	P,A,Y	68	NA
9	NA	39	Y	69	NA
10	NA	40	Y	70	NA
11	NA	41	A,Y,E	71	NA
12	NA	42	P,A	72	NA
13	NA	43	P,A	73	NA
14	NA	44	P,Y	74	NA
15	NA	45	A,Y,E	75	NA
16	NA	46	A,E	76	NA
17	NA	47	A,Y,E	77	NA
18	NA	48	A,Y,E	78	NA
19	NA	49	E	79	NA
20	NA	50	P,A,Y	80	NA
21	A	51	A,Y,E	81	NA
22	NA	52	Y,E	82	NA
23	NA	53	Y	83	NA
24	A,M	54	A,E	84	NA
25	NA	55	P,A,Y	85	NA
26	NA	56	A,Y,E	86	NA
27	NA	57	P,A	87	NA
28	A	58	A	88	NA
29	NA	59	A,E	89	NA
30	NA	60	A,Y,E	90	NA

¹ Gross lesion: A=atrophy, Y=yellowish color, E=edema, P=peribursal edema, M=macroscopic hemorrhage, NA=not applicable (no lesions)

Study Type	Efficacy
Pertaining to	Infectious Bursal Disease Virus (IBDV)
Study Purpose	To demonstrate effectiveness against IBDV in chickens with maternal antibodies to IBDV
Product Administration	<ol style="list-style-type: none"> 1. One dose administered by the <i>in ovo</i> route 2. One dose administered by the subcutaneous route
Study Animals	<ol style="list-style-type: none"> 1. 30 SPF chicken embryos per treatment group vaccinated at 18 days of incubation with product (<i>in ovo</i> vaccinate) 2. 30 SPF chicken embryos per treatment group vaccinated at day of age with product (SQ vaccinate) 3. 30 SPF chickens per treatment group vaccinated at day of age with placebo-matched vaccine (positive control) 4. 30 SPF chickens per treatment group vaccinated at day of age of incubation with placebo-matched vaccine (negative control)
Challenge Description	IBDV USDA Standard strain at nine weeks and four days of age except negative control group
Interval observed after challenge	Daily observation for 4 days post challenge; Tissues examined at 4 days post challenge
Results	<p>A chicken was considered affected by the challenge (positive) if grossly observable lesions caused by the IBDV USDA Standard challenge were present.</p> <p>In ovo vaccination: 0/30 (0%) vaccinates and 29/30 (97%) controls were affected by the challenge.</p> <p>SQ vaccination: 0/30 (0%) vaccinates and 29/30 (97%) controls were affected by the challenge.</p> <p>0/30 (0%) negative controls were affected by the challenge.</p> <p>Raw data are shown on the attached page.</p>
USDA Approval Date	May 2, 2016

In ovo Vaccinate ID	Infectious Bursal Disease Lesions ¹	SQ Vaccinate ID	Infectious Bursal Disease Lesions ¹	In ovo Positive Control ID	Infectious Bursal Disease Lesions ¹	SQ Positive Control ID	Infectious Bursal Disease Lesions ¹	Negative Control ID	Infectious Bursal Disease Lesions ¹
1	NA	31	NA	61	E	91	P,E	121	NA
2	NA	32	NA	62	P,Y,E	92	P,Y	122	NA
3	NA	33	NA	63	P,A,Y	93	P,A,E	123	NA
4	NA	34	NA	64	E	94	NA	124	NA
5	NA	35	NA	65	P,Y,E	95	P,A,Y	125	NA
6	NA	36	NA	66	P,Y	96	E	126	NA
7	NA	37	NA	67	P,Y	97	P,Y	127	NA
8	NA	38	NA	68	P,Y,E	98	P,Y,E	128	NA
9	NA	39	NA	69	P,Y	99	P,Y	129	NA
10	NA	40	NA	70	P,A,Y	100	P,Y	130	NA
11	NA	41	NA	71	E	101	P,Y	131	NA
12	NA	42	NA	72	NA	102	P	132	NA
13	NA	43	NA	73	A,Y	103	A,Y	133	NA
14	NA	44	NA	74	P,Y	104	P,Y	134	NA
15	NA	45	NA	75	E	105	P,Y	135	NA
16	NA	46	NA	76	P	106	P,Y	136	NA
17	NA	47	NA	77	P,Y	107	P,Y	137	NA
18	NA	48	NA	78	Y,E	108	P,Y	138	NA
19	NA	49	NA	79	P,Y	109	P,Y	139	NA
20	NA	50	NA	80	P,Y	110	P,A	140	NA
21	NA	51	NA	81	P	111	P,A,Y	141	NA
22	NA	52	NA	82	A,E	112	P,Y	142	NA
23	NA	53	NA	83	P	113	P,A	143	NA
24	NA	54	NA	84	P,A,Y,E	114	P	144	NA
25	NA	55	NA	85	P,A	115	P,E	145	NA
26	NA	56	NA	86	P,E	116	P,A	146	NA
27	NA	57	NA	87	Y	117	P,Y	147	NA
28	NA	58	NA	88	P,A	118	P,Y	148	NA
29	NA	59	NA	89	P,Y	119	P,E,Y	149	NA
30	NA	60	NA	90	P,Y	120	P	150	NA

¹ Gross lesion: A=atrophy, Y=yellowish color, E=edema, P=peribursal edema, M=macroscopic hemorrhage, NA=not applicable (no lesions)

Study Type	Efficacy
Pertaining to	Bursal Disease Virus (IBDV)
Study Purpose	To demonstrate effectiveness IBDV
Product Administration	One dose administered by the subcutaneous route (SQ)
Study Animals	<ol style="list-style-type: none"> 1. 30 SPF chicken embryos per treatment group vaccinated at day of age with product (SQ vaccinate) 2. 30 SPF chickens per treatment group vaccinated at 18 days of incubation with placebo-matched vaccine (positive control) 3. 20 SPF chickens per treatment group vaccinated at day of age of incubation with placebo-matched vaccine (negative control)
Challenge Description	IBDV USDA Standard at five weeks of age except the negative control group
Interval observed after challenge	Daily observation for four days post challenge; Tissues examined at four days post challenge
Results	<p>A chicken was considered affected by the challenge (positive) if gross lesions of Infectious Bursal Disease were present.</p> <p>1/30 (3%) vaccinates, 29/30 (97%) positive controls and 0/20 (0%) negative controls were affected by the challenge.</p> <p>Raw data are shown on the attached pages.</p>
USDA Approval Date	January 28, 2016

Vaccinate ID	Lesions of Infectious Bursal Disease ¹	Positive Control ID	Lesions of Infectious Bursal Disease ¹	Negative Control ID	Lesions of Infectious Bursal Disease ¹
61	NA	91	A	121	NA
62	A	92	A,P	122	NA
63	NA	93	A,P	123	NA
64	NA	94	A,E	124	NA
65	NA	95	Y,P	125	NA
66	NA	96	A,E	126	NA
67	NA	97	Y,P,E	127	NA
68	NA	98	A	128	NA
69	NA	99	A,M	129	NA
70	NA	100	A	130	NA
71	NA	101	A	131	NA
72	NA	102	A	132	NA
73	NA	103	A,E	133	NA
74	NA	104	P,Y,E	134	NA
75	NA	105	A,E	135	NA
76	NA	106	A,P	136	NA
77	NA	107	A,P	137	NA
78	NA	108	A,M	138	NA
79	NA	109	A	139	NA
80	NA	110	A,P	140	NA
81	NA	111	A,M,E		
82	NA	112	P,Y		
83	NA	113	P,M,E		
84	NA	114	NA		
85	NA	115	A,E		
86	NA	116	A,M		
87	NA	117	E		
88	NA	118	E		
89	NA	119	A,M		
90	NA	120	A,M,E		

¹ Gross lesion: A=atrophy, Y=yellowish color, E=edema, P=peribursal edema, M=macroscopic hemorrhage, NA=not applicable (no lesions)

Study Type	Efficacy
Pertaining to	Marek's Disease Virus (MDV)
Study Purpose	To demonstrate effectiveness against MDV
Product Administration	<ol style="list-style-type: none"> 1. One dose administered by the <i>in ovo</i> route 2. One dose administered by the subcutaneous route
Study Animals	<ol style="list-style-type: none"> 1. 45 SPF chicken embryos per treatment group vaccinated at 18 days of incubation with product (<i>in ovo</i> vaccinate) 2. 45 SPF chickens per treatment group vaccinated at day of age with product (SQ vaccinate) 3. 43 SPF chickens per treatment group vaccinated at day of age with Marek's Serotype 3 vaccine (Marek's Serotype 3 control) 4. 45 SPF chickens per treatment group vaccinated at at 18 days of incubation with placebo-matched vaccine (positive control) 5. 45 SPF chickens per treatment group vaccinated at at day of age with placebo-matched vaccine (negative control)
Challenge Description	MDV RB1/B at five days of age except negative control group
Interval observed after challenge	Daily observation for 44 days post challenge; Tissues examined at 44 days post challenge
Results	<p>A chicken was considered affected by the challenge (positive) if grossly observable lesions caused by the MDV RB1/B challenge were present.</p> <p>In ovo vaccination: 5/45 (11%) vaccinates were affected by the challenge.</p> <p>SQ vaccination: 7/45 (16%) vaccinates were affected by the challenge.</p> <p>Controls: 10/43 (23%) Marek's Serotype 3 controls were affected by the challenge. 42/45 (93%) positive controls were affected by the challenge. 0/45 (0%) negative controls were affected by the challenge.</p> <p>Raw data are shown on the attached page.</p>
USDA Approval Date	October 18, 2016

SQ Vaccinate ID	Marek's Lesions ¹	In ovo Vaccinate ID	Marek's Lesions ¹	Marek's Serotype 3 Control ID	Marek's Lesions ¹	Positive Control ID	Marek's Lesions ¹	Negative Control ID	Marek's Lesions ¹
1	NA	46	NA	91	NA	134	H,Sp	179	NA
2	L,K,G	47	NA	92	NA	135	NA	180	NA
3	NA	48	NA	93	H,L	136	H	181	NA
4	L,Sp,K,G	49	NA	94	K	137	H	182	NA
5	NA	50	NA	95	NA	138	H	183	NA
6	NA	51	NA	96	NA	139	H,G	184	NA
7	NA	52	NA	97	NA	140	L,G,K,H,M	185	NA
8	NA	53	H,G,K	98	NA	141	H,K	186	NA
9	NA	54	NA	99	NA	142	H	187	NA
10	NA	55	NA	100	NA	143	Sp,K	188	NA
11	G,K	56	NA	101	NA	144	H, Sp	189	NA
12	NA	57	NA	102	NA	145	L,H,K	190	NA
13	NA	58	H,L,Sp	103	NA	146	H	191	NA
14	NA	59	NA	104	NA	147	H,K,G	192	NA
15	NA	60	NA	105	NA	148	H,Sp	193	NA
16	NA	61	NA	106	NA	149	H,L,Sp,K,G	194	NA
17	NA	62	NA	107	NA	150	H,L	195	NA
18	L,Sp,K,G	63	NA	108	Sp,G,I	151	H,Sp	196	NA
19	NA	64	NA	109	NA	152	H,K	197	NA
20	NA	65	NA	110	NA	153	L,G,K,H	198	NA
21	NA	66	NA	111	NA	154	H,K	199	NA
22	NA	67	NA	112	NA	155	NA	200	NA
23	K	68	NA	113	NA	156	H	201	NA
24	NA	69	NA	114	L	157	H,L,Sp,K,G	202	NA
25	NA	70	NA	115	NA	158	H,L,Sp,K,G	203	NA
26	NA	71	H,L,G	116	NA	159	H,L,Sp,K,G	204	NA
27	K	72	NA	117	L,Sp,G,K	160	K,H	205	NA
28	NA	73	NA	118	H,K,Sp,M	161	H,M	206	NA
29	NA	74	NA	119	NA	162	H,K	207	NA
30	NA	75	NA	120	NA	163	H,Sp	208	NA
31	NA	76	NA	121	NA	164	H,G,K	209	NA
32	NA	77	NA	122	NA	165	H	210	NA
33	NA	78	NA	123	NA	166	H,Sp,K,G	211	NA
34	NA	79	NA	124	NA	167	H,L,Sp,K	212	NA
35	NA	80	NA	125	NA	168	H,Sp,K	213	NA
36	NA	81	L,K,G	126	NA	169	NA	214	NA
37	NA	82	NA	127	L,G,K,Sp	170	H,Sp	215	NA
38	NA	83	NA	128	H	171	H,K,Sp	216	NA
39	G,K,Sp	84	NA	129	NA	172	H,Sp	217	NA
40	NA	85	H,L,K	130	NA	173	H	218	NA
41	NA	86	NA	131	L,K,Sp,G	174	H,K	219	NA
42	NA	87	NA	132	NA	175	H,K	220	NA
43	NA	88	NA	133	K,M,I	176	H,K,G	221	NA
44	NA	89	NA			177	H,L,Sp,K	222	NA
45	NA	90	NA			178	H	223	NA

¹ Tissue with lesion: K=kidney, Sp=spleen, L=liver, H=heart, G=gonad, N=nerves, Sk=skin, E=eye, M=muscle, I=intestines, NA=not applicable (no lesions)

Study Type	Efficacy
Pertaining to	Newcastle Disease Virus (NDV)
Study Purpose	To demonstrate effectiveness against NDV
Product Administration	One dose administered by the <i>in ovo</i> route
Study Animals	<ol style="list-style-type: none"> 1. 30 SPF chicken embryos per treatment group vaccinated at 18 days of incubation with product (<i>in ovo</i> vaccinate) 2. 30 SPF chickens per treatment group vaccinated at at 18 days of incubation with placebo-matched vaccine (positive control) 3. 30 SPF chickens per treatment group vaccinated at at 18 days of incubation with placebo-matched vaccine (negative control)
Challenge Description	NDV Texas GB strain at 28 days of age except the negative control group
Interval observed after challenge	Daily observation for 14 days post challenge for clinical signs of NDV
Results	<p>A chicken was considered affected by the challenge (positive) if clinical signs of Newcastle disease caused by the NDV Texas GB challenge were present.</p> <p>3/30 (10%) vaccinates, 30/30 (100%) positive controls and 0/30 (0%) negative controls were affected by the challenge.</p> <p>Raw data are shown on the attached page.</p>
USDA Approval Date	February 26, 2016

Vaccinate ID	Clinical Signs of Newcastle Disease ¹	Positive Control ID	Clinical Signs of Newcastle Disease ¹	Negative Control ID	Clinical Signs of Newcastle Disease ¹
1	Neg	31	Pos	61	Neg
2	Neg	32	Pos	62	Neg
3	Neg	33	Pos	63	Neg
4	Neg	34	Pos	64	Neg
5	Neg	35	Pos	65	Neg
6	Neg	36	Pos	66	Neg
7	Pos	37	Pos	67	Neg
8	Pos	38	Pos	68	Neg
9	Neg	39	Pos	69	Neg
10	Neg	40	Pos	70	Neg
11	Neg	41	Pos	71	Neg
12	Neg	42	Pos	72	Neg
13	Neg	43	Pos	73	Neg
14	Neg	44	Pos	74	Neg
15	Neg	45	Pos	75	Neg
16	Neg	46	Pos	76	Neg
17	Neg	47	Pos	77	Neg
18	Neg	48	Pos	78	Neg
19	Pos	49	Pos	79	Neg
20	Neg	50	Pos	80	Neg
21	Neg	51	Pos	81	Neg
22	Neg	52	Pos	82	Neg
23	Neg	53	Pos	83	Neg
24	Neg	54	Pos	84	Neg
25	Neg	55	Pos	85	Neg
26	Neg	56	Pos	86	Neg
27	Neg	57	Pos	87	Neg
28	Neg	58	Pos	88	Neg
29	Neg	59	Pos	89	Neg
30	Neg	60	Pos	90	Neg

¹ Clinical Signs: Pos=death, Neg=negative for clinical signs, including death

Study Type	Efficacy
Pertaining to	Newcastle Disease Virus (NDV)
Study Purpose	To demonstrate effectiveness against NDV
Product Administration	One dose administered by the subcutaneous route (SQ)
Study Animals	<ol style="list-style-type: none"> 1. 30 SPF chicken embryos per treatment group vaccinated at day of age with product (SQ vaccinate) 2. 30 SPF chickens per treatment group vaccinated at 18 days of incubation with placebo-matched vaccine (positive control) 3. 32 SPF chickens per treatment group vaccinated at day of age of incubation with placebo-matched vaccine (negative control)
Challenge Description	NDV Texas GB at four weeks of age except the negative control group
Interval observed after challenge	Daily observation for 14 days post challenge for clinical signs of NDV
Results	<p>A chicken was considered affected by the challenge (positive) if clinical signs of Newcastle Disease were present.</p> <p>0/30 (0%) vaccinates, 30/30 (0%) positive controls and 0/32 (0%) negative controls were affected by the challenge.</p> <p>Raw data are shown on the attached pages.</p>
USDA Approval Date	January 28, 2016

Vaccinate ID	Clinical Signs of Newcastle Disease ¹	Positive Control ID	Clinical Signs of Newcastle Disease ¹	Negative Control ID	Clinical Signs of Newcastle Disease ¹
1	Neg	31	Pos	121	Neg
2	Neg	32	Pos	122	Neg
3	Neg	33	Pos	123	Neg
4	Neg	34	Pos	124	Neg
5	Neg	35	Pos	125	Neg
6	Neg	36	Pos	126	Neg
7	Neg	37	Pos	127	Neg
8	Neg	38	Pos	128	Neg
9	Neg	39	Pos	129	Neg
10	Neg	40	Pos	130	Neg
11	Neg	41	Pos	131	Neg
12	Neg	42	Pos	132	Neg
13	Neg	43	Pos	133	Neg
14	Neg	44	Pos	134	Neg
15	Neg	45	Pos	135	Neg
16	Neg	46	Pos	136	Neg
17	Neg	47	Pos	137	Neg
18	Neg	48	Pos	138	Neg
19	Neg	49	Pos	139	Neg
20	Neg	50	Pos	140	Neg
21	Neg	51	Pos	141	Neg
22	Neg	52	Pos	142	Neg
23	Neg	53	Pos	143	Neg
24	Neg	54	Pos	144	Neg
25	Neg	55	Pos	145	Neg
26	Neg	56	Pos	146	Neg
27	Neg	57	Pos	147	Neg
28	Neg	58	Pos	148	Neg
29	Neg	59	Pos	149	Neg
30	Neg	60	Pos	150	Neg
				151	Neg
				152	Neg

¹ Clinical Signs: Pos=death, Neg=negative for clinical signs, including death

Study Type	Efficacy
Pertaining to	Newcastle Disease Virus (NDV) Texas GB
Study Purpose	To demonstrate effectiveness against NDV Texas GB strain
Product Administration	1. One dose administered by the subcutaneous route (SQ) 2. One dose administered by the <i>in ovo</i> route
Study Animals	1. 30 maternal antibody-positive chickens per treatment group vaccinated at day of age by the SQ route 2. 30 maternal antibody-positive chicken embryos per treatment group vaccinated at 18 days of incubation by the <i>in ovo</i> route 3. 30 maternal antibody-positive chickens non-vaccinated as Positive Controls 4. 30 maternal antibody-positive chickens non-vaccinated and non-challenged as Negative Controls
Challenge Description	NDV Texas GB strain at 7 days of age for Positive Control group and 55 days of age for SQ and <i>in ovo</i> vaccinate groups. No challenge was administered to the Negative Controls group.
Interval observed after challenge	Daily observation for 14 days post challenge for clinical signs of New Castle Disease Virus.
Results	A chicken was considered affected by the challenge (positive) if clinical signs of Newcastle disease caused by the NDV Texas GB challenge were present. <i>In ovo</i> vaccination: 0/30 vaccinates and 30/30 Positive Controls were affected by the challenge, NDV Texas GB. SQ vaccination: 0/30 vaccinates and 30/30 Positive Controls were affected by the challenge, NDV Texas GB. 0/30 Negative Controls were affected by the challenge. Raw data are shown on the attached page.
USDA Approval Date	April 29, 2016

In ovo Vaccinate ID	Clinical Signs of Newcastle Disease ¹	SQ ² Vaccinate ID	Clinical Signs of Newcastle Disease ¹	In ovo Positive Control ID	Clinical Signs of Newcastle Disease ¹	SQ ² Positive Control ID	Clinical Signs of Newcastle Disease ¹	Negative Control ID	Clinical Signs of Newcastle Disease ¹
1	Neg	31	Neg	61	Pos	91	Pos	121	Neg
2	Neg	32	Neg	62	Pos	92	Pos	122	Neg
3	Neg	33	Neg	63	Pos	93	Pos	123	Neg
4	Neg	34	Neg	64	Pos	94	Pos	124	Neg
5	Neg	35	Neg	65	Pos	95	Pos	125	Neg
6	Neg	36	Neg	66	Pos	96	Pos	126	Neg
7	Neg	37	Neg	67	Pos	97	Pos	127	Neg
8	Neg	38	Neg	68	Pos	98	Pos	128	Neg
9	Neg	39	Neg	69	Pos	99	Pos	129	Neg
10	Neg	40	Neg	70	Pos	100	Pos	130	Neg
11	Neg	41	Neg	71	Pos	101	Pos	131	Neg
12	Neg	42	Neg	72	Pos	102	Pos	132	Neg
13	Neg	43	Neg	73	Pos	103	Pos	133	Neg
14	Neg	44	Neg	74	Pos	104	Pos	134	Neg
15	Neg	45	Neg	75	Pos	105	Pos	135	Neg
16	Neg	46	Neg	76	Pos	106	Pos	136	Neg
17	Neg	47	Neg	77	Pos	107	Pos	137	Neg
18	Neg	48	Neg	78	Pos	108	Pos	138	Neg
19	Neg	49	Neg	79	Pos	109	Pos	139	Neg
20	Neg	50	Neg	80	Pos	110	Pos	140	Neg
21	Neg	51	Neg	81	Pos	111	Pos	141	Neg
22	Neg	52	Neg	82	Pos	112	Pos	142	Neg
23	Neg	53	Neg	83	Pos	113	Pos	143	Neg
24	Neg	54	Neg	84	Pos	114	Pos	144	Neg
25	Neg	55	Neg	85	Pos	115	Pos	145	Neg
26	Neg	56	Neg	86	Pos	116	Pos	146	Neg
27	Neg	57	Neg	87	Pos	117	Pos	147	Neg
28	Neg	58	Neg	88	Pos	118	Pos	148	Neg
29	Neg	59	Neg	89	Pos	119	Pos	149	Neg
30	Neg	60	Neg	90	Pos	120	Pos	150	Neg

¹ Clinical Signs: Pos=death, Neg=negative for clinical signs, including death

²SQ = subcutaneous vaccination

Study Type	Safety																																									
Pertaining to	ALL																																									
Study Purpose	Field Safety																																									
Product Administration	One dose administered via the <i>in ovo</i> route.																																									
Study Animals	Broiler chickens at 18 or 19 days of embryonation. Two independent study sites consisting of two groups. One group received the product vaccine and the control group received vaccinations according to standard practices.																																									
Challenge Description	Not applicable																																									
Interval observed after challenge	Animals were observed daily for mortality through 21 days after vaccination.																																									
Results	<table border="1"> <thead> <tr> <th>Location</th> <th>Treatment</th> <th>% Hatchability</th> <th>Total Chicks Placed</th> <th>21 Day Mortality</th> <th>% Mortality</th> <th>% Condemnation</th> </tr> </thead> <tbody> <tr> <td>1</td> <td>Product Vaccine</td> <td>88.09</td> <td>33,300</td> <td>578</td> <td>1.74</td> <td>0.034</td> </tr> <tr> <td>1</td> <td>Control</td> <td>84.05</td> <td>34,500</td> <td>406</td> <td>1.18</td> <td>0.033</td> </tr> <tr> <td>2</td> <td>Product Vaccine</td> <td>89.69</td> <td>28,700</td> <td>544</td> <td>1.90</td> <td>0.14</td> </tr> <tr> <td>2</td> <td>Control</td> <td>88.00</td> <td>28,700</td> <td>462</td> <td>1.61</td> <td>0.18</td> </tr> </tbody> </table> <p>No adverse reactions attributable to the vaccine were recorded.</p>							Location	Treatment	% Hatchability	Total Chicks Placed	21 Day Mortality	% Mortality	% Condemnation	1	Product Vaccine	88.09	33,300	578	1.74	0.034	1	Control	84.05	34,500	406	1.18	0.033	2	Product Vaccine	89.69	28,700	544	1.90	0.14	2	Control	88.00	28,700	462	1.61	0.18
Location	Treatment	% Hatchability	Total Chicks Placed	21 Day Mortality	% Mortality	% Condemnation																																				
1	Product Vaccine	88.09	33,300	578	1.74	0.034																																				
1	Control	84.05	34,500	406	1.18	0.033																																				
2	Product Vaccine	89.69	28,700	544	1.90	0.14																																				
2	Control	88.00	28,700	462	1.61	0.18																																				
USDA Approval Date	August 1, 2017																																									

Study Type	Safety																																								
Pertaining to	ALL																																								
Study Purpose	Demonstrate safety of product under typical use conditions.																																								
Product Administration	One dose administered via the subcutaneous route.																																								
Study Animals	Commercial chickens at day of age. Chickens were observed daily for 22 days after vaccination.																																								
Challenge Description	Not applicable																																								
Interval observed after challenge	Not applicable																																								
Results	<table border="1"> <thead> <tr> <th rowspan="2">Location</th> <th rowspan="2">Vaccine Serial No./Treatment Group</th> <th rowspan="2">No. of Chickens Vaccinated</th> <th rowspan="2">No. of Birds Placed</th> <th colspan="2">Mortality</th> <th rowspan="2">Observations</th> </tr> <tr> <th>Total No. of Deaths</th> <th>Percent</th> </tr> </thead> <tbody> <tr> <td rowspan="2">PA</td> <td>377-001</td> <td>20,000</td> <td>20,000</td> <td>208</td> <td>1.04%</td> <td>No adverse reactions</td> </tr> <tr> <td>control</td> <td>19,998</td> <td>19,998</td> <td>155</td> <td>0.77%</td> <td>No adverse reactions</td> </tr> <tr> <td rowspan="2">MD</td> <td>377-002</td> <td>86,500</td> <td>86,500</td> <td>534</td> <td>0.62%</td> <td>No adverse reactions</td> </tr> <tr> <td>control</td> <td>86,600</td> <td>86,600</td> <td>512</td> <td>0.59%</td> <td>No adverse reactions</td> </tr> </tbody> </table> <p>No adverse reactions attributable to the vaccine were recorded.</p>						Location	Vaccine Serial No./Treatment Group	No. of Chickens Vaccinated	No. of Birds Placed	Mortality		Observations	Total No. of Deaths	Percent	PA	377-001	20,000	20,000	208	1.04%	No adverse reactions	control	19,998	19,998	155	0.77%	No adverse reactions	MD	377-002	86,500	86,500	534	0.62%	No adverse reactions	control	86,600	86,600	512	0.59%	No adverse reactions
Location	Vaccine Serial No./Treatment Group	No. of Chickens Vaccinated	No. of Birds Placed	Mortality		Observations																																			
				Total No. of Deaths	Percent																																				
PA	377-001	20,000	20,000	208	1.04%	No adverse reactions																																			
	control	19,998	19,998	155	0.77%	No adverse reactions																																			
MD	377-002	86,500	86,500	534	0.62%	No adverse reactions																																			
	control	86,600	86,600	512	0.59%	No adverse reactions																																			
USDA Approval Date	October 4, 2017																																								