

**Supplemental Assay Method for the Detection of Extraneous Hemadsorbing Agents in Master Seed Virus**

**United States Department of Agriculture  
Center for Veterinary Biologics  
Testing Protocol**

**SAM 313**

**Supplemental Assay Method for the Detection of Extraneous Hemadsorbing  
Agents in Master Seed Virus**

Date: August 23, 2023  
Number: SAM 313.05  
Supersedes: SAM 313.04, November 13, 2014  
Standard Requirement: 9 CFR, parts 113.46 and 113.55  
Contact: Email: [Methodsrequest.notification@usda.gov](mailto:Methodsrequest.notification@usda.gov)  
Phone: Center for Veterinary Biologics, 515-337-6100

United States Department of Agriculture  
Animal and Plant Health Inspection Service  
P. O. Box 844  
Ames, IA 50010

Mention of trademark or proprietary product does not constitute a guarantee or warranty of the product by USDA and does not imply its approval to the exclusion of other products that may be suitable.

**UNCONTROLLED COPY**

**Supplemental Assay Method for the Detection of Extraneous Hemadsorbing Agents in Master Seed Virus**

**Table of Contents**

- 1. Introduction**
- 2. Materials**
  - 2.1 Equipment/instrumentation**
  - 2.2 Reagents/supplies**
- 3. Preparation for the Test**
  - 3.1 Personnel qualifications/training**
  - 3.2 Preparation of RBC**
  - 3.3 Preparation of the sample**
- 4. Performance of the Test**
  - 4.1 Vessels incubated at each temperature separately**
  - 4.2 Separate vessels incubated simultaneously at each temperature**
- 5. Interpretation of the Test Results**
- 6. Report of Test Results**
- 7. References**
- 8. Summary of Revisions**

**Appendices**

- I. Macroscopic view of hemadsorption**
- II. Microscopic view of hemadsorption**

**UNCONTROLLED COPY**

**Supplemental Assay Method for the Detection of Extraneous Hemadsorbing Agents in Master Seed Virus**

**1. Introduction**

This Supplemental Assay Method (SAM) describes a method for detection of extraneous hemadsorbing (HAd) agents in master seeds (MS) used in the production of veterinary vaccines. Not all viruses cause a noticeable change in the cellular appearance, but can be detected by HAd of red blood cells (RBC) to virus-coded glycoproteins incorporated in the cell membrane. To perform the test, cell monolayers are washed and covered with a suspension of guinea pig and chicken RBC. Following incubation, uninfected cells will appear normal, while RBCs will adhere to cell monolayers that contain HAd surface viral antigens. Detection is accomplished by both macroscopic and microscopic reading methods.

**2. Materials**

**2.1 Equipment/instrumentation**

Equivalent equipment or instrumentation may be substituted for any brand name listed below. Equipment is calibrated and certified according to current standard operating procedures.

**2.1.1** Microscope, inverted bright light

**2.1.2** Illumination box

**2.1.3** Low speed, refrigerated centrifuge (Beckman J-6B centrifuge with JS 4.0 rotor or equivalent).

**2.1.4** Refrigerator, 2°-7°C

**2.1.5** Micropipettors and appropriate tips

**2.1.6** Vacuum source for aspirating liquids (vacuum pump with side arm flask of Chapman-type filter pump attached to a water line)

**2.1.7** Laminar Flow Biological Safety cabinet (NuAire, Inc. Labgard or equivalent )

**2.2 Reagents/supplies**

Equivalent reagents or supplies may be substituted for any brand name listed below. All reagents and supplies must be sterile.

**UNCONTROLLED COPY**

**Supplemental Assay Method for the Detection of Extraneous Hemadsorbing Agents in Master Seed Virus**

**2.2.1** Appropriate vessels at least 7 days from the last subculture containing:

- MS-inoculated
- negative control cell monolayers.

The preferred vessel is 25-cm<sup>2</sup> flasks, but any suitable alternative may be used if the surface area is  $\geq 6$  cm<sup>2</sup> each.

**2.2.2** A vessel containing a monolayer of Vero cells,  $\geq 6$  cm<sup>2</sup> to be used in preparation of positive control (PC).

**2.2.3** PC virus (current approved lot of Bovine Parainfluenza type 3)

**2.2.4** Alsever's solution

1. 8.0 g sodium citrate (C<sub>6</sub>H<sub>5</sub>Na<sub>3</sub>O<sub>7</sub>•2H<sub>2</sub>O)
2. 0.55 g citric acid (C<sub>6</sub>H<sub>8</sub>O<sub>7</sub>•H<sub>2</sub>O)
3. 4.2 g sodium chloride (NaCl)
4. 20.5 g glucose (C<sub>6</sub>H<sub>12</sub>O<sub>6</sub>)
5. Q.S. to 1000 mL with deionized water (DI).
6. Filter sterilize with a 0.22- $\mu$ m filter.
7. Store at 2°- 7°C.

**2.2.5** Guinea pig and chicken RBC in equal volumes of Alsever's Solution.  
Store at 2°- 7°C.

**2.2.6** 0.01 M Phosphate buffered saline (PBS)

1. 1.33 g sodium phosphate, dibasic, anhydrous (Na<sub>2</sub>HPO<sub>4</sub>)
2. 0.22 g sodium phosphate, monobasic, monohydrate (NaH<sub>2</sub>PO<sub>4</sub>•H<sub>2</sub>O)
3. 8.5 g NaCl
4. Q.S. to 1000 mL with DI.

**UNCONTROLLED COPY**

**Supplemental Assay Method for the Detection of Extraneous Hemadsorbing Agents in Master Seed Virus**

5. Adjust pH to 7.2-7.6 with 0.1 N sodium hydroxide (NaOH) or 1.0 N hydrochloric acid (HCl).
6. Sterilize by autoclaving at 15 psi,  $121^{\circ}\pm 2^{\circ}\text{C}$  for  $35 \pm 5$  minutes.
7. Store at  $2^{\circ}$ -  $7^{\circ}\text{C}$ .

**2.2.7** Pipettes: 1-mL, 5-mL, and 25-mL

**2.2.8** Conical tubes: 50-mL

**2.2.9** Cell culture flasks: 25-cm<sup>2</sup> (or appropriate vessel)

**2.2.10** Graduated cylinder

**3. Preparation for the Test**

**3.1 Personnel qualifications/training**

Laboratory personnel must be trained in the operation of instruments and equipment used prior to performing this test. Personnel must be familiar with proper use of test reagents and biological materials. Personnel must have knowledge of safe operating procedures and policies.

**3.2 Preparation of RBC**

**3.2.1 Wash guinea pig and/or chicken RBC**

1. Dispense 10-20 mL of guinea pig and chicken blood preserved in Alsever's solution into separate 50-mL conical centrifuge tubes.
2. Fill the tube with Alsever's/PBS and gently invert tube several times to wash RBC.
3. Centrifuge RBC for  $15 \pm 5$  minutes at 400-800 x g (e.g. 1500-1800 rpm, J6-B centrifuge with a JS-4.0 or JS-4.2 rotor) at  $4^{\circ}\pm 2^{\circ}\text{C}$ .
4. Aspirate PBS and buffy coat (bone colored layer of cells on top of RBC) using a pipette or vacuum type aspirator.

**UNCONTROLLED COPY**

### Supplemental Assay Method for the Detection of Extraneous Hemadsorbing Agents in Master Seed Virus

5. Repeat **Steps 2 through 4** for a total of 3 or more washes (until supernatant is clear). Store the packed guinea pig and chicken RBC at 2°-7°C for up to 1 week. RBC should be discarded if hemolysis is observed.

#### 3.2.2 Preparation of 0.2% RBC on day of test.

1. Dispense 49.8 mL of PBS into a suitable container.
2. Using 1-mL pipettes, add 100 µL of washed guinea pig RBC and 100 µL of chicken RBC to PBS. Rinse pipette(s) thoroughly to dispense all RBC. Gently swirl to mix.
3. RBC suspension should be discarded following completion of test.

#### 3.3 Preparation of the sample

For each cell line, seed at least one 25-cm<sup>2</sup> flask with MS inoculated cells and at least one 25-cm<sup>2</sup> flask with un-inoculated cells to be tested sequentially at 4°C and 25°C. In addition, seed two 25-cm<sup>2</sup> flasks of un-inoculated Vero cells to serve as positive controls, one flask for each temperature. The positive control flasks are seed-inoculated with bovine parainfluenza virus 3 at the time of the last subculture.

### 4. Performance of the Test

The HAd test is conducted at 2 temperatures: 4°± 2°C and room temperature (23°± 2°C) for MS and NC flasks of each cell type. Two PC flasks (Vero cell monolayers) are tested (one at 4° and one at room temperature). Alternatively, MS, NC and PC flasks of each cell type may be incubated simultaneously at each temperature (**Section 4.2**).

#### 4.1 Vessels incubated at each temperature separately

- 4.1.1 Decant medium from MS, NC and PC flasks.
- 4.1.2 Add 5 mL of PBS to each of the MS, NC and PC flasks, swirl gently and decant. Repeat for a total of 3 washes.
- 4.1.3 Gently swirl the 0.2% RBC mixture and add 5 mL into each MS, NC and PC flasks (**Section 3.2.2**).
- 4.1.4 Incubate all flasks except one PC flask at 4°± 2°C for 25 ± 5 minutes.
- 4.1.5 Decant the 0.2% RBC mixture.

**UNCONTROLLED COPY**

**Supplemental Assay Method for the Detection of Extraneous Hemadsorbing Agents in Master Seed Virus**

**4.1.6** Wash vessels 3 times as described in **Section 4.1.2**.

**4.1.7** Using an illumination box, compare each MS flask to the corresponding NC and PC control flasks. Look for areas of HAd on the cell monolayer (see **Appendix I**).

**4.1.8** Using an inverted microscope at 100X, compare each MS flask to the corresponding NC and PC flask(s). Look for HAd of individual RBC to cell membranes (see **Appendix II**).

**4.1.9** If HAd is not detected, add 0.2% RBC mixture to MS, NC cell monolayers and remaining PC flask as described in **Section 4.1.3**.

**4.1.10** Incubate all vessels at room temperature for  $25 \pm 5$  minutes.

**4.1.11** Repeat **Sections 4.1.6 through 4.1.9**.

**4.2 Separate vessels incubated simultaneously at each temperature**

**4.2.1** Repeat **Sections 4.1.1 through 4.1.3** for duplicate sets of vessels.

**4.2.2** Incubate MS and NC flasks for each cell line, as well as a PC flask (Vero) at  $4^{\circ} \pm 2^{\circ}\text{C}$  for  $25 \pm 5$  minutes.

**4.2.3** Incubate separate MS and NC flasks for each cell line, as well as a PC flask (Vero) at room temperature for  $25 \pm 5$  minutes.

**4.2.4** Repeat **Sections 4.1.6 through 4.1.9** for each set of vessels.

**5. Interpretation of the Test Results**

For a valid test, the PC cell culture must demonstrate HAd while the NC cell culture must be free of HAd activity.

HAd activity in the MS inoculated cell monolayers other than the non-neutralized MS is indicative of infection by extraneous viruses, and the MS is determined unsatisfactory. The MS may be retested to confirm original results if warranted. Additional investigative work may be warranted to determine cause of HAd.

**6. Report of Test Results**

All records are kept in accordance with current recordkeeping practices

**UNCONTROLLED COPY**

**Supplemental Assay Method for the Detection of Extraneous Hemadsorbing Agents in Master Seed Virus**

**7. References**

Title 9, *Code of Federal Regulations*, parts 113.46, 113.51, 113.52 and 113.55, U.S. Government Printing Office, Washington, DC.

**8. Summary of Revisions**

**Version .05**

- **The coversheet and contact information has been updated.**

**Version .04**

- The Contact information has been updated; however, the Virology Section has elected to keep the same next review date for the document.

**Version .03**

- The Contact information has been updated.

**Version .02**

This document was revised to clarify the practices currently in use at the Center for Veterinary Biologics and to provide additional detail. While no significant changes were made that impact the outcome of the test, the following changes were made to the document:

- **1:** An overview of the test method has been added.
- **2.1:** More current and appropriate equipment has been added.
- **2.2:** Surface area criteria to test and a Positive Control have been added to the test.
- **3.1:** Personnel qualifications and training have been clarified.
- **3.3:** Additional detail to washing and preparation of 0.2% RBC has been added.
- **3.4:** A Positive Control has been added to the test system.
- **4.2:** The Positive Control flask has been added to the incubation step.
- **5.1:** This section has been condensed and the roles of the Positive and Negative Control vessels have been clarified.

**UNCONTROLLED COPY**



**Supplemental Assay Method for the Detection of Extraneous Hemadsorbing Agents in Master Seed Virus**

- The refrigeration temperatures have been changed from  $4^{\circ} \pm 2^{\circ}\text{C}$  to  $2^{\circ}\text{-}7^{\circ}\text{C}$  throughout the document.
- Appendices containing macro- and microscopic images depicting hemadsorption have been added.

Supplemental Assay Method for the Detection of Extraneous Hemadsorbing Agents in Master Seed Virus

**Appendix I**

**Macroscopic view of hemadsorption**



**UNCONTROLLED COPY**

Supplemental Assay Method for the Detection of Extraneous Hemadsorbing Agents in Master Seed Virus

Appendix II

Microscopic view of hemadsorption

