



Summary of Studies Supporting USDA Product Licensure

Establishment Name	Boehringer Ingelheim Animal Health USA Inc.
USDA Vet Biologics Establishment Number	124
Product Code	49W5.20
True Name	Encephalomyelitis-Rhinopneumonitis-Influenza-West Nile Virus Vaccine, Eastern & Western & Venezuelan, Killed Virus, Tetanus Toxoid
Tradename(s) / Distributor or Subsidiary (if different from manufacturer)	Equi-Jec 7 - No distributor specified
Date of Compilation Summary	February 06, 2019

Disclaimer: Do not use the following studies to compare one product to another. Slight differences in study design and execution can render the comparisons meaningless.

Study Type	Efficacy
Pertaining to	Clostridium tetanus
Study Purpose	Demonstration of efficacy against Clostridium tetanus
Product Administration	One dose, administered intramuscularly
Study Animals	10 guinea pigs (10 vaccinates)
Challenge Description	Not applicable
Interval observed after challenge	Not applicable
Results	<p>Vaccinate serum samples were collected and pooled, then tested for antitoxin content by indirect Enzyme-Linked Immunosorbent Assay.</p> <p>A satisfactory value which met the requirements per 9 CFR 113.114(c) was achieved.</p>
USDA Approval Date	May 1, 2008

Study Type	Efficacy
Pertaining to	Eastern equine encephalomyelitis
Study Purpose	Demonstration of efficacy against Eastern equine encephalomyelitis
Product Administration	Two doses, administered intramuscularly, 14 to 21 days apart
Study Animals	12 guinea pigs (10 vaccinates, 2 controls)
Challenge Description	Not applicable
Interval observed after challenge	Not applicable
Results	<p>Serum samples were tested by a plaque reduction, serum neutralization test, 14 to 21 days after the second injection.</p> <p>Vaccinates and controls were evaluated in terms of Eastern equine encephalomyelitis per the criteria in 9 CFR 113.207(b) and the requirements were met.</p>
USDA Approval Date	May 1, 2008

Study Type	Efficacy
Pertaining to	Venezuelan equine encephalomyelitis
Study Purpose	Demonstration of efficacy against Venezuelan Equine Encephalomyelitis
Product Administration	Two doses, administered intramuscularly, 14 to 21 days apart
Study Animals	12 guinea pigs (10 vaccinates, 2 controls)
Challenge Description	Not applicable
Interval observed after challenge	Not applicable
Results	<p>Serum samples were tested by a plaque reduction, serum neutralization test, 14 to 21 days after the second injection.</p> <p>Vaccinates and controls were evaluated in terms of Venezuelan equine encephalomyelitis per the criteria in 9 CFR 113.207(b) and the requirements were met.</p>
USDA Approval Date	May 1, 2008

Study Type	Efficacy
Pertaining to	Western equine encephalomyelitis
Study Purpose	Demonstration of efficacy against Western equine encephalomyelitis
Product Administration	Two doses, administered intramuscularly, 14 to 21 days apart
Study Animals	12 guinea pigs (10 vaccinates, 2 controls)
Challenge Description	Not applicable
Interval observed after challenge	Not applicable
Results	<p>Serum samples were tested by a plaque reduction, serum neutralization test, 14 to 21 days after the second injection.</p> <p>Vaccinates and controls were evaluated in terms of Western equine encephalomyelitis per the criteria in 9 CFR 113.207(b) and the requirements were met.</p>
USDA Approval Date	May 1, 2008

Study Type	Efficacy																				
Pertaining to	Equine herpesvirus type 1 (EHV-1)																				
Study Purpose	Demonstration of efficacy against respiratory disease caused by EHV-1																				
Product Administration	Two doses, administered intramuscularly, 21 days apart																				
Study Animals	40 horses (20 vaccinates, 20 controls), 4-5 months of age																				
Challenge Description	Equine herpesvirus type 1 administered 15 days post-final vaccination																				
Interval observed after challenge	Horses were observed daily for 14 days post-challenge																				
Results	<p>See raw data on following pages.</p> <p>The horses were assessed for the presence of nasal discharge as signs of respiratory disease. The severity of nasal discharge was classified as “normal”, “mild”, or “moderate” according to the following classification of the nasal scores.</p> <table border="1"> <thead> <tr> <th>Disease status</th> <th>Maximum Nasal Score</th> </tr> </thead> <tbody> <tr> <td>Normal</td> <td>0 or 1</td> </tr> <tr> <td>Mild</td> <td>1.5 or 2</td> </tr> <tr> <td>Moderate</td> <td>4 or 6</td> </tr> </tbody> </table> <p>The number of horses in each category were:</p> <table border="1"> <thead> <tr> <th></th> <th>Normal</th> <th>Mild</th> <th>Moderate</th> </tr> </thead> <tbody> <tr> <td>Control</td> <td>0</td> <td>10</td> <td>10</td> </tr> <tr> <td>Vaccine</td> <td>6</td> <td>11</td> <td>3</td> </tr> </tbody> </table>	Disease status	Maximum Nasal Score	Normal	0 or 1	Mild	1.5 or 2	Moderate	4 or 6		Normal	Mild	Moderate	Control	0	10	10	Vaccine	6	11	3
Disease status	Maximum Nasal Score																				
Normal	0 or 1																				
Mild	1.5 or 2																				
Moderate	4 or 6																				
	Normal	Mild	Moderate																		
Control	0	10	10																		
Vaccine	6	11	3																		
USDA Approval Date	January 28, 2009																				

Nasal Discharge:

Treatment	ID	Day Postchallenge														
		0	1	2	3	4	5	6	7	8	9	10	11	12	13	14
Controls (20 horses)	1					1.5			1.5	1.5	1	1.5				
	2						1.5		1.5	1.5	1	1.5	1.5	1		
	3						1.5			1.5	2			1.5		
	4			1		2	1.5		1.5	1.5	1.5	1.5				1.5
	5				2	2	2	1	4	2	2	1.5	1.5		1.5	
	6			1		4	6	4	4	4	4	2	2	2		
	7					1.5	1.5	1.5	1.5	1.5	2	4	1		1	1
	8								1.5	2	2	4	1.5	4	2	1.5
	9				1	1.5	1.5	1.5	1.5	1.5	2	1.5				
	10			1			1		1.5	1.5	2	4			1.5	1.5
	11						1.5	1.5	1.5		2		1.5	1.5	1.5	
	12						1.5	1.5		2						1.5
	13						2	1.5	1.5	2	2	2	1.5	1.5	1.5	4
	14				1.5	2		1.5		1.5	1.5	1.5			2	2
	15				1	2	1.5	1	1.5		4		1		4	1.5
	16					1.5	1.5	2	2	2	2	1.5	1	1	4	2
	17					1.5		1			1.5	2		1.5	1.5	
	18						1	1.5	1.5	4	4	2	1.5	4	1.5	2
	19				1	2	1.5		1.5	2	4	1	1.5		1	
	20						1.5	1.5	2	1.5	2				1.5	
Vaccinates (20 horses)	1					1		1				1.5				
	2				1											
	3						1	1.5	4		1.5	1.5			1	
	4				1						2	1				
	5				1				1	1						
	6				1	1.5						1.5	2	2	2	1.5
	7							2					1.5			
	8															
	9					2	1.5	2	2	6	2	1.5		1.5	4	2
	10								1				1	1.5		
	11				1		1.5		2	2	1	1.5				
	12				1		1.5	2	1.5	2	2	2		2	2	1.5
	13				1.5						1.5	1.5			1.5	1.5
	14							1	1			1			1.5	
	15				1											
	16				1		1.5	1.5	1			1.5				
	17															
	18						1			1.5		1.5				
	19														6	2
	20															

Scoring:

Blank is 0 = none;

1 = slight serous, as may be observed in both normal and diseased horses;

1.5 = very slight mucopurulent discharge;

2 = moderate clear serous discharge, or slight mucopurulent discharge;

3 = abundant serous discharge;

4 = moderate mucopurulent discharge;

6 = heavy mucopurulent discharge

Study Type	Efficacy																										
Pertaining to	Equine herpesvirus (EHV) type 1																										
Study Purpose	Demonstration of efficacy against EHV-4																										
Product Administration	Two doses administered intramuscularly 21 days apart																										
Study Animals	37 horses (24 vaccinates, 13 controls), 4 to 5 months of age																										
Challenge Description	Equine herpesvirus type 4 was administered 15 days following second vaccination																										
Interval observed after challenge	Horses were observed daily for 14 days following challenge																										
Results	<p>The horses were assessed for the presence of nasal and ocular discharge as signs of respiratory disease. The severity of the nasal and ocular discharge resulted in the classification of the observed clinical signs as “mild” or “moderate” according to the following table:</p> <table border="1"> <thead> <tr> <th>Disease Status</th> <th>Nasal Score</th> <th>Ocular Score</th> </tr> </thead> <tbody> <tr> <td>Normal = 0</td> <td>0 or 1</td> <td>0 or 1</td> </tr> <tr> <td rowspan="2">Mild = 1</td> <td>0 or 1</td> <td>2</td> </tr> <tr> <td>1.5, 2, or 3</td> <td>Any</td> </tr> <tr> <td>Moderate = 2</td> <td>4 or 6</td> <td>Any</td> </tr> </tbody> </table> <p>Respiratory disease was observed as follows:</p> <table border="1"> <thead> <tr> <th></th> <th>Controls</th> <th>Vaccinates</th> </tr> </thead> <tbody> <tr> <td>Normal</td> <td>0 out of 13</td> <td>9 out of 24</td> </tr> <tr> <td>Mild</td> <td>12 out of 13</td> <td>14 out of 24</td> </tr> <tr> <td>Moderate</td> <td>1 out of 13</td> <td>1 out of 24</td> </tr> </tbody> </table> <p>See raw data on the following pages.</p>	Disease Status	Nasal Score	Ocular Score	Normal = 0	0 or 1	0 or 1	Mild = 1	0 or 1	2	1.5, 2, or 3	Any	Moderate = 2	4 or 6	Any		Controls	Vaccinates	Normal	0 out of 13	9 out of 24	Mild	12 out of 13	14 out of 24	Moderate	1 out of 13	1 out of 24
Disease Status	Nasal Score	Ocular Score																									
Normal = 0	0 or 1	0 or 1																									
Mild = 1	0 or 1	2																									
	1.5, 2, or 3	Any																									
Moderate = 2	4 or 6	Any																									
	Controls	Vaccinates																									
Normal	0 out of 13	9 out of 24																									
Mild	12 out of 13	14 out of 24																									
Moderate	1 out of 13	1 out of 24																									
USDA Approval Date	September 17, 2009																										

Ocular Discharge:

Treatment	Horse ID	Days Post-challenge															
		0	1	2	3	4	5	6	7	8	9	10	11	12	13	14	
Controls (13 horses)	8300																
	8861								1		1	2	1	1	1	1	
	0327										1	1				1	
	1029								1	1	1		1		1	1	
	2603											1					
	8822																
	5542												1	1	1	2	
	3280				1				1				1	1	1	1	
	5597														1	2	
	9331									2	2	2	2	1		2	1
	9339												1	1			1
	5103																
	6528																
Vaccinates (24 horses)	1278							1					1		1		
	1602							1	1	1	1	1	1	2			
	8026																
	3857									1	1	1				1	
	5560																
	5636																
	0261											1	1				
	0285						1	1								1	
	6051																
	6311							1			1						
	1310																
	5381																
	8023												1				
	8881				1	1		1					1				
	0019											1					
	1381																
	2333																1
	3086																
	3347																
	5379												1				
7297																	
7580								1				1					
7806								1	1								
8004																	

Scoring:

- No entry indicates 0 = horse is normal
- 1 = normal
- 2 = mild

Nasal Discharge:

Treatment	Horse #	Days Post-challenge															
		0	1	2	3	4	5	6	7	8	9	10	11	12	13	14	
Controls (13 horses)	8300							1.5	1.5	1.5	1.5			1.5			
	8861							1.5	1	1			1.5			1.5	
	0327							1	1		1.5	1.5		2		1	
	1029				1.5						1.5			1.5	1.5		
	2603				1.5			1.5			1.5						
	8822					2	1.5	1.5		2							
	5542								1.5	1.5	2	1.5	1		1.5	2	
	3280					1		2									
	5597					1.5		1	1.5	2	1.5		1		1.5	1.5	
	9331						1.5	1.5	1.5	4	1.5	1			1.5	1.5	
	9339							1.5	1.5	1.5	1.5	1.5	2	1			
	5103					1	1.5	2	1.5	2	1.5	2					
	6528							1.5	1.5			1.5		1			
Vaccinates (24 horses)	1278					1.5		1									
	1602							1.5	1.5	2	2	2	2	4			
	8026					2	1.5	1				2	1.5	2			
	3857									1.5	1.5						
	5560															1.5	
	5636									1.5							
	0261										1.5						
	0285										1.5						
	6051				1.5						1						1.5
	6311							1.5				2					
	1310																
	5381																
	8023																
	8881				1.5	1.5						1.5		1.5			
	0019																
	1381							1.5				1.5					
	2333																
	3086																
	3347				1.5												
	5379															2	
7297																	
7580					1											1.5	
7806																	
8004											1						

Scoring:

No entry is 0 = horse is normal

1 = slight clear serous, as may be observed in both normal and diseased horses;

1.5 = very slight mucopurulent discharge, one or both nostrils;

2 = moderate clear serous discharge or slightly mucopurulent discharge, in one or both nostrils;

3 = abundant clear serous discharge typically seen only in diseased horses;

4 = moderately mucopurulent discharge, in large quantities in both nostrils;

6 = heavy mucopurulent discharge in large amounts filling both nostrils

Study Type	Efficacy
Pertaining to	Equine influenza virus
Study Purpose	Demonstration of 6-month duration of immunity against respiratory disease caused by equine influenza
Product Administration	Two doses, administered intramuscularly, 21 days apart. Vaccinates received test product, and controls received adjuvanted diluent.
Study Animals	30 horses (20 vaccinates, 10 controls), 5-6 months of age
Challenge Description	Influenza A/eq/Ohio/2003 administered 184 days post-final vaccination
Interval observed after challenge	Horses were observed daily for 10 days post-challenge
Results	<p>See tables at the end of document for data.</p> <p>Clinical Signs: An animal was considered positive (affected by challenge) if the animal exhibited:</p> <ul style="list-style-type: none"> • Fever (temperature >102.5°F), OR • Nasal discharge (moderate serous discharge or mucopurulent discharge), OR • Ocular discharge <p>A total of 9/10 (90%) controls were positive as compared to only 9/20 (45%) vaccinates.</p> <p>There were no adverse reactions to vaccine administration at any timepoint.</p>
USDA Approval Date	September 7, 2010

Treatment	Clinical Sign	Days Post-challenge										
		0	1	2	3	4	5	6	7	8	9	10
Controls												
1	Fever											
	Nasal discharge							+	+	+	+	
	Ocular discharge							+			+	+
2	Fever											
	Nasal discharge			+				+		+	+	+
	Ocular discharge							+	+			+
3	Fever											
	Nasal discharge									+		
	Ocular discharge			+				+			+	+
4	Fever											
	Nasal discharge											
	Ocular discharge							+	+	+		+
5	Fever											
	Nasal discharge						+	+	+	+	+	
	Ocular discharge											
6	Fever											
	Nasal discharge						+			+		+
	Ocular discharge											+
7	Fever											
	Nasal discharge			+				+		+		+
	Ocular discharge			+					+			
8	Fever									+		
	Nasal discharge							+	+	+		+
	Ocular discharge			+	+			+	+			+
9	Fever											
	Nasal discharge											
	Ocular discharge											
10	Fever											
	Nasal discharge							+	+	+	+	+
	Ocular discharge						+	+		+	+	+

Treatment	Clinical Sign	Days Post-challenge												
		0	1	2	3	4	5	6	7	8	9	10		
Vaccinates														
1	Fever													
	Nasal discharge													
	Ocular discharge													
2	Fever													
	Nasal discharge													
	Ocular discharge													
3	Fever													
	Nasal discharge													
	Ocular discharge							+			+	+		
4	Fever													
	Nasal discharge									+				
	Ocular discharge													
5	Fever													
	Nasal discharge													
	Ocular discharge													
6	Fever													
	Nasal discharge													
	Ocular discharge													
7	Fever													
	Nasal discharge													
	Ocular discharge													
8	Fever													
	Nasal discharge													
	Ocular discharge													
9	Fever													
	Nasal discharge													
	Ocular discharge													
10	Fever							+						
	Nasal discharge								+	+				
	Ocular discharge										+		+	
11	Fever													
	Nasal discharge							+			+	+	+	
	Ocular discharge													
12	Fever													
	Nasal discharge										+			
	Ocular discharge													

Treatment	Clinical Sign	Days Post-challenge											
		0	1	2	3	4	5	6	7	8	9	10	
Vaccinates													
13	Fever												
	Nasal discharge						+						+
	Ocular discharge												
14	Fever												
	Nasal discharge												
	Ocular discharge												
15	Fever												
	Nasal discharge												
	Ocular discharge							+		+			
16	Fever												
	Nasal discharge								+				
	Ocular discharge												
17	Fever												
	Nasal discharge												
	Ocular discharge												
18	Fever												
	Nasal discharge												
	Ocular discharge												
19	Fever												
	Nasal discharge								+		+		
	Ocular discharge												
20	Fever												
	Nasal discharge												
	Ocular discharge												

Study Type	Efficacy
Pertaining to	Equine influenza
Study Purpose	Demonstration of efficacy against respiratory disease caused by equine influenza A2 strain Kentucky 95
Product Administration	Two doses, administered intramuscularly, 21 days apart
Study Animals	20 horses (20 vaccinates), 12 months of age
Challenge Description	Not applicable
Interval observed after challenge	Not applicable
Results	This product class allows the manufacturer to update micro-organisms in this vaccine under expedited procedures to respond to emerging needs. Abbreviated data to support influenza strain updates to the product composition were evaluated by USDA-APHIS and found to be acceptable based on regulations and policies at the time of approval. Full vaccination-challenge studies may not have been required for these updates.
USDA Approval Date	February 2, 2012

Study Type	Efficacy
Pertaining to	Equine influenza
Study Purpose	Demonstration of efficacy against respiratory disease caused by equine influenza A2 strain New Market 2/93
Product Administration	Two doses, administered intramuscularly, 21 days apart
Study Animals	20 horses (20 vaccinates), 12 months of age
Challenge Description	Not applicable
Interval observed after challenge	Not applicable
Results	This product class allows the manufacturer to update micro-organisms in this vaccine under expedited procedures to respond to emerging needs. Abbreviated data to support influenza strain updates to the product composition were evaluated by USDA-APHIS and found to be acceptable based on regulations and policies at the time of approval. Full vaccination-challenge studies may not have been required for these updates.
USDA Approval Date	February 2, 2012

Study Type	Efficacy									
Pertaining to	West Nile Virus (WNV)									
Study Purpose	Demonstration of twelve month duration of immunity against disease caused by WNV									
Product Administration	Two doses, administered intramuscularly, 25 days apart									
Study Animals	30 horses (20 vaccinates, 10 placebo controls) 4-5 months of age									
Challenge Description	West Nile Virus was administered at 380 days (10 vaccinated and 5 placebo control animals) or 408 days (10 vaccinated and 5 placebo control animals) post-final vaccination.									
Interval observed after challenge	Horses were observed twice daily for 14 days post-challenge and once daily for an additional 7 days post-challenge.									
Results	<p>An animal was considered affected by challenge if it developed neurological disease, as measured by mortality and microscopic evidence of virus-induced brain disease (histopathology).</p> <p>Animals were also monitored for viremia (detection of WNV in the blood).</p> <p>Results are summarized as follows:</p> <table border="1"> <thead> <tr> <th>Outcome</th> <th>Controls</th> <th>Vaccinates</th> </tr> </thead> <tbody> <tr> <td>Mortality</td> <td>7/10 (70%)</td> <td>1/20 (5%)</td> </tr> <tr> <td>Viremia at least one day</td> <td>10/10 (100%)</td> <td>2/20 (10%)</td> </tr> </tbody> </table> <p>See raw data on following pages.</p>	Outcome	Controls	Vaccinates	Mortality	7/10 (70%)	1/20 (5%)	Viremia at least one day	10/10 (100%)	2/20 (10%)
Outcome	Controls	Vaccinates								
Mortality	7/10 (70%)	1/20 (5%)								
Viremia at least one day	10/10 (100%)	2/20 (10%)								
USDA Approval Date	September 3, 2010									

Treatment	#	Died or Euthanized due to disease severity	Severity Histopathological lesions	
			Medulla	Pons
Controls (10 horses)	1	Yes	3	3
	2	Yes	3	3
	3	Yes	3	3
	4	Yes	3	3
	5	Yes	3	3
	6	Yes	2	2
	7	Yes	1	1
	8	No	1	1
	9	No	1	1
	10	No	1	0.5
Vaccinates (20 horses)	1	Yes	3	3
	2	No	2	0.5
	3	No	1	1
	4	No	1	0.5
	5	No	1	0.5
	6	No	1	0.5
	7	No	0.5	0.5
	8	No	0.5	0.5
	9	No	0.5	0
	10	No	0	0.5
	11	No	0	0
	12	No	0	0
	13	No	0	0
	14	No	0	0
15	No	0	0	
16	No	0	0	
17	No	0	0	
18	No	0	0	
19	No	0	0	
20	No	0	0	

Scoring of histopathological lesions:	
0 =	No significant lesions.
0.5 =	Rare, small, multifocal glial nodules scattered throughout the parenchyma.
1 =	Mild, nonsuppurative encephalitis characterized by mild multifocal perivascular cuffs with lymphocytes and plasma cells and a rare neutrophil and scattered multifocal glial nodules composed of glial cells with a few mononuclear inflammatory cells. Occasionally within this grade, there may be minimal perivascular cuffing and more moderate scattered glial nodules.
2 =	Moderate nonsuppurative encephalitis characterized by moderate lymphoplasmacytic perivascular cuffs around many vessels and multifocal accumulations of glial nodules scattered throughout the parenchyma.
3 =	Severe nonsuppurative encephalitis characterized by severe and thick lymphoplasmacytic perivascular cuffing with multiple scattered glial nodules throughout the parenchyma.

Study Type	Efficacy																																												
Pertaining to	West Nile Virus (WNV)																																												
Study Purpose	Demonstration of seven month duration of immunity against WNV																																												
Product Administration	Two doses, administered intramuscularly 22 days apart																																												
Study Animals	30 horses (20 vaccinates, 10 placebo controls) 4-5 months of age																																												
Challenge Description	Challenged with West Nile Virus at 201 days (Group 1: 10 vaccinated and 5 placebo control animals) or 222 days (Group 2: 10 vaccinated and 5 placebo control animals) after the second vaccination.																																												
Interval observed after challenge	Horses were bled on the day of challenge, twice daily for 6 days post-challenge, once daily for an additional 4 days post-challenge, and on day 14 post-challenge																																												
Results	<p>The primary outcome was viremia (detection of WNV in the blood). An animal was considered to be positive if virus was detected in the blood on one or more occasions post-challenge.</p> <p>The number of animals positive for viremia at least once is summarized for as follows:</p> <table border="1"> <thead> <tr> <th>Challenge Group</th> <th>Controls</th> <th>Vaccinates</th> </tr> </thead> <tbody> <tr> <td>1</td> <td>5/5 (100%)</td> <td>1/10 (10%)</td> </tr> <tr> <td>2</td> <td>5/5 (100%)</td> <td>3/10 (30%)</td> </tr> </tbody> </table> <p>The outcome for viremia is as follows for the first group of horses challenged 201 days following the second vaccination:</p> <table border="1"> <thead> <tr> <th></th> <th>Horse ID</th> <th>Challenge Group 1</th> </tr> </thead> <tbody> <tr> <td rowspan="5">Controls (5 horses)</td> <td>S16</td> <td>Positive</td> </tr> <tr> <td>S21</td> <td>Positive</td> </tr> <tr> <td>S23</td> <td>Positive</td> </tr> <tr> <td>S26</td> <td>Positive</td> </tr> <tr> <td>S30</td> <td>Positive</td> </tr> <tr> <td rowspan="10">Vaccinates (10 horses)</td> <td>S17</td> <td>Negative</td> </tr> <tr> <td>S18</td> <td>Negative</td> </tr> <tr> <td>S19</td> <td>Negative</td> </tr> <tr> <td>S20</td> <td>Negative</td> </tr> <tr> <td>S22</td> <td>Positive</td> </tr> <tr> <td>S24</td> <td>Negative</td> </tr> <tr> <td>S25</td> <td>Negative</td> </tr> <tr> <td>S27</td> <td>Negative</td> </tr> <tr> <td>S28</td> <td>Negative</td> </tr> <tr> <td>S29</td> <td>Negative</td> </tr> </tbody> </table> <p>Positive = WNV detected in blood on one or more occasions post-challenge Negative = WNV detected in blood on zero occasions post-challenge</p>	Challenge Group	Controls	Vaccinates	1	5/5 (100%)	1/10 (10%)	2	5/5 (100%)	3/10 (30%)		Horse ID	Challenge Group 1	Controls (5 horses)	S16	Positive	S21	Positive	S23	Positive	S26	Positive	S30	Positive	Vaccinates (10 horses)	S17	Negative	S18	Negative	S19	Negative	S20	Negative	S22	Positive	S24	Negative	S25	Negative	S27	Negative	S28	Negative	S29	Negative
Challenge Group	Controls	Vaccinates																																											
1	5/5 (100%)	1/10 (10%)																																											
2	5/5 (100%)	3/10 (30%)																																											
	Horse ID	Challenge Group 1																																											
Controls (5 horses)	S16	Positive																																											
	S21	Positive																																											
	S23	Positive																																											
	S26	Positive																																											
	S30	Positive																																											
Vaccinates (10 horses)	S17	Negative																																											
	S18	Negative																																											
	S19	Negative																																											
	S20	Negative																																											
	S22	Positive																																											
	S24	Negative																																											
	S25	Negative																																											
	S27	Negative																																											
	S28	Negative																																											
	S29	Negative																																											

The outcome for **viremia** is as follows for the second group of horses challenged 222 days following the second vaccination:

	Horse ID	Challenge Group 2
Controls (5 horses)	S32	Positive
	S36	Positive
	S39	Positive
	S40	Positive
	S43	Positive
Vaccinates (10 horses)	S31	Negative
	S33	Positive
	S34	Negative
	S35	Positive
	S37	Negative
	S38	Negative
	S41	Negative
	S42	Negative
	S44	Negative
	S45	Positive

Positive = WNV detected in blood on one or more occasions post-challenge
 Negative = WNV detected in blood on zero occasions post-challenge

USDA Approval Date

November 2, 2009

Study Type	Safety
Pertaining to	All fractions
Study Purpose	To demonstrate safety in pregnant mares under field conditions at two different test sites
Product Administration	Two intramuscular doses, given 16-28 days apart. 54 pregnant mares were injected with placebo and 325 pregnant mares were vaccinated with test product.
Study Animals	Three hundred seventy-nine pregnant mares at two locations were included in the study. The mares were confirmed to be pregnant by serum hormonal evaluation on the day of the first vaccination.
Challenge Description	Not applicable
Interval observed after vaccination	1 st and 2 nd trimester: Mares observed immediately after vaccination and daily for overall health and for abortion. Resulting foals were observed daily for 7 days following birth. 3 rd trimester: Mares observed immediately after vaccination and daily for overall health and for abortion. Resulting foals were observed daily for 30 days following birth.
Results	Results shown on next page

Results**Study 2013-PM-1009****North Dakota Site:**

Group	Vaccinated	Confirmed Pregnant	Foals	Parturition Rate
1 st trimester/ product	143	127	114	90%
1st trimester/ placebo	59	54	49	91%
2 nd trimester/ product	6	6	6	100%
3 rd trimester/ product	140	117	117	100%
Total – all animals	348	304	286	94%
Total – product only	289	250	237	95%
Total – placebo only	59	54	49	91%

Study 2013-PM-1009**Missouri Site:**

Group	Vaccinated	Confirmed Pregnant	Foals	Parturition Rate
2011 3 rd trimester	5	5	5	100%
2012 1 st trimester	1	1	1	100%
2012 2 nd trimester	53	43	39	91%
2012 3 rd trimester	26	26	25	96%
Total – product	85	75	70	93%

Study 2014-PM-1009**North Dakota Site:**

Group	Vaccinated	Confirmed Pregnant	Foaled	Parturition Rate	Foals Survived to End of Observation Period
2 nd trimester vaccinated	52	52	52	100%	51*
3 rd trimester vaccinated	69	69	67**	97.1%	67

*Lost foal affirmed by study cooperator to be due to causes other than vaccination.

**One mare died due to causes other than vaccination, as affirmed by study cooperator.

All other foals were normal and healthy

USDA Approval Date

September 12, 2014

Study Type	Safety																																																			
Pertaining to	All fractions																																																			
Study Purpose	To demonstrate safety under field conditions																																																			
Product Administration	Two doses, administered intramuscularly approximately 3 – 4 weeks apart																																																			
Study Animals	880 horses, including 218 foals 3 months of age and 52 foals 5 months of age																																																			
Challenge Description	Not applicable																																																			
Interval observed after challenge	Not applicable																																																			
Results	<p>Horses were observed at least daily following each vaccination, until resolution of any observed reactions.</p> <p>There were no systemic reactions observed at any of the sites. Two foals and one horse died from causes affirmed by licensee not attributed to vaccination.</p> <p>Adverse events were limited to transient, non-painful swellings at the injection site that resolved without treatment.</p> <p>Local injection site reactions are summarized below across the four sites:</p> <table border="1" data-bbox="432 1077 1445 1585"> <thead> <tr> <th rowspan="2">Site</th> <th rowspan="2">Total Number Of Vaccinates</th> <th rowspan="2">Number Of Vaccinates Administered 2 doses</th> <th colspan="2">Vaccinates With Transient Injection Site Swelling</th> <th colspan="2">Number Of Normal Vaccinates</th> </tr> <tr> <th>After 1st dose</th> <th>After 2nd dose</th> <th>After 1st dose</th> <th>After 2nd dose</th> </tr> </thead> <tbody> <tr> <td>North Dakota</td> <td>378</td> <td>378</td> <td>4</td> <td>0</td> <td>374</td> <td>378</td> </tr> <tr> <td>California</td> <td>43</td> <td>43</td> <td>4</td> <td>3</td> <td>39</td> <td>40</td> </tr> <tr> <td>Missouri</td> <td>292</td> <td>290</td> <td>0</td> <td>0</td> <td>292</td> <td>290</td> </tr> <tr> <td>Texas</td> <td>170</td> <td>169</td> <td>6</td> <td>1</td> <td>164</td> <td>168</td> </tr> <tr> <td>Total</td> <td>883</td> <td>880</td> <td>14 (1.6%)</td> <td>4 (0.5%)</td> <td>869 (98.4%)</td> <td>876 (99.5%)</td> </tr> </tbody> </table> <p>Results from each site are summarized on the following pages.</p>						Site	Total Number Of Vaccinates	Number Of Vaccinates Administered 2 doses	Vaccinates With Transient Injection Site Swelling		Number Of Normal Vaccinates		After 1 st dose	After 2 nd dose	After 1 st dose	After 2 nd dose	North Dakota	378	378	4	0	374	378	California	43	43	4	3	39	40	Missouri	292	290	0	0	292	290	Texas	170	169	6	1	164	168	Total	883	880	14 (1.6%)	4 (0.5%)	869 (98.4%)	876 (99.5%)
Site	Total Number Of Vaccinates	Number Of Vaccinates Administered 2 doses	Vaccinates With Transient Injection Site Swelling		Number Of Normal Vaccinates																																															
			After 1 st dose	After 2 nd dose	After 1 st dose	After 2 nd dose																																														
North Dakota	378	378	4	0	374	378																																														
California	43	43	4	3	39	40																																														
Missouri	292	290	0	0	292	290																																														
Texas	170	169	6	1	164	168																																														
Total	883	880	14 (1.6%)	4 (0.5%)	869 (98.4%)	876 (99.5%)																																														

North Dakota Site:

Summary	Number Of Vaccinates	Number Of Vaccinates Administered 2 doses	Vaccinates With Transient Injection Site Swelling		Number Of Normal Vaccinates	
			After 1 st dose	After 2 nd dose	After 1 st dose	After 2 nd dose
2-4 months	179	179	0	0	179	179
5-7 months	0	0	n/a	n/a	n/a	n/a
8-11 months	0	0	n/a	n/a	n/a	n/a
1-5 years*	121	121	2	0	119	121
6-15 years*	78	78	2	0	76	78
>16 years	0	0	n/a	n/a	n/a	n/a
Total	378	378	4	0	374	378

*Swellings were 3cm in size observed 1-3 days post vaccination that resolved within 3 days.

California Site:

Summary	Number Of Vaccinates	Number Of Vaccinates Administered 2 doses	Vaccinates With Transient Injection Site Swelling		Number Of Normal Vaccinates	
			After 1 st dose	After 2 nd dose	After 1 st dose	After 2 nd dose
2-4 months*	7	7	0	2	7	5
5-7 months**	1	1	1	0	0	1
8-11 months	0	0	n/a	n/a	n/a	n/a
1-5 years***	19	19	2	0	17	19
6-15 years****	15	15	1	1	14	14
>16 years	1	1	0	0	1	1
Total	43	43	4	3	39	40

*Swellings were 3cm in size observed within hours post vaccination that resolved within several hours.

**Swelling was 3cm in size observed immediately post vaccination that resolved within several hours.

***1 horse had a swelling 1cm in size observed immediately post vaccination that resolved within several hours. 1 horse had a swelling observed on day 1 that increased in size to 9cm on day 3 post vaccination and resolved by day 5.

****Same horse had a swelling after each vaccination that resolved within 3 weeks. Size after the first vaccination was 24cm. Size after the second vaccination was 10cm.

Missouri Site:

Summary	Number Of Vaccinates	Number Of Vaccinates Administered 2 doses	Vaccinates With Transient Injection Site Swelling		Number Of Normal Vaccinates	
			After 1 st dose	After 2 nd dose	After 1 st dose	After 2 nd dose
2-4 months	33	32	0	0	33	32
5-7 months	0	0	n/a	n/a	n/a	n/a
8-11 months	0	0	n/a	n/a	n/a	n/a
1-5 years	225	224	0	0	225	224
6-15 years	32	32	0	0	32	32
>16 years	2	2	0	0	2	2
Total	292	290	0	0	292	290

Texas Site:

Summary	Number Of Vaccinates	Number Of Vaccinates Administered 2 doses	Vaccinates With Transient Injection Site Swelling		Number Of Normal Vaccinates	
			After 1 st dose	After 2 nd dose	After 1 st dose	After 2 nd dose
2-4 months	0	0	n/a	n/a	n/a	n/a
5-7 months	52	51	1	1	51	50
8-11 months	0	0	n/a	n/a	n/a	n/a
1-5 years	114	114	5	0	109	114
6-15 years	0	0	n/a	n/a	n/a	n/a
>16 years	4	4	0	0	4	4
Total	170	169	6*	1**	164	168

*Swellings were <1.5cm were observed 4-7 days post vaccination and resolved within 6 days.

**Swelling was 5cm observed 1 day post vaccination that resolved within 2 days.

**USDA
Approval Date**

November 1, 2010