



Summary of Studies Supporting USDA Product Licensure

| | |
|---|--|
| Establishment Name | Intervet Inc. |
| USDA Vet Biologics Establishment Number | 165A |
| Product Code | 44B1.20 |
| True Name | Bovine Rhinotracheitis-Virus Diarrhea-Parainfluenza 3-Respiratory Syncytial Virus Vaccine, Modified Live Virus, Campylobacter Fetus-Leptospira Canicola-Grippotyphosa-Hardjo-Icterohaemorrhagiae-Pomona Bacterin |
| Tradename(s) / Distributor or Subsidiary (if different from manufacturer) | Bovilis Vista 5 VL5 SQ - Merck Animal Health Bovilis Vista 5 VL5 SQ - No distributor specified Vista 5 L5 SQ - Merck Animal Health Vista 5 VL5 SQ - Merck Sharp & Dohme Saude Animal Ltda. - Merck Animal Health Vista 5 VL5 SQ - No distributor specified |
| Date of Compilation Summary | December 15, 2020 |

Disclaimer: Do not use the following studies to compare one product to another. Slight differences in study design and execution can render the comparisons meaningless.

| | |
|--|---|
| Study Type | Efficacy |
| Pertaining to | Bovine Viral Diarrhea Virus Type 1 (BVDV1) |
| Study Purpose | To demonstrate efficacy against respiratory disease caused by BVDV1. |
| Product Administration | |
| Study Animals | Bovine |
| Challenge Description | BVDV Type 1b NY-1 strain |
| Interval observed after challenge | |
| Results | Study data were evaluated by USDA-APHIS prior to product licensure and met regulatory standards for acceptance at the time of submission. No data are published because this study was submitted to USDA-APHIS prior to January 1, 2007, and APHIS only requires publication of data submitted after that date. |
| USDA Approval Date | February 20, 2004 |

| | |
|--|---|
| Study Type | Efficacy |
| Pertaining to | Bovine Viral Diarrhea Virus Type 1 (BVDV1) |
| Study Purpose | To demonstrate efficacy against persistent infection of calves caused by BVDV1. |
| Product Administration | |
| Study Animals | Bovine |
| Challenge Description | BVDV Type 1b, strain SD02 BVD09 |
| Interval observed after challenge | |
| Results | Study data were evaluated by USDA-APHIS prior to product licensure and met regulatory standards for acceptance at the time of submission. No data are published because this study was submitted to USDA-APHIS prior to January 1, 2007, and APHIS only requires publication of data submitted after that date. |
| USDA Approval Date | June 23, 2005 |

| | |
|--|---|
| Study Type | Efficacy |
| Pertaining to | Bovine Viral Diarrhea Virus Type 1 (BVDV1) |
| Study Purpose | To demonstrate efficacy against fetal infection caused by BVDV1 206 days after vaccination. |
| Product Administration | |
| Study Animals | Bovine |
| Challenge Description | BVDV Type 1b strain SD02 BVD09 |
| Interval observed after challenge | |
| Results | Study data were evaluated by USDA-APHIS prior to product licensure and met regulatory standards for acceptance at the time of submission. No data are published because this study was submitted to USDA-APHIS prior to January 1, 2007, and APHIS only requires publication of data submitted after that date. |
| USDA Approval Date | October 6, 2005 |

| | |
|--|---|
| Study Type | Efficacy |
| Pertaining to | Bovine Viral Diarrhea Virus Type 1 (BVDV1) |
| Study Purpose | To demonstrate efficacy against respiratory disease caused by BVDV1 206 days after vaccination. |
| Product Administration | |
| Study Animals | Bovine |
| Challenge Description | |
| Interval observed after challenge | |
| Results | Study data were evaluated by USDA-APHIS prior to product licensure and met regulatory standards for acceptance at the time of submission. No data are published because this study was submitted to USDA-APHIS prior to January 1, 2007, and APHIS only requires publication of data submitted after that date. |
| USDA Approval Date | October 6, 2005 |

| | |
|--|---|
| Study Type | Efficacy |
| Pertaining to | Bovine Viral Diarrhea Virus Type 2 (BVDV2) |
| Study Purpose | To demonstrate efficacy against respiratory disease caused by BVDV2. |
| Product Administration | |
| Study Animals | Bovine |
| Challenge Description | BVDV2a strain 1373 |
| Interval observed after challenge | |
| Results | Study data were evaluated by USDA-APHIS prior to product licensure and met regulatory standards for acceptance at the time of submission. No data are published because this study was submitted to USDA-APHIS prior to January 1, 2007, and APHIS only requires publication of data submitted after that date. |
| USDA Approval Date | December 10, 2003 |

| | |
|--|---|
| Study Type | Efficacy |
| Pertaining to | Bovine Viral Diarrhea Virus Type 1 (BVDV2) |
| Study Purpose | To demonstrate efficacy against persistent infection of calves caused by BVDV2 |
| Product Administration | |
| Study Animals | Bovine |
| Challenge Description | BVDV Type 2 strain SD02 BVD05 |
| Interval observed after challenge | |
| Results | Study data were evaluated by USDA-APHIS prior to product licensure and met regulatory standards for acceptance at the time of submission. No data are published because this study was submitted to USDA-APHIS prior to January 1, 2007, and APHIS only requires publication of data submitted after that date. |
| USDA Approval Date | April 25, 2005 |

| Study Type | Efficacy | | | | | | | | | | | | | | | | | | | | | | | | |
|--|---|------------|--------------|------------|-------------|------------|----|---|---|----------|----|----|-----|-------|--------------|------------|-------------|------------|----|---|---|----------|----|----|----|
| Pertaining to | Bovine Viral Diarrhea Virus Type 2 (BVDV2) | | | | | | | | | | | | | | | | | | | | | | | | |
| Study Purpose | To demonstrate efficacy against fetal infection caused by BVDV2 206 days after vaccination. | | | | | | | | | | | | | | | | | | | | | | | | |
| Product Administration | 1 dose administered by the subcutaneous route 28 days prior to breeding. | | | | | | | | | | | | | | | | | | | | | | | | |
| Study Animals | 46 seronegative heifers, 28 vaccinates and 18 controls. | | | | | | | | | | | | | | | | | | | | | | | | |
| Challenge Description | All heifers were challenge with BVDV2 strain IV809-04 at 164-178 days of gestation. | | | | | | | | | | | | | | | | | | | | | | | | |
| Interval observed after challenge | Blood samples were collected on days 0, 5 through 10 post challenge for virus isolation. Fetuses were collected on day 60 after challenge. | | | | | | | | | | | | | | | | | | | | | | | | |
| Results | <p><u>Virus Isolation on heifers:</u></p> <table border="1"> <thead> <tr> <th>Group</th> <th># of Animals</th> <th># Affected</th> <th>Percent (%)</th> </tr> </thead> <tbody> <tr> <td>Vaccinates</td> <td>28</td> <td>2</td> <td>7</td> </tr> <tr> <td>Controls</td> <td>18</td> <td>18</td> <td>100</td> </tr> </tbody> </table> <p><u>Virus Isolation from fetal samples:</u> Calves (fetuses) were considered positive if virus was isolated from any fetal tissue (lung, spleen, thymus, kidney, buffy coat).</p> <table border="1"> <thead> <tr> <th>Group</th> <th># of Animals</th> <th># Affected</th> <th>Percent (%)</th> </tr> </thead> <tbody> <tr> <td>Vaccinates</td> <td>28</td> <td>2</td> <td>4</td> </tr> <tr> <td>Controls</td> <td>18</td> <td>17</td> <td>94</td> </tr> </tbody> </table> <p>Raw data shown on attached pages.</p> | Group | # of Animals | # Affected | Percent (%) | Vaccinates | 28 | 2 | 7 | Controls | 18 | 18 | 100 | Group | # of Animals | # Affected | Percent (%) | Vaccinates | 28 | 2 | 4 | Controls | 18 | 17 | 94 |
| Group | # of Animals | # Affected | Percent (%) | | | | | | | | | | | | | | | | | | | | | | |
| Vaccinates | 28 | 2 | 7 | | | | | | | | | | | | | | | | | | | | | | |
| Controls | 18 | 18 | 100 | | | | | | | | | | | | | | | | | | | | | | |
| Group | # of Animals | # Affected | Percent (%) | | | | | | | | | | | | | | | | | | | | | | |
| Vaccinates | 28 | 2 | 4 | | | | | | | | | | | | | | | | | | | | | | |
| Controls | 18 | 17 | 94 | | | | | | | | | | | | | | | | | | | | | | |
| USDA Approval Date | October 4, 2007 | | | | | | | | | | | | | | | | | | | | | | | | |

Viremia of Challenged Heifers

(Vaccinate Group)

| Heifer Number | Isolation of BVDV from Buffy-coats at Days post-challenge | | | | | | | |
|---------------|---|---|---|---|---|---|----|--|
| | 0 | 5 | 6 | 7 | 8 | 9 | 10 | |
| 1536 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | |
| 1538 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | |
| 1541 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | |
| 1544 | 0 | 0 | 1 | 0 | 0 | 0 | 0 | |
| 1545 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | |
| 1556 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | |
| 1558 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | |
| 1562 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | |
| 1566 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | |
| 1567 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | |
| 1569 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | |
| 1570 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | |
| 1575 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | |
| 1581 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | |
| 1582 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | |
| 1585 | 0 | 0 | 1 | 0 | 0 | 0 | 0 | |
| 1594 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | |
| 1596 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | |
| 1597 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | |
| 1598 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | |
| 1599 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | |
| 1601 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | |
| 1605 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | |
| 1606 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | |
| 1607 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | |
| 1608 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | |
| 1609 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | |
| 1614 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | |

0=negative; 1=positive.

(Control Group)

| Heifer Number | Isolation of BVDV from Buffy-coats at Days post-challenge | | | | | | | |
|---------------|---|---|---|---|---|---|----|--|
| | 0 | 5 | 6 | 7 | 8 | 9 | 10 | |
| 1540 | 0 | 1 | 1 | 1 | 1 | 0 | 0 | |
| 1542 | 0 | 1 | 1 | 0 | 1 | 0 | 0 | |
| 1543 | 0 | 1 | 1 | 1 | 1 | 0 | 1 | |
| 1546 | 0 | 0 | 1 | 1 | 1 | 0 | 0 | |
| 1549 | 0 | 1 | 1 | 1 | 1 | 1 | 0 | |
| 1553 | 0 | 1 | 1 | 1 | 1 | 1 | 0 | |
| 1557 | 0 | 0 | 1 | 1 | 1 | 0 | 0 | |
| 1571 | 0 | 1 | 1 | 1 | 1 | 0 | 0 | |
| 1572 | 0 | 1 | 1 | 0 | 1 | 0 | 0 | |
| 1573 | 0 | 1 | 1 | 1 | 1 | 0 | 0 | |
| 1574 | 0 | 1 | 1 | 1 | 0 | 1 | 0 | |
| 1577 | 0 | 1 | 1 | 1 | 0 | 0 | 0 | |
| 1586 | 0 | 1 | 1 | 1 | 0 | 0 | 0 | |
| 1590 | 0 | 1 | 1 | 1 | 0 | 0 | 0 | |
| 1591 | 0 | 1 | 1 | 1 | 1 | 1 | 0 | |
| 1593 | 0 | 1 | 1 | 1 | 1 | 1 | 0 | |
| 1595 | 0 | 1 | 1 | 1 | 1 | 1 | 0 | |
| 1615 | 0 | 1 | 0 | 0 | 0 | 1 | 0 | |

0=negative; 1=positive.

Virus Isolation from Fetal Samples

| Groups | Heifer ID | Virus isolations | | | | | VI Results |
|-----------|-----------|------------------|--------|------|--------|-------------|------------|
| | | Thymus | Spleen | Lung | Kidney | Buffy-coats | |
| Vaccinate | 1536 | 0 | 0 | 0 | 0 | 0 | 0 |
| | 1538 | 0 | 0 | 0 | 0 | 0 | 0 |
| | 1541 | 0 | 0 | 0 | 0 | 0 | 0 |
| | 1544 | 0 | 0 | 0 | 0 | 0 | 0 |
| | 1545 | 1 | 1 | 1 | 0 | 1 | 1 |
| | 1556 | 0 | 0 | 0 | 0 | 0 | 0 |
| | 1558 | 0 | 0 | 0 | 0 | 0 | 0 |
| | 1562 | 0 | 0 | 0 | 0 | 0 | 0 |
| | 1566 | 0 | 0 | 0 | 0 | 0 | 0 |
| | 1567 | 0 | 0 | 0 | 0 | 0 | 0 |
| | 1569 | 0 | 0 | 0 | 0 | 0 | 0 |
| | 1570 | 0 | 0 | 0 | 0 | 0 | 0 |
| | 1575 | 0 | 0 | 0 | 0 | 0 | 0 |
| | 1581 | 0 | 0 | 0 | 0 | 0 | 0 |
| | 1582 | 0 | 0 | 0 | 0 | 0 | 0 |
| | 1585 | 0 | 0 | 0 | 0 | 0 | 0 |
| | 1594 | 0 | 0 | 0 | 0 | 0 | 0 |
| | 1596 | 0 | 0 | 0 | 0 | 0 | 0 |
| | 1597 | 0 | 0 | 0 | 0 | 0 | 0 |
| | 1598 | 0 | 0 | 0 | 0 | 0 | 0 |
| 1599 | 0 | 0 | 0 | 0 | 0 | 0 | |
| 1601 | 0 | 0 | 0 | 0 | 1 | 1 | |
| 1605 | 0 | 0 | 0 | 0 | 0 | 0 | |
| 1606 | 0 | 0 | 0 | 0 | 0 | 0 | |
| 1607 | 0 | 0 | 0 | 0 | 0 | 0 | |
| 1608 | 0 | 0 | 0 | 0 | 0 | 0 | |
| 1609 | 0 | 0 | 0 | 0 | 0 | 0 | |
| 1614 | 0 | 0 | 0 | 0 | 0 | 0 | |

| Groups | Heifer ID | Virus isolations | | | | | VI Results |
|----------|-----------|------------------|--------|------|--------|-------------|------------|
| | | Thymus | Spleen | Lung | Kidney | Buffy-coats | |
| Controls | 1540 | 1 | 1 | 1 | 1 | 1 | 1 |
| | 1542 | 1 | 1 | 1 | 1 | 0 | 1 |
| | 1543 | 0 | 0 | 0 | 0 | 1 | 1 |
| | 1546 | 0 | 0 | 0 | 0 | 1 | 1 |
| | 1549 | 1 | 1 | 1 | 1 | 1 | 1 |
| | 1553 | 1 | 1 | 1 | 1 | 1 | 1 |
| | 1557 | 1 | 1 | 1 | 1 | 1 | 1 |
| | 1571 | 0 | 0 | 0 | 0 | 0 | 0 |
| | 1572 | 0 | 0 | 0 | 0 | 1 | 1 |
| | 1573 | 0 | 0 | 0 | 0 | 1 | 1 |
| | 1574 | 1 | 1 | 1 | 1 | 0 | 1 |
| | 1577 | 0 | 0 | 0 | 0 | 1 | 1 |
| | 1586 | 1 | 1 | 1 | 1 | 0 | 1 |
| | 1590 | 1 | 1 | 1 | 1 | 0 | 1 |
| | 1591 | 1 | 1 | 1 | 1 | 0 | 1 |
| | 1593 | 0 | 1 | 0 | 1 | 1 | 1 |
| | 1595 | 1 | 1 | 1 | 1 | 1 | 1 |
| | 1615 | 0 | 0 | 0 | 0 | 1 | 1 |

| | |
|--|---|
| Study Type | Efficacy |
| Pertaining to | Bovine Viral Diarrhea Virus Type 2 (BVDV2) |
| Study Purpose | To demonstrate efficacy against respiratory disease caused by BVDV2 200 days after vaccination. |
| Product Administration | |
| Study Animals | Bovine |
| Challenge Description | |
| Interval observed after challenge | |
| Results | Study data were evaluated by USDA-APHIS prior to product licensure and met regulatory standards for acceptance at the time of submission. No data are published because this study was submitted to USDA-APHIS prior to January 1, 2007, and APHIS only requires publication of data submitted after that date. |
| USDA Approval Date | June 22, 2006 |

| | |
|--|---|
| Study Type | Efficacy |
| Pertaining to | <i>Campylobacter fetus</i> |
| Study Purpose | To demonstrate effectiveness against infertility caused by <i>C. fetus</i> |
| Product Administration | Subcutaneous |
| Study Animals | Bovine |
| Challenge Description | |
| Interval observed after challenge | |
| Results | Study data were evaluated by USDA-APHIS prior to product licensure and met regulatory standards for acceptance at the time of submission. No data are published because this study was submitted to USDA-APHIS prior to January 1, 2007, and APHIS only requires publication of data submitted after that date. |
| USDA Approval Date | March 26, 2004 |

| | |
|--|---|
| Study Type | Efficacy |
| Pertaining to | Infectious Bovine Rhinotracheitis (IBR) |
| Study Purpose | To demonstrate effectiveness against disease caused by IBR |
| Product Administration | Subcutaneous |
| Study Animals | Bovine |
| Challenge Description | |
| Interval observed after challenge | |
| Results | Study data were evaluated by USDA-APHIS prior to product licensure and met regulatory standards for acceptance at the time of submission. No data are published because this study was submitted to USDA-APHIS prior to January 1, 2007, and APHIS only requires publication of data submitted after that date. |
| USDA Approval Date | March 29, 2004 |

| | |
|--|---|
| Study Type | Efficacy |
| Pertaining to | Infectious Bovine Rhinotracheitis (IBR) |
| Study Purpose | To demonstrate effectiveness against abortions caused by IBR |
| Product Administration | Subcutaneous |
| Study Animals | Bovine |
| Challenge Description | |
| Interval observed after challenge | |
| Results | Study data were evaluated by USDA-APHIS prior to product licensure and met regulatory standards for acceptance at the time of submission. No data are published because this study was submitted to USDA-APHIS prior to January 1, 2007, and APHIS only requires publication of data submitted after that date. |
| USDA Approval Date | June 27, 2005 |

| | |
|--|---|
| Study Type | Efficacy |
| Pertaining to | Infectious Bovine Rhinotracheitis (IBR) |
| Study Purpose | To demonstrate efficacy against abortions caused by IBR at 217 days post vaccination. |
| Product Administration | Subcutaneous |
| Study Animals | Bovine |
| Challenge Description | |
| Interval observed after challenge | |
| Results | Study data were evaluated by USDA-APHIS prior to product licensure and met regulatory standards for acceptance at the time of submission. No data are published because this study was submitted to USDA-APHIS prior to January 1, 2007, and APHIS only requires publication of data submitted after that date. |
| USDA Approval Date | March 16, 2006 |

| | |
|--|---|
| Study Type | Efficacy |
| Pertaining to | Infectious Bovine Rhinotracheitis (IBR) |
| Study Purpose | To demonstrate efficacy against respiratory disease caused by IBR 182 days after vaccination. |
| Product Administration | |
| Study Animals | Bovine |
| Challenge Description | |
| Interval observed after challenge | |
| Results | Study data were evaluated by USDA-APHIS prior to product licensure and met regulatory standards for acceptance at the time of submission. No data are published because this study was submitted to USDA-APHIS prior to January 1, 2007, and APHIS only requires publication of data submitted after that date. |
| USDA Approval Date | March 30, 2004 |

| | |
|--|---|
| Study Type | Efficacy |
| Pertaining to | <i>Leptospira canicola</i> |
| Study Purpose | To demonstrate effectiveness against disease caused by <i>L. canicola</i> |
| Product Administration | Subcutaneous |
| Study Animals | Bovine |
| Challenge Description | |
| Interval observed after challenge | |
| Results | Study data were evaluated by USDA-APHIS prior to product licensure and met regulatory standards for acceptance at the time of submission. No data are published because this study was submitted to USDA-APHIS prior to January 1, 2007, and APHIS only requires publication of data submitted after that date. |
| USDA Approval Date | March 10, 2005 |

| | |
|--|---|
| Study Type | Efficacy |
| Pertaining to | <i>Leptospira grippotyphosa</i> |
| Study Purpose | To demonstrate effectiveness against disease caused by <i>L. grippotyphosa</i> |
| Product Administration | Subcutaneous |
| Study Animals | Bovine |
| Challenge Description | |
| Interval observed after challenge | |
| Results | Study data were evaluated by USDA-APHIS prior to product licensure and met regulatory standards for acceptance at the time of submission. No data are published because this study was submitted to USDA-APHIS prior to January 1, 2007, and APHIS only requires publication of data submitted after that date. |
| USDA Approval Date | March 10, 2005 |

| | |
|--|---|
| Study Type | Efficacy |
| Pertaining to | <i>Leptospira hardjo</i> |
| Study Purpose | To demonstrate effectiveness against disease caused by <i>L. hardjo</i> |
| Product Administration | Subcutaneous |
| Study Animals | Bovine |
| Challenge Description | |
| Interval observed after challenge | |
| Results | Study data were evaluated by USDA-APHIS prior to product licensure and met regulatory standards for acceptance at the time of submission. No data are published because this study was submitted to USDA-APHIS prior to January 1, 2007, and APHIS only requires publication of data submitted after that date. |
| USDA Approval Date | May 24, 2004 |

| Study Type | Efficacy | | | | | | | | | | | | |
|--|--|------------|--------------|------------|-------------|------------|----|---|---|----------|----|----|-----|
| Pertaining to | <i>Leptospira hardjo</i> | | | | | | | | | | | | |
| Study Purpose | To demonstrate effectiveness against urinary shedding of <i>L. hardjo</i> | | | | | | | | | | | | |
| Product Administration | 1 dose administered by the subcutaneous route | | | | | | | | | | | | |
| Study Animals | 29 calves, > 6 months of age, 14 vaccinates and 15 controls | | | | | | | | | | | | |
| Challenge Description | All calves were challenged with <i>L. hardjo</i> 21 days after vaccination. | | | | | | | | | | | | |
| Interval observed after challenge | All calves were monitored daily for 30 days for clinical signs of disease. Urine samples were taken on days 16 (Week 2), 23 (Week 3) and 30 (Week 4) post challenge for <i>Leptospira</i> isolation. | | | | | | | | | | | | |
| Results | <p><u>Leptospira Isolation:</u> A calf was considered positive if leptospire were isolated from the urine during any of the four post-challenge weeks.</p> <table border="1"> <thead> <tr> <th>Group</th> <th># of Animals</th> <th># Positive</th> <th>Percent (%)</th> </tr> </thead> <tbody> <tr> <td>Vaccinates</td> <td>14</td> <td>0</td> <td>0</td> </tr> <tr> <td>Controls</td> <td>15</td> <td>15</td> <td>100</td> </tr> </tbody> </table> <p>Raw data shown on the attached page.</p> | Group | # of Animals | # Positive | Percent (%) | Vaccinates | 14 | 0 | 0 | Controls | 15 | 15 | 100 |
| Group | # of Animals | # Positive | Percent (%) | | | | | | | | | | |
| Vaccinates | 14 | 0 | 0 | | | | | | | | | | |
| Controls | 15 | 15 | 100 | | | | | | | | | | |
| USDA Approval Date | March 27, 2012 | | | | | | | | | | | | |

L. hardjo Urine Isolations

| Group | Animal No. | Post-challenge Week | | |
|---------|------------|---------------------|---|---|
| | | 2 | 3 | 4 |
| Control | 507 | + | - | - |
| | 509 | + | + | - |
| | 510 | + | + | + |
| | 517 | + | + | + |
| | 518 | + | + | - |
| | 519 | + | - | - |
| | 520 | - | + | - |
| | 521 | - | + | - |
| | 523 | + | - | - |
| | 526 | + | - | - |
| | 527 | + | + | - |
| | 529 | + | - | - |
| | 531 | - | + | - |
| | 533 | + | + | - |
| | 540 | + | + | - |
| | Vaccinated | 508 | - | - |
| 511 | | - | - | - |
| 512 | | - | - | - |
| 514 | | - | - | - |
| 516 | | - | - | - |
| 522 | | - | - | - |
| 524 | | - | - | - |
| 525 | | - | - | - |
| 532 | | - | - | - |
| 536 | | - | - | - |
| 537 | | - | - | - |
| 541 | | - | - | - |
| 542 | | - | - | - |
| 543 | - | - | - | |

| | |
|--|---|
| Study Type | Efficacy |
| Pertaining to | <i>Leptospira icterohaemorrhagiae</i> |
| Study Purpose | To demonstrate effectiveness against disease caused by <i>L. icterohaemorrhagiae</i> |
| Product Administration | Subcutaneous |
| Study Animals | Bovine |
| Challenge Description | |
| Interval observed after challenge | |
| Results | Study data were evaluated by USDA-APHIS prior to product licensure and met regulatory standards for acceptance at the time of submission. No data are published because this study was submitted to USDA-APHIS prior to January 1, 2007, and APHIS only requires publication of data submitted after that date. |
| USDA Approval Date | June 22, 2004 |

| | |
|--|---|
| Study Type | Efficacy |
| Pertaining to | <i>Leptospira pomona</i> |
| Study Purpose | To demonstrate effectiveness against disease caused by <i>L. pomona</i> |
| Product Administration | Subcutaneous |
| Study Animals | Bovine |
| Challenge Description | |
| Interval observed after challenge | |
| Results | Study data were evaluated by USDA-APHIS prior to product licensure and met regulatory standards for acceptance at the time of submission. No data are published because this study was submitted to USDA-APHIS prior to January 1, 2007, and APHIS only requires publication of data submitted after that date. |
| USDA Approval Date | May 24, 2004 |

| | |
|--|---|
| Study Type | Efficacy |
| Pertaining to | Parainfluenza ₃ Virus (PI3) |
| Study Purpose | To demonstrate effectiveness against shedding caused by PI3 |
| Product Administration | Subcutaneous |
| Study Animals | Bovine |
| Challenge Description | |
| Interval observed after challenge | |
| Results | Study data were evaluated by USDA-APHIS prior to product licensure and met regulatory standards for acceptance at the time of submission. No data are published because this study was submitted to USDA-APHIS prior to January 1, 2007, and APHIS only requires publication of data submitted after that date. |
| USDA Approval Date | April 12, 2005 |

| | |
|--|---|
| Study Type | Efficacy |
| Pertaining to | Bovine Respiratory Syncytial Virus (BRSV) |
| Study Purpose | To demonstrate effectiveness against BRSV |
| Product Administration | Subcutaneous |
| Study Animals | Bovine |
| Challenge Description | |
| Interval observed after challenge | |
| Results | Study data were evaluated by USDA-APHIS prior to product licensure and met regulatory standards for acceptance at the time of submission. No data are published because this study was submitted to USDA-APHIS prior to January 1, 2007, and APHIS only requires publication of data submitted after that date. |
| USDA Approval Date | June 3, 2004 |

| Study Type | Safety | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
|--|--|--------------------------|--|----------------------------------|------------------------|--|--|------------|---------------------------------|----------------------------------|------------------------|---|-----------------|-----|---|---|---|---|--------------|-----|---|---|---|---|-----------------|-----|---|---|---|---|--------------|-----|---|---|---|---|-----------------|-----|---|---|---|---|--------------|-----|---|---|---|-----------|----------------------------------|------------------|--|--------------------------|--|---|-----------------|-----|---|---|--------------|-----|---|---|-----------------|-----|---|---|--------------|-----|---|---|-----------------|-----|----|---|--------------|-----|---|
| Pertaining to | NA | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Study Purpose | Demonstrate safety in pregnant cows or in calves nursing pregnant cows if previously vaccinated pre-breeding with same product | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Product Administration | <p>One dose of test product was administered subcutaneously 30-60 days before breeding to all study animals. Then the test or control product was given during the targeted trimester of gestation.</p> <ul style="list-style-type: none"> • <u>Test product</u>: Contains modified live virus fractions • <u>Control product</u>: Product of similar composition except virus fraction(s) are killed | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Study Animals | Pregnant cows: separate groups vaccinated at each trimester of gestation. Similar sized groups in each trimester were maintained as controls. | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Challenge Description | NA | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Interval observed after challenge | No challenge. After vaccination, observed daily through birth of calves. Calves monitored for 48 hours (bull calves) or 4 weeks (heifer calves). | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Results | <table border="1"> <thead> <tr> <th rowspan="2">Trimester</th> <th rowspan="2">Vaccinated During Gestation With</th> <th colspan="4">Number Pregnant Cows</th> </tr> <tr> <th>Vaccinated</th> <th>Removed from study^a</th> <th>Unrelated Abortions^b</th> <th>Abortions^c</th> </tr> </thead> <tbody> <tr> <td>1</td> <td>Control Product</td> <td>233</td> <td>7</td> <td>2</td> <td>2</td> </tr> <tr> <td>1</td> <td>Test Product</td> <td>235</td> <td>4</td> <td>3</td> <td>1</td> </tr> <tr> <td>2</td> <td>Control Product</td> <td>230</td> <td>0</td> <td>2</td> <td>1</td> </tr> <tr> <td>2</td> <td>Test Product</td> <td>231</td> <td>2</td> <td>6</td> <td>1</td> </tr> <tr> <td>3</td> <td>Control Product</td> <td>224</td> <td>2</td> <td>5</td> <td>1</td> </tr> <tr> <td>3</td> <td>Test Product</td> <td>216</td> <td>1</td> <td>8</td> <td>0</td> </tr> </tbody> </table> <p>^aRemovals affirmed by study cooperators to be unrelated to vaccination. ^bAbortions affirmed by study cooperators to have known cause, unrelated to infectious bovine rhinotracheitis or bovine virus diarrhea viruses ^cAbortions with unknown causes OR related to infectious bovine rhinotracheitis or bovine virus diarrhea viruses</p> <table border="1"> <thead> <tr> <th rowspan="2">Trimester</th> <th rowspan="2">Vaccinated During Gestation With</th> <th colspan="2">Number of Calves</th> </tr> <tr> <th>Live Births^a</th> <th>Mortality Prior to End of Observation Period</th> </tr> </thead> <tbody> <tr> <td>1</td> <td>Control Product</td> <td>224</td> <td>4</td> </tr> <tr> <td>1</td> <td>Test Product</td> <td>229</td> <td>6</td> </tr> <tr> <td>2</td> <td>Control Product</td> <td>233</td> <td>5</td> </tr> <tr> <td>2</td> <td>Test Product</td> <td>230</td> <td>2</td> </tr> <tr> <td>3</td> <td>Control Product</td> <td>223</td> <td>14</td> </tr> <tr> <td>3</td> <td>Test Product</td> <td>221</td> <td>5</td> </tr> </tbody> </table> <p>^aLive births include twins</p> | Trimester | Vaccinated During Gestation With | Number Pregnant Cows | | | | Vaccinated | Removed from study ^a | Unrelated Abortions ^b | Abortions ^c | 1 | Control Product | 233 | 7 | 2 | 2 | 1 | Test Product | 235 | 4 | 3 | 1 | 2 | Control Product | 230 | 0 | 2 | 1 | 2 | Test Product | 231 | 2 | 6 | 1 | 3 | Control Product | 224 | 2 | 5 | 1 | 3 | Test Product | 216 | 1 | 8 | 0 | Trimester | Vaccinated During Gestation With | Number of Calves | | Live Births ^a | Mortality Prior to End of Observation Period | 1 | Control Product | 224 | 4 | 1 | Test Product | 229 | 6 | 2 | Control Product | 233 | 5 | 2 | Test Product | 230 | 2 | 3 | Control Product | 223 | 14 | 3 | Test Product | 221 | 5 |
| Trimester | Vaccinated During Gestation With | | | Number Pregnant Cows | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| | | Vaccinated | Removed from study ^a | Unrelated Abortions ^b | Abortions ^c | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| 1 | Control Product | 233 | 7 | 2 | 2 | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| 1 | Test Product | 235 | 4 | 3 | 1 | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| 2 | Control Product | 230 | 0 | 2 | 1 | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| 2 | Test Product | 231 | 2 | 6 | 1 | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| 3 | Control Product | 224 | 2 | 5 | 1 | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| 3 | Test Product | 216 | 1 | 8 | 0 | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Trimester | Vaccinated During Gestation With | Number of Calves | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| | | Live Births ^a | Mortality Prior to End of Observation Period | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| 1 | Control Product | 224 | 4 | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| 1 | Test Product | 229 | 6 | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| 2 | Control Product | 233 | 5 | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| 2 | Test Product | 230 | 2 | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| 3 | Control Product | 223 | 14 | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| 3 | Test Product | 221 | 5 | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| USDA Approval Date | May 9, 2013 | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |

| | |
|--|---|
| Study Type | Safety |
| Pertaining to | ALL |
| Study Purpose | To demonstrate safety under field conditions. |
| Product Administration | |
| Study Animals | Bovine |
| Challenge Description | |
| Interval observed after challenge | |
| Results | Study data were evaluated by USDA-APHIS prior to product licensure and met regulatory standards for acceptance at the time of submission. No data are published because this study was submitted to USDA-APHIS prior to January 1, 2007, and APHIS only requires publication of data submitted after that date. |
| USDA Approval Date | June 16, 2004 |

| | |
|--|---|
| Study Type | Safety |
| Pertaining to | ALL |
| Study Purpose | To demonstrate safety in pregnant animals under field conditions when cows or heifers are vaccinated prior to breeding, within the previous 12 months, with a modified live Infectious Bovine Rhinotracheitis Virus (IBRV) and Bovine Viral Diarrhea Virus (BVDV) product |
| Product Administration | Two doses, administered subcutaneously. First vaccination given 14 to 60 days prior to breeding. Second vaccination given during a specified trimester of pregnancy. |
| Study Animals | 1 st Trimester Study: 468 pregnant heifers (52 – 86 days pregnant) 2 years of age and older. 2 nd Trimester Study: 461 pregnant heifers (100 – 180 days pregnant) 2 – 14 years of age. 3 rd Trimester Study: 440 pregnant heifers (≥190 days pregnant) 2 years of age and older. |
| Challenge Description | Not applicable |
| Interval observed after challenge | All cows were observed from pre-breeding vaccination through calving. |
| Results | Summary of the results listed in the table below |
| USDA Approval Date | May 9, 2013 |

Summary of the results as follows:

| Trimester | Group | No. of Cows | | Fetal Loss (%) related to vaccination | Fetal Loss (%) unrelated to vaccination as affirmed by licensee |
|-----------------|------------|-------------|----------|---------------------------------------|---|
| | | Entered | Removed* | | |
| 1 st | Vaccinates | 235 | 4 | 1 (0.4 %) | 3 (1.3%) |
| | Controls | 233 | 7 | 2 (0.9%) | 2 (0.9%) |
| 2 nd | Vaccinates | 231 | 2 | 1 (0.4%) | 6 (2.5%) |
| | Controls | 230 | 0 | 1 (0.4%) | 2 (0.8%) |
| 3 rd | Vaccinates | 216 | 1 | 0 (0%) | 8 (3.7%) |
| | Controls | 224 | 2 | 1 (0.5%) | 5 (2.2%) |

*Number of cows removed from the study results due to death serious illness considered unrelated to vaccination as affirmed by licensee