



Summary of Studies Supporting USDA Product Licensure

| | |
|---|--|
| Establishment Name | Boehringer Ingelheim Animal Health USA Inc. |
| USDA Vet Biologics Establishment Number | 124 |
| Product Code | 48W5.A3 |
| True Name | Encephalomyelitis-West Nile Virus Vaccine, Eastern & Western, Killed Virus, Tetanus Toxoid |
| Tradename(s) / Distributor or Subsidiary (if different from manufacturer) | Equi-Jec WNV+EWT - No distributor specified Vetera EWT + WNV - No distributor specified |
| Date of Compilation Summary | December 02, 2020 |

Disclaimer: Do not use the following studies to compare one product to another. Slight differences in study design and execution can render the comparisons meaningless.

| | |
|--|--|
| Study Type | Efficacy |
| Pertaining to | Clostridium tetanus |
| Study Purpose | Demonstration of efficacy against Clostridium tetanus |
| Product Administration | One dose, administered intramuscularly |
| Study Animals | 10 guinea pigs (10 vaccinates) |
| Challenge Description | Not applicable |
| Interval observed after challenge | Not applicable |
| Results | <p>Vaccinate serum samples were collected and pooled, then tested for antitoxin content by indirect Enzyme-Linked Immunosorbent Assay.</p> <p>A satisfactory value which met the requirements per 9 CFR 113.114(c) was achieved.</p> |
| USDA Approval Date | April 18, 2008 |

| | |
|--|---|
| Study Type | Efficacy |
| Pertaining to | Eastern equine encephalomyelitis |
| Study Purpose | Demonstration of efficacy against Eastern equine encephalomyelitis |
| Product Administration | Two doses, administered intramuscularly, 14 to 21 days apart |
| Study Animals | 12 guinea pigs (10 vaccinates, 2 controls) |
| Challenge Description | Not applicable |
| Interval observed after challenge | Not applicable |
| Results | <p>Serum samples were tested by a plaque reduction, serum neutralization test, 14 days after the second injection.</p> <p>Vaccinates and controls were evaluated in terms of Eastern equine encephalomyelitis per the criteria in 9 CFR 113.207(b) and the requirements were met.</p> |
| USDA Approval Date | April 18, 2008 |

| | |
|--|---|
| Study Type | Efficacy |
| Pertaining to | Western equine encephalomyelitis |
| Study Purpose | Demonstration of efficacy against Western equine encephalomyelitis |
| Product Administration | Two doses, administered intramuscularly, 14 to 21 days apart |
| Study Animals | 12 guinea pigs (10 vaccinates, 2 controls) |
| Challenge Description | Not applicable |
| Interval observed after challenge | Not applicable |
| Results | <p>Serum samples were tested by a plaque reduction, serum neutralization test, 14 days after the second injection.</p> <p>Vaccinates and controls were evaluated in terms of Western equine encephalomyelitis per the criteria in 9 CFR 113.207(b) and the requirements were met.</p> |
| USDA Approval Date | April 18, 2008 |

| Study Type | Efficacy | | | | | | | | | |
|--|---|-------------------|-----------------|-------------------|-----------|------------|-----------|--------------------------|--------------|------------|
| Pertaining to | West Nile Virus (WNV) | | | | | | | | | |
| Study Purpose | Demonstration of twelve month duration of immunity against disease caused by WNV | | | | | | | | | |
| Product Administration | Two doses, administered intramuscularly, 25 days apart | | | | | | | | | |
| Study Animals | 30 horses (20 vaccinates, 10 placebo controls) 4-5 months of age | | | | | | | | | |
| Challenge Description | West Nile Virus was administered at 380 days (10 vaccinated and 5 placebo control animals) or 408 days (10 vaccinated and 5 placebo control animals) post-final vaccination. | | | | | | | | | |
| Interval observed after challenge | Horses were observed twice daily for 14 days post-challenge and once daily for an additional 7 days post-challenge. | | | | | | | | | |
| Results | <p>An animal was considered affected by challenge if it developed neurological disease, as measured by mortality and microscopic evidence of virus-induced brain disease (histopathology).</p> <p>Animals were also monitored for viremia (detection of WNV in the blood).</p> <p>Results are summarized as follows:</p> <table border="1"> <thead> <tr> <th>Outcome</th> <th>Controls</th> <th>Vaccinates</th> </tr> </thead> <tbody> <tr> <td>Mortality</td> <td>7/10 (70%)</td> <td>1/20 (5%)</td> </tr> <tr> <td>Viremia at least one day</td> <td>10/10 (100%)</td> <td>2/20 (10%)</td> </tr> </tbody> </table> <p>See raw data on following pages.</p> | Outcome | Controls | Vaccinates | Mortality | 7/10 (70%) | 1/20 (5%) | Viremia at least one day | 10/10 (100%) | 2/20 (10%) |
| Outcome | Controls | Vaccinates | | | | | | | | |
| Mortality | 7/10 (70%) | 1/20 (5%) | | | | | | | | |
| Viremia at least one day | 10/10 (100%) | 2/20 (10%) | | | | | | | | |
| USDA Approval Date | September 3, 2010 | | | | | | | | | |

| Treatment | # | Died or Euthanized due to disease severity | Severity Histopathological lesions | |
|------------------------|----|--|------------------------------------|------|
| | | | Medulla | Pons |
| Controls (10 horses) | 1 | Yes | 3 | 3 |
| | 2 | Yes | 3 | 3 |
| | 3 | Yes | 3 | 3 |
| | 4 | Yes | 3 | 3 |
| | 5 | Yes | 3 | 3 |
| | 6 | Yes | 2 | 2 |
| | 7 | Yes | 1 | 1 |
| | 8 | No | 1 | 1 |
| | 9 | No | 1 | 1 |
| | 10 | No | 1 | 0.5 |
| Vaccinates (20 horses) | 1 | Yes | 3 | 3 |
| | 2 | No | 2 | 0.5 |
| | 3 | No | 1 | 1 |
| | 4 | No | 1 | 0.5 |
| | 5 | No | 1 | 0.5 |
| | 6 | No | 1 | 0.5 |
| | 7 | No | 0.5 | 0.5 |
| | 8 | No | 0.5 | 0.5 |
| | 9 | No | 0.5 | 0 |
| | 10 | No | 0 | 0.5 |
| | 11 | No | 0 | 0 |
| | 12 | No | 0 | 0 |
| | 13 | No | 0 | 0 |
| | 14 | No | 0 | 0 |
| 15 | No | 0 | 0 | |
| 16 | No | 0 | 0 | |
| 17 | No | 0 | 0 | |
| 18 | No | 0 | 0 | |
| 19 | No | 0 | 0 | |
| 20 | No | 0 | 0 | |

| Scoring of histopathological lesions: | |
|---------------------------------------|--|
| 0 = | No significant lesions. |
| 0.5 = | Rare, small, multifocal glial nodules scattered throughout the parenchyma. |
| 1 = | Mild, nonsuppurative encephalitis characterized by mild multifocal perivascular cuffs with lymphocytes and plasma cells and a rare neutrophil and scattered multifocal glial nodules composed of glial cells with a few mononuclear inflammatory cells. Occasionally within this grade, there may be minimal perivascular cuffing and more moderate scattered glial nodules. |
| 2 = | Moderate nonsuppurative encephalitis characterized by moderate lymphoplasmacytic perivascular cuffs around many vessels and multifocal accumulations of glial nodules scattered throughout the parenchyma. |
| 3 = | Severe nonsuppurative encephalitis characterized by severe and thick lymphoplasmacytic perivascular cuffing with multiple scattered glial nodules throughout the parenchyma. |

| Treatment | # | Days Post-challenge | | | | | | | | | | | | | | | | | | | | |
|---------------------------|----|---------------------|----|-----|-----|-----|-----|-----|-----|-----|-----|----|----|---|--|---|---|---|----|----|----|---|
| | | 0 | | 1 | | 2 | | 3 | | 4 | | 5 | | 6 | | 7 | 8 | 9 | 10 | 14 | 21 | |
| | | AM | PM | AM | PM | AM | PM | AM | PM | AM | PM | AM | PM | | | | | | | | | |
| Controls (10 horses) | 1 | | | | | | | | | | | | | | | | | | | | | |
| | 2 | | | 15 | 25 | 30 | 5 | 20 | | | | | | | | | | | | | D | D |
| | 3 | | | 25 | 470 | 160 | 210 | 175 | 75 | 100 | | | | | | | | | | | D | D |
| | 4 | | | 5 | 20 | 205 | 175 | 90 | 75 | 50 | | | | | | | | | | | D | D |
| | 5 | | | 40 | 130 | 30 | 25 | 50 | 10 | | | | | | | | | | | | D | D |
| | 6 | | | 165 | 110 | 65 | 55 | 10 | | | | | | | | | | | | | D | D |
| | 7 | | | 10 | 330 | 200 | 180 | 225 | 115 | 15 | | | | | | | | | | | N | D |
| | 8 | | | | | 50 | 90 | 35 | 105 | 20 | 10 | | | | | | | | | | | |
| | 9 | | | | | 5 | 30 | 95 | 25 | 135 | 240 | 40 | 20 | | | | | | | | | |
| | 10 | | | | | | | 80 | 70 | 40 | 10 | | | | | | | | | | | |
| Vaccinates (20 horses) | 1 | | | | | | | | | | | | | | | | | | | | | |
| | 2 | | | | | | | | | | | | | | | | | | | | | |
| | 3 | | | | | | | | | | | | | | | | | | | | | |
| | 4 | | | | | | | | | | | | | | | | | | | | | |
| | 5 | | | | | | | | | | | | | | | | | | | | | |
| | 6 | | | | | | 95 | | | | | | | | | | | | | | | |
| | 7 | | | | | | | | | | | | | | | | | | | | | |
| | 8 | | | | | | 40 | | | | | | | | | | | | | | | |
| | 9 | | | | | | | | | | | | | | | | | | | | | |
| | 10 | | | | | | | | | | | | | | | | | | | | | |
| 11 | | | | | | | | | | | | | | | | | | | | | | |
| 12 | | | | | | | | | | | | | | | | | | | | | | |
| 13 | | | | | | | | | | | | | | | | | | | | | | |
| 14 | | | | | | | | | | | | | | | | | | | | | | |
| 15 | | | | | | | | | | | | | | | | | | | | | | |
| 16 | | | | | | | | | | | | | | | | | | | | | | |
| 17 | | | | | | | | | | | | | | | | | | | | | | |
| 18 | | | | | | | | | | | | | | | | | | | | | | |
| 19 | | | | | | | | | | | | | | | | | | | | | | |
| 20 | | | | | | | | | | | | | | | | | | | | | | |

Actual value in plaque-forming units per milliliter equivalents (PFUeq/mL) = Positive for virus isolation

Blank = Negative for virus isolation (<5 PFUeq/mL)

D = Dead

N = Not recorded; horse was circling with sporadic head / neck tremors.

| Study Type | Efficacy | | | | |
|--|---|----------|------------|-----------|-----------|
| Pertaining to | West Nile Virus | | | | |
| Study Purpose | Demonstration of efficacy against WNV | | | | |
| Product Administration | Two doses, administered intramuscularly 21 days apart | | | | |
| Study Animals | 28 horses (19 vaccinates, 9 placebo controls) 4-5 months of age | | | | |
| Challenge Description | West Nile Virus was administered intrathecally at 14 days (to 10 vaccinated and 5 placebo control animals) and 28 days (to 9 vaccinated and 4 placebo control animals) after the second vaccination | | | | |
| Interval observed after challenge | Horses were bled on the day of challenge, twice daily for 6 days post-challenge, once daily for an additional 4 days post-challenge, and on day 14 post-challenge | | | | |
| Results | <p>The primary outcome was viremia (detection of WNV in the blood). While the test method was quantitative, an animal was considered to be positive (affected by challenge) if any virus was detected in the blood on one or more occasions post-challenge.</p> <p>The number of animals positive for (affected by) viremia at least once is summarized as follows:</p> <table border="1" data-bbox="805 1070 1177 1146"> <thead> <tr> <th>Controls</th> <th>Vaccinates</th> </tr> </thead> <tbody> <tr> <td>8/9 (89%)</td> <td>1/19 (5%)</td> </tr> </tbody> </table> <p>See raw data on the following page.</p> | Controls | Vaccinates | 8/9 (89%) | 1/19 (5%) |
| Controls | Vaccinates | | | | |
| 8/9 (89%) | 1/19 (5%) | | | | |
| USDA Approval Date | August 25, 2008 | | | | |

| Viremia: | | Days Post-Challenge | | | | | | | | | | | | | | | | | | |
|------------------|----|---------------------|-----|------|-----|-----|-----|-----|----|----|----|----|----|----|----|----|---|---|----|----|
| | | 0 | | 1 | | 2 | | 3 | | 4 | | 5 | | 6 | | 7 | 8 | 9 | 10 | 14 |
| | | Horse ID | AM | PM | AM | PM | AM | PM | AM | PM | AM | PM | AM | PM | AM | PM | | | | |
| Treatment | 13 | 15 | 20 | 390 | 280 | 135 | | 20 | | | | | | | | | | | | |
| | 19 | | 5 | | 40 | | | 5 | 20 | 65 | 45 | | | | | | | | | |
| | 20 | | 125 | 1475 | 645 | 355 | 495 | 120 | 15 | | | | | | | | | | | |
| | 45 | | 20 | 85 | 235 | 140 | 235 | 145 | 80 | 15 | | | | | | | | | | |
| | 67 | | | 165 | | | | | | | | | | | | | | | | |
| | 71 | | | 110 | 675 | 110 | 70 | 120 | 70 | | | | | | | | | | | |
| | 72 | | | | | | | | | | | | | | | | | | | D |
| | 74 | | | 5 | 60 | 30 | 15 | 20 | 15 | | | | | | | | | | | |
| | 79 | | | 10 | 15 | 20 | | 15 | | | 10 | 5 | | | | | | | | |
| 14 | | | | | | | | | | | | | | | | | | | | |
| 16 | | | | | | | | | | | | | | | | | | | | |
| 21 | | | | | | | | | | | | | | | | | | | | |
| 22 | | | | | | | | | | | | | | | | | | | | |
| 23 | | | | | | | | | | | | | | | | | | | | |
| 24 | | | | | | | | | | | | | | | | | | | | |
| 25 | | | | | | | | | | | | | | | | | | | | |
| 26 | | | | | | | | | | | | | | | | | | | | |
| 27 | | | | | | | | | | | | | | | | | | | | |
| 37 | | | | | | | | | | | | | | | | | | | | |
| 73 | | | | | | | | | | | | | | | | | | | | |
| 75 | | | | | | | | | | | | | | | | | | | | |
| 77 | | | | | | | | | | | | | | | | | | | | |
| 80 | | | | | | | | | | | | | | | | | | | | |
| 81 | | | | | | | | | | | | | | | | | | | | |
| 82 | | | | | | | | | | | | | | | | | | | | |
| 83 | | | | | | | | | | | | | | | | | | | | |
| 84 | | | | | | | | | | | | | | | | | | | | |
| 85 | | | | 5 | | 5 | | 5 | | | | | | | | | | | | |

Actual value in plaque-forming units per milliliter equivalents (PFUeq/mL) = Positive for virus isolation
Blank = Negative for virus isolation (<5 PFUeq/mL)
D = Dead (euthanized on Day 11 due to West Nile Virus)

| Study Type | Efficacy | | | | | | | | | | | | |
|--|--|-----------------|----------|------------|---|------------|------------|---|------------|------------|----------|--------------|------------|
| Pertaining to | West Nile Virus (WNV) | | | | | | | | | | | | |
| Study Purpose | Demonstration of six month duration of immunity against WNV | | | | | | | | | | | | |
| Product Administration | Two doses, administered intramuscularly 21 days apart | | | | | | | | | | | | |
| Study Animals | 30 horses (20 vaccinates, 10 placebo controls) 4-5 months of age | | | | | | | | | | | | |
| Challenge Description | Challenge by West Nile Virus was administered at 180 days (Challenge Group 1: 10 vaccinated and 5 placebo control animals) or 243 days (Challenge Group 2: 10 vaccinated and 5 placebo control animals) after second vaccination | | | | | | | | | | | | |
| Interval observed after challenge | Horses were bled on the day of challenge, twice daily for 6 days post-challenge, once daily for an additional 4 days post-challenge, and on day 14 post-challenge | | | | | | | | | | | | |
| Results | <p>The primary outcome was viremia (detection of WNV in the blood). While the test method was quantitative, an animal was considered to be positive (affected by challenge) if any virus was detected in the blood on one or more occasions post-challenge.</p> <p>The number of animals positive for viremia at least once (affected) is summarized as follows:</p> <table border="1"> <thead> <tr> <th>Challenge Group</th> <th>Controls</th> <th>Vaccinates</th> </tr> </thead> <tbody> <tr> <td>1</td> <td>5/5 (100%)</td> <td>2/10 (20%)</td> </tr> <tr> <td>2</td> <td>5/5 (100%)</td> <td>4/10 (40%)</td> </tr> <tr> <td>Combined</td> <td>10/10 (100%)</td> <td>6/20 (30%)</td> </tr> </tbody> </table> <p>See raw data on the following page.</p> | Challenge Group | Controls | Vaccinates | 1 | 5/5 (100%) | 2/10 (20%) | 2 | 5/5 (100%) | 4/10 (40%) | Combined | 10/10 (100%) | 6/20 (30%) |
| Challenge Group | Controls | Vaccinates | | | | | | | | | | | |
| 1 | 5/5 (100%) | 2/10 (20%) | | | | | | | | | | | |
| 2 | 5/5 (100%) | 4/10 (40%) | | | | | | | | | | | |
| Combined | 10/10 (100%) | 6/20 (30%) | | | | | | | | | | | |
| USDA Approval Date | October 21, 2009 | | | | | | | | | | | | |

| Viremia: | | Days Post-challenge | | | | | | | | | | | | | | Final Outcome | | | | | | | | | | | | | |
|---------------------------------------|----------|---------------------|----|-----|-----|-----|-----|-----|----|----|----|----|----|----|----|---------------|---|---|---|----|----|--|--|--|--|--|----------|----------|----------|
| | | 0 | | 1 | | 2 | | 3 | | 4 | | 5 | | 6 | | | 7 | 8 | 9 | 10 | 14 | | | | | | | | |
| | | AM | PM | AM | PM | AM | PM | AM | PM | AM | PM | AM | PM | AM | PM | | | | | | | | | | | | | | |
| Treatment | Horse ID | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Controls (5 horses) Challenge 1 | S2 | | | 170 | 170 | 165 | 55 | 105 | 45 | | | | | | | | | | | | | | | | | | Positive | | |
| | S4 | | | 425 | 20 | 30 | 40 | 85 | 60 | 65 | | | | | | | | | | | | | | | | | N | Positive | |
| | S10 | | | 10 | 300 | 125 | 125 | 80 | 45 | | | | | | | | | | | | | | | | | | | Positive | |
| | S11 | | | | 50 | 30 | 40 | 40 | 25 | | | | | | | | | | | | | | | | | | | Positive | |
| | S13 | | | | 410 | 110 | 135 | 110 | 55 | 15 | | | | | | | | | | | | | | | | | N | Positive | |
| Vaccinates (10 horses) Challenge 1 | S1 | | | | | | | | | | | | | | | | | | | | | | | | | | | Negative | |
| | S3 | | | | | | | | | | | | | | | | | | | | | | | | | | | | Negative |
| | S5 | | | | | | | | | | | | | | | | | | | | | | | | | | | | Negative |
| | S6 | | | | | | | | | | | | | | | | | | | | | | | | | | | | Negative |
| | S7 | | | 470 | | 45 | 5 | | | | | | | | | | | | | | | | | | | | | Positive | |
| | S8 | | | 15 | | | | | | | | | | | | | | | | | | | | | | | | Positive | |
| | S9 | | | | | | | | | | | | | | | | | | | | | | | | | | N | Negative | |
| | S12 | | | | | | | | | | | | | | | | | | | | | | | | | | | | Negative |
| | S14 | | | | | | | | | | | | | | | | | | | | | | | | | | | | Negative |
| | S15 | | | | | | | | | | | | | | | | | | | | | | | | | | | | Negative |
| Controls (5 horses) Challenge 2 | S50 | | | 5 | 535 | 500 | 80 | 100 | 50 | 10 | | | | | | | | | | | | | | | | | | Positive | |
| | S53 | | | 20 | 320 | 380 | 100 | 135 | 45 | 10 | | | | | | | | | | | | | | | | | | Positive | |
| | S54 | | | | | | | | 5 | | 5 | 5 | 5 | | | | | | | | | | | | | | N | Positive | |
| | S55 | | | 10 | 95 | 70 | 30 | 25 | 40 | | | | | | | | | | | | | | | | | | | Positive | |
| | S59 | | | | 90 | 265 | 20 | 70 | 45 | 45 | 5 | | | | | | | | | | | | | | | | | Positive | |
| Vaccinates (10 horses) Challenge 2 | S46 | | | | | | | | | | | | | | | | | | | | | | | | | | | Negative | |
| | S47 | | | | | | | | | | | | | | | | | | | | | | | | | | | Positive | |
| | S48 | | | | | | | | | 5 | | | | | | | | | | | | | | | | | | Negative | |
| | S49 | | | | | | | | | | | | | | | | | | | | | | | | | | | Positive | |
| | S51 | | | | | | | | 15 | | | | | | | | | | | | | | | | | | | Negative | |
| | S52 | | | | | | | | | | | | | | | | | | | | | | | | | | | Negative | |
| | S56 | | | | | | | | | | | | | | | | | | | | | | | | | | | Negative | |
| | S57 | | | | | | | | | 5 | 5 | | | | | | | | | | | | | | | | | Positive | |
| | S58 | | | | | | | | | | 5 | | | | | | | | | | | | | | | | | Positive | |
| | S60 | | | | | | | | | | | | | | | | | | | | | | | | | | | | Negative |

Actual value in plaque-forming units per milliliter equivalents (PFU eq/mL) = Positive for virus isolation

Blank = Negative for virus isolation (<5 PFU eq/mL)

N = Not recorded

Positive = affected by challenge if virus was detected in the blood on one or more occasions post-challenge.

Negative = virus was detected in the blood on zero occasions post-challenge.

| | |
|--|---|
| Study Type | Safety |
| Pertaining to | All fractions |
| Study Purpose | To demonstrate safety in pregnant mares under field conditions at two different test sites |
| Product Administration | Two intramuscular doses, given 16-28 days apart. 54 pregnant mares were injected with placebo and 325 pregnant mares were vaccinated with test product. |
| Study Animals | Three hundred seventy-nine pregnant mares at two locations were included in the study. The mares were confirmed to be pregnant by serum hormonal evaluation on the day of the first vaccination. |
| Challenge Description | Not applicable |
| Interval observed after vaccination | 1 st and 2 nd trimester: Mares observed immediately after vaccination and daily for overall health and for abortion. Resulting foals were observed daily for 7 days following birth. 3 rd trimester: Mares observed immediately after vaccination and daily for overall health and for abortion. Resulting foals were observed daily for 30 days following birth. |
| Results | Results shown on next page |

Results

Study 2013-PM-1009

North Dakota Site:

| Group | Vaccinated | Confirmed Pregnant | Foals | Parturition Rate |
|---------------------------------------|------------|--------------------|------------|------------------|
| 1 st trimester/ product | 143 | 127 | 114 | 90% |
| 1st trimester/ placebo | 59 | 54 | 49 | 91% |
| 2 nd trimester/ product | 6 | 6 | 6 | 100% |
| 3 rd trimester/ product | 140 | 117 | 117 | 100% |
| Total – all animals | 348 | 304 | 286 | 94% |
| Total – product only | 289 | 250 | 237 | 95% |
| Total – placebo only | 59 | 54 | 49 | 91% |

Study 2013-PM-1009

Missouri Site:

| Group | Vaccinated | Confirmed Pregnant | Foals | Parturition Rate |
|-----------------------------------|------------|--------------------|-----------|------------------|
| 2011 3 rd trimester | 5 | 5 | 5 | 100% |
| 2012 1 st trimester | 1 | 1 | 1 | 100% |
| 2012 2 nd trimester | 53 | 43 | 39 | 91% |
| 2012 3 rd trimester | 26 | 26 | 25 | 96% |
| Total – product | 85 | 75 | 70 | 93% |

Study 2014-PM-1009

North Dakota Site:

| Group | Vaccinated | Confirmed Pregnant | Foaled | Parturition Rate | Foals Survived to End of Observation Period |
|--|------------|--------------------|--------|------------------|---|
| 2 nd trimester vaccinated | 52 | 52 | 52 | 100% | 51* |
| 3 rd trimester vaccinated | 69 | 69 | 67** | 97.1% | 67 |

*Lost foal affirmed by study cooperator to be due to causes other than vaccination.

**One mare died due to causes other than vaccination, as affirmed by study cooperator.

All other foals were normal and healthy

USDA Approval Date

September 12, 2014

| Study Type | Safety | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
|--|--|----------------------------|----------------------------|---|---|---|-----------------------------|-----------------------------|----------------------------|----------------------------|----------------------------|----------------------------|----------|-----|-----|-------------|-------------|----------------|----------------|----------|-----|-----|-------------|-------------|----------------|----------------|-------|-----|-----|-----------|-----------|---------------|---------------|--------------|------------|------------|---------------------|----------------------|------------------------|------------------------|
| Pertaining to | All fractions | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Study Purpose | To demonstrate safety under field conditions | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Product Administration | Two doses, administered intramuscularly 3 – 4 weeks apart | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Study Animals | 556 horses, including 438 foals between 2 months and approximately 1 year of age | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Challenge Description | Not applicable | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Interval observed after challenge | Not applicable | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Results | <p>Horses were observed at least daily following each vaccination, until resolution of any observed reactions. Observations ended 14 days after the second vaccination.</p> <p>There were no systemic reactions observed at any of the sites. One horse died from causes affirmed by licensee not associated with vaccination.</p> <p>Scoring Method for Injection Site Reactions: 0 = No reaction 1 = Localized swelling at or near the injection site which is not visible; detectable only by palpation. Not painful. 2 = Localized visible swelling at or near the injection site. Not painful. 3 = Localized visible swelling at or near the injection site. Raised, circumscribed and painful when palpated.</p> <p>Local injection site reactions are summarized below across the sites:</p> <table border="1"> <thead> <tr> <th rowspan="2">Site</th> <th rowspan="2">Total Number Of Vaccinates</th> <th rowspan="2">Number Of Vaccinates Administered 2 doses</th> <th colspan="2">Vaccinates With Transient Injection Site Swelling</th> <th colspan="2">Number Of Normal Vaccinates</th> </tr> <tr> <th>After 1st dose</th> <th>After 2nd dose</th> <th>After 1st dose</th> <th>After 2nd dose</th> </tr> </thead> <tbody> <tr> <td>Missouri</td> <td>315</td> <td>314</td> <td>3 (1.0%)</td> <td>9 (2.9%)</td> <td>312 (99.0%)</td> <td>305 (97.1%)</td> </tr> <tr> <td>Oklahoma</td> <td>110</td> <td>110</td> <td>1 (0.9%)</td> <td>2 (1.8%)</td> <td>109 (99.1%)</td> <td>108 (98.2%)</td> </tr> <tr> <td>Texas</td> <td>131</td> <td>131</td> <td>0 (0%)</td> <td>0 (0%)</td> <td>131 (100%)</td> <td>131 (100%)</td> </tr> <tr> <td>Total</td> <td>556</td> <td>555</td> <td>4 (0.7%)</td> <td>11 (2.0%)</td> <td>552 (99.3%)</td> <td>544 (98.0%)</td> </tr> </tbody> </table> <p>Results from each site are summarized on the following page.</p> | Site | Total Number Of Vaccinates | Number Of Vaccinates Administered 2 doses | Vaccinates With Transient Injection Site Swelling | | Number Of Normal Vaccinates | | After 1 st dose | After 2 nd dose | After 1 st dose | After 2 nd dose | Missouri | 315 | 314 | 3 (1.0%) | 9 (2.9%) | 312 (99.0%) | 305 (97.1%) | Oklahoma | 110 | 110 | 1 (0.9%) | 2 (1.8%) | 109 (99.1%) | 108 (98.2%) | Texas | 131 | 131 | 0 (0%) | 0 (0%) | 131 (100%) | 131 (100%) | Total | 556 | 555 | 4 (0.7%) | 11 (2.0%) | 552 (99.3%) | 544 (98.0%) |
| Site | Total Number Of Vaccinates | | | | Number Of Vaccinates Administered 2 doses | Vaccinates With Transient Injection Site Swelling | | Number Of Normal Vaccinates | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| | | After 1 st dose | After 2 nd dose | After 1 st dose | | After 2 nd dose | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Missouri | 315 | 314 | 3 (1.0%) | 9 (2.9%) | 312 (99.0%) | 305 (97.1%) | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Oklahoma | 110 | 110 | 1 (0.9%) | 2 (1.8%) | 109 (99.1%) | 108 (98.2%) | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Texas | 131 | 131 | 0 (0%) | 0 (0%) | 131 (100%) | 131 (100%) | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Total | 556 | 555 | 4 (0.7%) | 11 (2.0%) | 552 (99.3%) | 544 (98.0%) | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |

Missouri Site:

| Summary | Number Of Vaccinates | Number Of Vaccinates Administered 2 doses | Vaccinates With Transient Injection Site Swelling | | Number Of Normal Vaccinates | |
|--------------|----------------------|---|---|----------------------------|-----------------------------|----------------------------|
| | | | After 1 st dose | After 2 nd dose | After 1 st dose | After 2 nd dose |
| 2-4 months | 55 | 55 | 0 | 0 | 55 | 55 |
| 5-7 months | 8 | 8 | 0 | 0 | 8 | 8 |
| 8-11 months | 1 | 1 | 0 | 0 | 1 | 1 |
| 1 year | 170 | 170 | 1 | 2 | 169 | 168 |
| ≥ 2 years | 81 | 80 | 2 | 7 | 79 | 73 |
| Total | 315 | 314 | 3 | 9 | 312 | 305 |

| Horse No. | Age | Reaction Description | Injection # | Day | Score | Resolution Day |
|-----------|------|---|-------------|-----|-------|----------------|
| 10 | 11 y | Swelling on day 3, 5.5 cm x 2.25 cm x 5mm | 2 | 3 | 2 | 7 |
| 22 | 8 y | Swelling on day 3, 12 cm circle, raised 1.5 cm, painful, | 2 | 3 | 3 | 7 |
| 129 | 1 y | Swelling on day 3, 2.3 cm circle, raised 4 mm, painful but no heat | 2 | 3 | 3 | 7 |
| 183 | 1 y | Swelling on day 7, raised lesion 1.5 cm circle, height 0.2 cm | 1 | 7 | 2 | 14 |
| 183 | 1 y | Swelling on day 1, 3 cm lesion, not raised but palpable | 2 | 1 | 1 | 3 |
| 222 | 9 y | Swelling on day 3, 6 cm x 7 cm x 1.2 cm, raised lesion hard and painful | 2 | 3 | 3 | 7 |
| 266 | 10 y | Swelling localized in several places unsure if related to vaccine | 1 | 1 | 2 | 3 |
| 266 | 10 y | Swelling small palpable mass ~ 2 cm size, still present day 3 no worse | 2 | 1 | 2 | 3 |
| 271 | 13 y | Swelling 5cm circle, raised 5 mm, solid and painful | 2 | 3 | 3 | 7 |
| 288 | 8 y | Swelling < 2 cm, raised lesion ~ 1 mm deep | 1 | 3 | 2 | 7 |
| 288 | 8 y | Swelling ~ 8.5 cm circle raised ~ 1.3 cm, painful, not hot to touch | 2 | 3 | 3 | 7 |
| 300 | 10 y | Swelling 6 cm circle, solid swelling not painful | 2 | 3 | 2 | 7 |

Oklahoma Site:

| Summary | Number Of Vaccinates | Number Of Vaccinates Administered 2 doses | Vaccinates With Transient Injection Site Swelling | | Number Of Normal Vaccinates | |
|--------------|----------------------|---|---|----------------------------|-----------------------------|----------------------------|
| | | | After 1 st dose | After 2 nd dose | After 1 st dose | After 2 nd dose |
| 4-6 months | 49 | 49 | 1 | 1 | 48 | 48 |
| 1 year | 25 | 25 | 0 | 1 | 25 | 24 |
| ≥ 2 years | 36 | 36 | 0 | 0 | 36 | 36 |
| Total | 110 | 110 | 1 | 2 | 109 | 108 |

| Horse No. | Age | Reaction Description | Injection # | Day | Score | Resolution Day |
|-----------|-----|--|-------------|-----|-------|----------------|
| 19-A | 4 m | Swelling redness painful injection area 6 cm in diameter, reaction subsided in 10 days | 1 | 7 | 3 | 17 |
| 33-A | 5 m | Small swelling, 3 cm diameter, subsided in 3 days | 2 | 1 | 2 | 4 |
| 43 | 1 y | Mid-sized swelling, 5 cm diameter, reduced to 2.5 cm in 6 days; small, hard 2 cm at 10 days, probable subcutaneous leakage | 2 | 1 | 2 | Study End |

Texas Site:

| Summary | Number Of Vaccinates | Number Of Vaccinates Administered 2 doses | Vaccinates With Transient Injection Site Swelling | | Number Of Normal Vaccinates | |
|--------------|----------------------|---|---|----------------------------|-----------------------------|----------------------------|
| | | | After 1 st dose | After 2 nd dose | After 1 st dose | After 2 nd dose |
| 7-9 months | 130 | 130 | 0 | 0 | 130 | 130 |
| ≥ 2 years | 1 | 1 | 0 | 0 | 1 | 1 |
| Total | 131 | 131 | 0 | 0 | 131 | 131 |

USDA Approval Date

September 14, 2009