



## Summary of Studies Supporting USDA Product Licensure

Establishment Name	Boehringer Ingelheim Animal Health USA Inc.
USDA Vet Biologics Establishment Number	124
Product Code	4865.A1
True Name	Encephalomyelitis Vaccine, Eastern & Western, Killed Virus, Tetanus Toxoid
Tradename(s) / Distributor or Subsidiary (if different from manufacturer)	Vetera EWT - No distributor specified
Date of Compilation Summary	October 28, 2020

**Disclaimer: Do not use the following studies to compare one product to another. Slight differences in study design and execution can render the comparisons meaningless.**

<b>Study Type</b>	Efficacy
<b>Pertaining to</b>	Clostridium tetanus
<b>Study Purpose</b>	Demonstration of efficacy against Clostridium tetanus
<b>Product Administration</b>	One dose, administered intramuscularly
<b>Study Animals</b>	10 guinea pigs (10 vaccinates)
<b>Challenge Description</b>	Not applicable
<b>Interval observed after challenge</b>	Not applicable
<b>Results</b>	<p>Vaccinate serum samples were collected and pooled, then tested for antitoxin content by indirect Enzyme-Linked Immunosorbent Assay.</p> <p>A satisfactory value which met the requirements per 9 CFR 113.114(c) was achieved.</p>
<b>USDA Approval Date</b>	May 1, 2008

<b>Study Type</b>	Efficacy
<b>Pertaining to</b>	Eastern equine encephalomyelitis
<b>Study Purpose</b>	Demonstration of efficacy against Eastern equine encephalomyelitis
<b>Product Administration</b>	Two doses, administered intramuscularly, 14 to 21 days apart
<b>Study Animals</b>	12 guinea pigs (10 vaccinates, 2 controls)
<b>Challenge Description</b>	Not applicable
<b>Interval observed after challenge</b>	Not applicable
<b>Results</b>	<p>Serum samples were tested by a plaque reduction, serum neutralization test, 14 to 21 days after the second injection.</p> <p>Vaccinates and controls were evaluated in terms of Eastern equine encephalomyelitis per the criteria in 9 CFR 113.207(b) and the requirements were met.</p>
<b>USDA Approval Date</b>	May 1, 2008

<b>Study Type</b>	Efficacy
<b>Pertaining to</b>	Western equine encephalomyelitis
<b>Study Purpose</b>	Demonstration of efficacy against Western equine encephalomyelitis
<b>Product Administration</b>	Two doses, administered intramuscularly, 14 to 21 days apart
<b>Study Animals</b>	12 guinea pigs (10 vaccinates, 2 controls)
<b>Challenge Description</b>	Not applicable
<b>Interval observed after challenge</b>	Not applicable
<b>Results</b>	<p>Serum samples were tested by a plaque reduction, serum neutralization test, 14 to 21 days after the second injection.</p> <p>Vaccinates and controls were evaluated in terms of Western equine encephalomyelitis per the criteria in 9 CFR 113.207(b) and the requirements were met.</p>
<b>USDA Approval Date</b>	May 1, 2008

<b>Study Type</b>	Safety
<b>Pertaining to</b>	All fractions
<b>Study Purpose</b>	To demonstrate safety in pregnant mares under field conditions at two different test sites
<b>Product Administration</b>	Two intramuscular doses, given 16-28 days apart. 54 pregnant mares were injected with placebo and 325 pregnant mares were vaccinated with test product.
<b>Study Animals</b>	Three hundred seventy-nine pregnant mares at two locations were included in the study. The mares were confirmed to be pregnant by serum hormonal evaluation on the day of the first vaccination.
<b>Challenge Description</b>	Not applicable
<b>Interval observed after vaccination</b>	1 <sup>st</sup> and 2 <sup>nd</sup> trimester: Mares observed immediately after vaccination and daily for overall health and for abortion. Resulting foals were observed daily for 7 days following birth. 3 <sup>rd</sup> trimester: Mares observed immediately after vaccination and daily for overall health and for abortion. Resulting foals were observed daily for 30 days following birth.
<b>Results</b>	Results shown on next page

**Results****Study 2013-PM-1009****North Dakota Site:**

<b>Group</b>	<b>Vaccinated</b>	<b>Confirmed Pregnant</b>	<b>Foals</b>	<b>Parturition Rate</b>
1 <sup>st</sup> trimester/ product	143	127	114	90%
1st trimester/ placebo	59	54	49	91%
2 <sup>nd</sup> trimester/ product	6	6	6	100%
3 <sup>rd</sup> trimester/ product	140	117	117	100%
<b>Total – all animals</b>	<b>348</b>	<b>304</b>	<b>286</b>	<b>94%</b>
<b>Total – product only</b>	<b>289</b>	<b>250</b>	<b>237</b>	<b>95%</b>
<b>Total – placebo only</b>	<b>59</b>	<b>54</b>	<b>49</b>	<b>91%</b>

**Study 2013-PM-1009****Missouri Site:**

<b>Group</b>	<b>Vaccinated</b>	<b>Confirmed Pregnant</b>	<b>Foals</b>	<b>Parturition Rate</b>
2011 3 <sup>rd</sup> trimester	5	5	5	100%
2012 1 <sup>st</sup> trimester	1	1	1	100%
2012 2 <sup>nd</sup> trimester	53	43	39	91%
2012 3 <sup>rd</sup> trimester	26	26	25	96%
<b>Total – product</b>	<b>85</b>	<b>75</b>	<b>70</b>	<b>93%</b>

**Study 2014-PM-1009****North Dakota Site:**

<b>Group</b>	<b>Vaccinated</b>	<b>Confirmed Pregnant</b>	<b>Foaled</b>	<b>Parturition Rate</b>	<b>Foals Survived to End of Observation Period</b>
2 <sup>nd</sup> trimester vaccinated	52	52	52	100%	51*
3 <sup>rd</sup> trimester vaccinated	69	69	67**	97.1%	67

\*Lost foal affirmed by study cooperator to be due to causes other than vaccination.

\*\*One mare died due to causes other than vaccination, as affirmed by study cooperator.

All other foals were normal and healthy

**USDA Approval Date**

September 12, 2014

<b>Study Type</b>	Safety																																																			
<b>Pertaining to</b>	All fractions																																																			
<b>Study Purpose</b>	To demonstrate safety under field conditions																																																			
<b>Product Administration</b>	Two doses, administered intramuscularly approximately 3 – 4 weeks apart																																																			
<b>Study Animals</b>	880 horses, including 218 foals 3 months of age and 52 foals 5 months of age																																																			
<b>Challenge Description</b>	Not applicable																																																			
<b>Interval observed after challenge</b>	Not applicable																																																			
<b>Results</b>	<p>Horses were observed at least daily following each vaccination, until resolution of any observed reactions.</p> <p>There were no systemic reactions observed at any of the sites. Two foals and one horse died from causes affirmed by licensee not attributed to vaccination.</p> <p>Adverse events were limited to transient, non-painful swellings at the injection site that resolved without treatment.</p> <p>Local injection site reactions are summarized below across the four sites:</p> <table border="1"> <thead> <tr> <th rowspan="2">Site</th> <th rowspan="2">Total Number Of Vaccinates</th> <th rowspan="2">Number Of Vaccinates Administered 2 doses</th> <th colspan="2">Vaccinates With Transient Injection Site Swelling</th> <th colspan="2">Number Of Normal Vaccinates</th> </tr> <tr> <th>After 1<sup>st</sup> dose</th> <th>After 2<sup>nd</sup> dose</th> <th>After 1<sup>st</sup> dose</th> <th>After 2<sup>nd</sup> dose</th> </tr> </thead> <tbody> <tr> <td>North Dakota</td> <td>378</td> <td>378</td> <td>4</td> <td>0</td> <td>374</td> <td>378</td> </tr> <tr> <td>California</td> <td>43</td> <td>43</td> <td>4</td> <td>3</td> <td>39</td> <td>40</td> </tr> <tr> <td>Missouri</td> <td>292</td> <td>290</td> <td>0</td> <td>0</td> <td>292</td> <td>290</td> </tr> <tr> <td>Texas</td> <td>170</td> <td>169</td> <td>6</td> <td>1</td> <td>164</td> <td>168</td> </tr> <tr> <td><b>Total</b></td> <td><b>883</b></td> <td><b>880</b></td> <td><b>14</b> <b>(1.6%)</b></td> <td><b>4</b> <b>(0.5%)</b></td> <td><b>869</b> <b>(98.4%)</b></td> <td><b>876</b> <b>(99.5%)</b></td> </tr> </tbody> </table> <p>Results from each site are summarized on the following pages.</p>						Site	Total Number Of Vaccinates	Number Of Vaccinates Administered 2 doses	Vaccinates With Transient Injection Site Swelling		Number Of Normal Vaccinates		After 1 <sup>st</sup> dose	After 2 <sup>nd</sup> dose	After 1 <sup>st</sup> dose	After 2 <sup>nd</sup> dose	North Dakota	378	378	4	0	374	378	California	43	43	4	3	39	40	Missouri	292	290	0	0	292	290	Texas	170	169	6	1	164	168	<b>Total</b>	<b>883</b>	<b>880</b>	<b>14</b> <b>(1.6%)</b>	<b>4</b> <b>(0.5%)</b>	<b>869</b> <b>(98.4%)</b>	<b>876</b> <b>(99.5%)</b>
Site	Total Number Of Vaccinates	Number Of Vaccinates Administered 2 doses	Vaccinates With Transient Injection Site Swelling		Number Of Normal Vaccinates																																															
			After 1 <sup>st</sup> dose	After 2 <sup>nd</sup> dose	After 1 <sup>st</sup> dose	After 2 <sup>nd</sup> dose																																														
North Dakota	378	378	4	0	374	378																																														
California	43	43	4	3	39	40																																														
Missouri	292	290	0	0	292	290																																														
Texas	170	169	6	1	164	168																																														
<b>Total</b>	<b>883</b>	<b>880</b>	<b>14</b> <b>(1.6%)</b>	<b>4</b> <b>(0.5%)</b>	<b>869</b> <b>(98.4%)</b>	<b>876</b> <b>(99.5%)</b>																																														

North Dakota Site:

Summary	Number Of Vaccinates	Number Of Vaccinates Administered 2 doses	Vaccinates With Transient Injection Site Swelling		Number Of Normal Vaccinates	
			After 1 <sup>st</sup> dose	After 2 <sup>nd</sup> dose	After 1 <sup>st</sup> dose	After 2 <sup>nd</sup> dose
2-4 months	179	179	0	0	179	179
5-7 months	0	0	n/a	n/a	n/a	n/a
8-11 months	0	0	n/a	n/a	n/a	n/a
1-5 years*	121	121	2	0	119	121
6-15 years*	78	78	2	0	76	78
>16 years	0	0	n/a	n/a	n/a	n/a
<b>Total</b>	<b>378</b>	<b>378</b>	<b>4</b>	<b>0</b>	<b>374</b>	<b>378</b>

\*Swellings were 3cm in size observed 1-3 days post vaccination that resolved within 3 days.

California Site:

Summary	Number Of Vaccinates	Number Of Vaccinates Administered 2 doses	Vaccinates With Transient Injection Site Swelling		Number Of Normal Vaccinates	
			After 1 <sup>st</sup> dose	After 2 <sup>nd</sup> dose	After 1 <sup>st</sup> dose	After 2 <sup>nd</sup> dose
2-4 months*	7	7	0	2	7	5
5-7 months**	1	1	1	0	0	1
8-11 months	0	0	n/a	n/a	n/a	n/a
1-5 years***	19	19	2	0	17	19
6-15 years****	15	15	1	1	14	14
>16 years	1	1	0	0	1	1
<b>Total</b>	<b>43</b>	<b>43</b>	<b>4</b>	<b>3</b>	<b>39</b>	<b>40</b>

\*Swellings were 3cm in size observed within hours post vaccination that resolved within several hours.

\*\*Swelling was 3cm in size observed immediately post vaccination that resolved within several hours.

\*\*\*1 horse had a swelling 1cm in size observed immediately post vaccination that resolved within several hours. 1 horse had a swelling observed on day 1 that increased in size to 9cm on day 3 post vaccination and resolved by day 5.

\*\*\*\*Same horse had a swelling after each vaccination that resolved within 3 weeks. Size after the first vaccination was 24cm. Size after the second vaccination was 10cm.



Missouri Site:

Summary	Number Of Vaccinates	Number Of Vaccinates Administered 2 doses	Vaccinates With Transient Injection Site Swelling		Number Of Normal Vaccinates	
			After 1 <sup>st</sup> dose	After 2 <sup>nd</sup> dose	After 1 <sup>st</sup> dose	After 2 <sup>nd</sup> dose
2-4 months	33	32	0	0	33	32
5-7 months	0	0	n/a	n/a	n/a	n/a
8-11 months	0	0	n/a	n/a	n/a	n/a
1-5 years	225	224	0	0	225	224
6-15 years	32	32	0	0	32	32
>16 years	2	2	0	0	2	2
<b>Total</b>	<b>292</b>	<b>290</b>	<b>0</b>	<b>0</b>	<b>292</b>	<b>290</b>

Texas Site:

Summary	Number Of Vaccinates	Number Of Vaccinates Administered 2 doses	Vaccinates With Transient Injection Site Swelling		Number Of Normal Vaccinates	
			After 1 <sup>st</sup> dose	After 2 <sup>nd</sup> dose	After 1 <sup>st</sup> dose	After 2 <sup>nd</sup> dose
2-4 months	0	0	n/a	n/a	n/a	n/a
5-7 months	52	51	1	1	51	50
8-11 months	0	0	n/a	n/a	n/a	n/a
1-5 years	114	114	5	0	109	114
6-15 years	0	0	n/a	n/a	n/a	n/a
>16 years	4	4	0	0	4	4
<b>Total</b>	<b>170</b>	<b>169</b>	<b>6*</b>	<b>1**</b>	<b>164</b>	<b>168</b>

\*Swellings were <1.5cm were observed 4-7 days post vaccination and resolved within 6 days.

\*\*Swelling was 5cm observed 1 day post vaccination that resolved within 2 days.

**USDA  
Approval Date**

November 1, 2010