

**Table F. Wildlife Disease Sampling Conducted by
Wildlife Services in Michigan - FY 2007**

| <i>State</i> | <i>Disease/Condition</i> | <i>Species Sampled</i> | <i>Samples</i> |
|--------------|----------------------------|---------------------------------------|----------------|
| MI | avian influenza | | |
| | | ducks, american black | 5 |
| | | ducks, bufflehead | 9 |
| | | ducks, canvasback | 16 |
| | | ducks, gadwall | 3 |
| | | ducks, goldeneye, common | 1 |
| | | ducks, mallards | 146 |
| | | ducks, northern pintail | 92 |
| | | ducks, ruddy | 4 |
| | | ducks, scaup, lesser | 26 |
| | | ducks, teal, green-winged | 122 |
| | | ducks, wigeon, american | 1 |
| | | ducks, wood | 1 |
| | | geese, canada | 2 |
| | | grebes, horned | 2 |
| | | herons, great blue | 4 |
| | | non-wildlife | 749 |
| | | swans, mute | 230 |
| | | swans, trumpeter | 30 |
| | | swans, tundra | 16 |
| | | swans, whooper | 30 |
| | | <i>Subtotal of Samples by Disease</i> | 1,489 |
| | bovine tuberculosis | | |
| | | coyotes | 1 |
| | | opossums, virginia | 30 |
| | | raccoons | 68 |
| | | <i>Subtotal of Samples by Disease</i> | 99 |
| | pseudorabies | | |
| | | hogs, feral | 1 |
| | | <i>Subtotal of Samples by Disease</i> | 1 |
| | rabies | | |
| | | coyotes | 4 |
| | | foxes, gray | 40 |
| | | raccoons | 308 |
| | | skunks, striped | 10 |
| | | <i>Subtotal of Samples by Disease</i> | 362 |
| | swine brucellosis | | |
| | | hogs, feral | 1 |

Entries of "bird, unidentifiable" or "non-wildlife" usually refer to samples of matter, such as fecal material, found in the environment. "Bird, unidentifiable" is a name assigned to material when WS wants to know it is associated with birds.

| <i>State</i> | <i>Disease/Condition</i> | <i>Species Sampled</i> | <i>Samples</i> |
|--------------|--------------------------|---|----------------|
| MI | swine brucellosis | | |
| | | <i>Subtotal of Samples by Disease</i> | 1 |
| | | <i>Total of All Samples of All Types This State</i> | 1,952 |