

# Influenza A Subtype H3 Viruses in Feral Swine, United States, 2011–2012

Zhixin Feng, John A. Baroch, Li-Ping Long, Yifei Xu, Frederick L. Cunningham, Kerri Pedersen, Mark W. Lutman, Brandon S. Schmit, Andrew S. Bowman, Thomas J. DeLiberto, and Xiu-Feng Wan

To determine whether, and to what extent, influenza A subtype H3 viruses were present in feral swine in the United States, we conducted serologic and virologic surveillance during October 2011–September 2012. These animals were periodically exposed to and infected with A(H3N2) viruses, suggesting they may threaten human and animal health.

Swine are proposed as “mixing vessels” to generate novel influenza A viruses (IAVs) by facilitating reassortment among IAVs and providing a potential pathway in which these viruses can move from wild birds to humans (1). Subtype H3N2 is one of the most common subtypes in the US domestic swine population, which possibly resulted from spillover of human seasonal A(H3N2) virus (2,3). Since its introduction in the mid-1990s, A(H3N2) virus has evolved genetically and antigenically in domestic swine. Four genetic groups (so-called clusters I–IV) were identified, and these 4 clusters were also antigenically distinct (4). The viruses in cluster IV formed at least 2 antigenic subclusters, H3N2- $\alpha$  and H3N2- $\beta$  (5). Both subclusters are co-circulating in pigs, and subcluster H3N2- $\beta$  predominated among the isolates obtained from domestic swine at Ohio county fairs during 2010–2011 (5). During July and August 2011, two children were infected with novel

reassortant H3N2 variant (H3N2v), 1 in Indiana and 1 in Pennsylvania (6). This H3N2v virus is antigenically similar to the viruses in subcluster H3N2- $\beta$  and has the matrix gene of influenza A(H1N1)pdm09 virus. It caused illness in  $\approx$ 2,055 persons during August 2011–April 2012 (7).

The role of feral swine in IAV ecology has not been adequately addressed. Feral swine could be a reservoir of IAVs or, possibly, a spatially dynamic mixing vessel, given their free-ranging habits. Such unrestricted movement provides the potential for exchange of IAVs among subpopulations of feral swine and the opportunity for exposure to different IAVs through contact with a variety of habitats and species. Also, feral swine can live up to 8 years, which provides ample opportunities for reinfection with the same subtype IAVs, especially those with antigenically distinct hemagglutinins. The IAVs can be transmitted bidirectionally between feral and domestic swine because contact between them is not unusual (8). Ultimately the IAVs emerging in feral swine potentially could be transmitted to humans.

The United States has  $\approx$ 4–5 million feral swine (9) throughout at least 38 states (10). Feral swine are expanding their range because of a lack of natural predators and intentional introductions for hunting. Our goals in this study were to determine through virologic and serologic surveillance whether, and to what extent, subtype H3 IAVs were present in the US feral swine metapopulation.

## The Study

We collected 1,983 nasal swab samples from swine during October 2011–September 2012 (online Technical Appendix, [wwwnc.cdc.gov/EID/article/20/5/13-1578-Techapp1.pdf](http://wwwnc.cdc.gov/EID/article/20/5/13-1578-Techapp1.pdf)). Matrix gene-based quantitative reverse transcription PCR showed that 9 swabs samples were IAV positive; 1 A(H3N2) feral swine isolate, A/swine/Texas/A01104013/2012(H3N2), was recovered. Phylogenetic analyses showed that all genes of this feral swine isolate are genetically similar to those of A(H3N2v) viruses isolated from humans, and other contemporary subtype H3N2 isolates from swine from county fairs farms. The matrix gene of this feral swine IAV was genetically close to that of A(H1N1)pdm09 and the human A(H3N2v) viruses (online Technical Appendix Figure) (11). Similar to other viruses in the antigenic cluster, H3N2- $\beta$  A/swine/Texas/A01104013/2012(H3N2) had a R189K mutation at the antibody-binding site of the hemagglutinin protein, which caused a recent antigenic drift in subtype H3N2 IAVs (5,12). The 8 genes of A/swine/Texas/A01104013/2012(H3N2) have a minimal 99.59% nt sequence identity to those of the human subtype H3N2v isolate A/Indiana/10/2011(H3N2).

We also collected 1,989 serum samples from swine in 31 states; these samples were tested by using an IAV-specific ELISA (Figure 1). We identified 182 samples as

Author affiliations: Mississippi State University, Mississippi State, Mississippi, USA (Z. Feng, L.-P. Long, Y. Xu, X.-F. Wan); Jiangsu Academy of Agricultural Sciences, Nanjing, China (Z. Feng); Animal and Plant Health Inspection Service of the US Department of Agriculture, Fort Collins, Colorado, USA (J.A. Baroch, K. Pedersen, M.W. Lutman, B.S. Schmit, T.J. DeLiberto); Animal and Plant Health Inspection Service of the US Department of Agriculture, Mississippi State (F.L. Cunningham); and The Ohio State University, Columbus, Ohio, USA (A.S. Bowman)

DOI: <http://dx.doi.org/10.3201/eid2005.131578>




Figure 1. Geographic distributions of serum samples from feral swine, United States, 2011–2012. A) Of 1,989 samples tested by ELISA, 182 were positive (red) and 1,807 were negative (blue). B) The 76 samples (blue) were selected for hemagglutination-inhibition and microneutralization subtyping. C) The distributions of feral swine (green) and domestic swine (orange) were also marked (A and C).

IAV positive, from swine that were broadly distributed over 19 states. The average IAV seropositive rate was 9.15% but it varied by month. The highest positive rate (22.9%) was in June 2012 (Figure 2). Although no clear temporal pattern was found in the IAV seropositive rate, the rate was relatively higher in the summer than in other seasons. One explanation could be that noncommercial swine farmers might give their animals more pasture time during the summer, thereby increasing the chance of contact between domestic and feral swine. In addition, our results showed adult feral swine ( $\geq 1$  year of age) had the highest rate of IAV positivity (11.1% [of 1,380 animals]),

followed by subadults (2 months–1 year of age) (5.1% [of 494 animals]), and juveniles (<2 months of age) (3.8% [of 105 animals]) (age was not determined for 10 animals). (Dentition patterns were used to determine the age of feral swine [13].) Female and male pigs were equally as likely to be seropositive (102 [9.6%] of 1,058 vs. 80 [9.7%] of 821).

Of the 182 IAV-positive serum samples, 76 were randomly selected for influenza subtyping. We used hemagglutination-inhibition (HI) and microneutralization (MN) assays for subtyping against A(H1N1)pdm09 virus and 22 H3 IAVs, which represent a wide range of antigenically




Figure 2. Epidemiologic analyses of feral swine serum samples seropositive for influenza A virus by ELISA, United States, 2011–2012. A) Temporal distribution. B) Distribution of feral pigs, by age. C) Distribution of feral pigs, by sex. The numbers on the bar indicate the influenza A virus–seropositive percentile.

Table. Cross-reactive antibody responses against H3 and influenza A(H1N1)pdm09 virus in 76 influenza-positive serum samples from feral swine, United States, 2011–2012\*

| Source, virus                       | Antigenic cluster† | No. (%) seropositive‡ |           | GMT (95% CI)§ |                | Overall no. (%) seropositive¶ |
|-------------------------------------|--------------------|-----------------------|-----------|---------------|----------------|-------------------------------|
|                                     |                    | HI                    | MN        | HI            | MN             |                               |
| <b>Feral swine IAV</b>              |                    |                       |           |               |                |                               |
| A/swine/Texas/A011040013/2012(H3N2) | H3N2-β             | 36 (47.4)             | 36 (47.4) | 163 (40–640)  | 259 (40–1,280) | 25 (32.9)                     |
| <b>Domestic swine IAV</b>           |                    |                       |           |               |                |                               |
| A/swine/Ohio/11SW64/2009 (H3N2)     | H3N2-α             | 27 (35.5)             | 30 (39.5) | 65 (40–160)   | 121 (40–1,280) | 17 (22.4)                     |
| A/swine/Iowa/12627/2009(H3N2)       | H3N2-α             | 30 (39.5)             | 31 (40.8) | 73 (40–320)   | 183 (40–1,280) | 15 (19.7)                     |
| A/swine/Ohio/11SW347/2011(H3N2)     | H3N2-β             | 31 (40.8)             | 31 (40.8) | 122 (40–640)  | 112 (40–1,280) | 17 (22.4)                     |
| A/swine/Iowa/6368/2012(H3N2)        | H3N2-β             | 26 (34.2)             | 36 (47.4) | 94 (0–640)    | 240 (40–1,280) | 20 (26.3)                     |
| <b>Human H3N2v IAV</b>              |                    |                       |           |               |                |                               |
| A/IA/07/2011                        |                    | 32 (42.1)             | 31 (40.8) | 109 (40–640)  | 313 (40–1,280) | 21 (27.6)                     |
| <b>Human seasonal IAV</b>           |                    |                       |           |               |                |                               |
| A/Perth/16/2009                     | H3N2-β             |                       |           |               |                |                               |
| A/Perth/16/2009                     | H3N2-β             | 12 (15.8)             | 5 (6.6)   | 76 (40–320]   | 121 (40–1,280) | 1 (1.3)                       |
| A/Victoria/361/2011                 | H3N2-β             | 11 (14.5)             | 2 (2.6)   | 141 (40–640)  | 226 (40–1,280) | 0                             |
| A/California/7/2009                 | H3N2-β             | 5 (6.6)               | 3 (3.9)   | 70 (40–640)   | 80 (40–320)    | 1 (1.3)                       |

\*Influenza detected by ELISA. GMT, geometric mean titer; HI, hemagglutination-inhibition; MN, microneutralization; IAV, influenza A virus.

†The antigenic cluster H3N2-α and H3N2-β were defined in (5).

‡Seropositive was defined as the HI or MN titer  $\geq 40$ .

§Only serum with an HI or MN titer  $\geq 40$  are used to calculate the GMT.

¶Overall seroconversion was defined as both HI and MN titer  $\geq 40$ .

distinct H3 IAVs (online Technical Appendix Table). Serum was defined as seropositive if its titer was  $\geq 40$ .

HI results showed that 46 (60.5%) of 76 feral swine samples were positive against at least 1 of the 22 H3 IAVs tested, of which A/swine/Texas/A011040013/2012(H3N2) had the highest seropositive rate (47.4%), followed by 4 human A(H3N2v) isolates (Table). MN results were consistent with those of HI. The geometric mean titers for HI and MN against the feral swine isolate were 163 and 259, respectively. The maximum MN titer among the serum and A/swine/Texas/A011040013/2012(H3N2) was 1,280. The seropositive rates of these serum samples varied from 34.2% to 42.1% against the other viruses from subclusters H3N2-α and H3N2-β (Table), but the viruses in subcluster H3N2-β had significantly higher geometric mean titers than did those in H3N2-α ( $p < 0.001$ ).

HI results also showed that 12 samples were seropositive against A/Perth/16/2009(H3N2); 11 were seropositive against A/Victoria/361/2011(H3N2), and 5 were seropositive against A/California/7/2009(H1N1). These 3 viruses do not cross-react with the IAVs from subclusters H3N2-α and H3N2-β. Those 5 subtype H1N1-positive samples were also seropositive against subtype H3N2 IAVs, indicating potential previous exposures of these feral swine to both H3 and H1 IAVs.

The HI results demonstrated that only 2 serum samples had a low-level cross-reaction with avian influenza A(H3N2) viruses, and the HI titers for both were 40. This result is consistent with findings in an earlier report (14). Further studies are needed to determine whether additional antigenic clusters of H3 IAVs are present in migratory waterfowl. The discrepancies in the cross-reactivity of these serum samples against the IAVs tested in this study suggested that these feral swine had different exposure histories against antigenically diverse IAVs.

## Conclusions

Our study demonstrated that subtype H3N2 IAVs are periodically infecting feral swine in the United States. Feral swine are a potential source of IAVs with bidirectional transmission to domestic swine or humans. Detection of an H3N2v-like IAV in the feral swine population demonstrates a potential threat to human health. Continued surveillance is recommended to monitor the distribution and the genomic and antigenic diversities of IAVs in feral swine to better assess the risk.

## Acknowledgments

We are grateful to Jianqiang Ye and Brigitte Martin for technical support and discussion and Dena Pruett for editorial assistance. We also thank Michael Shaw providing subtype H3N2v isolates and Kyoung-Jin Yoon for providing swine influenza isolates.

This work was supported by the US Department of Agriculture, Wildlife Services, National Wildlife Research Center (grant no. 13-7428-0961-CA).

Dr Feng is a visiting scholar at the Influenza Systems Biology Laboratory, Department of Basic Sciences, College of Veterinary Medicine, Mississippi State University. His research focuses on swine influenza serology and ecology.

## References

- Hinshaw VS, Webster RG, Easterday BC, Bean WJ Jr. Replication of avian influenza A viruses in mammals. *Infect Immun*. 1981;34:354–61.
- Zhou NN, Senne DA, Landgraf JS, Swenson SL, Erickson G, Rossow K, et al. Genetic reassortment of avian, swine, and human influenza A viruses in American pigs. *J Virol*. 1999;73:8851–6.
- Zhou NN, Senne DA, Landgraf JS, Swenson SL, Erickson G, Rossow K, et al. Emergence of H3N2 reassortant influenza A viruses in North American pigs. *Vet Microbiol*. 2000;74:47–58. [http://dx.doi.org/10.1016/S0378-1135\(00\)00165-6](http://dx.doi.org/10.1016/S0378-1135(00)00165-6)

4. Hause BM, Oleson TA, Bey RF, Stine DL, Simonson RR. Antigenic categorization of contemporary H3N2 swine influenza virus isolates using a high-throughput serum neutralization assay. *J Vet Diagn Invest.* 2010;22:352–9. <http://dx.doi.org/10.1177/104063871002200302>
5. Feng Z, Gomez J, Bowman AS, Ye J, Long LP, Nelson SW, et al. Antigenic characterization of H3N2 influenza A viruses from Ohio agricultural fairs. *J Virol.* 2013;87:7655–67. <http://dx.doi.org/10.1128/JVI.00804-13>
6. Centers for Disease Control and Prevention. Swine-origin influenza A (H3N2) virus infection in two children—Indiana and Pennsylvania, July–August 2011. *MMWR Morb Mortal Wkly Rep.* 2011;60:1213–5.
7. Biggerstaff M, Reed C, Epperson S, Jhung MA, Gambhir M, Bresee JS, et al. Estimates of the number of human infections with influenza A(H3N2) variant virus, United States, August 2011–April 2012. *Clin Infect Dis.* 2013;57(Suppl 1):S12–5. <http://dx.doi.org/10.1093/cid/cit273>
8. Wyckoff AC, Henke SE, Campbell TA, Hewitt DG, VerCauteren KC. Feral swine contact with domestic swine: a serologic survey and assessment of potential for disease transmission. *J Wildl Dis.* 2009;45:422–9. <http://dx.doi.org/10.7589/0090-3558-45.2.422>
9. Pimentel D. Environmental and economic costs of vertebrate species invasions into the United States. In: Witmer GW, Pitt WC, Fagerstone KA, editors. *Managing vertebrate invasive species: proceedings of an international symposium; 2007 August 7–9, 2007.* Fort Collins (CO): USDA/APHIS/National Wildlife Research Center; 2007. p. 1–8.
10. Fogarty EP. National distribution of and stakeholder attitudes toward feral hogs. Mississippi State (MS): Mississippi State University; 2007.
11. Shu B, Garten R, Emery S, Balish A, Cooper L, Sessions W, et al. Genetic analysis and antigenic characterization of swine origin influenza viruses isolated from humans in the United States, 1990–2010. *Virology.* 2012;422:151–60. <http://dx.doi.org/10.1016/j.virol.2011.10.016>
12. Ye J, Xu Y, Harris J, Sun H, Bowman AS, Cunningham F, et al. Mutation from arginine to lysine at the position 189 of hemagglutinin contributes to the antigenic drift in H3N2 swine influenza viruses. *Virology.* 2013;446:225–9. <http://dx.doi.org/10.1016/j.virol.2013.08.004>
13. Matschke GH. Aging European wild hogs by dentition. *J Wildl Manage.* 1967;31:109–13. <http://dx.doi.org/10.2307/3798365>
14. Hall JS, Minnis RB, Campbell TA, Barras S, Deyoung RW, Pabilonia K, et al. Influenza exposure in United States feral swine populations. *J Wildl Dis.* 2008;44:362–8. <http://dx.doi.org/10.7589/0090-3558-44.2.362>

Address for correspondence: Xiu-Feng Wan, Department of Basic Sciences, College of Veterinary Medicine, Mississippi State University, 240 Wise Center Dr, Mississippi State, MS 39762, USA; email: wan@cvm.msstate.edu

