

**Table F. Wildlife Disease Sampling Conducted by
Wildlife Services in Illinois - FY 2010**

| <i>Disease/Condition</i> | <i>Species Sampled</i> | <i># of Samples</i> |
|--------------------------------|---|---------------------|
| Avian Influenza | | |
| | Coots, American | 3 |
| | Ducks, Canvasback | 1 |
| | Ducks, Gadwall | 16 |
| | Ducks, Mallards | 91 |
| | Ducks, Merganser Common | 1 |
| | Ducks, Merganser, Hooded | 4 |
| | Ducks, Northern Pintail | 4 |
| | Ducks, Northern Shoveler | 5 |
| | Ducks, Redhead | 5 |
| | Ducks, Ring-Necked | 7 |
| | Ducks, Scaup, Lesser | 2 |
| | Ducks, Teal, Blue-Winged | 75 |
| | Ducks, Teal, Green-Winged | 36 |
| | Ducks, Wigeon, American | 1 |
| | Ducks, Wood | 15 |
| | Geese, Canada | 126 |
| | Gulls, Bonaparte`s | 1 |
| | Gulls, Ring-Billed | 43 |
| | Killdeers | 26 |
| | Sandpipers, Least | 1 |
| | Sandpipers, Solitary | 2 |
| | Sandpipers, Spotted | 7 |
| | <i>Subtotal of # of Samples for Avian Influenza</i> | 472 |
| Bluetongue | | |
| | Deer, White-Tailed (Wild) | 23 |
| | <i>Subtotal of # of Samples for Bluetongue</i> | 23 |
| Chronic Wasting Disease | | |
| | Deer, White-Tailed (Wild) | 156 |
| | <i>Subtotal of # of Samples for Chronic Wasting Disease</i> | 156 |
| Classical Swine Fever | | |
| | Swine, Feral | 18 |
| | <i>Subtotal of # of Samples for Classical Swine Fever</i> | 18 |
| Disease (Non-Specific) | | |
| | Swine, Feral | 2 |
| | <i>Subtotal of # of Samples for Disease (Non-Specific)</i> | 2 |

Entries of "bird, unidentifiable" or "non-wildlife" usually refer to samples of matter, such as fecal material, found in the environment. "Bird, unidentifiable" is a name assigned to material when WS wants to know it is associated with birds.

| <i>Disease/Condition</i> | <i>Species Sampled</i> | <i># of Samples</i> |
|--------------------------------------|---|---------------------|
| Epizootic Hemorrhagic Disease | Deer, White-Tailed (Wild) | 23 |
| | <i>Subtotal of # of Samples for Epizootic Hemorrhagic Disease</i> | 23 |
| Genetic | Coyotes | 14 |
| | <i>Subtotal of # of Samples for Genetic</i> | 14 |
| Leptospirosis | Raccoons | 80 |
| | <i>Subtotal of # of Samples for Leptospirosis</i> | 80 |
| Pseudorabies | Swine, Feral | 18 |
| | <i>Subtotal of # of Samples for Pseudorabies</i> | 18 |
| Q Fever | Pigeons, Feral (Rock) | 16 |
| | Starlings, European | 212 |
| | <i>Subtotal of # of Samples for Q Fever</i> | 228 |
| Rabies | Skunks, Striped | 20 |
| | <i>Subtotal of # of Samples for Rabies</i> | 20 |
| Raccoon Roundworm | Raccoons | 98 |
| | <i>Subtotal of # of Samples for Raccoon Roundworm</i> | 98 |
| Swine Brucellosis | Swine, Feral | 18 |
| | <i>Subtotal of # of Samples for Swine Brucellosis</i> | 18 |
| Ticks:All | Deer, White-Tailed (Wild) | 10 |
| | <i>Subtotal of # of Samples for Ticks:All</i> | 10 |
| Toxoplasmosis | Skunks, Striped | 2 |
| | <i>Subtotal of # of Samples for Toxoplasmosis</i> | 2 |
| Tularemia | Coyotes | 4 |
| | Opossums, Virginia | 3 |
| | Raccoons | 19 |
| | Skunks, Striped | 2 |
| | Swine, Feral | 4 |

Entries of "bird, unidentifiable" or "non-wildlife" usually refer to samples of matter, such as fecal material, found in the environment. "Bird, unidentifiable" is a name assigned to material when WS wants to know it is associated with birds.

| <i>Disease/Condition</i> | <i>Species Sampled</i> | <i># of Samples</i> |
|--------------------------|---|---------------------|
| Tularemia | | |
| | <i>Subtotal of # of Samples for Tularemia</i> | 32 |
| West Nile Virus | | |
| | Turtles Other | 22 |
| | Turtles, Common Snapping | 19 |
| | Turtles, Painted | 28 |
| | <i>Subtotal of # of Samples for West Nile Virus</i> | 69 |
| | <i>Grand Total of Samples For All Diseases/Conditions</i> | 1,283 |
| | <i>Total # Diseases/Conditions Sampled In IL</i> | 17 |

Entries of "bird, unidentifiable" or "non-wildlife" usually refer to samples of matter, such as fecal material, found in the environment. "Bird, unidentifiable" is a name assigned to material when WS wants to know it is associated with birds.